

Perennial grain on a Midwest Alfisol shows no sign of early soil carbon gain

Christine D. Sprunger^{1,2*}, Steve W. Culman³, G. Philip Robertson^{1,4} and Sieglinde S. Snapp⁴

¹W.K. Kellogg Biological Station, Michigan State University, Hickory Corners, Michigan 49060, USA.

²Agriculture and Food Security Center, The Earth Institute, Columbia University, New York city, New York 10025, USA.

³School of Environment and Natural Resources, Ohio State University, Wooster, Ohio 44691, USA.

⁴Department of Plant, Soil, and Microbial Sciences, Michigan State University, East Lansing, Michigan 48824, USA.

*Corresponding author: cs3476@columbia.edu

Accepted 22 February 2017; First published online 23 March 2017

Research Paper

Abstract

Perennial grain crops are expected to sequester soil carbon (C) and improve soil health due to their large and extensive root systems. To examine the rate of initial soil C accumulation in a perennial grain crop, we compared soil under perennial intermediate wheatgrass (IWG) with that under annual winter wheat 4 years after the crops were first planted. In addition, we tested the effect of three nitrogen (N) sources on C pools: Low available N (Low N (Organic N); 90 kg N ha⁻¹ poultry litter), moderately available N (Mid N; 90 kg N ha⁻¹ urea) and high available N (High N; 135 kg N ha⁻¹ urea). We measured aboveground C (grain + straw), and coarse and fine root C to a depth of 1 m. Particulate organic matter (POM-C), fractionated by size, was used to indicate labile and more stabilized soil C pools. At harvest, IWG had 1.9 times more straw C and up to 15 times more root C compared with wheat. There were no differences in the size of the large (6 mm–250 µm) or medium (250–53 µm) POM-C fractions between wheat and IWG ($P > 0.05$) in surface horizons (0–10 cm). Large POM-C under IWG ranged from 3.6 ± 0.3 to 4.0 ± 0.7 g C kg soil⁻¹ across the three N rates, similar to wheat under which large POM-C ranged from 3.6 ± 1.4 to 4.7 ± 0.7 g C kg soil⁻¹. Averaged across N level, medium POM-C was 11.1 ± 0.8 and 11.3 ± 0.7 g C kg soil⁻¹ for IWG and wheat, respectively. Despite IWG's greater above and belowground biomass (to 70 cm), POM-C fractions in IWG and wheat were similar. Post-hoc power analysis revealed that in order to detect differences in the labile C pool at 0–10 cm with an acceptable power (~80%) a 15% difference would be required between wheat and IWG. This demonstrates that on sandy soils with low cation exchange capacity, perennial IWG will need to be in place for longer than 4 years in order to detect an accumulated soil C difference > 15%.

Key words: perennial grain crops, annual winter wheat, roots, soil carbon sequestration, sustainable agriculture

Introduction

Intensive agricultural practices have depleted soil carbon (C) pools by up to 75% and contributed ~124 Pg C to the atmosphere over the past 140 years (Houghton and Hackler, 2001; Lal, 2011). Several management practices can replenish the soil C pool (West and Post, 2002; Jarecki and Lal, 2003) and one of the most effective is to convert annual row crops to perennial vegetation (Post and Kwon, 2000; McLauchlan et al., 2006; Syswerda et al., 2011). For example, Post and Kwon (2000) reported average C accumulation rates following conversion to grasslands of 33.2 g C m⁻² yr⁻¹ and Gebhart et al. (1994) found rates as high as 110 g C m⁻² yr⁻¹ 12 years post conversion. Similar estimates have been reported for row crop

conversion to forest although rates vary between tropical and temperate stands (Post and Kwon, 2000). Evidence of C accrual in abandoned agricultural fields have also been reported with annual increases of 20 g C m⁻² yr⁻¹ (Knops and Tilman, 2000) to 60 g C m⁻² yr⁻¹ (Robertson et al., 2000) in surface soils.

One of the most important drivers of C accumulation following conversion to perennials is an increase in soil organic matter inputs (Post and Kwon, 2000). Perennial systems often have between 3 and 10 times more belowground biomass compared with annual row crops (Zan et al., 2001; Culman et al., 2010; Dupont et al., 2014). Furthermore, C accrual will occur faster in perennial vegetation because perennial crops are usually not tilled and are typically planted for longer intervals compared

with annual crops, thus minimizing disturbance and reducing soil C loss (Huggins et al., 1998). Other important factors that lead to C accumulation under perennials are the inputs of soil organic matter deeper in the soil profile and enhanced physical protection of soil C through aggregation (Six et al., 1998; Grandy and Robertson, 2006; Syswerda et al., 2011; Tiemann and Grandy, 2015).

Less is known about the length of time required before increases in soil C are detectable or before soil C stabilizes post conversion (McLauchlan et al., 2006). In some cases, C accrual is detectable within the first few years post conversion (McLauchlan et al., 2006; Rehbein et al., 2015) while in other cases C accumulation among croplands and perennial systems may not be significantly different after 12 years (Syswerda et al., 2011). Different C accrual rates between experiments likely result from a number of interacting factors including soil type, climate, fertilization practices, soil spatial variability (Anderson-Teixerira et al., 2009; Kravchenko and Robertson, 2011; Tiemann and Grandy, 2015) and how close a system is to reaching C equilibrium (Six et al., 2002). Further, soil C comprises of different pools that turn over at different rates (Wander, 2004). For example, organic matter that is readily decomposed by microbial activity is typically associated with labile or active pools of C compared with organic material that is inaccessible to microbial activity either due to inherent chemical recalcitrance (Rasse et al., 2005) or physical protection (Schmidt et al., 2011). The proportion of C in the labile pools compared with more processed C pools is rarely determined and thus C stabilization potential post conversion is poorly understood.

Particulate organic matter (POM) is commonly used to decipher labile and stable soil C pools through physical size fractionation (Cambardella and Elliott, 1992). The large fraction mainly comprises organic materials that are in the initial phases of decomposition that can turn-over anywhere from a few days to a year. The medium fraction consists of more processed organic materials that can take a few decades to fully decompose. Given the more stable nature of the medium fraction, it has a greater influence on long-term soil carbon sequestration compared with the labile C pool (Rehbein et al., 2015).

There is widespread interest to accumulate soil C in both labile and stable pools within agricultural systems for both farm-scale and global benefits. One option could be to cultivate perennial grain crops in place of annual grain crops (Asbjornsen et al., 2013). Perennial wheat and perennial intermediate wheatgrass (IWG) are being developed to achieve the high yields of annual wheat and have extensive root systems that could potentially increase soil C (Glover et al., 2010; Kell, 2011). For example, compared with annual wheat, perennial grains developed by DeHaan et al. (2005) have significantly more coarse and fine roots to 40 cm and 70 cm depths, respectively (Sprunger, 2015). However, yields

from perennial grain crops seem to peak after 2 or 3 years (Wagoner, 1990; Culman et al., unpublished), at which point a farmer would need to replant or rotate to another crop. Understanding whether initial gains in soil C can occur within this period is important for determining the value of perennial grains as a plausible strategy for C sequestration.

Here we examine root C and large and medium POM-C fractions under an experimental perennial grain crop (IWG) 4 years post conversion compared with annual winter wheat. We hypothesize that (1) IWG will accumulate more C in both large and medium POM-C fractions compared with wheat because of greater C inputs from both above and belowground sources; and (2) POM-C fractions will be greatest in systems receiving higher levels of more available nitrogen (N).

Methods

Site description

This study was conducted at the W.K. Kellogg Biological Station (KBS) Long-term Ecological Research site, located in southwest Michigan, USA (42°24'N, 85°24'W, elevation 288 m). Mean annual precipitation and temperature are 1005 mm and 10.1°C. KBS soils are co-mingled Kalamazoo soil series (fine loamy) and Oshtemo (coarse loamy) soil series, mixed, mesic Typic Hapludalfs. These soils typically have an A horizon to a depth of 30 cm, a deep Bt horizon that reaches 80+ cm and a BC horizon that extends to 140 cm (Table 1). Prior to the establishment of this experiment, this field was under a conventionally managed corn (*Zea mays* L.)-soybean [*Glycine max* (L.) Meer.]-wheat (*Triticum aestivum* L.) rotation.

Experimental design

The experiment was established in 2009 as a split plot randomized complete block design with four replicated blocks. The main factor was N level and the sub-factor was crop type for a total of 24 plots (3 N levels × 2 crops × 4 blocks). Each plot was 3.0 m × 4.6 m, with 2.4 m buffers in between the plots and 0.9 m buffers on the perimeter. Three N levels included (1) Low N (Organic N), which contained approximately 21 kg of plant available N ha⁻¹ from poultry litter, and will hereafter be referred to as Low N; (2) Mid N, which received 90 kg N ha⁻¹ of urea; and (3) High N, which received 135 kg N ha⁻¹ of urea. Pelletized poultry litter (2.24 Mg ha⁻¹) was applied each October at a total rate of 90 kg N ha⁻¹, with 41% of N available in the first year (A&L Great Lakes Laboratory). The applied poultry litter had an NPK formula of 4–3–2. Manure subsamples were analyzed for C content using Loss on Ignition at 500 °C and analyses revealed that 0.83 Mg C ha⁻¹ was added to the soil each year (A&L Great Lakes Laboratory). The Mid

Table 1. Baseline soil profile characteristics taken in the fall of 2009 prior to planting.

Soil depth cm	Texture			Bulk density Mg m ⁻³	pH	CEC cmol _c kg ⁻¹	Total soil carbon g C kg ⁻¹ soil	Total soil nitrogen g N kg ⁻¹ soil
	sand	Silt %	Clay					
0–10	56	37	7	1.4	5.5	6.7	9.0	0.9
10–20	54	39	7	1.6	5.6	6.2	7.0	0.8
20–40	56	31	13	1.6	5.9	6.8	3.0	0.4
40–70	73	14	13	1.8	5.6	7.7	1.0	0.3
70–100	85	7	8	1.5	5.6	5.3	0.5	0.2

N level is the recommended rate for conventionally grown wheat in the state of Michigan, while the High N level received 50% more N than the Mid N level. In both the Mid N and High N treatments, a starter of 33.6 kg N ha⁻¹ as pelletized urea and 44 kg K₂O ha⁻¹ of KCL fertilizer were applied immediately before fall planting. The following spring, typically at the beginning of April and again at the end of May, plots were top-dressed with 28 and 50.4 kg N ha⁻¹ for Mid N and High N, respectively.

Prior to the start of the experiment in fall 2009, the site was chisel plowed to 20 cm and in subsequent years the wheat plots were rototilled to 15 cm depth. *Thinopyrum intermedium* (Host) Barworkth and D.R. Dewey, “*Kernza*TM” (IWG) was developed through bulk breeding and mass selection at the Land Institute in Salina, KS (DeHaan *et al.*, 2005; Cox *et al.*, 2010). IWG was planted in mid November 2009 at a seeding rate of 310 seeds m⁻² (1.25 million seeds acre⁻¹) at 15 cm row spacing (Culman *et al.*, 2013). Winter wheat (*Triticum aestivum* L. var. *Caledonia*) was planted in early October of 2009, 2010, 2011 and 2012 at a seeding rate of 432 seeds m⁻² (1.75 million seeds acre⁻¹). More details regarding planting and chemical application can be found in Culman *et al.* (2013).”

Aboveground biomass sampling

We sampled aboveground biomass (seed and straw) during the summer of 2013, when the IWG was in its fourth year and fully established. Aboveground biomass was measured at maturity for both crops. Wheat was harvested on July 15, 2013 and the IWG was harvested on August 26, 2013. Aboveground biomass was determined by randomly placing two 0.25-m² quadrats in every plot and clipping crop biomass to 10 cm above the soil. The aboveground biomass was separated into seed heads and straw then dried at 60°C for 48 h and weighed. Seeds were separated from their hulls using a custom-made tabletop thresher.

Belowground biomass and soil sampling

Belowground biomass and soil samples were collected on June 7 and 8, 2013, which was near peak aboveground

biomass for both wheat and IWG. We used a hydraulic direct-push soil sampler (Geoprobe, Salina, KS) to extract three 6 cm diameter soil cores to 1 m depth from each plot. The three cores per plot were subsequently divided into five depths (0–10 cm, 10–20 cm, 20–40 cm, 40–70 cm and 70–100 cm) and composited by depth interval. From each depth interval a sub-sample of 400 g was taken for root analysis. Roots were separated into two size classes, coarse (>6 mm) and fine (1–6 mm). We obtained coarse roots by gently sieving field moist soil through 6 mm sieves. We obtained fine roots by wet sieving the remaining soil through a 1 mm sieve. We then used tweezers to separate fine roots from plant residues that remained on the 1 mm sieve. We made no attempt to separate live and dead roots. To ensure that roots were soil-free, we hand-washed roots by soaking them in deionized water. Both coarse and fine roots were dried at 60°C for 48 h and then weighed.

Crop C and N analysis

Dried grain and stems were ground separately to a fine powder. Dried roots were frozen in liquid N and then immediately ground to a fine powder using a mortar and pestle. We analyzed both above and belowground crop parts for C and N in a CHNS analyzer (Costech Analyzer ECS 4010, Costech Analytical Technologies, Valencia, CA).

Particulate organic matter C

We used physical size fractionation to determine POM-C, which has been shown to reflect both labile and more stabilized C pools (Cambardella and Elliott, 1992; Culman *et al.*, 2012). First, we gently sieved 100 g of soil through a 6 mm screen so as not to disturb soil aggregates (Ontl *et al.*, 2013). Next, 10 g of air-dried soil and 30 mL of 0.05 sodium hexametaphosphate were combined in a 50 mL centrifuge tube and placed on a shaker for 8 h at 120 oscillations min⁻¹. Using deionized water, we passed the solution of soil and sodium hexametaphosphate through a 250 µm sieve (large POM-C), which was placed over a 0.053 µm mesh sieve (medium POM-C). The large POM-C fraction is associated with coarser material and

reflects the labile C pool, while the medium POM-C comprises silt and clay particles and is associated with more stabilized C pools. The materials that were retained on both sieves were oven dried at 55°C until samples reached a constant weight. Dried samples were then ground using a mortar and pestle and analyzed for C and N as above. POM-C on an areal basis was determined by multiplying POM-C concentration (g C kg^{-1} soil), dry weight of POM-C fraction (g)/proportion of total soil sample (g), bulk density (g cm^{-3}) and depth interval (cm).

Statistical analyses

All above and belowground biomass as well as labile soil C data were analyzed separately with the Proc Mixed procedure in SAS (version 9.3; SAS Institute, Cary, NC, USA). Crop, N level, depth and treatment interactions were treated as fixed effects. While, block and treatment \times block interactions were treated as random effects. Significant differences were determined at $\alpha = 0.05$. For POM-C and roots, depth was analyzed as a repeated measure. The covariance structure assigned in the repeated measures statement was compound symmetry. Means were compared with an adjusted Tukey's pairwise means comparison.

We used a post-hoc statistical power analysis to determine if a type II error occurred during the POM-C statistical analysis. Power analyses have been widely used in soil science to determine if the lack of significance is more likely due to insufficient sampling (number of replications) or an absence of biogeochemical differences between treatments (Kravchenko and Robertson, 2011; Ladoni et al., 2015). Detailed explanations of power analyses that have been used for soil C studies can be found in Garten and Wullschleger (1999); Poussart et al. (2004), and Kravchenko and Robertson (2011). In brief, we conducted a post-hoc power analysis that included (1) hypothesizing a size difference in the large POM-C between wheat and IWG; (2) estimating the variability (estimates of variance); (3) specifying a significance level of $\alpha = 0.05$; (4) specifying the probability of detecting statistical differences (power); and (5) calculating a proposed number of replications. The power analysis was conducted using the PROC MIXED procedure in SAS (version 9.3; SAS Institute, Cary, NC, USA).

Results

Aboveground C

Grain C differed by crop (Table 2; $F = 289$, $P < 0.05$), for example, grain C for wheat ranged from 1.28 ± 0.14 (mean \pm standard error of the mean) to 1.39 ± 0.15 Mg C ha^{-1} across N levels and was up to 25 times greater than for IWG, where grain C ranged from 0.07 ± 0.002 to 0.54 ± 0.1 Mg C ha^{-1} . There was no overall N level effect as both crops had statistically similar grain C

across N levels ($F = 0.5$, $P = 0.6$). Straw C was significantly greater in IWG compared with wheat (Table 2; $F = 110$, $P < 0.05$). Averaging across N levels, IWG had 1.9 times greater straw C compared with wheat. Neither the wheat nor IWG aboveground C differed across the three N levels.

Root C content and depth distribution

IWG coarse root C for the entire soil profile was up to 15 times greater than that of wheat, such that coarse root C for IWG ranged from 1.70 ± 0.30 to 2.42 ± 0.13 Mg C ha^{-1} and for wheat ranged from 0.11 ± 0.05 to 0.29 ± 0.07 Mg C ha^{-1} (Table 3). Despite no overall N level effect ($F = 1.4$, $P = 0.3$), pairwise comparisons revealed that IWG coarse root C under high N was significantly greater than coarse root C under Mid N and Low N ($P < 0.03$). Wheat coarse root C was statistically similar across N levels. The majority of coarse root C was concentrated at the surface for both crops (Fig. 1). Averaging across N levels, 60% of IWG total coarse root C was in the top 10 cm and 81% was in the top 20 cm. On average 79% of wheat total coarse root C was in the top 0–10 cm and 96% was in the top 20 cm. IWG had significantly greater root C compared with wheat to 40 cm depths under Low N and Mid N levels and to 20 cm depths under the High N level.

Differences between wheat and IWG were also apparent for fine root C (Table 3), which was four times greater in IWG compared with wheat ($F = 34.6$, $P = 0.0002$). Total fine root C for IWG was 0.24 ± 0.02 , 0.47 ± 0.09 , and 0.47 ± 0.10 Mg C ha^{-1} for Low N, Mid N and High N, respectively. In contrast, total fine root C for wheat was 0.063 ± 0.001 , 0.01 ± 0.01 , and 0.11 ± 0.05 Mg C ha^{-1} for Low N, Mid N and High N, respectively. There was a marginal overall N level effect on fine root C ($F = 3.0$, $P = 0.1$). In addition, pairwise comparisons showed that IWG under High N and Mid N had significantly greater root C content compared with IWG under Low N (Table 3).

IWG fine root C was more evenly distributed throughout the soil profile compared with coarse root C, but still a large portion was in the top 20 cm (Fig. 2). For example, averaging across N levels, 48% of IWG fine root C was in the top 10 cm and 72% was in the top 20 cm. Fine root distributions in the wheat systems mirrored the coarse root biomass distributions, with 67% found in the top 10 cm and 92% found in the top 90 cm. IWG had significantly more fine root C compared with wheat to 70 cm depth in Mid N and High N levels (Fig. 2). Within the Low N level, IWG had significantly greater fine root C compared with wheat down to only 20 cm.

Root C and N concentrations and C:N ratios

There was a significant crop effect for root C concentrations ($F = 98.9$, $P < 0.0001$), but differences between

Table 2. Grain and straw C Content for wheat and IWG across three N levels (Low N (Organic N), Mid N and High N) at the Kellogg Biological Station in 2013.


N Level	Grain		Mg C ha ⁻¹	Straw	
	Wheat	IWG		Wheat	IWG
Low N	1.28 (0.14) ^a	0.07 (0.002) ^b		2.34 (0.28) ^b	4.59 (0.41) ^a
Mid N	1.43 (0.19) ^a	0.11 (0.02) ^b		2.73 (0.24) ^b	5.19 (0.40) ^a
High N	1.39 (0.15) ^a	0.54 (0.01) ^b		2.42 (0.18) ^b	5.36 (0.25) ^a

Comparisons of means within rows (between crops) followed by same lowercase letters are not significantly different. Different lowercase letters denote significant differences between crops and across N levels. Within a column (crop), different lowercase letters denote a significant difference amongst N Levels at $P = 0.05$. Values in parentheses are standard errors. See Supplementary Materials for ANOVA tables.

Table 3. Total coarse and fine root C contents for the entire soil profile (0–100 cm) between wheat and IWG across three N levels (Low N (Organic N), Mid N and High N) at the Kellogg Biological Station in 2013.

N Level	Coarse roots		Mg C ha ⁻¹	Fine roots	
	Wheat	IWG		Wheat	IWG
Low N	0.29 (0.07) ^c	1.77 (0.29) ^b		0.06 (0.001) ^c	0.24 (0.02) ^b
Mid N	0.19 (0.06) ^c	1.74 (0.13) ^b		0.10 (0.01) ^c	0.47 (0.09) ^a
High N	0.16 (0.03) ^c	2.42 (0.13) ^a		0.11 (0.05) ^c	0.47 (0.1) ^a

Comparisons of means within rows (between crops) followed by same lowercase letters are not significantly different. Different lowercase letters denote significant differences between crops and across N levels. Within a column (crop), different lowercase letters denote a significant difference amongst N Levels at $P = 0.05$. Values in parentheses are standard errors. See Supplementary Materials for ANOVA tables.

**Figure 1.** Coarse root C content for annual winter wheat (triangles) and IWG (circles) for three N levels (Low N (Organic N), Mid N and High N) at five different soil depths. Error bars represent the standard error of the mean and asterisks denote significance at $P < 0.05$ and t denotes significance at $P < 0.1$.




Figure 2. Fine root C in annual winter wheat (triangles) and IWG (circles) for three N levels (Low N (Organic N), Mid N and High N) at five different depths throughout the soil profile. Error bars represent the standard error of the mean and asterisks denote significance at $P < 0.05$ and t denotes significance at $P < 0.1$.

wheat and IWG mainly occurred in the top 10 cm of soil (Table 4). At the surface depth interval, IWG root C concentrations ranged from 28.9 to 33.1% and were greater than in wheat, which ranged from 16.8 to 22.3%. Coarse root C did not significantly differ across N levels ($F = 3.1$, $P = 0.07$). Despite significant overall crop and N level effects on fine root C concentrations ($F = 8.2$, $P = 0.01$ and $F = 3.8$, $P = 0.05$), distinct trends between the two crops for fine root C concentrations were not as apparent compared with those in coarse roots. In general, greater C concentrations were found under the Low N level compared with the Mid N and High N levels (Table 4).

Coarse root N concentrations were significantly greater in the wheat systems compared with IWG (Table 5; $F = 130$, $P < 0.0001$) and decreased significantly by depth ($F = 77$, $P < 0.0001$). There was also a strong N level effect, where coarse root N concentrations were typically greatest in the High N level ($F = 16$, $P < 0.0001$). There was a significant N level \times crop \times depth interaction ($F = 5.2$, $P = 0.0002$), most likely caused by lack of differences across N level and between crops at depths below 40 cm.

Fine root N concentrations differed by crop ($F = 21$, $P = 0.0002$) but not by N level ($F = 4.5$, $P = 0.2$). Wheat had greater N concentrations compared with IWG at almost every depth (Table 5). On average, fine root N

concentrations were 36% greater than coarse root N concentrations for both crops.

The C:N ratio for coarse roots was significantly greater in IWG systems compared with wheat at almost every depth (Fig. 3; $F = 269$, $P < 0.0001$). There was also a strong overall N level effect ($F = 74.8$, $P < 0.0001$), where the coarse root C:N ratio was greater under Low N, especially at lower depths. Similarly, there was an overall crop ($F = 62.5$, $P < 0.001$) and N level ($F = 10.4$, $P < 0.002$) effect for fine root C:N ratio, where IWG had a significantly greater C:N ratio at all depths under Low N and greater C:N ratio in subsurface depths under Mid N and High N (Fig. 4). In addition, there was a significant crop \times N level interaction because IWG was more affected by N level compared with wheat (Fig. 4).

Particulate organic matter C

There were no significant differences in large or medium POM-C concentrations between the two crops (Fig. 5; $F = 0.5$ and $P = 0.5$ and $F = 0$, $P = 0.9$, respectively) or across N levels ($F = 0.3$, $P = 0.8$ and $F = 0.6$, $P = 0.9$, respectively). Large POM-C concentrations were greatest in the top 0–10 cm of soil beneath both crops compared with other depth intervals. Mean IWG large POM-C concentrations at the surface depth were 3.6 ± 0.4 , 3.8 ± 0.8 ,

Table 4. The total amount of C in coarse and fine root biomass at five depths associated with wheat and IWG across N levels (Low N (Organic N), Mid N and High N) at the Kellogg Biological Station in 2013.

N Level	Soil depth Cm	Coarse Root C concentration		Fine Root C concentration	
		Wheat g C kg ⁻¹	IWG g C kg ⁻¹	Wheat g C kg ⁻¹	IWG g C kg ⁻¹
Low N (Organic N)	0–10	22.3 (0.9) ^{Ab}	31.1 (2.8) ^{Bb}	29.8 (1.5) ^{Aa}	28.7 (2.2) ^{Aa}
	10–20	27.6 (1.4) ^{Ab}	31.8 (0.4) ^{Ab}	29.1 (0.9) ^{Aa}	29.9 (1.7) ^{Aa}
	20–40	30.3 (0.8) ^{Aa}	36.1 (1.2) ^{Aa}	30.3 (2.5) ^{Aa}	35.5 (0.3) ^{Ab}
	40–70	21.9 (7.4) ^{Ab}	38.4 (0.6) ^{Bab}	30.8 (1.9) ^{Aa}	33.6 (1.1) ^{Ab}
	70–100	No Roots	28.7 (4.3)	26.5 (4.9) ^{Aa}	28.8 (1.3) ^{Ab}
Mid N	0–10	16.8 (1.2) ^{Ab}	29.8 (2.8) ^{Bb}	28.9 (3.9) ^{Aa}	21.8 (3.2) ^{Bb}
	10–20	25.4 (2.1) ^{Ab}	30.9 (5.10) ^{Bb}	30.2 (1.2) ^{Aa}	17.4 (3.0) ^{Bb}
	20–40	24.2 (2.4) ^{Ab}	25.64 (2.4) ^{Ab}	29.4 (1.2) ^{Aa}	26.5 (3.2) ^{Ac}
	40–70	27.9 (5.6) ^{Ab}	29.7 (0.6) ^{Ab}	31.1 (1.0) ^{Aa}	32.7 (0.4) ^{Ab}
	70–100	No Roots	28.4 (1.4)	32.8 (5.0) ^{Aa}	30.1 (1.1) ^{Ab}
High N	0–10	18.5 (2.5) ^{Ab}	33.3 (1.9) ^{Bb}	31.1 (1.6) ^{Aa}	16.9 (0.8) ^{Ba}
	10–20	21.9 (1.3) ^{Ab}	26.8 (3.3) ^{Ab}	24.4 (3.3) ^{Aa}	17.9 (1.6) ^{Ba}
	20–40	24.8 (2.7) ^{Ab}	33.7 (3.0) ^{Aa}	31.1 (2.3) ^{Aa}	29.1 (3.4) ^{Abc}
	40–70	20.1 (0.6) ^{Ab}	27.5 (2.8) ^{Abc}	28.9 (2.2) ^{Aa}	32.4 (1.1) ^{Ab}
	70–100	12.9 (0.8) ^{Ab}	22.6 (4.9) ^{Ab}	34.0 (1.7) ^{Aa}	30.4 (2.3) ^{Ab}

Comparisons of crop means (between rows) that are followed by different uppercase letters are significantly different at ($P = 0.05$). Within crop and depth interval (columns), different lowercase letters denote a significant difference among N Level at $P = 0.05$. Values in parentheses are standard errors. See Supplementary Materials for ANOVA tables.

Table 5. Total amount of N in coarse and fine root biomass at five depths associated with wheat and IWG across N levels (Low N (Organic N), Mid N and High N) at the Kellogg Biological Station in 2013.

N Level	Soil depth cm	Coarse root N concentrations		Fine root N concentrations	
		Wheat g N kg ⁻¹	IWG g N kg ⁻¹	Wheat g N kg ⁻¹	IWG g N kg ⁻¹
Low N (Organic N)	0–10	0.89 (0.06) ^{Aa}	0.57 (0.03) ^{Bc}	1.30 (0.08) ^{Aa}	0.86 (0.09) ^{Ba}
	10–20	0.48 (0.1) ^{Ab}	0.43 (0.01) ^{Ab}	0.98 (0.02) ^{Aa}	0.77 (0.03) ^{Ba}
	20–40	0.59 (0.06) ^{Abc}	0.30 (0.03) ^{Ab}	0.95 (0.06) ^{Ab}	0.65 (0.06) ^{Ba}
	40–70	0.44 (0.05) ^{Ab}	0.26 (0.04) ^{Ab}	0.74 (0.09) ^{Aa}	0.49 (0.02) ^{Bb}
	70–100	No roots	0.33 (0.06) ^a	0.66 (0.08) ^{Ab}	0.55 (0.06) ^{Aa}
Mid N	0–10	0.8 (0.04) ^{Aa}	0.80 (0.02) ^{Ab}	0.99 (0.2) ^{Aa}	0.68 (0.1) ^{Ba}
	10–20	0.86 (0.04) ^{Aa}	0.62 (0.07) ^{Ba}	1.10 (0.05) ^{Aa}	0.64 (0.09) ^{Bab}
	20–40	0.7 (0.04) ^{Aab}	0.44 (0.02) ^{Aab}	1.0 (0.04) ^{Aab}	0.6 (0.04) ^{Ba}
	40–70	0.77 (0.06) ^{Aa}	0.46 (0.04) ^{Ba}	0.89 (0.08) ^{Aa}	0.66 (0.03) ^{Bab}
	70–100	No roots	0.48 (0.03) ^a	0.92 (0.09) ^{Aa}	0.64 (0.03) ^{Ba}
High N	0–10	0.85 (0.06) ^{Aa}	0.93 (0.01) ^{Aa}	1.09 (0.2) ^{Aa}	0.8 (0.02) ^{Ba}
	10–20	0.84 (0.07) ^{Aa}	0.68 (0.04) ^{Aa}	1.0 (0.09) ^{Aa}	0.59 (0.1) ^{Bb}
	20–40	0.81 (0.05) ^{Aa}	0.52 (0.05) ^{Aa}	1.1 (0.09) ^{Aa}	0.7 (0.06) ^{Ba}
	40–70	0.77 (0.05) ^{Aa}	0.54 (0.09) ^{Aa}	0.9 (0.05) ^{Aa}	0.76 (0.1) ^{Aa}
	70–100	0.38 (0.04) ^{Aa}	0.41 (0.04) ^{Aa}	0.84 (0.2) ^{Aab}	0.67 (0.06) ^{Aa}

Comparisons of crop means (between rows) that are followed by different uppercase letters are significantly different at ($P = 0.05$). Within crop and depth interval (columns), different lowercase letters denote a significant difference among N Level at $P = 0.05$. Values in parentheses are standard errors. See Supplementary Materials for ANOVA tables.

and 4.0 ± 0.7 g C kg soil⁻¹ for Low N, Mid N and High N, respectively. Wheat POM-C concentrations at 0–10 cm depth ranged from 3.6 ± 1.3 to 4.7 ± 0.7 g C kg soil⁻¹, with greater concentrations found in the Low N treatment.

Medium POM-C was greater than the large POM-C across all N levels. At 0–10 cm, IWG medium POM-C ranged from 10.9 ± 1.4 to 11.2 ± 0.6 g C kg soil⁻¹ across N levels, with the Mid N treatment having the lowest concentrations. Surface soil concentrations were very similar




Figure 3. Coarse root C:N ratios for annual winter wheat (triangles) and IWG (circles) for three N levels (Low N (Organic N), Mid N and High N) at five different depths throughout the soil profile. Error bars represent the standard error of the mean and asterisks denotes significance at <0.05, and *t* denotes significance at <0.1.




Figure 4. Fine root C:N ratios for annual winter wheat (triangles) and IWG (circles) for three N levels (Low N (Organic N), Mid N and High N) at five different depths throughout the soil profile. Error bars represent the standard error of the mean and asterisks denote significance at $P < 0.05$.

in wheat systems where concentrations ranged from 9.8 ± 1.6 to 12.6 ± 1.9 g C kg soil⁻¹, again with concentrations slightly higher in the Low N (Organic N) treatment. Pairwise comparisons reveal that large POM-C

concentrations below the 10 cm depth interval were statistically similar to one another ($p > 0.05$). In contrast, medium POM-C fractions significantly decreased by depth to 40 cm ($P < 0.0001$).




Figure 5. Large and Medium POM-C concentrations for annual winter wheat (triangles) and IWG (circles) for three N levels (Low N (Organic N), Mid N and High N) at five different depths throughout the soil profile. Error bars represent the standard error of the mean.

POM-C content accounts for the weight of the fraction, C concentration, bulk density and length of depth interval. There was no difference in large or medium POM-C content between the two crops throughout the soil profile to 1 m (Fig. 6, $F=0$ and $P=0.9$ and $F=0.11$, $P=0.7$, respectively). Approximately 40% of POM-C was found in the top 0–10 cm for both crops. POM-C below 20 cm was evenly distributed throughout the soil profile in the large fraction, but steadily decreased by depth in the medium fraction (Fig. 6). In addition, POM-C content was statistically similar across N levels for both large and medium fractions ($F=0.8$ and $P=0.5$, $F=1.6$, $P=0.2$; respectively).

Power analysis

We conducted a post-hoc power analysis for two different scenarios. First, we used the observed difference between wheat and IWG in the large POM-C at 0–10 cm depth and increased the number of replicates. Secondly, we hypothesized a 15% difference in C between wheat and IWG, while keeping the number of replicates at $n=4$. For scenario one, a total of 52 replicates were needed in order to achieve 78% power. For scenario two, a 15% difference between wheat and IWG large POM-C with four replicates was needed to achieve 84% power.

Discussion

Perennial IWG had significantly greater above and below-ground biomass compared with annual wheat. Despite this considerable difference in biomass, we did not

detect significant differences in large or medium POM-C fractions between wheat and IWG 4 years after establishment.

Crop C and POM-C fractions

Perennial crops are often touted for their greater and more extensive root systems compared with annual crops (Glover *et al.*, 2007), which we found here with total coarse and fine root C stores of IWG between 6 and 15 times greater than root C stores of wheat. The magnitude of differences in root C between IWG and wheat is on par with other studies comparing annual and perennial crops (Jarchow and Liebman, 2012; Anderson-Teixeira *et al.*, 2013). Our findings are also consistent with expectations that perennial grains will have greater root C at subsurface depths (Glover *et al.*, 2010). Significant root C differences between the two crops were detectable to 70 cm depth and demonstrate that perennials are capable of placing greater amounts of root C deeper in the soil profile compared with annual crops. Although the majority of aboveground biomass is removed in both IWG and wheat, there is a portion of aboveground C that is left on the soil surface. Given that IWG has significantly greater straw C than wheat, there could potentially be more aboveground C contributing to soil C stores in IWG systems compared with wheat.

However, despite greater overall aboveground C and up to 15 times more root C within the IWG systems compared with wheat, we did not find significant differences in the large or medium POM-C fractions between the two crops at any depth. Furthermore, we found similar POM-C concentrations across the three N levels, even




Figure 6. Large and Medium POM-C content for annual winter wheat (triangles) and IWG (circles) for three N levels (Low N (Organic N), Mid N and High N) at five different depths throughout the soil profile. Error bars represent the standard error of the mean.

though IWG grown under High N had more root C than IWG grown with Low N and Mid N levels.

In our study, the 0–10 cm large POM-C fraction consisted of recently deposited organic material, such as roots and residue and had a C:N of 19 compared with the 0–10 cm medium POM-C fraction, which comprised finer mineral associated particles and had a C:N of 14. These findings are corroborated by others who associate fractions that have high C:N ratios with more labile C pools compared with fractions with lower C:N ratios, that tend to reflect a more stabilized pool of C (Willson et al., 2001).

Ultimately, these findings show that IWG does not accumulate more C compared with wheat in labile or stabilized pools, which goes against our initial hypothesis. This is surprising given the widespread evidence for gains in soil C under perennial systems compared with annual row-crops (Zan et al., 2001; McLauchlan et al., 2006; and Anderson-Teixeira et al., 2009).

Similar soil C under IWG and annual wheat

One reason for similar soil POM-C status in IWG and in annual wheat systems might be the limited time since conversion and/or establishment. Forest and grassland systems that had increased soil C, reviewed by Post and Kwon (2000), were between 8 and 126 years post-conversion from cropland. In studies that reported soil C accumulation in perennial grasses or cellulosic biofuels compared with annual crops, perennial systems were typically 4–15 years old (Collins et al., 2010 and Rehbein et al., 2015). However not all studies report consistent C gains within this time frame. For example, at a site

within 1 km of our experiment, Syswerda et al. (2011) found greater (10.3%) surface soil C concentrations 12 years post establishment in alfalfa compared with a conventionally managed corn-soybean-wheat system, but no differences between alfalfa and an organic or no-till corn-soybean-wheat system.

An important consideration here is that our study site has relatively sandy soils with low cation exchange capacity (CEC) relative to most Midwestern soils, resulting in an overall lower capacity to accumulate soil C (Table 1). Sandy soils often require more time to build soil C compared with soils with greater clay content (Schneckenberger and Kuzyakov, 2007) and has been shown to be a significant factor in predicting soil C accrual rates (Anderson-Teixeira et al., 2009). In addition, our perennial sites received annual applications of N, which can influence soil C accrual rates both positively and negatively (Anderson-Teixeira et al., 2009). Thus, more time may be needed for soil C to accumulate in these coarse loam soils.

The labile C pool comprises recent inputs from above-ground litter and or root rhizodeposition; thus, if we were to see a difference in C between the two crops, we would expect to see it first in the large POM-C fraction. That we did not detect an increase in soil C within the large POM-C fraction could be because the labile soil C pool under IWG may be lower quality due to slower root decomposition compared with wheat. Thus, IWG roots may be adding C into the large POM-C fraction slower than roots of annual wheat. The C:N ratios of both coarse and fine IWG roots were significantly greater than those for wheat throughout the entire profile, which could lead to reduced turnover and smaller C contributions to

the large POM-C within the initial years of establishment. A higher C:N ratio within perennial roots compared with annual roots is common (Craine *et al.*, 2003) and the greater C content could lead to longer root persistence. Over time, as the roots higher in C content turn over, gains in soil C might be detected under IWG.

Another plausible explanation for similar soil C between the two crops is priming under IWG. Priming occurs when increased root exudates stimulate microbial activity, causing an increase in decomposition rates of older soil C (Cheng, 1999). Strickland *et al.* (2015) found a 21% decline in total soil C under established switchgrass stands compared with total C prior to planting, mainly due to losses in POM-C. They attributed this loss to priming that occurred due to increased microbial activity. In the present study, omnivore nematodes were greater under IWG (Culman *et al.*, unpublished), which could have led to increased decomposition and ultimately less C accumulation. In addition, Culman *et al.* (2013) found significantly greater C mineralization under IWG compared with wheat in the second year of this study, while we found no difference in C mineralization (data not shown) between the two crops 4 years after establishment. An initial pulse in C mineralization under IWG that has since subsided could be due to priming that has resulted in less C accumulation under IWG than was expected.

Results from a power analysis further help explain the similarity of soil C in the wheat and IWG systems. The power analysis for the large POM-C at the 0–10 cm depth interval, where we most expected to see differences between the two systems, revealed that 52 replicates would likely be needed to reach an acceptable probability (78%) of detecting a significant difference ($P = 0.05$) in large POM-C at that time. With four replicates in this study, alternatively, a 15% difference in surface soil C between wheat and IWG would be needed to achieve 84% power. Over time, then, soil C might accumulate sufficiently to reveal a 15% difference in POM-C. However, a long-term experiment would be required to capture such differences and in any case would take longer than the expected 3 year perennial grain rotation now projected (Wagoner, 1990; Culman *et al.*, unpublished). Thus, the power analysis reinforces our inference that a longer rotation interval would be needed in order to detect a difference in C between wheat and IWG.

Vision of perennial grains as a means to accumulate soil C

The concept of perennial grains as a means to increase yields while providing ecosystem services within agricultural landscapes has garnered much attention (Wagoner, 1990; Glover *et al.*, 2007). In particular, proponents of perennial wheat development argue that a perennial version of wheat could lead to crops that are more productive with less need for fertilizers, that ameliorate

erosion and reduce nitrate leaching, and that possess greater water use efficiency (Glover *et al.*, 2010; Kell, 2011; Culman *et al.*, 2013). Proponents especially tout the potential for soil C accrual throughout the soil profile due to deep roots (Asbjornsen *et al.*, 2013; Crews and DeHaan, 2015).

Perennial wheat and IWG yields at KBS are 50 and 70% lower than annual winter wheat yields (Jaikumar *et al.*, 2012; Culman *et al.*, 2013). However, proponents argue that it could still be valuable to farmers who want to improve soil health and other ecosystem services (Adebisi *et al.*, 2016). Four years post establishment, however, we were unable to detect any gains in C accumulation under IWG compared with wheat in either labile or stabilized pools. Four years was an insufficient amount of time to detect gains in C under IWG, especially given the coarse loam soils at this site. Soil C gains could eventually occur because of the large amount of belowground C content found within IWG. That said, since grain yields often decline after 3 or 4 years, gains in C that fail to show up in this time period may never be realized if farmers till and rotate to another crop.

Nevertheless, soil C sequestration is not the only ecosystem service that IWG can provide. For example, Culman *et al.* (2013) found that IWG reduced nitrate leaching up to 99% compared with wheat and Sprunger (2015) found that IWG improved crop-level N use efficiency by up to 42%. In addition, there is evidence that perennial roots persist even after a new annual crop is established (Dupont *et al.*, 2014). Perennial roots could, therefore, contribute to soil C pools after conversion to an annual system. However, the lack of detectable soil C accumulation in a coarse loam soil, 4 years post establishment, weakens the appeal of perennial wheat and IWG.

Conclusions

We measured large and medium POM-C fractions in annual wheat and fourth year perennial IWG across three N levels differing in rates and types of N. Coarse and fine root C were up to 15 times greater under IWG compared with wheat. However, we did not detect any soil C gains under IWG in either labile or stabilized pools. Post-hoc power analysis suggests a much longer time is needed to detect significant soil C accumulation in labile C pools. Since grain yields of perennial IWG often decline after 3 or 4 years, it seems unlikely that detectable (>15%) C gains would be achieved in these low CEC soils before IWG is rotated to a different crop.

Supplementary Material

The supplementary material for this article can be found at <https://doi.org/10.1017/S1742170517000138>

Acknowledgements. We thank M. Freeman, J. Green, and many others for assistance both in the field and laboratory. We thank Lee DeHaan and the Land Institute for supplying Kernza seed and valuable guidance on management. We also thank S. K. Hamilton for suggestions and extensive comments at all stages of this research. In addition, we thank Jane Schuette for help with figure preparation. This work was funded by a USDA Organic Research and Education Initiative grant (2009-51300-05513) and by the NSF Long-Term Ecological Research Program (DEB 1027253) and Michigan State AgBioResearch.

References

- Adebiyi, J., Schmitt Olabisi, L., and Snapp, S. 2016. Understanding perennial wheat adoption as a transformative technology: Evidence from the literature and farmers. *Renewable Agriculture and Food Systems* 31:1–10.
- Anderson-Teixeira, K.J., Davis, S.C., Masters, M.D., and Delucia, E.H. 2009. Changes in soil organic carbon under biofuel crops. *Global Change Bioenergy* 1:75–96.
- Anderson-Teixeira, K.J., Masters, M.D., Black, C.K., Zeri, M., Hussain, M.Z., Bernacchi, C.J., and DeLucia, E.H. 2013. Altered belowground carbon cycling following land-use change to perennial bioenergy crops. *Ecosystems* 16: 508–520.
- Ashbjornsen, H., Hernandez-Santana, V., Liebman, M.Z., Bayala, J., Chen, J., Helmers, M., Ong, C.K., and Schulte, L.A. 2013. Targeting perennial vegetation in agricultural landscapes for enhancing ecosystem services. *Renewable Agriculture and Food Systems* 29:101–125.
- Cambardella, C.A. and Elliott, E.T. 1992. Particulate soil organic matter changes across a grassland cultivation sequence. *Soil Science Society of America Journal* 56:777–783.
- Cheng, W. 1999. Rhizosphere feedbacks in elevated CO₂. *Tree Physiology* 19:313–320.
- Collins, H.P., Smith, J.L., Fransen, S., Alva, A.K., Kruger, C.E., and Granatstein, D.M. 2010. Carbon sequestration under irrigated switchgrass (L.) production. *Soil Science Society of America Journal* 74(6):2049–2058.
- Cox, T.S., Van Tassel, D.L., Cox, C.M., and DeHaan, L.R. 2010. Progress in breeding perennial grains. *Crop and Pasture Science* 61:513–521.
- Craine, J.M., Wedin, D.A., Chapin, F.S., and Reich, P.B. 2003. The dependence of root system properties on root system biomass of 10 North American grassland species. *Plant and Soil* 250:39–47.
- Crews, T.E. and DeHaan, L.R. 2015. The strong perennial vision: A response. *Agroecology and Sustainable Food Systems* 39:500–515.
- Culman, S.W., DuPont, S.T., Glover, J.D., Buckley, D.H., Fick, G.W., Ferris, H., and Crews, T.E. 2010. Long-term impacts of high-input annual cropping and unfertilized perennial grass production on soil properties and belowground food webs in Kansas, USA. *Agriculture, Ecosystems & Environment* 137:13–24.
- Culman, S.W., Snapp, S.S., Freeman, M.A., Schipanski, M.E., Beniston, J., Lal, R., Drinkwater, L.E., Franzluebbers, A.J., Glover, J.D., Grandy, A.S., Lee, J., Six, J., Maul, J.E., Mirksy, S.B.J., Spargo, T., and Wander, M.M. 2012. Permanganate oxidizable carbon reflects a processed soil fraction that is sensitive to management. *Soil Science Society of America Journal* 76:494–506.
- Culman, S.W., Snapp, S.S., Ollenburger, M., Basso, B. and DeHaan, L.R. 2013. Soil and water quality rapidly responds to the perennial grain Kernza Wheatgrass. *Agronomy Journal* 105:735–744.
- Culman, S.W., Snapp, S. S., Sprunger, C.S., Peralta, A. L., DeHaan, L.R. unpublished. In prep. Enhanced ecosystem services under perennial intermediate wheatgrass compared to annual winter wheat.
- DeHaan, L.R., Van Tassel, D.L., and Cox, T.S. 2005. Perennial grain crops: A synthesis of ecology and plant breeding. *Renewable Agriculture and Food Systems* 20:5–14.
- DuPont, S., Beniston, T.J., Glover, J.D., Hodson, A., Culman, S. W., Lal, R., and Ferris, H. 2014. Root traits and soil properties in harvested perennial grassland, annual wheat, and never-tilled annual wheat. *Plant and Soil* 381: 405–420.
- Garten, C.T. and Wulschleger, S.D. 1999. Soil carbon inventories under a bioenergy crop (switchgrass): Measurement limitations. *Journal of Environmental Quality* 28:1359–1365.
- Gebhart, D.L., Johnson, H.B., Mayeux, H.S., and Polley, H.W. 1994. The CRP increases soil organic carbon. *Journal of Soil and Water Conservation* 49:488–492.
- Glover, J.D., Cox, C.M., and Reganold, J.P. 2007. Future farming: A return to roots? *Scientific American* 83:82–89.
- Glover, J.D., Reganold, J.P., Bell, L.W., Borevitz, J., Brummer, E.C., Buckler, E.S., Cox, C.M., Cox, T.S., Crews, T.E., Culman, S.W., DeHaan, L.R., Eriksson, D., Gill, B.S., Holland, J., Hu, F., Hulke, B.S., Ibrahim, A.M. H., Jackson, W., Jones, S.S., and Murray, S.C. 2010. Increased food and ecosystem security via perennial grains. *Science* 328:1638–1639.
- Grandy, A.S., and Robertson, G.P. 2006. Aggregation and organic matter protection following tillage of a previously uncultivated soil. *Soil Science Society of America Journal* 70:1398–1406.
- Houghton, R.A. and Hackler, J.L. 2001. Carbon Flux to the Atmosphere from Land-Use Changes: 1850 to 1990. In R. M. Cushman (ed.). *Trends: A Compendium of Data on Global Change. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, TN.* pp. 305–318.
- Huggins, D.R., Buyanovsky, G.A., Wagner, G.H., Brown, J.R., Darmody, R.G., Peck, T.R., Lesoing, G.W., Vanotti, M.B., and Bundy, L.G. 1998. Soil organic C in the tallgrass prairie-derived region of the corn belt: Effects of long-term crop management. *Soil and Tillage Research* 47:219–234.
- Jaikumar, N.S., Snapp, S.S., Murphy, K., and Jones, S.S. 2012. Agronomic assessment of perennial wheat and perennial rye as cereal crops. *Agronomy Journal* 104:1716–1726.
- Jarchow, M.E. and Liebman, M. 2012. Tradeoffs in biomass and nutrient allocation in prairies and corn managed for bioenergy production. *Crop Science* 52:1330–1342.
- Jarecki, M.K. and R. Lal. 2003. Crop management and soil carbon sequestration. *Critical Reviews in Plant Sciences* 22: 471–502.
- Kell, D.B. 2011. Breeding crop plants with deep roots: Their role in sustainable carbon, nutrient and water sequestration. *Annals of Botany* 108:407–18.
- Knops, J.M.H. and Tilman, D. 2000. Dynamics of soil nitrogen and carbon accumulation for 61 years after agricultural abandonment. *Ecology* 81:88–98.

- Kravchenko, A.N. and Robertson, G.P.** 2011. Whole-profile soil carbon stocks: The danger of assuming too much from analyses of too little. *Soil Science Society of America Journal* 75: 235–240.
- Ladoni, M., Basir, A., and Kravchenko, A.N.** 2015. Which soil carbon fraction is the best for assessing management differences? A statistical power perspective. *Soil Science Society of America Journal* 79:848–857.
- Lal, R.** 2011. Sequestering carbon in soils of agro-ecosystems. *Food Policy* 36:S33–S39.
- McLauchlan, K.K., Hobbie, S.E., and Post, W.M.** 2006. Conversion from agriculture to grassland builds soil organic matter on decadal timescales. *Ecological Applications* 16:143–153.
- Ontl, T.A., Hofmocker, K.S., Cambardella, C.A., Schulte, L.A., and Kolka, R.K.** 2013. Topographic and soil influences on root productivity of three bioenergy cropping systems. *New Phytologist* 199(3):727–737.
- Post, W.M. and Kwon, K.** 2000. Soil carbon sequestration and land-use change: Processes and potential. *Global Change Biology* 6:317–327.
- Poussart, J.N., Adro, J. and Olsson, L.** 2004. Verification of soil carbon sequestration: Sample requirements. *Environmental Management* 33:S416–S425.
- Rasse, D.P., Rumpel, C., and Dignac, M.F.** 2005. Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant and Soil* 269(1–2):341–356.
- Rehbein, K., Sandhage-Hofmann, A., and Amelung, A.** 2015. Soil carbon accrual in particle-size fractions under *Miscanthus* × *giganteus* cultivation. *Biomass and Bioenergy* 78:80–91.
- Robertson, G.P., Paul, E.A., and Harwood, R.R.** 2000. Greenhouse gases in intensive agriculture: Contributions of individual gases to the radiative forcing of the atmosphere. *Science* 289:1922–1925.
- Schmidt, M.W., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A., Kleber, M., Kogel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse, D.P., Weiner, S., Trumbore, S.** 2011. Persistence of soil organic matter as an ecosystem property. *Nature* 478:41–56.
- Schneckenberger, K. and Kuzyakov, Y.** 2007. Carbon sequestration under *Miscanthus* in sandy and loamy soils estimated by natural ^{13}C abundance. *Journal of Plant Nutrition and Soil Science* 170:538–542.
- Six, J., Elliott, E.T., Paustian, K., and Doran, J.W.** 1998. Aggregation and soil organic matter accumulation in cultivated and native grassland soils. *Soil Science Society of America Journal* 62:1367–1377.
- Six, J., Conant, R.T., Paul, E.A., and Paustian, K.** 2002. Stabilization mechanisms of soil organic matter: Implications for c-saturation of soils. *Plant and Soil* 241: 155–176.
- Sprunger, C.D.** 2015. Root production and soil carbon accumulation in annual, perennial, and diverse cropping systems. Dissertation. Michigan State University. ProQuest.
- Strickland, M.S., Leggett, Z.H., Sucre, E.B., and Bradford, M. A.** 2015. Biofuel intercropping effects on soil carbon and microbial activity. *Ecological Applications* 25: 140–150.
- Syswerda, S.P., Corbin, A.T., Mokma, D.L., Kravchenko, A.N., and Robertson, G.P.** 2011. Agricultural management and soil carbon storage in surface vs. deep layers. *Soil Science Society of America Journal* 75:92–101.
- Tiemann, L.K., and Grandy, A.S.** 2015. Mechanisms of soil carbon accrual and storage in bioenergy cropping systems. *Global Change Biology Bioenergy* 7(2):161–174.
- Wagoner, P.** 1990. New use for intermediate wheatgrass. *Journal of Soil and Water Conservation* 45:81–82.
- Wander, M.** 2004. Soil organic matter fractions and their relevance to soil function. In F. Magdoff and R. Weil (ed.). *Soil Organic Matter in Sustainable Agriculture*. CRC Press, Boca Raton. pp. 67–102.
- West, T.O. and Post, W.M.** 2002. Soil organic carbon sequestration rates by tillage and crop rotation: A global data analysis. *Soil Science Society of America Journal* 66:1930–1946.
- Willson, T.C., Paul, E.A., and Harwood, R.R.** 2001. Biologically active soil organic matter fractions in sustainable cropping systems. *Applied Soil Ecology* 16:63–76.
- Zan, C.S., Fyles, J.W., Girouard, P., and Samson, R.A.** 2001. Carbon sequestration in perennial bioenergy, annual corn and uncultivated systems in southern Quebec. *Agriculture, Ecosystems & Environment* 86:135–144.