

## Original Article

**Cite this article:** Soykan O, Gülşahin A, Cerim H (2020). Contribution to some biological aspects of invasive marbled spinefoot (*Siganus rivulatus*, Forsskål 1775) from the Turkish coast of southern Aegean Sea. *Journal of the Marine Biological Association of the United Kingdom* **100**, 453–460. <https://doi.org/10.1017/S0025315420000351>

Received: 13 November 2019

Revised: 19 March 2020

Accepted: 23 March 2020

First published online: 27 April 2020


### Key words:

Aegean Sea; Lessepsian; marbled spinefoot; small-scale fishery

### Author for correspondence:

Hasan Cerim, E-mail: [hasancerim@gmail.com](mailto:hasancerim@gmail.com)

# Contribution to some biological aspects of invasive marbled spinefoot (*Siganus rivulatus*, Forsskål 1775) from the Turkish coast of southern Aegean Sea

Ozan Soykan<sup>1</sup>, Anıl Gülşahin<sup>2,3</sup> and Hasan Cerim<sup>2</sup> 

<sup>1</sup>Ege University, Faculty of Fisheries, İzmir, 35100, Turkey; <sup>2</sup>Mugla University, Faculty of Fisheries, Mugla, 48000, Turkey and <sup>3</sup>Mugla University, Underwater Research Center, Mugla, 48000, Turkey

## Abstract

Some biological aspects of marbled spinefoot (*Siganus rivulatus*) were determined from a southern Aegean Sea small-scale fishery. A total of 721 individuals were captured (335 females, 279 males and 87 immature) and sex ratio was found to be 1:1.27, female to male. Total lengths and ages of individuals ranged between 9.0–26.1 cm and 1–9 years. Asymptotic length ( $L_{\infty}$ ), growth coefficient ( $K$ ) and the theoretical age at zero length ( $t_0$ ) were estimated to be 27.7 cm,  $K = 0.162 \text{ y}^{-1}$  and  $t_0 = -1.87 \text{ y}$ . Length at maturity value was determined as 16.6 cm for females and 16.84 cm for males. Although *S. rivulatus* is one of the target species for small-scale and recreational fishery in the Turkish coast of the southern Aegean Sea, especially in Gökova Bay, official landing data is lacking for Turkey. Results of the present study are of crucial importance for further fisheries management of this invasive species.

## Introduction

Members of the family Siganidae have 29 species worldwide (Froese & Pauly, 2019) and they are known as rabbit fishes. Genus *Siganus* has been represented by two species along the Turkish coast; dusky spinefoot, *Siganus luridus* (Rüppell, 1829) and marbled spinefoot, *Siganus rivulatus* (Forsskål 1775). *Siganus rivulatus* is mostly found in the western Indian Ocean, several localities in East Africa and from the Red Sea to the eastern Mediterranean. It inhabits shallow waters, usually in schools of 50 to several hundred individuals, as a grazer by feeding on algae and generally prefers protected areas (Froese & Pauly, 2019). It is one of the Lessepsian migrants (Ben-Tuvia, 1985) which invaded the Mediterranean and became one of the commercial alien fish for capture fishery (Saoud & Ghanawi, 2010). Additionally, there have been limited studies in terms of the culture potential of this species in the Mediterranean Sea (Stephanou & Georgiou, 2000).

Due to commercial importance of the marbled spinefoot as a food fish, many fishing methods either commercially or recreationally such as spears, gill nets, beach seines, small-mesh nets, cast nets, seine nets, baited hooks and lines (Pacific Community, 2019) have been applied to the species. Global Siganidae capture production from the eastern and western Indian Ocean, Mediterranean Sea and Western Central Pacific was reported to be 157,679 tons in 2017. Furthermore, total capture production of the Siganids in the Mediterranean countries – Cyprus, Egypt, Israel, Lebanon, Libya, Palestine and Syrian Arab Republic – was 1368 tons in 2017 (Fisheries & Aquaculture software, 2019). *Siganus rivulatus* is one of the target species for small-scale fishery and captured by gill nets and baited hooks in the Turkish Coast of the southern Aegean Sea, especially in Gökova Bay. However, there is no official landing data of *S. rivulatus* for Turkey.

There have been some scientific studies on the biology and fisheries of marbled spinefoot in the Red Sea and the eastern Mediterranean. Growth, mortality and yield per recruit of marbled spinefoot were reported by El-Gammal (1988), El-Ganainy (2003), Mehanna & Abdallah (2002) and Gabr *et al.* (2018) in the Red Sea. Moreover, Ben-Tuvia (1986) and Hussein (1986) focused on the reproduction of the species, Lundberg & Lipkin (1993) and Bariche (2006) studied the feeding habits of *S. rivulatus* in the eastern Mediterranean. Besides, different biological aspects including age, growth, mortality and yield per recruit of this species in the eastern Mediterranean were studied by Mohammed (1991), El-Okda (1998), Bariche (2005) and El-Far (2008). Only three scientific studies were conducted on the eastern Mediterranean coast of Turkey by Yeldan & Avşar (2000), Bilecenoglu & Kaya (2002) and Ergenler (2016) and they reported the growth and reproduction of the species. Therefore, information regarding the growth, length at reproduction and age are scarce for the southern Aegean Sea.

The purpose of the current study was to present new and original data on the growth and reproduction of marbled spinefoot from the Turkish coast of the southern Aegean Sea. These data are lacking in the scientific literature for the Aegean coast of Turkey and this study is the first to address this deficiency.





**Fig. 1.** Sampling locations of *Siganus rivulatus* from Gökova Bay, southern Aegean Sea (black points indicate the sampling stations).

## Materials and methods

Fish samples ( $n = 721$ ) were collected between July 2016 and June 2018 from small-scale (trammel nets) and recreational fisheries (by traps and angling) in Gökova Bay, southern Aegean Sea (Figure 1).

Trammel nets were made of polyamide material with 50 mm stretched mesh size and total length of 500–600 m. They were used on the rocky and seagrass substrates between the time of sunset and sunrise (nets stayed underwater for one night). Fishing tackle had a simple design and angling was performed on the rocky shores of Gökova Bay with crumb as the main bait. Fishing trap (locally named as kirtil) was used on the seagrass and rocky substrates and crumb, purslane and lettuce were the baits. Depth of the fishing operations ranged from 1–20 m. Technical characteristics of fishing gears used in the study were given in Figure 2.

Total length (TL) was measured in the natural body position to the nearest millimetre. Total weight ( $W$ ) and gonad weight ( $GW$ ) were measured to the nearest 0.01 g, and sex was recorded after dissection of the specimen. Pooled data were used to draw annual length frequency diagrams. Studies regarding sex and maturity were carried out by macroscopic analysis of the gonads. Maturity stages were assigned in a 5-stage classification scheme: stage I immature, stage II resting, stage III developing, stage IV ripe and stage V spent (Gunderson, 1993). The female to male (F:M) ratio was calculated using only mature individuals, and a chi-square ( $\chi^2$ ) test was applied for determining the significance of the female to male ratio.

The length–weight relationship was calculated with the formula  $W = a \times L^b$ , where  $W$  refers to total body weight (g),  $L$  is total length (cm) where  $a$  and  $b$  are coefficients (Ricker, 1973). The parameters ( $a$  and  $b$ ) of the length–weight relationship were estimated according to linear regression analysis of log-transformed data. Significant difference of  $b$  values from 3, which represent isometric growth, was examined with the  $t$ -test (Pauly, 1993). The degree of association between variables was calculated by the correlation coefficient ( $R^2$ ). Fulton's Condition Factor (CF) values were calculated for males and females separately with the following formulae:

$$CF = \left( \frac{\text{Body Weight} - \text{Gonad Weight}}{\text{Total Length}^3} \right) \times 100$$

Age estimations ( $n = 360$ ) were made by two experienced independent readers who did not have prior access to information on size, sex or date of capture while they were counting growth

increments. The dataset agreed by the independent readers was considered for the estimations. If the readings did not coincide, the vertebra was rejected. Therefore, vertebrae of 307 individuals, covering all size classes, were used for determining the age. Otolith examination process of *S. rivulatus*, especially the removal, requires too much time and effort due to small-sized otoliths. Therefore, age determination was conducted on vertebrae (Campana, 2001; Yalcin et al., 2002). All vertebrae from 13th to 16th were taken out, kept in boiling water for 10 min, cleaned from soft tissues and air dried for age estimations and they were examined in xylol by stereoscopic zoom microscope under reflected light against a black background (Yalcin et al., 2002). The above-mentioned sections of vertebra were bigger than the others which made age reading easier under the binocular and more practical during the cleaning process. Opaque and transparent rings were counted; one opaque zone together with one translucent zone was considered to be the annual growth indicator.

For growth estimation, standard non-linear optimization methods (Sparre & Venema, 1998) were used and von Bertalanffy growth function was fitted to size-at-age data. The function  $L_t = L_\infty [1 - e^{-K(t-t_0)}]$  was fitted to the data, where  $L_t$  is the fish length (cm) at time  $t$  (year),  $L_\infty$  is the asymptotic length (cm),  $K$  is the growth coefficient ( $\text{year}^{-1}$ ), and  $t_0$  (year) is the hypothetical time at when the length is equal to zero. In addition, accuracy of the growth parameters was examined using Munro's growth performance index ( $\phi = \log(K) + 2\log(L_\infty)$ ) and the  $t$ -test (Pauly & Munro, 1984).

The spawning period was established based on monthly variation in the gonadosomatic index (GSI, %) using the equation  $GSI = [GW/(W-GW)] \times 100$ , where  $GW$  is the gonad weight (g) and  $W$  is the total weight (g) of the fish (Ricker, 1975). Length at first maturity ( $L_m$ ) was defined as the length at which 50% of the population investigated was near spawning (King, 1996). The length at 50% maturity was determined with the formula  $L_m = 1/(1 + \exp(-(r \times (TL - TL_{50}))))$  where  $r$  is the slope of the curve,  $TL$  is the mean length of length classes,  $TL_{50}$  is model constant and  $L_m$  is the mean length at sexual maturity or the length which corresponds to a proportion of 0.5 (or 50%) in reproductive condition.

## Results

A total of 721 marbled spinefoot individuals were sampled during the study, including 355 females, 279 males and 87 immature




Fig. 2. Fishing gears for *Siganus rivulatus* capture used in the southern Aegean Sea (Out of scale, A; Trammel net, B; Handline, C; Trap).




Fig. 3. Length-frequency graph of *Siganus rivulatus* from the south-eastern Aegean Sea.

individuals that were excluded from the sex ratio determination. Female:male ratio was calculated to be 1:1.27. Chi-square analysis indicated a significant difference between number of males and females ( $\chi^2$ ,  $P < 0.05$ ). The lengths of fish ranged from 9.0 cm (April) to 26.1 cm (August) TL. Males ranged from 9.4–25.2 cm and females varied from 10.8–26.1 cm (Figure 3). The mean length of the whole sample was calculated to be  $17.9 \pm 0.1$  cm, while  $17.5 \pm 0.2$  and  $18.7 \pm 0.2$  cm were determined for males and females, respectively.

The weights of the samples varied from 8.81–291.26 g. Males were in the range of 11.57–196.08 g and females 14.98–293.42 g. The mean weight of all samples was determined as  $79.0 \pm 1.87$  g, while  $71.0 \pm 2.54$  g was obtained for males and  $91.1 \pm 2.93$  g for females. The length-weight relationship was  $W = 0.00941L^{3.0828}$  ( $R^2 = 0.958$ ) for all individuals. While isometric growth was

observed for males and all individuals, positive allometric growth was detected for females ( $t$ -test,  $P < 0.05$ ; Table 1).



Our results on condition factor (CF) revealed that the values fluctuated between 0.783–1.674 with an average of 1.154 for males and it was between 0.131–2.101 with an average of 1.204 for females.

Age of the stock ranged from 1 to 9 years. Length of the individuals included in the age estimations varied between 9.0 and 26.1 cm TL with a mean length of  $18.2 \pm 0.2$  cm. Regarding the growth, length at infinity was calculated as  $L_\infty = 27.7$  cm. In addition, growth coefficient was found to be  $K = 0.162 \text{ y}^{-1}$  and  $t_0 = -1.87 \text{ y}$  ( $a = 4.14$ ,  $b = -0.15$ ,  $R^2 = 0.985$ ) is the value calculated for the age before birth. The last result on the growth of the species was the growth performance index which was determined to be  $\phi = 2.09$ .

**Table 1.** Length–weight relationship of marbled spinefoot from southern Aegean Sea

	N	<i>a</i>	<i>b</i>	SE <sub><i>b</i></sub>	CL	<i>R</i> <sup>2</sup>	GT
Total	721	0.0094	3.0828	0.06	3.04–3.13	0.958	0
Male	279	0.0133	2.9539	0.06	2.87–3.03	0.950	0
Female	355	0.0066	3.2091	0.06	3.13–3.27	0.956	A+

N, number of individuals; *a* and *b*, intercept and slope of the relationship; *R*<sup>2</sup>, coefficient of determination; SE<sub>*b*</sub>, standard error of slope; CL, 95% confidence limits; GT, growth type (A+: positive allometry, 0: Isometry).

**Fig. 4.** Monthly gonadosomatic index (GSI) for females and males of *Siganus rivulatus* in the southern Aegean Sea between 2016–2018.**Fig. 5.** First reproduction length of female *Siganus rivulatus*. TL, total length.


High GSI values were detected in June and July, but were at the minimum level in September for each sex, whilst individuals ready for spawning were observed mostly between May and August, peaking in June (Figure 4). While gonad formation occurred at 10.5 cm in males and 12.2 cm for females, mature gonads were detected at 15.9 and 16.0 cm for males and females respectively. Moreover, length and age at first reproduction were:  $L_m = 16.6 \pm 0.44$  cm and 1 year ( $a = 216.517$ ,  $b = 0.769$ ,  $R^2 = 0.907$ ) for females,  $L_m = 16.8 \pm 0.48$  cm and 2 years ( $a = 216.543$ ,  $b = 0.645$ ,  $R^2 = 0.833$ ) for males (Figure 5 & 6).

## Discussion

This study addressed first data on age, growth, reproduction, spawning season and some other related biological parameters

of *S. rivulatus* for the Turkish coast of the Aegean Sea. Updated information on population parameters is of crucial importance for alien species such as marbled spinefoot due to their potential ecological impacts. This study is considered to provide useful information for further fisheries management of marbled spinefoot by filling the gap on length at maturity, lifespan and spawning season which are lacking in the Turkish fishery regulations.

*Siganus rivulatus* is considered to be a difficult fish to collect due to its habitat preference. Sampling gears of most studies (Yeldan & Avşar, 2000; Bilecenoğlu & Kaya, 2002; Gökçe et al., 2010) are gill nets, pots and fishing line which limits the total number of samples in comparison to other fishing techniques such as trawl and purse seine nets. The number of samples ( $n = 721$ ,  $L_{\min} - L_{\max} = 9.0 - 26.1$  cm) collected in this study is slightly higher than that of Bilecenoğlu & Kaya (2002) ( $n = 521$ ,



**Fig. 6.** First reproduction length of male *Siganus rivulatus*. TL, total length.

$L_{\min}-L_{\max}=7.0-21.5$  cm), Yeldan & Avşar (2000) ( $n=473$ ) and Ergenler (2016) ( $n=531$ ,  $L_{\min}-L_{\max}=8.5-29.5$  cm). Therefore, the present study includes the highest number of representatives in each length group conducted so far around the surrounding waters of Turkey.

Our data indicated that females dominated the stock in terms of number. While El-Drawany (2015) and Yeldan & Avşar (2000) reported similar results on sex ratio, Ergenler (2016) revealed the dominance of males in İskenderun Bay, eastern Mediterranean coast of Turkey. The range of minimum and maximum lengths ( $TL_{\min}=9.0$  cm,  $TL_{\max}=26.1$  cm) of this study is in between the range of the previous study results (minimum 7.0 cm by Bilecenoğlu & Kaya, 2002; maximum 30.2 by Gabr *et al.*, 2018). Furthermore, the biggest individual on the Turkish Mediterranean coast was reported from İskenderun Bay with a total length of 29.5 cm by Ergenler (2016). Similarity within the length ranges of the reported studies is attributable to sampling gears as most of the studies were performed by artisanal fishery which includes small-scale fishing gears such as trammel nets, pots, traps and handlines.

Length–weight relationship (LWR) is not only an important tool in fisheries management, but also an important growth indicator. LWR parameters vary for same species according to regional and sampling differences and accuracy of the measurements (Gülşahin & Soykan, 2017). The LWR in fish is affected by a number of factors including gonad maturity, sex, diet, stomach fullness, differences in the observed length ranges of the caught specimen and health as well as season and habitat (Froese, 2006). The growth of marbled spinefoot in the southern Aegean Sea was isometric ( $b=3.0828$ ) and it was found that females grew in weight more rapidly than males (Table 2) similar to those inhabiting the Jeddah coast (Saudi Arabia) (Gabr *et al.*, 2018), the İskenderun Bay (Turkey) (Ergenler, 2016) and Bitter Lakes in Egypt (El-Drawany, 2015). On the other hand, Bilecenoğlu & Kaya (2002) stated allometric growth for marbled spinefoot from Antalya Bay (eastern Mediterranean). ‘ $b$ ’ values of previous studies ranged from 2.59 (El-Far, 2008) to 3.32 (Bariche, 2005). Variations on  $b$  values are attributable to availability or lack of food supply and spatio-temporal differences between regions.

Age is a required parameter to assess population dynamics and to determine the state of exploited resources (Allain & Lorange, 2000). For age determination, bony structures such as otoliths and vertebra are considered to be more suitable and satisfactory than scales due to formation timing in the lifespan (Gabr *et al.*,

2018). Clay (1982) also offered that reading age from vertebrae is better than the other methods. Therefore, age estimation of *S. rivulatus* in the present study was based on the counting and interpretation of growth zones formed on the vertebrae. Marbled spinefoot was reported as a long life species (Bilecenoğlu & Kaya, 2002) and our result on maximum age (9 years) confirmed this. Yeldan & Avşar (1998) stated the highest age to be 7 years for all samples from İskenderun Bay. The maximum observed lifespan around Turkish seas was reported as 10 years by Ergenler (2016) from İskenderun Bay. Although aged individuals were announced from the Turkish coast of the Mediterranean, this situation differs in the Red Sea. El Gammal (1988), Mehanna & Abdallah (2002) and Gabr *et al.* (2018) revealed 5 years as the maximum age in the Egyptian coast of the Red Sea. The difference in maximum age between the eastern Mediterranean and Red Sea may be attributed to the sampling and ageing methods used (Bariche, 2005), fishing pressure, different growth coefficients, various biotic and abiotic factors such as temperature, predation and food availability. The presented maximum theoretical length of *S. rivulatus* is slightly higher than the results from comparable studies (Yeldan & Avşar, 1998; Bilecenoğlu & Kaya, 2002) in the Mediterranean Sea. However, Ergenler (2016) reported the  $L_{\infty}$  to be more than twice the results of the abovementioned studies. On the other hand, the maximum theoretical length values reported from the Red Sea are clearly higher than our finding (Table 2). The differences between the growth parameters from different locations (Mediterranean and Red Sea) given in Table 2 are considered due to different factors such as sampling, methodology of ageing or even geographic differences of habitats and condition of the fish samples. Phi-prime index values of the published studies were calculated using reported parameters and they ranged between 2.02 and 2.80. These values were compared with our study’s phi-prime index value of 2.09 and a statistically significant difference was observed between the growth performance index values of our study and the published papers.

Gonad formation in females and males was determined to occur at 10.5 and 12.2 cm total length, respectively. Furthermore, 16.6 cm (females) and 16.8 cm (males) total length values were calculated to be the length at first reproduction. El-Drawany’s (2015) examination of male and female maturity stages indicated that males and females of *S. rivulatus* matured at about 15.5 cm total length in the Bitter Lakes of Egypt. Bariche *et al.* (2003) estimated the length at maturity to be 13.2 cm for males and 13.6 cm for females on the Lebanese

**Table 2.** Length–weight relationship and the growth parameters of previous studies for *S. rivulatus* in the Red Sea and Eastern Mediterranean

Sex	L	LR (Mean ± SE)	a	b	n <sub>1</sub>	L <sub>∞</sub>	K	t <sub>0</sub>	n <sub>2</sub>	φ'	Year(s)	AM	Area	Author
♀	TL	10.8–26.1 (18.7 ± 0.2)	0.0066	3.2091	355						2016–2018	V	Southern Aegean Sea, Turkey	Present study
♂		9.4–25.2 (17.5 ± 0.2)	0.0133	2.9539	279									
Σ		9.0–26.1 (17.9 ± 0.1)	0.0094	3.0828	721	27.7	0.162	−1.87		2.09				
♀	TL	11.2–30.2	0.0123	3.029	1094	38.2	0.27	−0.27		2.59	2013–2014	O	Red Sea coast of Jeddah, Saudi Arabia	Gabr et al. (2018)
♂		11.3–28.3	0.12	3.049	1082	37.9	0.268	−0.24		2.58				
Σ		11.2–30.2	0.011	3.06	2176	38.1	0.275	−0.24		2.60				
Σ	TL	17.2–30.0	0.0077	3.166	2000	34.4	0.38	−0.41		2.65 <sup>a</sup>	2017–2018	O	Southern Red Sea, Egypt	Mehanna et al. (2018)
♀	TL	8.5–29.5	0.013	2.982	166	49.0	0.054	−3.495		2.426	2014–2015	O	İskenderun Bay, Eastern Mediterranean, Turkey	Ergenler (2016)
♂		8.5–26.9	0.015	2.934	365	43.5	0.059	−3.646		2.409				
Σ		8.5–29.5	0.015	2.948	531	47.2	0.054	−3.615		2.412				
♀	TL	9.0–28.9	0.009	3.042		35.5	0.084	−0.843		2.02 <sup>a</sup>	2011	O	Bitter Lakes, Egypt	El-Drawany (2015)
♂		9.0–28.5	0.010	3.01		36.5	0.078	−1.003		2.02 <sup>a</sup>				
Σ		9.0–28.9			420									
♀	TL	6.2–27.0 (19.0 ± 2.9)									2005–2006	O	Libyan coast of Mediterranean	Shakman et al. (2008)
♂		6.3–26.5 (18.8 ± 2.7)												
Σ		6.2–27.4	0.023	2.818	1672	35.0	0.16	−1.04		2.29 <sup>a</sup>				
♀	TL	12.6–26.7(19.1 ± 3.3)	0.00001	3.037	93						1999–2000	O	Lebanese coastal waters	Bariche (2005)
♂		12.7–13.2(17.7 ± 2.5)	0.000002	3.322	98									
Σ		2.5–26.7 (12.1 ± 5.7)	0.00001	3.037	781	31.9	0.225	−1.307	78	2.36 <sup>a</sup>				
♀	TL	7.0–21.5 (15.1 ± 0.3)	0.0064	3.221	292	22.5	0.267	−0.47		2.13 <sup>a</sup>	1996–1998	S	Antalya Bay, Eastern Mediterranean, Turkey	Bilecenoğlu and Kaya (2002)
♂		7.1–20.6 (16.1 ± 0.3)	0.0079	3.135	229	21.6	0.343	−0.54		2.20 <sup>a</sup>				
Σ		7.0–21.5	0.0071	3.179	521	22.3	0.279	−0.503		2.14 <sup>a</sup>				
Σ	TL	9.4–33.7	0.0122	3.018		37.1	0.397	−0.19	425	2.74 <sup>a</sup>	2000–2001	O	Egyptian sector of the Red Sea	Mehanna and Abdallah (2002)
Σ	TL				600	24.4	0.735	−0.22		2.803	2000–2001	–	Gulf of Suez, Egypt	El-Ganainy and Ahmed (2002)
♀	TL	9.0–24.0	0.0061	3.220	224	23.6	0.28	−0.11		2.19 <sup>a</sup>	1995–1996	S	Mersin and İskenderun Bays, Eastern Mediterranean, Turkey	Yeldan and Avşa, (1998)
♂		10.0–22.0	0.0088	3.089	190	20.0	0.40	−0.05		2.20 <sup>a</sup>				
Σ		9.0–24.0 (17.1)	0.0082	3.118	414	21.5	0.34	−0.09		2.20 <sup>a</sup>				

–, not given; ♂, ♀ and å, male, female and total; L, length type (TL: total length); LR, length range (cm); SE, standard error of mean length (cm); a, b and n<sub>1</sub>, intercept and slope of the length–weight relationship, and sample size; L<sub>∞</sub>, k, t<sub>0</sub>, n<sub>2</sub> and φ', asymptotic length (cm), growth coefficient (y<sup>−1</sup>), hypothetical time at which the length is equal to zero (y), sample size and Munro's phi prime index value; Year(s), sampling year/years; AM, Ageing methods; V, vertebrae; S, scales; O, otoliths.

<sup>a</sup>Phi-primes of the studies were calculated by the authors.

coast. Ergenler (2016) reported the sexual maturity length to be 13.91 and 14.1 cm for females and males respectively in İskenderun Bay (eastern Mediterranean coast of Turkey). Yeldan & Aşar (2000) revealed the existence of sexually mature individuals bigger than 15.5 cm in the north-eastern Mediterranean coast of Turkey. Conversely, 20.0 cm TL was declared to be the length of first sexual maturation from the southern Red Sea, Egypt by Mehanna *et al.* (2018). Differences between the previous studies about the length at maturity are relatively small and discrepancies may be attributed to sampling techniques, geographic disparity and the sampling year of the studies. The biggest effect of geographic disparity is undoubtedly on the water temperature. While average seawater surface temperature was 17.2°C in the Mediterranean (southern Aegean Sea – sampling area of the present study), it was reported to be 22.7°C in the Red Sea (World Sea Temperatures, 2020). Therefore, differences in length at maturity values in different locations may be attributable to the water temperature. Although spawning period of the species was reported to be seven months (from March to September) in the Red Sea (Amin, 1985), El-Drawany (2015) stated that the spawning season of the species was between April and August in the Egyptian Bitter Lakes. Bariche *et al.* (2003) revealed the spawning season as June on the Lebanese coast and they announced that high temperature in the summer period was the limiting factor in gonadal development. Yeldan & Aşar (2000) indicated the reproduction period to be July and August in the Turkish coast of the north-eastern Mediterranean. Our results are in accordance with Yeldan & Aşar (2000) as we detected an increase in GSI values starting from May, a peak in June and a decrease in July and August showing that spawning occurred in between June and August in the southern Aegean Sea. It is clear that *S. rivulatus* is a summer spawner species.

One of the major concerns about invasive marbled spinefoot is the lack of information on its stock assessment in the Turkish Seas. Many studies (Stergiou, 1988; Bariche *et al.*, 2004; Moleana *et al.*, 2016) reported that Siganids are strong competitors in terms of feeding and adaptation and *S. rivulatus* was reported as an herbivorous species (Stergiou & Karpouzi, 2002; Bariche, 2006; Pickholtz *et al.*, 2018). Therefore, identifying the probable competition scenarios of the species against other local herbivores such as *Sarpa salpa* and juvenile fish is of crucial importance in terms of fisheries management and preserving the biodiversity. Despite its commercial importance, no regulative tool such as minimum landing size or fishing period limitation is applied on the fishery of the species in Turkey. This discussion raises two complex questions, i.e. ‘Should we somehow limit the fishery of the species?’ or ‘Should we promote the catch pressure on the species?’ Whatever the answer is, the above-mentioned cases indicate that stock assessment studies and long-term monitoring of the Siganid stocks in the Turkish seas should be the first attempt for ecosystem-based fishery management.

The aims of the present study were to investigate the length distribution, length–weight relationship, age, growth, spawning period and length at maturity of invasive marbled spinefoot in the Turkish coast of the southern Aegean Sea and to compare these results with those of previous studies. Although small-scale fishing fleets have been targeting this valuable species for many years especially in the Mediterranean, more scientific information is needed for providing the sustainability of the fisheries. Therefore, additional comprehensive studies are required in order to expand our knowledge of the biology, ecology and fisheries on this species.

**Acknowledgements.** This study is supported by Ege University Scientific Research Projects Coordination Unit. Project Number: 2016/SÜF/014.

## References

- Allain V and Lorance P (2000) Age estimation and growth of some deep-sea fish from the northeast Atlantic Ocean. *Cybiurn* **24**, 7–16.
- Amin EM (1985) Seasonal development changes in the ovaries of *Siganus rivulatus*, Forsskal, from the Red Sea. *Bulletin of the Institute of Oceanography and Fisheries* **11**, 131–147.
- Bariche M (2005) Age and growth of Lessepsian rabbitfish from the eastern Mediterranean. *Journal of Applied Ichthyology* **21**, 141–145.
- Bariche M (2006) Diet of the Lessepsian fishes, *Siganus rivulatus* and *S. lurides* (Siganidae) in the eastern Mediterranean: a bibliographic analysis. *Cybiurn* **30**, 41–49.
- Bariche M, Harmelin-Vivien M and Quignard J (2003) Reproductive cycles and spawning periods of two Lessepsian siganid fishes on the Lebanese coast. *Journal of Fish Biology* **62**, 129–142.
- Bariche M, Letourneur Y and Harmelin-Vivien M (2004) Temporal fluctuations and settlement patterns of native and Lessepsian herbivorous fishes on the Lebanese coast (eastern Mediterranean). *Environment Biology of Fishes* **70**, 81–90.
- Ben-Tuvia A (1985) The impact of the Lessepsian (Suez Canal) fish migration on the eastern Mediterranean ecosystem. In Moraitou-Apostolopoulou M and Kiortsis V (eds), *Mediterranean Marine Ecosystems*. New York, NY: Plenum Press, pp. 367–375.
- Ben-Tuvia A (1986) Siganidae. In Whitehead PJP, Bauchot ML, Hureau JC, Nielsen J and Tortonese E (eds), *Fishes of the Northeastern Atlantic and the Mediterranean*, vol. II, Paris: UNESCO, pp. 964–966.
- Bilecenoğlu M and Kaya M (2002) Growth of marbled spinefoot, *Siganus rivulatus*, (Forsskal, 1775, Teleostei: Siganidae) introduced to Antalya Bay, eastern Mediterranean Sea (Turkey). *Journal of Fisheries Research* **54**, 279–285.
- Campana SE (2001) Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *Journal of Fish Biology* **59**, 197–242.
- Clay D (1982) A comparison of different methods of age determination in the sharp-toothed catfish *Clarias gariepinus*. *Journal of the Limnological Society of Southern Africa* **8**, 61–70.
- El-Drawany M (2015) On the biology of *Siganus rivulatus* inhabits Bitter Lakes in Egypt. *Journal of Aquaculture Research and Development* **6**, 342.
- El-Far A (2008) *Artisanal Fisheries Along Alexandria Coast Area With Special Reference to the Fishery Biology of Siganus spp.* M.Sc. thesis. Department of Zoology, Faculty of Science, Zagazig University.
- El-Gammal FI (1988) Age, growth and mortality of the rabbitfish *Siganus rivulatus* (Forsk., 1775) from the Red Sea. *Bulletin of the Institute of Oceanography and Fisheries* **14**, 13–21.
- El-Ganainy AA (2003) Assessment of the lizard fish *Saurida undosquamis* (Richardson) fishery in the Gulf of Suez, Red Sea. *Bulletin of the Institute of Oceanography and Fisheries* **29**, 1–13.
- El-Ganainy A and Ahmed A (2002) Growth, mortality and yield-per-recruit of the rabbit fish, *Siganus rivulatus*, from the eastern side of the Gulf of Suez, Sinai coast, Red Sea. *Egyptian Journal of Aquatic Biology and Fisheries* **6**, 67–81.
- El-Okda NI (1998) *Comparative Studies on Certain Biological Aspects of Siganus in Marine Waters of Egypt*. Ph.D. thesis. Fac. Sci., (Benha) Zagazig. Univ.
- Ergenler A (2016) *The Investigation of Some Biological Characteristics of Marbled Spinefoot (Siganus rivulatus (Forsskal 1775)) Living In İskenderun Bay*. M.Sc. thesis. İskenderun Teknik Üniversitesi Mühendislik ve Fen Bilimleri Enstitüsü, 53 pp.
- Fisheries and Aquaculture Software (2019) *FishStatJ – Software for Fishery Statistical Time Series*. Rome: FAO. Available at <http://www.fao.org/fishery/statistics/software/fishstatj/en> (Accessed 12 February 2019).
- Froese R (2006) Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology* **22**, 241–253.
- Froese R and Pauly D (2019) Fishbase (version Feb 2018). In Roskov Y, Ower G, Orrell T, Nicolson D, Bailly N, Kirk PM, Bourgoin T, DeWalt RE, Decock W, Nieukerken E, Zarucchi J and Penev L (eds) *Species 2000 & ITIS Catalogue of Life, 2019 Annual Checklist*. Leiden: Species 2000: Naturalis. Available at [www.catalogueoflife.org/col](http://www.catalogueoflife.org/col) (Accessed 12 February 2019).
- Gabr MH, Bakaili AS and Mal AO (2018) Growth, mortality and yield per recruit of the rabbitfish *Siganus rivulatus* (Forsskal 1775) in the Red Sea coast of Jeddah, Saudi Arabia. *International Journal of Fisheries and Aquatic Studies* **6**, 87–96.

- Gökce G, Mustafa C and Filiz H** (2010) Length weight relationship of marine fishes off Yumurtalik coast (Iskenderun Bay), Turkey. *Turkish Journal of Zoology* **34**, 101–104.
- Gülşahin A and Soykan O** (2017) Catch composition, length-weight relationship and discard ratios of commercial longline fishery in the Eastern Mediterranean. *Cahiers de Biologie Marine* **58**, 1–7.
- Gunderson DR** (1993) *Surveys of Fisheries Resources*. New York, NY: John Wiley and Sons, 248 pp.
- Hussein KA** (1986) The use of minute embedded scales for age determination of Mediterranean Sea siganid *Siganus rivulatus* (Forsskal). *Bulletin of the Institute of Oceanography and Fisheries Cairo* **12**, 187–198.
- King RP** (1996) Length-weight relationships of Nigerian freshwater fishes. *Naga: The ICLARM Quarterly* **19**, 49–52.
- Lundberg B and Lipkin Y** (1993) Seasonal, grazing site and fish size effects on patterns of algal consumption by the herbivorous fish, *Siganus rivulatus*, at Mikhmoret (Mediterranean, Israel). *Water Science and Technology* **27**, 413–419.
- Mehanna SF and Abdallah M** (2002) Population dynamics of the rabbitfish *Siganus rivulatus* from the Egyptian sector of the Red Sea. *Journal of King Abdulaziz University: Marine Sciences* **13**, 161–170.
- Mehanna SF, Mohammad AS, El-Mahdyand SM and Osman YA** (2018) Stock assessment and management of the rabbitfish *Siganus rivulatus* from the Southern Red Sea, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries* **22**, 323–329.
- Mohammed N** (1991) *Biological and Biochemical Studies of Some Siganid Fishes from the Mediterranean Waters off Alexandria*. M.Sc. thesis. Faculty of Science, Alexandria University.
- Moleana T, Patrona LD, Meziane T and Letourneur Y** (2016) First record of *Siganus randalli* (Teleost, Siganidae) in New Caledonia, and comments on its diet. *Marine Biodiversity Records* **9**, 1–5.
- Pacific Community** (2019) Coastal fisheries programme, information sheet 02: Rabbitfish (Siganidae). Available at <http://coastfish.spc.int/en/component/content/article/393-guide-and-information-sheets-for-fishing-communities.html> (Accessed 12 February 2019).
- Pauly D** (1993) Fishbyte Section Editorial. *Naga, the ICLARM Quarterly* **16**, 26.
- Pauly D and Munro JL** (1984) Length-converted catch curves: a powerful tool for fisheries research in the Tropics (Part 1). *ICLARM Fishbyte* **1**, 9–13.
- Pickholtz RSM, Kiflawi M and Friedlander AM** (2018) Habitat utilization by an invasive herbivorous fish (*Siganus rivulatus*) in its native and invaded range. *Biological Invasions* **20**, 3499–3512.
- Ricker WE** (1973) Linear regressions in fishery research. *Journal of the Fisheries Research Board of Canada* **30**, 409–434.
- Ricker WE** (1975) Computation and Interpretation of Biological Statistics of Fish Populations. *Bulletin of the Fisheries Research Board of Canada*.
- Saoud IP and Ghanawi J** (2010) Culture potential for marbled spinefoot rabbitfish (*Siganus rivulatus*). *The Practical: Asian Aquaculture* **1**, 15–17.
- Shakman E, Winkler H, Oeberst R and Kinzelbach R** (2008) Morphometry, age and growth of *Siganus luridus* Rüppell, 1828 and *Siganus rivulatus* Forsskal, 1775 (Siganidae) in the central Mediterranean (Libyan coast). *Revista de biología marina y oceanografía* **43**, 521–529.
- Sparre P and Venema SC** (1998) *Introduction to Tropical Fish Stock Assessment, Part I: Manual*. FAO Fisheries Technical Paper. Rome: FAO.
- Stephanou D and Georgiou G** (2000) Recent experiences on the culture of rabbitfish *Siganus rivulatus* in Cyprus. *Cahiers Options Mediterranennes* **47**, 295–301.
- Stergiou KI** (1988) Feeding habits of the Lessepsian migrant *Siganus luridus* in the eastern Mediterranean, its new environment. *Journal of Fisheries Biology* **33**, 531–543.
- Stergiou KI and Karpouzi VS** (2002) Feeding habits and trophic levels of Mediterranean fish. *Reviews in Fish Biology and Fisheries* **11**, 217–254.
- World Sea Temperatures** (2020) Sea surface temperature of Marmaris and Al Quşayr. Available at <https://www.seatemperature.org> (Accessed 04 February 2020).
- Yalcin ÖŞ, Solak K and Akyurt İ** (2002) Growth of the catfish *Clarias gariepinus* (Clariidae) in the River Asi (Orontes), Turkey. *Cybiurn* **26**, 163–172.
- Yeldan H and Avşar D** (1998) The identification of exploitation rate on the rabbitfish (*Siganus rivulatus* Forsskali 1775) stock from the northeastern Mediterranean (in Turkish). XIV. Ulusal Biyoloji Kongresi 7-10 Eylül 1998 Samsun. vol. II. pp. 152–162.
- Yeldan H and Avşar D** (2000) A preliminary study on the reproduction of the rabbitfish (*Siganus rivulatus* (Forsskal, 1775)) in the Northeastern Mediterranean. *Turkish Journal of Zoology* **24**, 173–182.