

# On the decay properties of solutions to the non-stationary Navier–Stokes equations in $\mathbb{R}^3$

**Cheng He**

Institute of Applied Mathematics, Academia Sinica, Beijing, 100080, People's Republic of China, and Institute of Mathematical Sciences, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong (cheng@math03.math.ac.cn)

**Zhouping Xin**

Courant Institute, New York University, 251 Mercer Street, New York, NY 10012, USA, and Department of Mathematics and Institute of Mathematical Sciences, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong (zpxin@ims.cuhk.edu.hk)

(MS received 30 April 1999; accepted 21 January 2000)

In this paper, we study the asymptotic decay properties in both spatial and temporal variables for a class of weak and strong solutions, by constructing the weak and strong solutions in corresponding weighted spaces. It is shown that, for the strong solution, the rate of temporal decay depends on the rate of spatial decay of the initial data. Such rates of decay are optimal.

## 1. Introduction

We consider the decay properties of solutions to the non-stationary Navier–Stokes equations in  $\mathbb{R}^3 \times [0, +\infty)$ :

$$\left. \begin{aligned} \frac{\partial u}{\partial t} - \nu \Delta u + (u \cdot \nabla)u &= -\nabla p, & \text{in } \mathbb{R}^3 \times (0, \infty), \\ \operatorname{div} u &= 0, & \text{in } \mathbb{R}^3 \times (0, \infty), \\ u &\rightarrow 0, & \text{as } |x| \rightarrow +\infty, \\ u(x, 0) &= a(x), & \text{in } \mathbb{R}^3. \end{aligned} \right\} \quad (1.1)$$

Here  $u = u(x, t) = (u_1, u_2, u_3)$  and  $p = p(x, t)$  denote the unknown velocity vector and the pressure of the fluid at point  $(x, t) \in \mathbb{R}^3 \times (0, \infty)$  respectively, while  $\nu > 0$  is the viscosity and  $a(x)$  is a given initial velocity vector field. For simplicity, let  $\nu = 1$ .

Since Leray [25] constructed the weak solutions for (1.1) in  $\mathbb{R}^3$ , and posed the question about the decay properties of his weak solution, there is large literature discussing the decay properties of weak solutions and strong solutions to the non-stationary Navier–Stokes equations [1–4, 7, 8, 10, 11, 14–20, 22, 23, 26–34, 40, 42]. The  $L^2$  decay properties have been extensively studied and the decay rates, similar to

that of heat equation, are also obtained. Their results show: for each  $a \in \mathring{J}^2(\mathbb{R}^3)$ , subspace of  $L^2(\mathbb{R}^3)$  consisting of all solenoidal vector fields, there exists a weak solution  $u$  such that

$$\lim_{t \rightarrow \infty} \|u(t)\|_2 = 0. \tag{1.2}$$

If  $a \in L^r(\mathbb{R}^3) (1 \leq r < 2)$ , then

$$\|u(t)\|_2 \leq Ct^{-(3/r-3/2)/2}. \tag{1.3}$$

Here and after,  $\|\cdot\|_r$  denotes the norm in  $L^r(\mathbb{R}^3)$ . Furthermore, if  $a \in \mathring{J}^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$ , then

$$\|u(t)\|_r \leq C(1+t)^{-(3-3/r)/2} \tag{1.4}$$

for  $1 < r \leq 2$  and

$$\lim_{t \rightarrow \infty} \|u(t)\|_1 = 0. \tag{1.5}$$

If  $\|e^{-tA}a\|_1 \leq C(1+t)^{-\beta}$  for some  $\beta > 0$  with  $A$  being the Stokes operator, then

$$\|u(t)\|_r \leq C(1+t)^{-(3-3/r+\gamma)/2}, \quad \gamma = \min\{1, 2\beta\}, \tag{1.6}$$

for  $1 \leq r \leq 2$ . See [2–4, 19, 26, 27, 29, 31–33, 41]. Schonbek gave the estimates of upper and the lower bound of decay rates for weak solutions in a series of works [32–34]. Moreover, the decay rates of high-order derivatives have been studied in [22, 23] for weak solutions after sufficient large time. Recently, Miyakawa studied the decay rates in Hardy space and established the decay results for weak solutions in some ‘ $L^r$ -like space’ with  $r < 1$ . For details, see [29, 30] and references therein. The time-decay properties are therefore well understood. However, there are few results on the spatial decay properties. Farwing and Sohr [11] showed a class of weighted ( $|x|^\alpha$ ) weak solutions with second derivatives about spatial various and one order derivatives about time variable in  $L^s(0, +\infty; L^q)$  for  $1 < q < 3/2, 1 < s < 2$  and  $0 \leq 3/q + 2/s - 4 \leq \alpha < \min\{1/2, 3 - 3/q\}$ , in the case of exterior domain. In [10], they also showed that there exists a class of weak solutions satisfying

$$\| |x|^{\alpha/2} u \|_2^2 + \int_0^t \| |x|^{\alpha/2} \nabla u \|_2^2 d\tau \leq \begin{cases} C(a, f, \alpha) & \text{if } 0 \leq \alpha < \frac{1}{2}, \\ C(a, f, \alpha', \alpha) t^{\alpha'/2-1/4} & \text{if } \frac{1}{2} \leq \alpha < \alpha' < 1, \\ C(a, f) (t^{1/4} + t^{1/2}) & \text{if } \alpha = 1. \end{cases} \tag{1.7}$$

While in [15], a class of weak solutions

$$(1 + |x|^2)^{1/4} u \in L^\infty(0, +\infty; L^p(\mathbb{R}^3)) \tag{1.8}$$

was constructed for  $6/5 \leq p < 3/2$ , which satisfies (1.7)<sub>3</sub> for  $f = 0$ .

The first purpose of this paper is to construct weak solutions in weighted spaces. We show that, if  $a \in L^1(\mathbb{R}^3) \cap \mathring{J}^2(\mathbb{R}^3)$  and  $|x|a \in L^2(\mathbb{R}^3)$ , there exists a class of weak solution  $u$  satisfying

$$\|(1 + |x|^2)^{1/2} u(t)\|_2^2 + \int_0^t \|(1 + |x|^2)^{1/2} \nabla u(\tau)\|_2^2 d\tau \leq C,$$

which improves the corresponding local weighted estimate as (1.7)<sub>3</sub> in [15]. The interpolation inequality implies that  $u$  satisfies

$$(1 + |x|)^{\alpha}u(t) \in L^{\infty}(0, +\infty; L^p(\mathbb{R}^3))$$

for  $1 < p \leq 2$  and  $\alpha = 2(p-1)/p$ , which also improves estimate (1.8). Furthermore, if  $|x|^{3/2}a \in L^2(\mathbb{R}^3)$ , then we show that there exists a class of weak solutions satisfying

$$\| (1 + |x|^2)^{\alpha/2}u(t) \|_2^2 + \int_0^t \| (1 + |x|^2)^{\alpha/2}\nabla u(\tau) \|_2^2 d\tau \leq C(1 + \log(1 + t)), \quad \forall t \geq 0, \tag{1.9}$$

for  $0 \leq \alpha \leq 3$ . If  $\|e^{-tA}a\|_1 \leq C(1 + t)^{-\gamma}$  for some  $\gamma > 0$ , then the right-hand side of (1.9) can be replaced by a constant independent of  $t$ . It should be noted that Schonbek and Schonbek [35] studied the decay properties of the moment estimate  $\| |x|^{\alpha/2}u \|_2$  for smooth solutions.

The second purpose of this paper is to establish the estimates of decay rate of weighted norm for strong solutions in  $L^p(\mathbb{R}^3)$ . For the existence and decay properties of strong solutions, there are many results, see [1, 5–9, 12–18, 20, 21, 28, 40, 42]. In particular, Heywood showed the existence and decay properties of strong solutions as  $\|a\|_{H^1}$  is small in [17]. Miyakawa [28] improved Heywood’s results by a different method. On the other hand, many results about the existence and decay properties of strong solution have been obtained in space  $L^p(0, T; L^q(\mathbb{R}^n))$  for  $q \geq n \geq 3$  and  $1/p + n/2q \leq 1/2$  in [1, 6–9, 12, 14, 16, 18, 20, 21, 42]. In particular, let  $BC(0, T)$  ( $T > 0$ ) denote the set of bounded and continuous functions defined in  $(0, T)$ , Kato [20] first obtains the following decay rates for strong solutions in  $L^p(\mathbb{R}^n)$

$$t^{(1-n/q)/2}u \in BC([0, +\infty; L^q(\mathbb{R}^n)) \quad \forall n \leq q \leq +\infty, \tag{1.10}$$

$$t^{(1-n/q)/2+1/2}\nabla u \in BC([0, +\infty; L^q(\mathbb{R}^n)) \quad \forall n \leq q < +\infty \tag{1.11}$$

provided that  $a \in L^n(\mathbb{R}^n)$  and  $\|a\|_n$  is small. In the case that  $a \in L^n(\mathbb{R}^n) \cap L^p(\mathbb{R}^n)$  ( $1 < p < n$ ), then

$$t^{(3/p-n/q)/2}u \in BC([1, +\infty; L^q(\mathbb{R}^n)) \quad \forall p \leq q \leq +\infty, \tag{1.12}$$

$$t^{(n/p-n/q)/2+1/2}\nabla u \in BC([1, +\infty; L^q(\mathbb{R}^n)) \quad \forall p \leq q < +\infty \tag{1.13}$$

provided the exponent of  $t$  in (1.12) and (1.13) is smaller than 1; otherwise, (1.12) and (1.13) are valid for any positive number less than 1 as the exponent of  $t$  is bigger than 1. By adding a correction term, Carpio [7] showed that, for initial data  $a$  satisfying (i)  $\|a\|_n$  ( $n \geq 3$ ) is small; (ii)  $a \in L^p(\mathbb{R}^n) \cap L^n(\mathbb{R}^n)$  ( $1 \leq p < n$ ), the solution to the Navier–Stokes equations behaves like the solutions of the heat equations taking the same initial data. Thus Carpio [7] removes Kato’s restriction on exponent of  $t$  and extends the decay estimate to reach the case  $p = 1$ . So this problem is also well understood. While for the spatial decay properties at large distance, a class of weighted strong solutions was constructed, which satisfies that  $(1 + |x|^2)u \in L^{\infty}(0, +\infty; L^2(\mathbb{R}^3))$  and  $t^{1/2}\nabla u \in L^{\infty}(0, +\infty; L^2(\mathbb{R}^3))$ , if  $\|a\|_2 + \|a\|_p$  small for some  $1 \leq p < 2$  in [15], and similar results for exterior domains in [16]. Recently, Takahashi [39] showed that if  $u$  is a smooth solution, then

$$t^{\beta}|x|^{\alpha}|u(x, t)| \leq C \quad \text{for } |x| + t \text{ large} \tag{1.14}$$

for all  $\alpha \geq 0$  and  $\beta \geq 0$  such that  $\alpha + 2\beta < 3$ . Meanwhile, Miyakawa [30] pointed out that if  $a = \partial b / \partial x_1$ , then

$$\sup_{t>0} |e^{-tA} a| \sim C|x|^{-1-n} \quad \text{and} \quad \sup_x |e^{-tA} a| \sim Ct^{-(n+1)/2},$$

as  $|x| \rightarrow \infty$  and  $t \rightarrow \infty$ . Thus he conjectured that if  $u$  is a smooth solution, then  $u$  should satisfy that

$$|u(x, t)| \sim C|x|^{-\alpha}t^{-\beta} \quad \text{as } |x| + t \rightarrow \infty, \tag{1.15}$$

for all  $\alpha \geq 0$  and  $\beta \geq 0$  such that  $\alpha + 2\beta = n + 1$ .

In this paper, we show that, with the hypothesis that  $\|a\|_1 + \|a\|_2$  is small, there exists a unique strong solution  $u$  such that

$$t^\beta \|(1 + |x|^2)^{\alpha/2} u(t)\|_p \leq C \quad \text{for } t \geq 0 \tag{1.16}$$

and

$$t^{\beta+1/2} \|(1 + |x|^2)^{\alpha/2} \nabla u(t)\|_p \leq C \quad \text{for } t \geq 0 \tag{1.17}$$

with  $\alpha = 3 - 3/p_0$ ,  $\beta = (3/p_0 - 3/p)/2$  for  $1 \leq p_0 \leq p \leq +\infty$  and  $3 < p$ . So  $\alpha + 2\beta = 3 - 3/p$ . Therefore, if  $p = \infty$ , our results improve (1.14). Moreover, the time decay rates of weighted norm of the solutions also improve the results of Kato's in the sense that there is no restriction on exponent of  $t$  and  $p_0$  can be taken to be 1. If  $a = \partial b / \partial x_i$  for some  $i = 1, 2, 3$ , we show that, if  $\|a\|_1 + \|a\|_2$  small, there exists a unique strong solution  $u$ , which satisfies

$$t^{\beta+1/2} \|(1 + |x|^2)^{\alpha/2} u(t)\|_p \leq C \quad \text{for } t \geq 0. \tag{1.18}$$

In the special case  $p = \infty$ , our result (1.18) shows that Miyakawa's conjecture (1.15) is right. Finally, it should be noted that in order to obtain the global strong solutions, we assume the smallness of the initial data, i.e.  $\|a\|_1 + \|a\|_2 \leq \delta$ , which is different from any previous known small assumptions.

The paper is organized as follows. In §2, we state our main results. A new class of approximate solutions is constructed and then the integral representations are delivered in §3. In §§4 and 5, we establish the main weighted estimates for weak and strong solutions.

We conclude this introduction by listing some notation used in the rest of the paper.

Let  $L^p(\mathbb{R}^3)$ ,  $1 \leq p \leq +\infty$ , represent the usual Lebesgue space of scalar functions as well as that of vector-valued functions with norm denoted by  $\|\cdot\|_p$ . Let  $C_{0,\sigma}^\infty(\mathbb{R}^3)$  denote the set of all  $C^\infty$  real vector-valued functions  $\phi = (\phi_1, \phi_2, \phi_3)$  with compact support in  $\mathbb{R}^3$ , such that  $\text{div } \phi = 0$ .  $\dot{J}^p(\mathbb{R}^3)$ ,  $1 \leq p < \infty$ , is the closure of  $C_{0,\sigma}^\infty(\mathbb{R}^3)$  with respect to  $\|\cdot\|_p$ .  $H^m(\mathbb{R}^3)$  denotes the usual Sobolev Space. Finally, given a Banach space  $X$  with norm  $\|\cdot\|_X$ , we denote by  $L^p(0, T; X)$ ,  $1 \leq p \leq +\infty$ , the set of function  $f(t)$  defined on  $(0, T)$  with values in  $X$  such that  $\int_0^T \|f(t)\|_X^p dt < +\infty$ . Let  $P$  be the Helmholtz projection from  $L^p(\mathbb{R}^3)$  to  $\dot{J}^p(\mathbb{R}^3)$ . Then the Stokes operator  $A$  is defined by  $A = -P\Delta$  with  $D(A) = H^2(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$ . Let

$$D^2 = \sum_{i,j} \left| \frac{\partial^2}{\partial x_i \partial x_j} \right| \quad \text{and} \quad D^3 = \sum_{j,k} \left| \frac{\partial^3}{\partial x_i \partial x_j \partial x_k} \right|.$$

At last, we denote a generic constant by  $C$ , the value of which is inessential to our aims, and which may change from line to line.

**2. The main results**

We first give the definitions of weak and strong solutions.

DEFINITION 2.1.  $u$  is called a weak solution to the Cauchy problem (1.1) if

- (i)  $u \in L^\infty(0, T; L^2(\mathbb{R}^3)) \cap L^2(0, T; H^1(\mathbb{R}^3))$  for any  $T > 0$ ;
- (ii)  $u$  satisfies the equations (1.1) in the sense of distribution, i.e.

$$\int_0^\infty \int_{\mathbb{R}^3} \left( -\frac{\partial \phi}{\partial \tau} u + \nabla u \cdot \nabla \phi + (u \cdot \nabla) u \cdot \phi \right) dx d\tau = \int_{\mathbb{R}^3} \phi(x, 0) a(x)$$

for every  $\phi \in C_{0,\sigma}^\infty(R \times \mathbb{R}^3)$ .

- (iii)  $\operatorname{div} u = 0$  in the sense of distribution, i.e.

$$\int_{\mathbb{R}^3} u(x, t) \nabla \psi(x) = 0$$

for every  $\psi \in C_0^\infty(\mathbb{R}^3)$ .

DEFINITION 2.2.  $u$  is called a strong solution to the Cauchy problem (1.1) if  $u \in L^\infty(0, T; L^p(\mathbb{R}^3))$  for  $3 < p \leq +\infty$  and any  $T > 0$ , and (ii) and (iii) in definition 2.1 hold for  $u$ .

The main results in this paper are described in the following theorems. Our first result concerns the existence and global estimates of weak solutions in a weighted  $L^2$ -space.

THEOREM 2.3. Let  $a \in L^1(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$  and  $(1 + |x|^2)^{1/2} a \in L^2(\mathbb{R}^3)$ . Then there exists a weak solution  $u$  in  $L^\infty(0, +\infty; L^2(\mathbb{R}^3))$  to (1.1) such that

$$\|u(t)\|_2^2 + \int_0^t \|\nabla u(\tau)\|_2^2 d\tau \leq 4\|a\|_2^2 \tag{2.1}$$

and

$$\|(1 + |x|^2)^{1/2} u\|_2^2 + \int_0^t \|(1 + |x|^2)^{1/2} \nabla u\|_2^2 d\tau \leq CA_1 \tag{2.2}$$

for any  $t \geq 0$ . Moreover, for  $t \geq 0$ ,

$$\|u(t)\|_2 \leq CN(1 + t)^{-3/4} \tag{2.3}$$

and

$$\|u(t)\|_1 \leq CB. \tag{2.4}$$

Here

$$\begin{aligned} A_1 &= e^{C\|a\|_2^2} (\|(1 + |x|^2)^{1/2} a\|_2^2 + N^2), \\ N &= \|a\|_1 + \|a\|_1^2 + \|a\|_2 + \|a\|_2^2, \\ B &= \|a\|_1 + \|a\|_2 N. \end{aligned}$$

In the case that the initial velocity field possesses higher-order moments, we have the following estimates.

**THEOREM 2.4.** *Let  $a \in L^2(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$  and  $|x|^{3/2}a \in L^2(\mathbb{R}^3)$ , then there exists a weak solution  $u$  in  $L^\infty(0, +\infty; L^2(\mathbb{R}^3))$  to (1.1), which satisfies that*

$$\int_{\mathbb{R}^3} (1+|x|^2)^{3/2}u^2 dx + \int_0^t \int_{\mathbb{R}^3} (1+|x|^2)^{3/2}|\nabla u|^2 dx d\tau \leq C(A_2 + B^{2/3}N^{4/3} \log(1+t)) \tag{2.5}$$

for any  $t \geq 0$ , (2.1)-(2.4) are valid for  $u$ . Moreover, if  $\|e^{-tA}a\|_1 \leq C(1+t)^{-\gamma}$  for some  $\gamma > 0$ , then

$$\int_{\mathbb{R}^3} (1+|x|^2)^{3/2}u^2 dx + \int_0^t \int_{\mathbb{R}^3} (1+|x|^2)^{3/2}|\nabla u|^2 dx d\tau \leq C(A_2 + B^{2/3}N^{4/3}) \tag{2.6}$$

for any  $t \geq 0$  with  $A_2 = A_1^{3/2}(\|a\|_2^{5/4}N^{3/4} + N\|a\|_2^{5/6})$ .

Next, the weighted norm (both in time and in space) estimates of strong solutions are established in the following theorem.

**THEOREM 2.5.** *Let  $a \in L^1(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$ ,  $(1+|x|^2)^{1/2}a \in L^2(\mathbb{R}^3)$  and  $(1+|x|^2)^{\alpha/2}a \in L^{p_0}(\mathbb{R}^3)$  for  $1 \leq p_0 \leq +\infty$  and  $\alpha = 3 - 3/p_0$ . Then there exists a constant  $\lambda > 0$ , such that if  $\|a\|_1 + \|a\|_2 \leq \lambda$ , then there exists a unique strong solution  $u \in L^\infty(0, +\infty; L^p(\mathbb{R}^3))$  for  $3 < p \leq +\infty$ , which satisfies the estimate*

$$\|t^\beta(1+|x|^2)^{\alpha/2}u\|_p + \|t^{1/2+\beta}(1+|x|^2)^{\alpha/2}\nabla u\|_p \leq C(\|(1+|x|^2)^{\alpha/2}a\|_{p_0} + N^2 + A_1^{5/2}B^{3/2} + B^{1/2}A_1N) \tag{2.7}$$

for any  $t \geq 0$  and  $\beta = (3/p_0 - 3/p)/2$ .

Finally, for a class of special initial data, the results in theorem 2.5 can be improved by the following theorem.

**THEOREM 2.6.** *Let  $a \in L^1(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$  and  $(1+|x|^2)^{1/2}a \in L^2(\mathbb{R}^3)$ . Let  $a = \partial b/\partial x_i$  for some  $i = 1, 2, 3$  with  $b \in L^1(\mathbb{R}^3)$  and  $(1+|x|^2)^{\alpha/2}b \in L^{p_0}(\mathbb{R}^3)$  for  $1 \leq p_0 \leq +\infty$  and  $\alpha = 3 - 3/p_0$ . Then there exists a constant  $\lambda_0 > 0$ , such that if  $\|a\|_1 + \|a\|_2 \leq \lambda_0$ , then there exists a unique strong solution  $u \in L^\infty(0, +\infty; L^p(\mathbb{R}^3))$  for  $3 < p \leq +\infty$ , which satisfies the estimate*

$$\|t^{1/2+\beta}(1+|x|^2)^{\alpha/2}u\|_p \leq C(\|(1+|x|^2)^{\alpha/2}b\|_{p_0} + \|b\|_1 + N^2 + A_1^{5/2}B^{3/2} + B^{1/2}A_1N) \tag{2.8}$$

for any  $t \geq 0$  and  $\beta = (3/p_0 - 3/p)/2$ .

**REMARK 2.7.**

- (i) By the interpolation inequality, (2.3) and (2.4) imply that the solution  $u$ , obtained in theorems 2.3 and 2.4, satisfies the decay property (1.4), which for weak solutions has already been obtained in, for example, [3, 19]. However, (1.5) is not a consequence.

(ii) The weak solution  $u$ , obtained in theorem 2.3, also satisfies that

$$(1 + |x|)^\alpha u \in L^\infty(0, +\infty; L^p(\mathbb{R}^3))$$

for  $1 < p \leq 2$  and  $\alpha = 2(p - 1)/p$ , which also improves the corresponding results in [10, 11, 15].

(iii) Schonbek and Schonbek [35] studied the decay properties of the moment estimate  $\| |x|^\alpha u \|_2$  for  $0 \leq \alpha \leq 3/2$ , when  $u$  is a smooth solution. While we get estimates (2.5) and (2.6) for the weak solution. Moreover, the weak solution  $u$ , obtained in theorem 2.4, satisfies that, for any  $0 \leq \alpha \leq 3$ ,

$$\int_{\mathbb{R}^3} |x|^\alpha |u|^2 dx + \int_0^t \int_{\mathbb{R}^3} |x|^\alpha |\nabla u|^2 dx d\tau$$

can be dominated by the terms on the right-hand sides of (2.5) or (2.6), respectively.

(iv) In theorem 2.4, the assumption  $\| e^{-tA} a \|_1 \leq C(1 + t)^{-\gamma}$  holds for some  $\gamma > 0$ , if  $a = A^\gamma b$  for some  $b \in L^1(\mathbb{R}^3)$ .

(v) In theorem 2.5,  $\alpha + 2\beta = 3 - 3/p$ . When  $p = +\infty$ , our result improves Takahashi’s result. Moreover, we obtain the decay rate similar to that of Kato, for weighted norm. At same time, we remove the restriction on the exponent of  $t$  and extend the decay rate to reach to the case  $p_0 = 1$ . By adding a correction term, Carpio [7] have showed the solution to the Navier–Stokes equations behaves like the solution of the heat equation, with the same initial data, in  $L^q(\mathbb{R}^n)$  ( $n \geq 3$ ) for  $q \geq p$ , as initial data  $a$  satisfying (i)  $\| a \|_n$  is small, (ii)  $a \in L^p(\mathbb{R}^n) \cap L^n(\mathbb{R}^n)$  for  $1 \leq p < n$ . But our results are different from that in [7].

(vi) Taking  $p = \infty$  in theorem 2.6, then estimate (2.8) yields (1.15), which gives an assertive answer to Miyakawa’s conjecture.

(vii) Let  $p_0 = 1$  and  $p = \infty$  in theorem 2.6, then

$$\| u(t) \|_\infty = O(t^{-2}),$$

which has been proved by Miyakawa under different assumptions on initial data  $a$ . See theorem 1.9(i) in [30].

Applying the weighted estimates obtained in §§ 4 and 5, the proof of theorems 2.3–2.6 are standard. So we will only deduce the necessary weighted estimates, and omit the details of the procedure of the proof.

### 3. The approximation solutions and their integral representations

In this section, we construct a sequence of approximate solutions by using the linearized Navier–Stokes equations in  $\mathbb{R}^3$ , and derive the integral representations of the approximate solutions. First, let  $a \in \dot{J}^p(\mathbb{R}^3) \cap \dot{J}^q(\mathbb{R}^3)$  ( $1 \leq p, q \leq +\infty$ ). We select  $a^k \in C_{0,\sigma}^\infty(\mathbb{R}^3)$ , such that

$$a^k \rightarrow a \quad \text{in } \dot{J}^p(\mathbb{R}^3) \cap \dot{J}^q(\mathbb{R}^3) \text{ strongly}$$

and

$$\|a^k\|_p \leq 2\|a\|_p, \quad \|a^k\|_q \leq 2\|a\|_q. \tag{3.1}$$

The approximate solutions are defined as follows: let  $(u^0, p^0)$  solve the Cauchy problem of the Stokes equations,

$$\left. \begin{aligned} \frac{\partial u^0}{\partial t} - \Delta u^0 &= -\nabla p^0, & \text{in } \mathbb{R}^3 \times (0, \infty), \\ \operatorname{div} u^0 &= 0, & \text{in } \mathbb{R}^3 \times (0, \infty), \\ u^0 &\rightarrow 0, & \text{as } |x| \rightarrow +\infty, \\ u^0(x, 0) &= a^0(x), & \text{in } \mathbb{R}^3, \end{aligned} \right\} \tag{3.2}$$

and  $(u^k, p^k)$  ( $k \geq 1$ ) solve the Cauchy problem for the linearized Navier–Stokes equations

$$\left. \begin{aligned} \frac{\partial u^k}{\partial t} - \Delta u^k + (u^{k-1} \cdot \nabla)u^k &= -\nabla p^k, & \text{in } \mathbb{R}^3 \times (0, \infty), \\ \operatorname{div} u^k &= 0, & \text{in } \mathbb{R}^3 \times (0, \infty), \\ u^k &\rightarrow 0, & \text{as } |x| \rightarrow +\infty, \\ u^k(x, 0) &= a^k(x), & \text{in } \mathbb{R}^3, \end{aligned} \right\} \tag{3.3}$$

for  $k \geq 1$ . It is well known (cf. [24]) that there exists a unique solution  $u^k$  ( $k \geq 0$ ) to (3.2) and (3.3) satisfying

$$\frac{\partial u^k}{\partial t}, \quad \frac{\partial u^k}{\partial x_i}, \quad \frac{\partial^2 u^k}{\partial x_i \partial x_j}, \quad \frac{\partial p^k}{\partial x_i} \in L^2(0, T; L^2(\mathbb{R}^3)) \tag{3.4}$$

for  $i, j = 1, 2, 3, k \geq 0$  and any  $T > 0$ .

In order to derive an integral expression for  $u^k$ , one can use the singular integral expression of the projection operator  $P : L^2(\mathbb{R}^3) \rightarrow \dot{J}^2(\mathbb{R}^3)$ , that is,

$$P\phi = \phi + \frac{1}{4\pi} \nabla \operatorname{div} \int_{\mathbb{R}^3} \frac{\phi(y)}{|x-y|} dy \tag{3.5}$$

for any  $\phi \in L^2(\mathbb{R}^3)$  (cf. [24]). Applying the fundamental solution of the heat equation, we can rewrite the solution to the Cauchy problem for the Stokes equations,

$$\left. \begin{aligned} \frac{\partial v}{\partial t} - \Delta v &= -\nabla p + f, \\ \operatorname{div} v &= 0, \\ v(x, 0) &= 0, \end{aligned} \right\}$$

as

$$v_i = \int_0^t \int_{\mathbb{R}^3} V^i(x-y, t-\tau) \cdot f(y, \tau) dy d\tau, \quad i = 1, 2, 3, \tag{3.6}$$

where

$$\left. \begin{aligned} V^i(x, t) &= \Gamma(x, t)e^i + \frac{1}{4\pi} \nabla \frac{\partial}{\partial x_i} \int_{\mathbb{R}^3} \frac{\Gamma(x-z, t)}{|z|} dz = P(\Gamma e^i), \\ \Gamma(x, t) &= (4\pi t)^{-3/2} e^{-|x|^2/4t}, \end{aligned} \right\} \tag{3.7}$$

and  $e^i$  is the unit vector along  $x_i$ -axis. It is easy to see that

$$V^i(x, t) = \text{curl}(\text{curl}\omega^i) = -\Delta\omega^i + \nabla \text{div}\omega^i, \quad i = 1, 2, 3$$

with

$$\left. \begin{aligned} \omega^i(x, t) &= \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\Gamma(x - z, t)}{|z|} dz e^i, \\ \bar{\theta}(x, t) &= \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{\Gamma(x - z, t)}{|z|} dz. \end{aligned} \right\} \quad (3.8)$$

For the detailed derivation of (3.7) and (3.8), see Ladyzhenskaya [24].

For simplicity, we drop the right upper label  $k$  of the solution  $u^k$  of (3.3) and use  $b$  to denote  $u^{k-1}$ . Let  $y$  and  $\tau$  denote the variables in equations (3.3). We multiply both sides of (3.3) by  $V^i(x - y, t - \tau)$ , then integrate for  $y \in \mathbb{R}^3$  and  $\tau \in [0, t - \varepsilon]$  for arbitrary  $0 < \varepsilon < t$ , to get

$$\begin{aligned} \int_0^{t-\varepsilon} \int_{\mathbb{R}^3} \left( \frac{\partial u}{\partial \tau} - \Delta u \right) (y, \tau) V^i(x - y, t - \tau) dy d\tau \\ = \int_0^{t-\varepsilon} \int_{\mathbb{R}^3} (-\nabla_y p - (b \cdot \nabla)u)(y, \tau) V^i(x - y, t - \tau) dy d\tau. \end{aligned}$$

Since  $(-\partial/\partial\tau - \Delta_y)V^i = 0$  and  $V^i = P(\Gamma e^i)$ , it follows that

$$\begin{aligned} \int_{\mathbb{R}^3} u(y, t - \varepsilon) V^i(x - y, \varepsilon) dy - \int_{\mathbb{R}^3} a(y) V^i(x - y, t) dy \\ = - \int_0^{t-\varepsilon} \int_{\mathbb{R}^3} (b \cdot \nabla)u(y, \tau) V^i(x - y, t - \tau) dy d\tau. \end{aligned}$$

Since  $u$  is divergence free, so it follows, by the structure of  $V^i(x - y, t - \tau)$ , that

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3} u(y, t - \varepsilon) V^i(x - y, \varepsilon) dy = u_i(x, t),$$

where  $u_i$  denotes the  $i$ th component of the vector  $u$ . Thus,

$$u_i = - \int_0^t \int_{\mathbb{R}^3} (b \cdot \nabla)u(y, \tau) V^i(x - y, t - \tau) dy d\tau + \int_{\mathbb{R}^3} a(y) V^i(x - y, t) dy.$$

Substituting (3.7) into above equation, we get that

$$\begin{aligned} u_i = - \int_0^t \int_{\mathbb{R}^3} (b \cdot \nabla)u(y, \tau) \Gamma(x - y, t - \tau) e^i dy d\tau \\ - \int_0^t \int_{\mathbb{R}^3} (b \cdot \nabla)u(y, \tau) \nabla \frac{\partial}{\partial y_i} \bar{\theta}(x - y, t - \tau) dy d\tau \\ + \int_{\mathbb{R}^3} a(y) \Gamma(x - y, t) e^i dy. \quad (3.9) \end{aligned}$$

By integration by parts, we arrive at the desired integral representation form:

$$u_i = \int_0^t \int_{\mathbb{R}^3} \sum_j b_j u_i(y, \tau) \frac{\partial}{\partial y_j} \Gamma(x - y, t - \tau) dy d\tau \tag{3.10}$$

$$+ \int_0^t \int_{\mathbb{R}^3} \sum_{l,k=1}^3 b_l u_k(y, \tau) \frac{\partial^3}{\partial y_i \partial y_l \partial y_k} \bar{\theta}(x - y, t - \tau) dy d\tau \tag{3.11}$$

$$+ \int_{\mathbb{R}^3} a(y) \Gamma(x - y, t) e^i dy. \tag{3.12}$$

Let

$$\left. \begin{aligned} I_1^k &= \int_{\mathbb{R}^3} |a^k|(y) \Gamma(x - y, t) dy, \\ I_2^k &= \int_0^{t/2} \int_{\mathbb{R}^3} |u^{k-1}| |u^k|(y, \tau) (|\nabla \Gamma| + |D^3 \bar{\theta}|)(x - y, t - \tau) dy d\tau, \\ I_3^k &= \int_{t/2}^t \int_{\mathbb{R}^3} |u^{k-1}| |u^k|(y, \tau) (|\nabla \Gamma| + |D^3 \bar{\theta}|)(x - y, t - \tau) dy d\tau. \end{aligned} \right\} \tag{3.13}$$

Thus

$$|u^k(x, t)| \leq C(I_1^k + I_2^k + I_3^k). \tag{3.14}$$

For  $\Gamma$  and  $\bar{\theta}$ , direct calculations show that

$$\left. \begin{aligned} |D^m \Gamma(x, t)| &\leq C_m (|x|^2 + t)^{-(m+3)/2}, \\ |D^m \bar{\theta}(x, t)| &\leq C_m (|x|^2 + t)^{-(m+1)/2}. \end{aligned} \right\} \tag{3.15}$$

for  $m \in N$ .

#### 4. Weighted estimates for the approximate solutions I

In this section, we establish some *a priori* estimates for the approximate solutions constructed in §3, which result in theorems 2.3 and 2.4 by standard compactness argument. First, standard energy estimates yield the following.

LEMMA 4.1. *Let  $a \in \dot{J}^2(\mathbb{R}^3)$ . Then the estimates,*

$$\left. \begin{aligned} \|u^k(t)\|_2 &\leq 2\|a\|_2 \quad \forall t > 0, \\ \int_0^\infty \|\nabla u^k(s)\|_2^2 ds &\leq 4\|a\|_2^2, \end{aligned} \right\} \tag{4.1}$$

hold uniformly for  $k \geq 0$ .

The next lemma follows from (3.12)–(3.15), (4.1) and standard convolution estimates.

LEMMA 4.2. *Let  $a \in L^1(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$ . Then we have*

$$\|u^k(t)\|_2 \leq CNt^{-3/4} \tag{4.2}$$

and

$$\|u^k(t)\|_1 \leq CB \tag{4.3}$$

holds uniformly for  $k \geq 0$  and  $t > 0$ . Furthermore, if  $\|e^{-tA}a\|_1 \leq C(1+t)^{-\gamma}$  for some  $\gamma > 0$ , then

$$\|u^k(t)\|_2 \leq Ct^{-3/4-\gamma_1/2}, \tag{4.4}$$

for  $\gamma_1 = \min\{1, 2\gamma\}$ , where  $N = \|a\|_1 + \|a\|_1^2 + \|a\|_2 + \|a\|_2^2$  and  $B = \|a\|_1 + \|a\|_2N$ .

We now turn to the main weighted norm estimates in theorems 2.3 and 2.4.

LEMMA 4.3. Let  $a \in L^1(\mathbb{R}^3)$  and  $(1 + |x|^2)^{1/2}a \in L^2(\mathbb{R}^3)$ . Then

$$\|(1 + |x|^2)^{1/2}u^k\|_2^2 + \int_0^t \|(1 + |x|^2)^{1/2}|\nabla u^k|\|_2^2 dx d\tau \leq CA_1 \tag{4.5}$$

for any  $k \geq 0$  and  $t \geq 0$  with  $A_1 = e^{C\|a\|_2^2}(\|(1 + |x|^2)^{1/2}a\|_2^2 + N^2)$ .

Proof. Taking the divergence of the first equations of (3.3) yields

$$-\Delta p^k = \sum_{ij=1}^3 \frac{\partial^2}{\partial x_i \partial x_j} (u_i^{k-1}u_j^k). \tag{4.6}$$

Then the standard Calderón–Zygmund estimate gives

$$\|p^k\|_r \leq C\|u^{k-1}\|_{2r}\|u^k\|_{2r}$$

for  $1 < r < +\infty$ .

If  $(1 + |x|^2)^{1/2}u^{k-1} \in L_{loc}^\infty(0, \infty; L^2(\mathbb{R}^3))$ , we can show that  $(1 + |x|^2)^{1/2}u^k \in L_{loc}^\infty(0, \infty; L^2(\mathbb{R}^3))$ , by (3.12)–(3.15). By induction,

$$\int_{\mathbb{R}^3} (1 + |x|^2)|u^k|^2 dx \quad \text{and} \quad \int_0^t \int_{\mathbb{R}^3} (1 + |x|^2)|\nabla u^k|^2 dx d\tau$$

are well defined. Multiplying the first equation in (3.3) by  $(1 + |x|^2)u^k$  and integrating over  $\mathbb{R}^3$ , we obtain

$$\begin{aligned} & \frac{d}{dt} \|(1 + |x|^2)^{1/2}u^k\|_2^2 + \int_{\mathbb{R}^3} (1 + |x|^2)|\nabla u^k|^2 dx \\ & \leq C\|u^k\|_2^2 + C\|(1 + |x|^2)^{1/2}u^k\|_2\|u^{k-1}\|_4\|u^k\|_4 \\ & \leq 2\|u^k\|_2^2 + C\|(1 + |x|^2)^{1/2}u^k\|_2\|u^{k-1}\|_2^{1/4}\|u^k\|_2^{1/4}\|\nabla u^{k-1}\|_2^{3/4}\|\nabla u^k\|_2^{3/4} \\ & \leq \|(1 + |x|^2)^{1/2}u^k\|_2^2\|\nabla u^{k-1}\|_2\|\nabla u^k\|_2 + C\|u^k\|_2^2 \\ & \quad + C\|u^{k-1}\|_2\|u^k\|_2 + \|\nabla u^{k-1}\|_2\|\nabla u^k\|_2. \end{aligned} \tag{4.7}$$

Now (4.5) follows from (4.1), (4.2) and (4.7), by Gronwall’s inequality. □

The higher-order moments are estimated as follows.

LEMMA 4.4. Let  $a \in L^1(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3)$  and  $|x|^{3/2}a \in L^2(\mathbb{R}^3)$ . Then

$$\int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |u^k|^2 dx + \int_0^t \int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |\nabla u^k|^2 dx d\tau \leq CA_2 + B^{3/2}N^{4/3} \log(1 + t) \tag{4.8}$$

hold uniformly for  $k \geq 0$  and  $t \geq 0$  with  $A_2 = A_1^{3/2}(\|a\|_2^{5/4}N^{3/4} + \|a\|_2^{5/6}N)$ .

Moreover, if  $\|e^{-tA}a\|_1 \leq C(1 + t)^{-\gamma}$ , then

$$\int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |u^k|^2 dx + \int_0^t \int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |\nabla u^k|^2 dx d\tau \leq CA_2 + N^2(\frac{3}{2}) \tag{4.9}$$

hold uniformly for  $k \geq 0$  and  $t \geq 0$ .

*Proof.* Applying estimate (3.15) and the inequality

$$(1 + |x|^2)^{\alpha/2} \leq 2^{\alpha/2}((1 + |y|^2)^{\alpha/2} + |x - y|^\alpha) \quad \text{for } \alpha \geq 0, \tag{4.10}$$

we can show by using (3.12)–(3.14) and lengthy calculation that

$$(1 + |x|^2)^{3/4} u^k \in L^\infty_{\text{loc}}(0, +\infty; L^2(\mathbb{R}^3))$$

and

$$(1 + |x|^2)^{3/4} |\nabla u^k| \in L^2_{\text{loc}}(0, +\infty; L^2(\mathbb{R}^3))$$

as long as  $(1 + |x|^2)^{3/4} u^{k-1} \in L^\infty_{\text{loc}}(0, +\infty; L^2(\mathbb{R}^3))$ . By induction,

$$\int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |u^k|^2 dx \quad \text{and} \quad \int_0^t \int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |\nabla u^k|^2 dx d\tau$$

are well defined.

We now multiply both sides of (3.3) by  $(1 + |x|^2)^{3/2} u^k$  and integrate over  $\mathbb{R}^3$  to get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |u^k|^2 dx + \int_{\mathbb{R}^3} (1 + |x|^2)^{3/2} |\nabla u^k|^2 dx \\ & \leq 3 \int_{\mathbb{R}^3} (1 + |x|^2) |u^k| |\nabla u^k| dx + 3 \int_{\mathbb{R}^3} (1 + |x|^2) |u^{k-1}| |u^k|^2 dx \\ & \quad + 3 \int_{\mathbb{R}^3} (1 + |x|^2) |u^k| |p^k| dx. \end{aligned} \tag{4.11}$$

Employing the weighted estimates on singular integral operators (cf. [36]), we deduce from (4.6) that

$$\|(1 + |x|^2)^{1/2} p^k\|_2 \leq C \|(1 + |x|^2)^{1/2} |u^{k-1}| |u^k|\|_2.$$

Thus

$$\begin{aligned}
 \int_{\mathbb{R}^3} (1 + |x|^2)|p^k||u^k| \, dx &\leq \|(1 + |x|^2)^{1/2}u^k\|_2\|(1 + |x|^2)^{1/2}p^k\|_2 \\
 &\leq C\|(1 + |x|^2)^{1/2}u^k\|_2\|(1 + |x|^2)^{1/2}|u^{k-1}||u^k\|_2 \\
 &\leq C\|(1 + |x|^2)^{1/2}u^k\|_2\|(1 + |x|^2)^{3/4}u^k\|_6^{2/3}\|u^k\|_{24/7}^{1/3}\|u^{k-1}\|_{24/7} \\
 &\leq CA_1\|\nabla\{(1 + |x|^2)^{3/4}u^k\}\|_2^{2/3}\|u^k\|_{24/7}^{1/3}\|u^{k-1}\|_{24/7} \\
 &\leq \frac{1}{4}\|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2^2 + CA_1^{3/2}\|u^k\|_{24/7}^{1/2}\|u^{k-1}\|_{24/7}^{3/2} \\
 &\quad + CA_1\|(1 + |x|^2)^{1/4}u^k\|_2\|u^k\|_{24/7}^{1/3}\|u^{k-1}\|_{24/7} \\
 &\leq \frac{1}{4}\|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2^2 + CA_1^{3/2}\|u^k\|_2^{3/16}\|u^{k-1}\|_2^{9/16}\|\nabla u^k\|_2^{5/16}\|\nabla u^{k-1}\|_2^{15/16} \\
 &\quad + CA_1^{3/2}\|u^k\|_2^{5/8}\|u^{k-1}\|_2^{3/8}\|\nabla u^k\|_2^{5/24}\|\nabla u^{k-1}\|_2^{5/8},
 \end{aligned}$$

where we have used lemma 4.3 and the inequality,

$$\|u\|_{24/7} \leq \|u\|_2^{3/8}\|\nabla u\|_2^{5/8}.$$

By (4.1) and (4.2), we obtain

$$\begin{aligned}
 \int_0^t \int_{\mathbb{R}^3} (1 + |x|^2)|p^k||u^k| \, dx \, d\tau &\leq \frac{1}{4} \int_0^t \|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2^2 \, d\tau + CA_1^{3/2}(N^{3/4}\|a\|_2^{5/4} + N\|a\|_2^{5/6}). \quad (4.12)
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \int_0^t \int_{\mathbb{R}^3} (1 + |x|^2)|u^{k-1}||u^k|^2 \, dx \, d\tau &\leq \frac{1}{4} \int_0^t \|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2^2 \, d\tau + CA_1^{3/2}(N^{3/4}\|a\|_2^{5/4} + N\|a\|_2^{5/6}). \quad (4.13)
 \end{aligned}$$

Finally, we estimate the first term on the right-hand side of (4.11). By Hölder’s inequality we get

$$\begin{aligned}
 \int_{\mathbb{R}^3} (1 + |x|^2)|u^k||\nabla u^k| \, dx &= \int_{\mathbb{R}^3} (1 + |x|^2)^{3/4}|\nabla u^k|(1 + |x|^2)^{1/4}|u^k| \, dx \\
 &\leq \|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2\|(1 + |x|^2)^{1/4}u^k\|_2 \\
 &\leq \frac{1}{8}\|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2^2 + C\|(1 + |x|^2)^{1/4}u^k\|_2^2.
 \end{aligned}$$

Since

$$\begin{aligned}
 \|(1 + |x|^2)^{1/4}u^k\|_2^2 &= \int_{\mathbb{R}^3} (1 + |x|^2)^{1/2}|u^k|^{2/3}|u^k|^{4/3} \, dx \\
 &\leq \|(1 + |x|^2)^{3/4}u^k\|_6^{2/3}\|u^k\|_{3/2}^{4/3} \\
 &\leq C\|\nabla\{(1 + |x|^2)^{3/4}u^k\}\|_2^{2/3}\|u^k\|_{3/2}^{4/3} \\
 &\leq \varepsilon\|(1 + |x|^2)^{3/4}|\nabla u^k|\|_2^2 + \frac{1}{2}\|(1 + |x|^2)^{1/4}u^k\|_2^2 + C\|u^k\|_{3/2}^2,
 \end{aligned}$$

for some  $\varepsilon > 0$ . Thus,

$$\|(1 + |x|^2)^{1/4} u^k\|_2^2 \leq 2\varepsilon \|(1 + |x|^2)^{3/4} |\nabla u^k|\|_2^2 + C \|u^k\|_1^{2/3} \|u^k\|_2^{4/3}.$$

Taking  $2\varepsilon = 1/8$  yields

$$\int_{\mathbb{R}^3} (1 + |x|^2) |u^k| |\nabla u^k| \, dx \leq \frac{1}{4} \|(1 + |x|^2)^{3/4} |\nabla u^k|\|_2^2 + CB^{2/3} \|u^k\|_2^{4/3}. \tag{4.14}$$

Substituting (4.1)–(4.14) into (4.11), we obtain estimates (4.8) and (4.9), by lemma 4.2. □

### 5. Weighted estimates for approximate solutions II

In this section, we establish the decay rates estimates of weighted norms for approximate solutions in  $L^p(\mathbb{R}^3)$  ( $p > 3$ ), which are used to show the existence of corresponding strong solutions. To this end, we first recall some basic estimates on  $\Gamma$  and  $\bar{\theta}$ . Applying the inequality  $\tau^\alpha e^{-C\tau} \leq C^\alpha e^{-1}$  for  $\alpha > 0$ , we can verify directly that

$$\left. \begin{aligned} \||x|^\alpha \Gamma\|_p &\leq C t^{\alpha/2 - (3-3/p)/2}, \\ \||x|^\alpha \nabla \Gamma\|_p &\leq C t^{(\alpha-1)/2 - (3-3/p)/2}, \end{aligned} \right\} \tag{5.1}$$

for  $1 \leq p \leq +\infty$  and  $\alpha \geq 0$ . By the weighted estimates about the singular integrals (cf. [36–38]) and (3.8), we have

$$\left. \begin{aligned} \||x|^\alpha D^2 \bar{\theta}\|_p &\leq C \||x|^\alpha \Gamma\|_p \leq C t^{\alpha/2 - (3/2)(1-1/p)}, \\ \||x|^\alpha D^3 \bar{\theta}\|_p &\leq C \||x|^\alpha |\nabla \Gamma|\|_p \leq C t^{(\alpha-1)/2 - (3/2)(1-1/p)}, \end{aligned} \right\} \tag{5.2}$$

for  $1 < p < +\infty$  and  $-1/p < \alpha < 3 - 3/p$ .

Now we can deduce the main estimates needed for theorems 2.5 and 2.6.

LEMMA 5.1. *Let*

$$a \in L^1(\mathbb{R}^3) \cap \dot{J}^2(\mathbb{R}^3), \quad (1 + |x|^2)^{1/2} a \in L^2(\mathbb{R}^3) \quad \text{and} \quad (1 + |x|^2)^{\alpha/2} a \in L^{p_0}(\mathbb{R}^3)$$

for  $1 \leq p_0 \leq +\infty$  and  $\alpha = 3 - 3/p_0$ . Then there exists a constant  $\lambda > 0$ , such that if  $N \leq \lambda$ , then, for  $\beta = (3/p_0 - 3/p)/2$ , the estimate

$$\|t^\beta (1 + |x|^2)^{\alpha/2} u^k\|_p \leq C (\|(1 + |x|^2)^{\alpha/2} a\|_{p_0} + N^2 + A_1^{5/2} + B^{1/2} A_1 N) \tag{5.3}$$

holds uniformly for  $k \geq 0$  and for  $3 < p \leq +\infty$ .

*Proof.* By (3.14) and Minkowski inequality, we obtain

$$\|(1 + |x|^2)^{\alpha/2} u^k\|_p \leq C \sum_{i=1}^3 \|(1 + |x|^2)^{\alpha/2} I_i^k\|_p. \tag{5.4}$$

By the Minkowski inequality, (4.10) and the basic  $L^p$ -estimates for convolutions, we have, for  $3 < p \leq +\infty$ ,

$$\begin{aligned} \|(1 + |x|^2)^{\alpha/2} I_1^k\|_p &\leq C \left\| \int_{\mathbb{R}^3} (1 + |y|^2)^{\alpha/2} |a(y)| \Gamma(x - y, t) \, dy \right\|_p \\ &\quad + C \left\| \int_{\mathbb{R}^3} |a(y)| |x - y|^\alpha \Gamma(x - y, t) \, dy \right\|_p \\ &\leq C \|(1 + |x|^2)^{\alpha/2} a\|_{p_0} t^{-(3/2)(1/p_0 - 1/p)} + C \|a\|_1 t^{-3/2 + 3/(2p) + \alpha/2} \\ &\leq C(\|a\|_1 + \|(1 + |x|^2)^{\alpha/2} a\|_{p_0}) t^{-(3/2)(1/p_0 - 1/p)}, \end{aligned} \tag{5.5}$$

where we have used the estimates (5.1) (if  $p = \infty$ , we use (3.15) to estimate the second term of the first step in (5.5)).

By (4.10) and the Minkowski inequality, we can get, for  $3 < p \leq +\infty$ ,

$$\begin{aligned} \|(1 + |x|^2)^{\alpha/2} I_2^k\|_p &\leq C \left\| \int_0^{t/2} \int_{\mathbb{R}^3} (1 + |y|^2)^{\alpha/2} |u^{k-1}| |u^k| (|\nabla \Gamma| + |D^3 \bar{\theta}|)(x - y, t - \tau) \, dy \, d\tau \right\|_p \\ &\quad + C \left\| \int_0^{t/2} \int_{\mathbb{R}^3} |u^{k-1}| |u^k| |x - y|^\alpha (|\nabla \Gamma| + |D^3 \bar{\theta}|)(x - y, t - \tau) \, dy \, d\tau \right\|_p \\ &\triangleq C(\|I_{21}^k\|_p + \|I_{22}^k\|_p). \end{aligned} \tag{5.6}$$

Similarly,

$$\begin{aligned} \|(1 + |x|^2)^{\alpha/2} I_3^k\|_p &\leq C \left\| \int_{t/2}^t \int_{\mathbb{R}^3} (1 + |y|^2)^{\alpha/2} |u^{k-1}| |u^k| (|\nabla \Gamma| + |D^3 \bar{\theta}|)(x - y, t - \tau) \, dy \, d\tau \right\|_p \\ &\quad + C \left\| \int_{t/2}^t \int_{\mathbb{R}^3} |u^{k-1}| |u^k| |x - y|^\alpha (|\nabla \Gamma| + |D^3 \bar{\theta}|)(x - y, t - \tau) \, dy \, d\tau \right\|_p \\ &\triangleq C(\|I_{31}^k\|_p + \|I_{32}^k\|_p). \end{aligned} \tag{5.7}$$

Let  $J_p^k \triangleq \|(1 + |x|^2)^{\alpha/2} u^k\|_p$ . In order to estimate  $I_2^k$  and  $I_3^k$ , we discuss three separated cases: (i)  $p = +\infty$  and  $1 \leq p_0 \leq +\infty$ ; (ii)  $1 \leq p_0 < p < \infty$  and  $p > 3$ ; (iii)  $3 < p_0 = p < \infty$ .

CASE I.  $p = +\infty$  and  $1 \leq p_0 \leq +\infty$ .

In order to establish the uniform estimates on  $t^\beta \|(1 + |x|^2)^{\alpha/2} u^k\|_\infty$  with  $\beta = 3/(2p_0)$ , the singular factor  $t^{-\beta}$  will later appear in the integral. So it seems necessary to treat two cases:  $p_0 > 3/2$  and  $1 \leq p_0 \leq 3/2$ . Similarly, to establish the uniform estimates on  $t^{1/2+\beta} \|(1 + |x|^2)^{\alpha/2} \nabla u^k\|_\infty$ , the singular factor  $t^{-1/2-\beta}$  will appear in this procedure, we need to distinguish three cases:  $p_0 > 3$ ,  $3/2 < p_0 \leq 3$  and  $1 \leq p_0 \leq 3$ .

First, it follows from (5.1), (5.2) and lemma 4.2, that

$$\begin{aligned} \|I_{21}^k\|_\infty &\leq C \int_0^{t/2} \|u^{k-1}\|_2 J_\infty^k \|\nabla \Gamma + |D^3 \bar{\theta}|\|_2 \, d\tau \\ &\leq C \int_0^{t/2} \|u^k\|_2 J_\infty^{k-1} (t - \tau)^{-5/4} \, d\tau \\ &\leq CN \int_0^{t/2} J_\infty^{k-1} (1 + \tau)^{-3/4} (t - \tau)^{-5/4} \, d\tau \end{aligned} \tag{5.8}$$

which yields the desired estimate for  $p_0 > 3$ . If  $1 \leq p_0 \leq 3/2$ , then  $\alpha \leq 1$  and  $1 \leq \beta \leq 3/2$ . By (3.15), we get, with the help of lemmas 4.2 and 4.3, that

$$\begin{aligned} \|I_{21}^k\|_\infty &\leq C \int_0^{t/2} \|u^k\|_{3/2} \|(1 + |y|^2)^\alpha u^{k-1}\|_2^{2/3} (J_\infty^{k-1})^{1/3} (t - \tau)^{-2} \, d\tau \\ &\leq CA_1^{2/3} \int_0^{t/2} \|u^k\|_1^{1/3} \|u^k\|_2^{2/3} (J_\infty^{k-1})^{1/3} (t - \tau)^{-2} \, d\tau \\ &\leq CA_1^{2/3} B^{1/3} N^{2/3} \int_0^{t/2} (J_\infty^{k-1})^{1/3} (1 + \tau)^{-1/2} (t - \tau)^{-2} \, d\tau. \end{aligned} \tag{5.9}$$

If  $3/2 < p_0 \leq 3$ , then  $\alpha - 1 \in (0, 1]$  and  $1/2 \leq \beta \leq 1$ . By (3.15), we obtain, with the aid of lemmas 4.2 and 4.3, that

$$\|I_{21}^k\|_\infty \leq CA_1^{5/3} B \int_0^{t/2} (J_\infty^{k-1})^{1/3} (t - \tau)^{-2} \, d\tau, \tag{5.10}$$

where we have used the inequality

$$\|(1 + |x|^2)^{\gamma_0/2} u\|_p \leq \|u\|_q \|(1 + |x|^2)^{1/2} u\|_2 \tag{5.11}$$

for  $1 \leq q < p < 2$  and  $\gamma_0 = 2(p - q)/(p(2 - q))$ , which follows from the Hölder inequality.

The  $\|I_{22}^k\|_\infty$  can be estimated as

$$\begin{aligned} \|I_{22}^k\|_\infty &\leq C \left\| \int_0^{t/2} |u^{k-1}| |u^k| |x - y|^\alpha (|x - y|^2 + (t - \tau))^{-2} \, dy \, d\tau \right\|_\infty \\ &\leq C \int_0^{t/2} \|u^{k-1}\|_2 \|u^k\|_2 (t - s)^{-2+\alpha/2} \, d\tau \\ &\leq CN^2 \int_0^{t/2} (1 + \tau)^{-3/2} (t - \tau)^{-1/2-3/(2p_0)} \, d\tau \\ &\leq CN^2 t^{-1-3/(2p_0)}, \end{aligned} \tag{5.12}$$

where one has used (3.15).

Next, we estimate  $I_3^k$ . By (5.1) and (5.2), we have

$$\begin{aligned} \|I_{31}^k\|_\infty &\leq C \int_{t/2}^t \|(1 + |x|^2)^{\alpha/2} |u^{k-1}| |u^k|\|_4 (t - \tau)^{-7/8} \, d\tau \\ &\leq C \int_{t/2}^t J_\infty^{k-1} (J_\infty^k)^{1/2} \|u^k\|_2^{1/2} (t - \tau)^{-7/8} \, d\tau \\ &\leq CN^{1/2} \int_{t/2}^t J_\infty^{k-1} (J_\infty^k)^{1/2} (1 + \tau)^{-3/8} (t - \tau)^{-7/8} \, d\tau. \end{aligned} \tag{5.13}$$

Note that in the estimate of  $I_{32}^k$ , the singular factor  $(t - \tau)^{-2+\alpha/2}$  will appear in the integral with  $\alpha = 3 - 3/p_0$ . In order to control this singularity, we will separate two cases:  $p_0 > 3$  and  $1 \leq p_0 \leq 3$ .

If  $p_0 > 3$ , it follows from the estimate (3.15) that

$$\|I_{32}^k\|_\infty \leq C \int_{t/2}^t \|u^{k-1}\|_2 \|u^k\|_2 (t - \tau)^{-1/2-3/(2p_0)} \, d\tau \leq CN^2 t^{-1-\beta}. \tag{5.14}$$

If  $1 \leq p_0 \leq 3$ , using (5.1) and (5.2), we can get

$$\|I_{32}^k\|_\infty \leq C \int_{t/2}^t \| |u^{k-1}| |u^k| \|_{(3p_0+1)/(3p_0-2)} (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} \, d\tau. \tag{5.15}$$

If  $5/3 \leq p_0 \leq 3$ , then  $(3p_0 + 1)/(3p_0 - 2) \leq 2$ . By lemma 4.2, we get

$$\begin{aligned} \| |u^{k-1}| |u^k| \|_{(3p_0+1)/(3p_0-2)} &\leq J_\infty^{k-1} \|u^k\|_1^{(3p_0-5)/(3p_0+1)} \|u^k\|_2^{6/(3p_0+1)} \\ &\leq J_\infty^{k-1} B^{(3p_0-5)/(3p_0+1)} N^{6/(3p_0+1)} (1 + \tau)^{-9/(6p_0+2)}. \end{aligned} \tag{5.16}$$

If  $1 \leq p_0 < 5/3$ , then  $(3p_0 + 1)/(3p_0 - 2) > 2$ . Then,

$$\begin{aligned} \| |u^{k-1}| |u^k| \|_{(3p_0+1)/(3p_0-2)} &\leq J_\infty^{k-1} (J_\infty^k)^{(5-3p_0)/(3p_0+1)} \|u^k\|_2^{2(3p_0-2)/(3p_0+1)} \\ &\leq J_\infty^{k-1} (J_\infty^k)^{(5-3p_0)/(3p_0+1)} N^{2(3p_0-2)/(3p_0+1)} (1 + \tau)^{-3(3p_0-2)/(2(3p_0+1))}. \end{aligned} \tag{5.17}$$

CASE II.  $1 \leq p_0 < p < +\infty$  and  $3 < p$ .

This can be achieved in a similar way as case I with slight modification. Indeed, we deal with three cases for  $p_0$ . For  $p_0 > 3$ , (5.1), (5.2) and lemma 4.2 then imply that

$$\begin{aligned} \|I_{21}^k\|_p &\leq C \int_0^{t/2} \|u^k\|_2 J_p^{k-1} \| |\nabla \Gamma| + |D^3 \bar{\theta}| \|_2 \, d\tau \\ &\leq C \int_0^{t/2} \|u^k\|_2 J_p^{k-1} (t - \tau)^{-5/4} \, d\tau \\ &\leq CN \int_0^{t/2} J_p^{k-1} (1 + \tau)^{-3/4} (t - \tau)^{-5/4} \, d\tau. \end{aligned} \tag{5.18}$$

If  $1 \leq p_0 \leq 3/2$ , then  $\alpha \leq 1$ . It now follows from (5.1), (5.2) and lemmas 4.2 and 4.3 that

$$\begin{aligned} \|I_{21}^k\|_p &\leq C \int_0^{t/2} \|u^k\|_{4p/(3p-2)} \|(1 + |y|^2)^{\alpha/2} u^{k-1}\|_2^{1/2} (J_p^{k-1})^{1/2} (t - \tau)^{-2+3/(2p)} \, d\tau \\ &\leq CA_1^{1/2} \int_0^{t/2} \|u^k\|_1^{(p-2)/(2p)} \|u^k\|_2^{(p+2)/(2p)} (J_p^{k-1})^{1/2} (t - \tau)^{-2+3/(2p)} \, d\tau \\ &\leq CA_1^{1/2} B^{(p-2)/(2p)} N^{(p+2)/(2p)} \\ &\quad \times \int_0^{t/2} (J_p^{k-1})^{1/2} (1 + \tau)^{-3/8-3/(4p)} (t - \tau)^{-2+3/(2p)} \, d\tau \end{aligned} \tag{5.19}$$

If  $3/2 < p_0 \leq 3$ , then  $\alpha - 1 \in (0, 1]$ . By (5.1) and (5.2), we obtain, with the help of lemmas 4.2 and 4.3 and inequality (5.11), that

$$\begin{aligned} \|I_{21}^k\|_p &\leq C \int_0^{t/2} \|(1 + |y|^2)^{1/2} u^{k-1}\|_2^{1/2} \|(1 + |y|^2)^{(\alpha-1)/4} u^k\|_{4p/(3p-2)} \\ &\quad \times (J_p^{k-1})^{1/2} (t - \tau)^{-2+3/(2p)} \, d\tau \\ &\leq CA_1^{1/2} \int_0^{t/2} \|u^k\|_1 \|(1 + |x|^2)^{1/2} u^k\|_2 (J_p^{k-1})^{1/2} (t - \tau)^{-2+3/(2p)} \, d\tau \\ &\leq CA_1^{3/2} B \int_0^{t/2} (J_p^{k-1})^{1/2} (t - \tau)^{-2+3/(2p)} \, d\tau. \end{aligned} \tag{5.20}$$

Applying (5.1) and (5.1), we can estimate the  $\|I_{22}^k\|_p$  as

$$\|I_{22}^k\|_p \leq C \int_0^{t/2} \|u^{k-1}\|_2 \|u^k\|_2 (t - s)^{-2+\alpha/2+3/(2p)} \, d\tau \leq CN^2 t^{-1-\beta}. \tag{5.21}$$

Now we estimate  $I_3^k$ . Let  $r = 6(p - 1)/(p + 1)$ . By (5.1) and (5.2), we have

$$\begin{aligned} \|I_{31}^k\|_p &\leq C \int_{t/2}^t \|(1 + |x|^2)^{\alpha/2} |u^{k-1}| |u^k|\|_{pr/(p+r)} (t - \tau)^{-1/2-3/(2r)} \, d\tau \\ &\leq C \int_{t/2}^t J_p^{k-1} (J_p^k)^{p(r-2)/(r(p-2))} \|u^k\|_2^{2(p-r)/(r(p-2))} (t - \tau)^{-1/2-3/(2r)} \, d\tau \\ &\leq CN^{(p-3)/(3(p-1))} \int_{t/2}^t J_p^{k-1} (J_p^k)^{2p/(3(p-1))} \\ &\quad \times (1 + \tau)^{-(p-3)/(4(p-1))} (t - \tau)^{-1/2-3/(2r)} \, d\tau. \end{aligned} \tag{5.22}$$

For the estimate on  $I_{32}^k$ , We also consider two cases. If  $p_0 \geq 3$ , we have, by (5.1) and (5.2),

$$\|I_{32}^k\|_p \leq C \int_{t/2}^t \|u^{k-1}\|_2 \|u^k\|_2 (t - \tau)^{-1/2-\beta} \, d\tau \leq CN^2 t^{-1-\beta}. \tag{5.23}$$

By Young's inequality and estimates (5.1) and (5.2), we get

$$\|I_{32}^k\|_p \leq C \int_{t/2}^t \| |u^{k-1}| |u^k| \|_l (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau$$

for  $1/l = 1/p + (3p_0 - 2)/(3p_0 + 1)$ . If  $5/3 \leq p_0 \leq 3$ , by lemma 4.2,

$$\begin{aligned} \|I_{32}^k\|_p &\leq C \int_{t/2}^t J_p^{k-1} \|u^k\|_{(3p_0+1)/(3p_0-2)} (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau \\ &\leq C \int_{t/2}^t J_p^{k-1} \|u^k\|_1^{(3p_0-5)/(3p_0+1)} \|u^k\|_2^{6/(3p_0+1)} (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau \\ &\leq CB^{(3p_0-5)/(3p_0+1)} N^{6/(3p_0+1)} \int_{t/2}^t J_p^{k-1} (1 + \tau)^{-9/(2(3p_0+1))} \\ &\quad \times (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau. \end{aligned} \tag{5.24}$$

If  $1 \leq p_0 < 5/3$ , then

$$\begin{aligned} \|I_{32}^k\|_p &\leq C \int_{t/2}^t J_p^{k-1} \|u^k\|_{(3p_0+1)/(3p_0-2)} (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau \\ &\leq C \int_{t/2}^t J_p^{k-1} (J_p^k)^{(5-3p_0)/(3p_0+1)} \|u^k\|_2^{2(3p_0-2)/(3p_0+1)} \\ &\quad \times (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau \\ &\leq CN^{2(3p_0-2)/(3p_0+1)} \int_{t/2}^t J_p^{k-1} (J_p^k)^{(5-3p_0)/(3p_0+1)} \\ &\quad \times (1 + \tau)^{-3(3p_0-2)/(2(3p_0+1))} (t - \tau)^{-1/2-3/(2p_0(3p_0+1))} d\tau. \end{aligned} \tag{5.25}$$

CASE III.  $3 < p_0 = p < +\infty$ .

The estimates of  $\|I_{21}^k\|_p$  and  $\|I_{31}^k\|_p$  in this case are same as (5.18) and (5.22). So we only give the estimate of  $\|I_{22}^k\|_p$  and  $\|I_{32}^k$ . Applying (3.15) and the theory of singular integral operator (cf. [37]), we get

$$\begin{aligned} \|I_{22}^k\|_p + \|I_{32}^k\|_p &\leq C \left\| \int_0^t |u^{k-1}| |u^k| |x - y|^\alpha (|x - y|^2 + (t - \tau))^{-2} dy d\tau \right\|_p \\ &\leq C \left\| \int_0^t |u^{k-1}| |u^k| |x - y|^{-1-3/p} dy d\tau \right\|_p \\ &\leq C \int_0^t \| |u^{k-1}| |u^k| \|_{3/2} d\tau \\ &\leq C \|a\|_2 N t^{-1/4}. \end{aligned} \tag{5.26}$$

Summarizing the above estimates, it is obvious that, if  $t^\beta J_p^{k-1} \leq C$ , then  $t^\beta J_p^k \leq C$ , by the Gronwall inequality. Therefore, it holds that  $t^\beta J_p^k \in L_{loc}^\infty(0, +\infty)$ , by

induction. Thus

$$\begin{aligned}
 t^\beta J_\infty^k &\leq C \left( \|a\|_1 + \|(1 + |x|^2)^{\alpha/2} a\|_{p_0} + N^2(1 + t)^{-1} \right. \\
 &\quad \left. + N^{1/2}(1 + t)^{-\beta/2-1/4} \max_{0 \leq t \leq \infty} (t^\beta J_\infty^{k-1})(t^\beta J_\infty^k)^{1/2} \right) \\
 &\quad \left\{ \begin{aligned}
 &C \left( N(1 + t)^{-1} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1}) + N^2(1 + t)^{-1} \right), \\
 &\hspace{10em} \text{as } p_0 > 3; \\
 &C \left( A_1^{5/3} B(1 + t)^{-1+2\beta/3} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{1/3} \right. \\
 &\quad \left. + B^{(3p_0-5)/(3p_0+1)} N^{6/(3p_0+1)} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1}) \right), \\
 &\hspace{10em} \text{as } \frac{5}{3} \leq p_0 \leq 3; \\
 &C \left( A_1^{5/3} B(1 + t)^{-1+2\beta/3} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{1/3} \right. \\
 &\quad \left. + N^{2(3p_0-2)/(3p_0+1)} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})(t^\beta J_\infty^k)^{(5-3p_0)/(3p_0+1)} \right), \\
 &\hspace{10em} \text{as } \frac{3}{2} < p_0 < \frac{5}{3}; \\
 &C \left( B^{1/3} A_1^{2/3} N^{2/3} (1 + t)^{-3/2+2\beta/3} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{1/3} \right. \\
 &\quad \left. + N^{2(3p_0-2)/(3p_0+1)} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})(t^\beta J_\infty^k)^{(5-3p_0)/(3p_0+1)} \right), \\
 &\hspace{10em} \text{as } 1 \leq p_0 \leq \frac{3}{2}.
 \end{aligned} \right.
 \end{aligned}$$

By Young’s inequality, we deduce that

$$\begin{aligned}
 \max_{t \in [0, \infty]} (t^\beta J_\infty^k) &\leq C \left( \|a\|_1 + \|(1 + |x|^2)^{\alpha/2} a\|_{p_0} + N^2 + N \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^2 \right) \\
 &\quad \left\{ \begin{aligned}
 &C \left( N \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1}) + N^2 \right), \\
 &\hspace{10em} \text{as } p_0 > 3; \\
 &C \left( A_1^{5/3} B \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{1/3} \right. \\
 &\quad \left. + B^{(3p_0-5)/(3p_0+1)} N^{6/(3p_0+1)} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1}) \right), \\
 &\hspace{10em} \text{as } \frac{5}{3} \leq p_0 \leq 3; \\
 &C \left( A_1^{5/3} B \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{1/3} \right. \\
 &\quad \left. + N \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{(3p_0+1)/(2(3p_0-2))} \right), \\
 &\hspace{10em} \text{as } \frac{3}{2} < p_0 < \frac{5}{3}; \\
 &C \left( B^{1/3} A_1^{2/3} N^{2/3} \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{1/3} \right. \\
 &\quad \left. + N \max_{t \in [0, \infty]} (t^\beta J_\infty^{k-1})^{(3p_0+1)/(2(3p_0-2))} \right), \\
 &\hspace{10em} \text{as } 1 \leq p_0 \leq \frac{3}{2}.
 \end{aligned} \right.
 \end{aligned}$$

Therefore, if  $N$  is suitably small, then

$$\max(t^\beta J_\infty^k) \leq C(\|a\|_1 + \|(1 + |x|^2)^{\alpha/2} a\|_{p_0} + N^2 + A_1^{5/2} B^{3/2} + B^{1/2} A_1 N),$$

which yields the desired estimate for the case  $p = \infty$ . The argument for  $3 < p < \infty$  is similar. □

By a lengthy but similar calculation, we can show the following lemma.

**LEMMA 5.2.** *Assume the conditions in lemma 5.1 are satisfied. Then if  $N \leq \lambda$ , the estimates, for  $t \geq 0$ ,*

$$t^{1/2+\beta} \|\nabla u^k\|_p \leq C(\|a\|_1 + \|(1 + |x|^2)^{\alpha/2} a\|_{p_0} + N^2 + A_1^{5/2} B^{3/2} + B^{1/2} A_1 N)$$

hold uniformly for  $k \geq 0$  and  $3 < p \leq +\infty$ .

If  $a = (\partial b)/(\partial x_i)$  for some  $i = 1, 2, 3$ , we can show, by similar discussion, the following.

**LEMMA 5.3.** *Assume the conditions in lemma 5.1 hold. If  $a = (\partial b)/(\partial x_i)$  for some  $i = 1, 2, 3$  with  $b \in L^1(\mathbb{R}^3)$  and  $(1 + |x|^2)^{\alpha/2} \in L^{p_0}(\mathbb{R}^3)$ , then there exists a constant  $\lambda_0 > 0$  such that if  $N \leq \lambda_0$ , the estimate*

$$t^{1/2+\beta} \|u^k\|_p \leq C(\|b\|_1 + \|(1 + |x|^2)^{\alpha/2} b\|_{p_0} + N^2 + A_1^{5/2} B^{3/2} + B^{1/2} A_1 N),$$

$t \in [0, \infty)$

holds uniformly for  $k \geq 0$  and  $3 < p \leq +\infty$ .

### Acknowledgments

This work was done when C.H. was visiting the Institute of Mathematical Sciences at the Chinese University of Hong Kong. The authors express their sincere thanks to Professor Zhimin Chen for helpful discussions. The research of C.H. was supported in part by the National Youth Foundation of China, and Zheng Ge Ru Foundation. The research of Z.X. was supported in part by the Zheng Ge Ru Foundation, NSF grant DMS-96-00137, and DOE grant De-FG02-88ER-25053.

### References

- 1 H. Beirao da Veiga. Existence and asymptotic behavior for strong solutions of the Navier–Stokes equations in the whole space. *Indiana Univ. Math. J.* **30** (1987), 149–166.
- 2 W. Borchers and T. Miyakawa.  $L^2$  decay for the Navier–Stokes flows in halfspace. *Math. Ann.* **282** (1988), 141–155.
- 3 W. Borchers and T. Miyakawa. Algebraic  $L^2$  decay for Navier–Stokes flows in exterior domains. *Acta Math.* **165** (1990), 189–227.
- 4 W. Borchers and T. Miyakawa.  $L^2$  decay for Navier–Stokes flows in unbounded domains, with application to exterior stationary flows. *Arch. Ration. Mech. Analysis* **118** (1992), 273–295.
- 5 C. P. Calderón. Existence of weak solutions for the Navier–Stokes equations with initial data in  $L^p$ . *Trans. Am. Math. Soc.* **318** (1990), 179–200.
- 6 C. P. Calderón. Addendum to the paper ‘Existence of weak solutions for the Navier–Stokes equations with initial data in  $L^p$ ’. *Trans. Am. Math. Soc.* **318** (1990), 201–207.

- 7 A. Carpio. Large-time behavior in incompressible Navier–Stokes equations. *SIAM J. Math. Analysis* **27** (1996), 449–475.
- 8 Z.-m. Chen. Solutions of the stationary and nonstationary Navier–Stokes equations in exterior domains. *Pac. J. Math.* **159** (1993), 227–242.
- 9 E. B. Fabes, B. F. Jones and N. M. Riviere. The initial value problem for the Navier–Stokes equations with data in  $L^p$ . *Arch. Ration. Mech. Analysis* **45** (1972), 222–242.
- 10 R. Farwig and H. Sohr. Weighted energy inequalities for the Navier–Stokes equations in exterior domains. *Appl. Analysis* **58**(1–2) (1995), 157–173.
- 11 R. Farwig and H. Sohr. Global estimates in weighted spaces of weak solutions of the Navier–Stokes equations in exterior domains. *Arch. Math.* **67** (1996), 319–330.
- 12 Y. Giga. Solutions for semilinear parabolic equations in  $L^p$  and the regularity of weak solutions of the Navier–Stokes equations. *J. Diff. Eqns* **62** (1986), 182–212.
- 13 Y. Giga and T. Miyakawa. Solutions in  $L^r$  of the Navier–Stokes initial value problem. *Arch. Ration. Mech. Analysis* **89** (1985), 267–281.
- 14 C. He. The Cauchy problem for the Navier–Stokes equations. *J. Math. Analysis Appl.* **209** (1997), 228–242.
- 15 C. He. Weighted estimates for nonstationary Navier–Stokes equations. *J. Diff. Eqns* **148** (1998), 422–444.
- 16 C. He and Z. Xin. Weighted estimates for nonstationary Navier–Stokes equations in exterior domains. *Meth. Applic. Analysis* **7**(3) (2000). (Special issue dedicated to Professor Morawetz.)
- 17 J. G. Heywood. The Navier–Stokes equations: on the existence, regularity and decay of solutions. *Indiana Univ. Math. J.* **29** (1980), 641–681.
- 18 H. Iwashita.  $L^p$ – $L^q$  estimates for solutions of the nonstationary Stokes equations in an exterior domain and the Navier–Stokes initial value problem in  $L^q$  space. *Math. Ann.* **285** (1989), 265–288.
- 19 R. Kajikiya and T. Miyakawa. On  $L^2$  decay of weak solutions of the Navier–Stokes equations in  $R^n$ . *Math. Z.* **192** (1986), 136–148.
- 20 T. Kato. Strong  $L^p$ -solutions of the Navier–Stokes equations in  $R^n$ , with application to weak solutions. *Math. Z.* **187** (1984), 471–480.
- 21 T. Kato. *Liapunov functions and monotonicity in the Navier–Stokes equations*, Lecture Notes in Mathematics, vol. 1450, pp. 53–64 (Springer, 1989).
- 22 H. Kozono and T. Ogawa. Some  $L^p$  estimate for the exterior Stokes flow and an application to the non-stationary Navier–Stokes equations. *Indiana Univ. Math. J.* **41** (1992), 789–808.
- 23 H. Kozono, T. Ogawa and H. Sohr. Asymptotic behavior in  $L^r$  for weak solutions of the Navier–Stokes equations in exterior domains. *Manuscr. Math.* **74** (1992), 253–275.
- 24 O. A. Ladyzhenskaya. *The mathematical theory of viscous incompressible flow* (Gordon and Breach, 1969).
- 25 J. Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.* **63** (1934), 193–248.
- 26 P. Maremonti. On the asymptotic behaviour of the  $L^2$ -norm of suitable weak solutions to the Navier–Stokes equations in three-dimensional exterior domains. *Commun. Math. Phys.* **118** (1988), 405–420.
- 27 K. Masuda.  $L^2$ -decay of solutions of the Navier–Stokes equations in the exterior domains. In *Proc. Symposia in Pure Mathematics*, vol. 45, part 2, pp. 179–182 (American Mathematical Society, 1986).
- 28 T. Miyakawa. On nonstationary solutions of the Navier–Stokes equations in an exterior domains. *Hiroshima Math. J.* **12** (1982), 115–142.
- 29 T. Miyakawa. Hardy spaces of solenoidal vector fields, with applications to the Navier–Stokes equations. *Kyushu J. Math.* **50** (1996), 1–64.
- 30 T. Miyakawa. Application of Hardy space techniques to the time-decay problem for incompressible Navier–Stokes flows in  $R^n$ . *Funkcialaj Ekvacioj* **41** (1998), 383–434.
- 31 T. Miyakawa and H. Sohr. On energy inequality, smoothness and large time behavior in  $L^2$  for weak solutions of the Navier–Stokes equations in exterior domains. *Math. Z.* **199** (1988), 455–478.
- 32 M. E. Schonbek.  $L^2$  decay for weak solutions of the Navier–Stokes equations. *Arch. Ration. Mech. Analysis* **88** (1985), 209–222.

- 33 M. E. Schonbek. Large time behavior of solutions to the Navier–Stokes equations. *Commun. Partial Diff. Eqns* **11** (1986), 733–763.
- 34 M. E. Schonbek. Lower bounds of rates of decay for solutions to the Navier–Stokes equations. *J. Am. Math. Soc.* **4** (1991), 423–449.
- 35 M. E. Schonbek and T. Schonbek. On the boundedness and decay of the moments of solutions of the Navier–Stokes equations. *Adv. Diff. Eqns* **5** (2000), 861–898.
- 36 E. M. Stein. Note on singular integrals. *Proc. Am. Math. Soc.* **8** (1957), 250–254.
- 37 E. M. Stein. *Singular integrals and differentiability properties of functions* (Princeton University Press, 1970).
- 38 E. M. Stein and G. Weiss. Fractional integral on  $n$ -dimensional Euclidean space. *J. Math. Mech.* **17** (1958), 503–514.
- 39 S. Takahashi. A weighted equations approach to decay rate estimates for the Navier–Stokes equations. *Nonlinear Analysis* **37** (1999), 751–789.
- 40 S. Ukai. A solution formula for the Stokes equations in  $R_+^n$ . *Commun. Pure Appl. Math.* **40** (1987), 611–621.
- 41 M. Weigner. Decay results for weak solutions of the Navier–Stokes equations in  $R^n$ . *J. Lond. Math. Soc.* **36** (1987), 303–313.
- 42 F. B. Weissler. Initial value problem in  $L^p$  for the Navier–Stokes fluids. *Arch. Ration. Mech. Analysis* **74** (1980), 219–229.

(Issued 15 June 2001)