



L^p -regularity of the Bergman projection on quotient domains

Chase Bender, Debraj Chakrabarti, Luke Edholm, and Meera Mainkar

Abstract. We obtain sharp ranges of L^p -boundedness for domains in a wide class of Reinhardt domains representable as sublevel sets of monomials, by expressing them as quotients of simpler domains. We prove a general transformation law relating L^p -boundedness on a domain and its quotient by a finite group. The range of p for which the Bergman projection is L^p -bounded on our class of Reinhardt domains is found to shrink as the complexity of the domain increases.

1 Introduction

1.1 Main result

Let $n \geq 2$, and for each $1 \leq j \leq n$, let $b^j = (b_1^j, \dots, b_n^j) \in \mathbb{Q}^n$ be an n -tuple of rational numbers. Let $\mathcal{U} \subset \mathbb{C}^n$ be a bounded domain (open connected subset) of the form

$$(1.1) \quad \mathcal{U} = \left\{ z \in \mathbb{C}^n : \text{for } 1 \leq j \leq n, \quad \prod_{k=1}^n |z_k|^{b_k^j} < 1 \right\},$$

where it is understood that a point $z \in \mathbb{C}^n$ does not belong to \mathcal{U} if for some $1 \leq j \leq n$, the quantity $\prod_{k=1}^n |z_k|^{b_k^j}$ is not defined due to division by zero. We call a domain such as \mathcal{U} a *monomial polyhedron*.

We refer the reader to Section 1.2 for a discussion of the significance of monomial polyhedra in complex analysis. Our main result is the following:

Theorem 1.2 *Suppose that the monomial polyhedron \mathcal{U} of (1.1) is bounded. Then there is a positive integer $\kappa(\mathcal{U})$ (the complexity of \mathcal{U} , whose computation is described below) such that the Bergman projection on \mathcal{U} is bounded on $L^p(\mathcal{U})$ if and only if*

$$(1.3) \quad \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) + 1} < p < \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) - 1}.$$

Received by the editors August 30, 2020; revised January 16, 2021; accepted January 25, 2021.

Published online on Cambridge Core February 8, 2021.

Chase Bender was supported by a Student Research and Creative Endeavors grant from Central Michigan University.

Debraj Chakrabarti was partially supported by National Science Foundation grant DMS-1600371.

AMS subject classification: 32A36.

Keywords: Bergman spaces, Bergman projection, Reinhardt domains, Generalized Hartogs triangle.

To compute $\kappa(\mathcal{U})$, first define for a vector $x \in \mathbb{Q}^n \setminus \{0\}$, the positive integer $h(x)$ (the *projective height* of x) as follows. If we think of x as the homogeneous coordinates of a point $[x]$ in the rational projective space $\mathbb{P}^{n-1}(\mathbb{Q})$, there is clearly an *integer* vector $y \in \mathbb{Z}^n$ such that $[y] = [x]$ (i.e., there is a $\lambda \in \mathbb{Q} \setminus \{0\}$ such that $y = \lambda x$), and we have additionally that $\gcd(y_1, \dots, y_n) = 1$. We then set

$$(1.4) \quad h(x) = \sum_{j=1}^n |y_j|.$$

We can think of h as a *height function* on $\mathbb{P}^{n-1}(\mathbb{Q})$ in the sense of Diophantine geometry, uniformly comparable to the standard multiplicative height function (see [HS00, pp. 174 ff.]).

Let B be the $n \times n$ matrix whose entry in the j th row and k th column is b_k^j , i.e., the j th row of B is the multi-index b^j in (1.1). It will follow from our work below (Proposition 3.2) that the matrix $B \in M_n(\mathbb{Q})$ is invertible. We define

$$(1.5) \quad \kappa(\mathcal{U}) = \max_{1 \leq k \leq n} h(B^{-1}e_k),$$

where e_k denotes the $n \times 1$ column vector all whose entries are zero, except the k th, which is 1. Notice that $B^{-1}e_k$ is simply the k th *column* of the matrix B^{-1} , that is, the arithmetic complexity of the monomial polyhedron \mathcal{U} is the maximum projective height of the columns of B^{-1} , where B is the rational matrix whose rows are the multi-indices occurring in the n inequalities that define \mathcal{U} in (1.1). It will be shown below in Proposition 3.5 that the integer $\kappa(\mathcal{U})$ is determined only by the domain \mathcal{U} and not the particular representation on the right hand side of (1.1).

1.2 Singular Reinhardt domains in complex analysis

Except in the degenerate case when it reduces to a polydisc (e.g., when $b^j = e_j$, the j th natural basis vector of \mathbb{Q}^n), the domain \mathcal{U} is a Reinhardt pseudoconvex domain (with center of symmetry at the origin) such that the origin is a boundary point. These *singular Reinhardt domains* (their boundaries are not Lipschitz at 0) display pathological holomorphic extension phenomena: the best-known example is that of the Hartogs triangle $\{|z_1| < |z_2| < 1\} \subset \mathbb{C}^2$, corresponding to a \mathcal{U} with $b^1 = (1, -1)$, $b^2 = (0, 1)$ (see [Beh33, Sib75]). For example, on a singular Reinhardt domain, each holomorphic function smooth up to the closure extends holomorphically to a fixed neighborhood of the closure (see [Cha19]), something which is impossible for smoothly bounded pseudoconvex domains [HS80, Cat80]. Therefore, a profound understanding of function theory on these domains is an important step in extending classical results on the regularity of the $\bar{\partial}$ -problem (and associated operators such as the Bergman projection) to new and more general settings (see [JP08]). Monomial polyhedra are an interesting class of such singular Reinhardt pseudoconvex domains with tractable geometry and some very interesting properties. They can be compared to analytic polyhedra in the classical theory of pseudoconvex domains: model exhausting domains where explicit computations are possible (see [Vla66, Section 24]).

The striking phenomenon observed in Theorem 1.2 was first noticed (see [EM17]) in the setting of the so-called *generalized Hartogs triangles*, defined for coprime positive integers k_1, k_2 as

$$(1.6) \quad H_{k_1/k_2} = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^{k_1/k_2} < |z_2| < 1\},$$

which corresponds to $b^1 = \left(\frac{k_1}{k_2}, -1\right)$ and $b^2 = (0, 1)$. It is striking that the range (1.3) should depend, not on the shape of the domain as a subset of \mathbb{C}^2 (which is determined in the case of H_{k_1/k_2} by the “fatness exponent” $\frac{k_1}{k_2}$), but on the complexity (which, for H_{k_1/k_2} is $k_1 + k_2$), the range becoming narrower as the complexity rises. As a limiting case, if $\gamma > 0$ is irrational, on the domain $\{|z_1|^\gamma < |z_2| < 1\} \subset \mathbb{C}^2$ (a domain of infinite complexity!), the Bergman projection is bounded in the L^p -norm, only if $p = 2$. The proofs of the precise range of L^p -boundedness of the Bergman projection on H_{k_1/k_2} and its generalizations in earlier work [EM17, Che17, Huo18, Zha20, Zha19] consist of an explicit computation of the Bergman kernel, followed by an application of Schur’s theorem on L^p boundedness of operators defined by integral kernels to determine the range of L^p -boundedness. Other authors have used techniques of classical harmonic analysis, such as weak-type endpoint estimates along with interpolation in L^p -spaces, Muckenhoupt A_p weights etc. to study related questions. See [CZ16, Edh16, EM16, CEM19, HW19, CKY20, EM20, CJY20] for other results in this circle of ideas.

Theorem 1.2 not only encompasses the known examples of domains on which the relation between the regularity of the Bergman projection and arithmetic complexity has been observed, but also substantially extends this class of domains. Its proof is based on an understanding of the geometry of monomial polyhedra as quotient domains. It is hoped that this will eventually lead to a deeper understanding of this mysterious notion of complexity, and its extension to other contexts.

1.3 Ingredients in the proof of Theorem 1.2

The range of L^p -boundedness of the Bergman projection on a domain is a function theoretic property determined by its Hermitian geometry, but the full extent of this relationship is yet to be understood. This article brings to bear a new perspective on this problem in the case of monomial polyhedra: one in which the domain is realized as a *quotient* of a simpler domain under the action of a group of biholomorphic automorphisms Γ (Theorem 3.12 below).

Our approach to the geometry of \mathcal{U} in Section 3 is inspired by the observation that in the “log-absolute coordinates” $\xi_k = \log |z_k|$, it is represented as

$$(1.7) \quad \sum_{k=1}^n b_k^j \xi_k < 0, \quad \text{for each } 1 \leq j \leq n,$$

which is an *open polyhedral cone* (intersection of open half-spaces) in the sense of convex geometry. By a classical result (see [Zie95, Theorem 1.3, p. 30], also [Grü03, Section 3.1]), such a polyhedral cone can also be represented as the cone generated by its extreme points, i.e., it is the image of an orthant (in a possibly higher dimensional Euclidean space) under a linear map. In Section 3, we prove an analogous statement

for \mathcal{U} : there is a domain $\mathbb{D}_{L(B)}^n \subset \mathbb{C}^n$, which is the product of a certain number of unit discs with a certain number of punctured unit discs, and a proper holomorphic map $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$ which is of the “quotient type” (see Definition 3.10), i.e., off some small analytic sets in the source and target, it is essentially a quotient map by a group Γ of automorphisms of the source. It turns out that the map Φ_A is of “monomial type” in the sense of [NP20], i.e., an n -dimensional analog of the branched covering map from the disc to itself given by $z \mapsto z^a$ for an integer $a > 0$. This fact has several pleasant consequences and facilitates computations.

One of the consequences of the existence of the map Φ_A is that it allows us to compute the Bergman kernel of \mathcal{U} explicitly. This has been a crucial step in the study of the Bergman projection on such domains in all prior investigations. In this paper, however, we avoid computing Bergman kernels and directly study the transformation properties of L^p -Bergman spaces. However, we do show in Proposition 3.22, using Theorem 3.12, that the Bergman kernel of \mathcal{U} is a rational function.

In Section 4, we study how L^p -Bergman spaces and the Bergman projection acting on L^p -Bergman spaces transform under proper holomorphic maps of quotient type. This point of view leads to a transformation law (Theorem 4.15 below) relating the L^p -Bergman spaces and Bergman projections of the source and target—one that is closely connected to the well-known Bell’s transformation law relating the Bergman kernels. One new ingredient here is the use of subspaces *invariant* under the action of the deck transformation group of the proper holomorphic map, which allows us to state a sharp result which can be used in the Proof of Theorem 1.2. We believe that the considerations of Section 4 have an independent interest beyond their application here.

The transformation law Theorem 4.15 can be used to pull back the problem from \mathcal{U} to the well-understood domain $\mathbb{D}_{L(B)}^n$, which is the polydisc, except for a missing analytic hypersurface. The pulled-back problem can be solved using classical estimates on the polydisc. This is achieved in Sections 5 and 6, completing the proof of Theorem 1.2.

The germ of the idea of relating the L^p -regularity of the Bergman projection with the properties of a “uniformizing” map from a simpler domain may already be found in [CZ16, CKY20]. The sharper version of this technique presented in this paper may be thought of as a step toward a unified understanding of the way in which boundary singularities affect the mapping properties of the Bergman projection.

1.4 Examples in \mathbb{C}^2

It is not difficult to see that a monomial polyhedron in \mathbb{C}^n is bounded by n Levi-flat “faces” (in the log-absolute representation (1.7), these faces are linear hyperplanes; see the proof of Proposition 3.5). The generalized Hartogs triangles of (1.6) are special in that one of the faces is a “coordinate face”, i.e., represented by a coordinate hyperplane in log-absolute coordinates. In (1.6), this is, $\{|z_2| = 1\}$ which corresponds to $\{\xi_2 = 0\}$ in the log-absolute representation. In two dimensions, the generic monomial polyhedron has two noncoordinate faces, and can be thought of as an intersections of two generalized Hartogs triangles. The Reinhardt shadows in \mathbb{R}^2 (the image of




Figure 1: Reinhardt shadows of some monomial polyhedra.

$z \mapsto (|z_1|, |z_2|)$ of three monomial polyhedra in \mathbb{C}^2 can be seen in Figure 1. The domains corresponding to (a), (b), and (c) are respectively given by:

$$\{|z_1|^4 < |z_2| < |z_1|^{1/3}\}, \quad \{|z_1|^{1/2} < |z_2| < |z_1|^{1/4}\}, \quad \text{and} \quad \{|z_1|^{1/2} < |z_2| < 1\}.$$

1.5 L^p -theory of the Bergman projection

We collect here some general information about the Bergman projection, and set up notation for later use.

Let Ω be a domain (an open connected set) in \mathbb{C}^n . The Bergman space $A^2(\Omega)$ is the Hilbert space of holomorphic functions on Ω which are square integrable with respect to the Lebesgue measure; see [Kra13] for a modern treatment. The space $A^2(\Omega)$ is a closed subspace of $L^2(\Omega)$, the usual Hilbert space of measurable functions square integrable with respect to the Lebesgue measure. The Bergman projection is the orthogonal projection

$$B_\Omega : L^2(\Omega) \rightarrow A^2(\Omega).$$

The construction of Bergman spaces has a contravariant functorial character. If $\phi : \Omega_1 \rightarrow \Omega_2$ is a suitable holomorphic map of domains, we can associate a continuous linear mapping of Hilbert spaces $\phi^\sharp : L^2(\Omega_2) \rightarrow L^2(\Omega_1)$ defined for each $f \in L^2(\Omega_2)$ by

$$(1.8) \quad \phi^\sharp(f) = f \circ \phi \cdot \det \phi',$$

where $\phi'(z) : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is the complex derivative of the map ϕ at $z \in \Omega_1$. It is clear that ϕ^\sharp restricts to a map $A^2(\Omega_2) \rightarrow A^2(\Omega_1)$. We will refer to ϕ^\sharp as the pullback induced by ϕ . It is not difficult to see that if ϕ is a biholomorphism, then the pullback ϕ^\sharp is an isometric isomorphism of Hilbert spaces $L^2(\Omega_2) \cong L^2(\Omega_1)$, and restricts to an isometric isomorphism $A^2(\Omega_2) \cong A^2(\Omega_1)$. This biholomorphic invariance of Bergman spaces can be understood intrinsically by interpreting the Bergman space as a space of top-degree holomorphic forms (see [Kob59] or [Kra13, p. 178 ff.]), and the map ϕ^\sharp as the pullback map of forms induced by the holomorphic map ϕ . This invariance can be extended to proper holomorphic mappings via Bell’s transformation formula, and lies at the heart of classical applications of Bergman theory to the boundary regularity of holomorphic maps; see [Bel81, Bel82, DF82, BC82].

For $0 < p < \infty$, define L^p -Bergman spaces $A^p(\Omega)$ of p th power integrable holomorphic functions on Ω . For $p \geq 1$, these are Banach spaces when equipped with the L^p -norm. An extensive theory of these spaces on the unit disc has been developed, in analogy with the theory of Hardy spaces (cf. [DS04, HKZ00]). Unlike the L^2 -Bergman space, the general L^p -Bergman space is not invariantly determined by the complex structure alone, but also depends on the Hermitian structure of the domain as a subset of \mathbb{C}^n . An important question about these spaces is the boundedness of the Bergman projection in the L^p -norm. After initial results were obtained for discs and balls ([ZJ64, FR75]) the problem was studied on various classes of smoothly bounded pseudoconvex domains using estimates on the kernel (e.g., [PS77, MS94]). On these domains the Bergman projection is bounded in L^p for $1 < p < \infty$. Many examples have been given which show that there are domains on which the Bergman projection fails to be bounded in L^p for certain p . See [Bar84, BŞ12, KP08, Hed02], in addition to the singular Reinhardt domains already mentioned in [CZ16, EM16, EM17, Chel7, CKY20, Huo18, HW19]. This paper proves sharp L^p -regularity results on a large class of singular Reinhardt domains.

1.6 Two examples of Theorem 1.2

We illustrate the application of Theorem 1.2 to two families of domains generalizing the Hartogs triangle to higher dimensions, recapturing the results of [Zha20, Zha19].

Let $k = (k_1, \dots, k_n)$ be an n -tuple of positive integers. The domain

$$(1.9) \quad \mathcal{H}_k = \left\{ z \in \mathbb{D}^n : |z_1|^{k_1} < \prod_{j=2}^n |z_j|^{k_j} \right\},$$

was introduced in [CKMM20] where it was called an *elementary Reinhardt domain of signature 1*, and its Bergman kernel was computed explicitly. We see that \mathcal{H}_k is a monomial polyhedron as in (1.1), since

$$(1.10) \quad \mathcal{H}_k = \left\{ z \in \mathbb{C}^n : |z_1|^{k_1} |z_2|^{-k_2} \dots |z_n|^{-k_n} < 1, \text{ and } |z_j| < 1 \text{ for } 2 \leq j \leq n \right\}.$$

The matrix B whose rows are multi-indices occurring in the inequalities in (1.10) is then given by

$$B = \left(\begin{array}{c|ccc} k_1 & -k_2 & \dots & -k_n \\ \hline 0 & & & \\ \vdots & & I_{n-1} & \\ 0 & & & \end{array} \right), \text{ so that } B^{-1} = \frac{1}{k_1} \left(\begin{array}{c|ccc} 1 & k_2 & \dots & k_n \\ \hline 0 & & & \\ \vdots & & k_1 \cdot I_{n-1} & \\ 0 & & & \end{array} \right),$$

where I_{n-1} is the identity matrix of size $n - 1$. The projective height of the first column of B^{-1} is 1, and for $2 \leq j \leq n$ that of the j th column is $\frac{k_1 + k_j}{\gcd(k_1, k_j)}$, using (1.4). Therefore, we get the complexity of \mathcal{H}_k as (see (1.5)):

$$\kappa(\mathcal{H}_k) = \max \left\{ 1, \frac{k_1 + k_2}{\gcd(k_1, k_2)}, \dots, \frac{k_1 + k_n}{\gcd(k_1, k_n)} \right\} = \max_{2 \leq j \leq n} \left\{ \frac{k_1 + k_j}{\gcd(k_1, k_j)} \right\}.$$

Noting that the function $x \mapsto \frac{x}{x-1}$ is decreasing for $x > 1$, and the function $x \mapsto \frac{x}{x+1}$ is increasing, by Theorem 1.2, the Bergman projection on \mathcal{H}_k is bounded on $L^p(\mathcal{H}_k)$ if and only if

$$\max_{2 \leq j \leq n} \frac{2(k_1 + k_j)}{k_1 + k_j + \gcd(k_1, k_j)} < p < \min_{2 \leq j \leq n} \frac{2(k_1 + k_j)}{k_1 + k_j - \gcd(k_1, k_j)},$$

in consonance with the result of [Zha19].

The second family of domains, which we denote by \mathcal{S}_k , gives a different type of generalization of the Hartogs triangle. For an n -tuple of positive integers (k_1, \dots, k_n) we define:

$$(1.11) \quad \mathcal{S}_k = \{z \in \mathbb{C}^n : |z_1|^{k_1} < |z_2|^{k_2} < \dots < |z_n|^{k_n} < 1\}.$$

In [Par18], the Bergman kernel of \mathcal{S}_k was explicitly computed for $n = 3$, and in [Chel7] the special case $k = (1, 1, \dots, 1)$ was considered and it was shown that the Bergman projection is L^p -bounded on $\mathcal{S}_{(1,1,\dots,1)}$ if and only if $\frac{2n}{n+1} < p < \frac{2n}{n-1}$. In [Zha20] the Bergman kernel of \mathcal{S}_k was computed in general, and the range of L^p -boundedness of the Bergman projection was determined. Since \mathcal{S}_k is a monomial polyhedron given by

$$(1.12) \quad \mathcal{S}_k = \{z \in \mathbb{C}^n : \text{for } 1 \leq j \leq n-1, |z_j|^{k_j} |z_{j+1}|^{-k_{j+1}} < 1, \text{ and } |z_n|^{k_n} < 1\},$$

the matrix B in the definition of complexity and its inverse B^{-1} are given as below:

$$B = \begin{pmatrix} k_1 & -k_2 & 0 & 0 & \dots & 0 \\ 0 & k_2 & -k_3 & 0 & \dots & 0 \\ \vdots & & \ddots & \ddots & & \vdots \\ \vdots & & & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & k_{n-1} & -k_n \\ 0 & 0 & 0 & \dots & 0 & k_n \end{pmatrix}, \quad B^{-1} = \frac{1}{K} \begin{pmatrix} \ell_1 & \ell_1 & \ell_1 & \dots & \dots & \ell_1 \\ 0 & \ell_2 & \ell_2 & \dots & \dots & \ell_2 \\ 0 & 0 & \ell_3 & \dots & \dots & \ell_3 \\ \vdots & \vdots & & \ddots & & \vdots \\ 0 & 0 & \dots & \dots & 0 & \ell_n \end{pmatrix},$$

where $K = \prod_{j=1}^n k_j$ and $\ell_j = \frac{K}{k_j}$ for $1 \leq j \leq n$. Note that the height of the m th column of B^{-1} is

$$h_m = \frac{\sum_{j=1}^m \ell_j}{\gcd(\ell_1, \dots, \ell_m)}.$$

As m increases, the numerator $\sum_{j=1}^m \ell_j$ of h_m increases and the denominator $\gcd(\ell_1, \dots, \ell_m)$ decreases, so h_m increases with m . Therefore, the complexity of \mathcal{S}_k is

$$\kappa(\mathcal{S}_k) = \max_{1 \leq m \leq n} h_m = h_n = \frac{\sum_{j=1}^n \ell_j}{\gcd(\ell_1, \dots, \ell_n)}.$$

Therefore, Theorem 1.2 shows that the Bergman projection on \mathcal{S}_k is bounded on $L^p(\mathcal{S}_k)$ if and only if

$$\frac{2 \sum_{j=1}^n \ell_j}{\sum_{j=1}^n \ell_j + \gcd(\ell_1, \dots, \ell_n)} < p < \frac{2 \sum_{j=1}^n \ell_j}{\sum_{j=1}^n \ell_j - \gcd(\ell_1, \dots, \ell_n)},$$

recapturing the main result of [Zha20].

2 Notation

2.1 Elementwise operations on matrices

Let A be an $m \times n$ matrix. We denote the entry of A at the j th row and k th column by a_k^j , where $1 \leq j \leq m, 1 \leq k \leq n$. We use two kinds of products of matrices: one is the standard matrix multiplication, denoted by simple juxtaposition AB , or sometimes a dot for clarity: $A \cdot B$.

We also need a second type of multiplication, the *elementwise*, or *Hadamard–Schur* product, in which the product of two $m \times n$ matrices $A = (a_k^j)$ and $B = (b_k^j)$ is the $m \times n$ matrix C , for which we have $c_k^j = a_k^j b_k^j$, i.e., the element at a certain position of the product is the product of the corresponding entries of the factors. We denote this by

$$(2.1) \quad C = A \odot B.$$

It will be important to distinguish between column and row vectors. We denote the R -module of $n \times 1$ column vectors by R^n , where R can be one of $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$. The R -module of $1 \times n$ row vectors is denoted by $(R^n)^\dagger$. We write the entries of the row vector a (resp. the column vector b) as (a_1, \dots, a_n) (resp. as $(b_1, \dots, b_n)^T$). We let $\mathbb{1}$ denote a $1 \times n$ row vector, all whose entries are 1. The positive integer n will be clear from the context:

$$(2.2) \quad \mathbb{1} = (1, \dots, 1). \quad (n \text{ ones})$$

Similarly, $\mathbb{1}^T$ is the $n \times 1$ column vector, all whose entries are 1. Notice that these are identity elements for the elementwise multiplication of row and column vectors.

For real matrices of the same size A, B , the notations

$$(2.3) \quad A > B, A \geq B, A < B, A \leq B$$

stand for the natural elementwise order, e.g., $A > B$ denotes that $a_k^j > b_k^j$. Note that this elementwise ordering of matrices is distinct from other notions of matrix ordering, such as positive (semi-)definiteness.

2.2 Vector and matrix powers

If α is a row vector of size n , and z is column vector of the same size, we will denote

$$z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n} = \prod_{j=1}^n z_j^{\alpha_j},$$

whenever the powers $z_j^{\alpha_j}$ make sense, and where we use the convention $0^0 = 1$. For example, we could have $\alpha \in (\mathbb{Z}^n)^\dagger$ and $z \in \mathbb{C}^n$ such that for each j such that $\alpha_j < 0$, we have $z_j \neq 0$. In this context α is typically called a *multi-index* and z^α a *Laurent monomial*. We also set

$$(2.4) \quad \varphi_\alpha(z) = z^\alpha.$$

Informally, we think of a monomial function as a “vector power” of a vector variable.

We will also use “matrix powers.” Recall from the previous subsection that for an $n \times n$ matrix A , we denote the element at the j th row and k th column of A by a_k^j . Let a^j denote the j th row of A , so that each row can be thought of as a multi-index. We then define for a column vector z of size n ,

$$(2.5) \quad z^A = \begin{pmatrix} z^{a^1} \\ \vdots \\ z^{a^n} \end{pmatrix} = \begin{pmatrix} z_1^{a_1^1} z_2^{a_2^1} \cdots z_n^{a_n^1} \\ \vdots \\ z_1^{a_1^n} z_2^{a_2^n} \cdots z_n^{a_n^n} \end{pmatrix}$$

provided each of the monomials is defined. We will also set

$$(2.6) \quad \Phi_A(z) = z^A,$$

which is called a *monomial map*.

For row vectors α, β , square matrices A, B and column vectors z, w such that the vector and matrix powers are well-defined, the following simple algebraic properties of monomials and monomial mappings can be easily verified. Not unexpectedly, these mirror familiar rules of elementary algebra for exponents.

$$(2.7a) \quad (z \odot w)^\alpha = z^\alpha \cdot w^\alpha$$

$$(2.7b) \quad (z \odot w)^A = z^A \odot w^A$$

$$(2.7c) \quad z^{\alpha+\beta} = z^\alpha \cdot z^\beta$$

$$(2.7d) \quad z^{A+B} = z^A \odot z^B$$

$$(2.7e) \quad (z^A)^\alpha = z^{\alpha A} \quad \text{i.e.,} \quad \varphi_\alpha \circ \Phi_A = \varphi_{\alpha A}$$

$$(2.7f) \quad (z^A)^B = z^{BA} \quad \text{i.e.,} \quad \Phi_B \circ \Phi_A = \Phi_{BA}$$

2.3 Two other maps

(1) The (elementwise) *exponential map*

$$(2.8) \quad \exp : \mathbb{C}^n \rightarrow (\mathbb{C}^*)^n, \quad \exp(z) = (e^{z_1}, \dots, e^{z_n})^T$$

can be thought of as the exponential map associated to the abelian Lie group $(\mathbb{C}^*)^n$. Notice that for $\alpha \in (\mathbb{Z}^n)^\dagger$ and $A \in M_n(\mathbb{Z})$ we have:

$$(2.9) \quad \exp(z)^\alpha = e^{\alpha z} \text{ and } \exp(z)^A = \exp(Az).$$

(2) The *multi-radius map*

$$(2.10) \quad \rho : \mathbb{C}^n \rightarrow \mathbb{R}^n \quad z \mapsto (|z_1|, \dots, |z_n|)^T$$

restricts to a surjective group homomorphism $(\mathbb{C}^*)^n \rightarrow (\mathbb{R}^+)^n$, and satisfies, for $\alpha \in (\mathbb{Z}^n)^\dagger$ and $A \in M_n(\mathbb{Z})$

$$(2.11) \quad \rho(z)^\alpha = |z^\alpha| \text{ and } \rho(z)^A = \rho(z^A).$$

3 Geometry of monomial polyhedra

The main goal of this section is Theorem 3.12, which gives a “uniformization” of a monomial polyhedron by a proper holomorphic monomial map. This construction is crucial for everything that follows. We also show that the Bergman kernel of a monomial polyhedron is rational, although this fact is not used in the sequel.

3.1 The matrix *B*

The definition of the domain \mathcal{U} as given in (1.1) can be succinctly rewritten, using the notation introduced in the previous Section 2 as

$$(3.1) \quad \mathcal{U} = \{z \in \mathbb{C}^n : \rho(z)^B \text{ is defined and } \rho(z)^B < \mathbb{1}^T\},$$

where $B \in M_n(\mathbb{Q})$ is the matrix whose j th row is b^j . In the next proposition, we show that the matrix B in the definition of \mathcal{U} can always be taken to be an integer matrix which is *monotone* in the sense of Collatz (see [Col66, p. 376 ff.]):

Proposition 3.2 *In the representation (3.1) of the bounded monomial polyhedron \mathcal{U} of Theorem 1.2, we may assume without loss of generality (after switching two rows, if necessary) that*

$$(3.3) \quad B \in M_n(\mathbb{Z}), \quad \det B > 0, \quad \text{and} \quad B^{-1} \geq 0.$$

Proof Let $B \in M_n(\mathbb{Q})$ be the matrix whose rows are b^1, \dots, b^n , where for $1 \leq j \leq n$, the vector $b^j \in (\mathbb{Q}^n)^\dagger$, $b^j = (b_1^j, \dots, b_n^j)$ is as in (1.1), which can be written in the form (3.1) using the notation introduced above. Notice that if any one of the vectors b^j is 0, then \mathcal{U} is empty, since the inequality $\rho(z)^{b^j} < 1$ becomes $1 < 1$.

If $\delta_j > 0$ is a common denominator for the rational numbers b_1^j, \dots, b_n^j , then notice that for a $z \in \mathbb{C}^n$, the quantity $\rho(z)^{b^j}$ is defined and less than 1 if and only if $\rho(z)^{\delta_j b^j}$ is defined and less than 1. Therefore, we can assume without loss of generality that the matrix B has integer entries. Note also that interchanging the rows of B simply corresponds renumbering the equations in (1.1), so we can further assume without loss of generality that $\det B \geq 0$.

Now, assuming that $B \in M_n(\mathbb{Z})$ and $\det B \geq 0$, suppose for a contradiction that $\det B = 0$. We will show that \mathcal{U} is unbounded, which will contradict the hypothesis of Theorem 1.2. Since $\det B = 0$, there is a nonzero vector $x \in \mathbb{R}^n$ such that $Bx = 0$. Let $r \in \mathcal{U}$ be such that $r > 0$ (such an r exists since \mathcal{U} is nonempty, open, and Reinhardt). Consider the curve $f : \mathbb{R} \rightarrow \mathbb{C}^n$ parametrized by

$$f(t) = r \odot \exp(tx),$$

where \exp is as in (2.8). Observe that

$$f(t)^B = r^B \odot \exp(tx)^B = r^B \odot \exp(tBx) = r^B,$$

since $Bx = 0$. Therefore,

$$\rho(f(t))^B = \rho(f(t)^B) = r^B.$$

But since $r \in \mathcal{U}$, it follows that $\rho(f(t))^B = r^B < \mathbb{1}^T$, so in fact f is a curve in \mathcal{U} . As $x \in \mathbb{R}^n$ is nonzero, some component x_k of x is nonzero, so that the k th component of the curve f , given by $f_k(t) = r_k e^{tx_k}$ is unbounded as $t \rightarrow \pm\infty$. Thus, \mathcal{U} contains the unbounded image of the curve f , showing that \mathcal{U} is unbounded if $\det B = 0$. Since by assumption \mathcal{U} is bounded, we have $\det B \neq 0$. So $\det B > 0$.

To show that $B^{-1} \geq 0$, we let A be the adjugate of B , i.e., $A = \text{adj } B$, and we show that

$$\Phi_A((\mathbb{D}^*)^n) \subset \mathcal{U},$$

where Φ_A is the monomial map as in (2.6), and $(\mathbb{D}^*)^n$ is the product of n punctured unit disks. Since \mathcal{U} is bounded this implies that $\Phi_A((\mathbb{D}^*)^n)$ is bounded. Let $w \in \Phi_A((\mathbb{D}^*)^n)$ and let $z \in (\mathbb{D}^*)^n$ be such that $z^A = w$. Notice that $z \in (\mathbb{D}^*)^n$ is equivalent to $0 < \rho(z) < \mathbb{1}^T$, and we also have $z^A = w \in (\mathbb{C}^*)^n$. By applying (2.11) and Cramer’s rule, we find

$$\rho(w)^B = (\rho(z)^A)^B = \rho(z)^{BA} = \rho(z)^{(\det B)I} = (|z_1|^{\det B}, \dots, |z_n|^{\det B})^T.$$

As $\det B > 0$ we find that $\rho(w)^B < \mathbb{1}^T$, i.e., $w \in \mathcal{U}$, yielding the desired inclusion $\Phi_A((\mathbb{D}^*)^n) \subset \mathcal{U}$.

If $A \not\geq 0$ then there exist some $1 \leq j, k \leq n$ such that $a_k^j < 0$. Let $z \in (\mathbb{D}^*)^n$ be such that all its components, except the k th, are $1/2$. Then the j th component of $\Phi_A(z)$ is given by

$$z^{a_j^j} = z_k^{a_k^j} \cdot \frac{1}{2^m}, \quad \text{where} \quad m = \sum_{\substack{1 \leq \ell \leq n \\ \ell \neq k}} a_\ell^j.$$

As $a_k^j < 0$ we see that as $z_k \rightarrow 0$, the monomial $z^{a_j^j}$ is unbounded, showing that $\Phi_A((\mathbb{D}^*)^n)$ is unbounded. Therefore, the boundedness of \mathcal{U} guarantees that $A = \text{adj } B \geq 0$.

To complete the proof note that $A \geq 0$ implies that $B^{-1} = (\det B)^{-1} \text{adj } B \geq 0$. ■

Due to Proposition 3.2, we can easily establish a more concrete bound on monomial polyhedra:

Corollary 3.4 *A bounded monomial polyhedron is contained in the unit polydisc \mathbb{D}^n and its boundary contains the unit torus $\mathbb{T}^n = \{|z_j| = 1, 1 \leq j \leq n\}$.*

Proof For a bounded monomial polyhedron \mathcal{U} , let $z \in \mathcal{U}$ so that $\rho(z)^B < 1$, where B is as in Proposition 3.2. Since $B^{-1} \geq 0$, we have for any $r \in \mathbb{R}^n$ with, $0 \leq r < 1$ that $r^{B^{-1}} < 1$. Therefore, we have $(\rho(z)^B)^{B^{-1}} = \rho(z) < 1$, i.e., $z \in \mathbb{D}^n$. The second assertion follows on noting that in the log absolute representation (1.7) of \mathcal{U} , the origin (which corresponds to the unit torus of \mathbb{C}^n) is a boundary point. ■

In the Definition (3.1), it is clear that the matrix B is not unique: permuting the order of inequalities, does not change to domain \mathcal{U} , and multiplying for each j the row b^j by the same rational $\delta_j > 0$ also gives exactly the same \mathcal{U} . Therefore, for any permutation matrix P and any positive diagonal matrix D with rational entries, the domains corresponding to B and DPB are the same. The following proposition shows that the complexity $\kappa(\mathcal{U})$ is independent of the matrix B and depends only on \mathcal{U} .

Proposition 3.5 *The complexity $\kappa(\mathcal{U})$ is independent of the choice of the representing matrix B in (3.1).*

Proof Notice that switching two rows of B does not change the complexity of \mathcal{U} , we can assume that the conditions (3.3) hold for the representing matrix. In the log-absolute coordinates $\xi_k = \log |z_k|$ the domain \mathcal{U} is represented by the equations (1.7), which define the open polyhedral cone $\mathcal{C}_{\mathcal{U}} = \{B\xi < 0\} \subset \mathbb{R}^n$. We claim that for each $1 \leq j \leq n$, the hyperplane $H_j = \{b^j \xi = 0\}$ determined by the j th row of the matrix B is a face of $\mathcal{C}_{\mathcal{U}}$, i.e., the intersection $\overline{\mathcal{C}_{\mathcal{U}}} \cap H_j \neq \emptyset$. Indeed, since by Proposition 3.2, $B^{-1} > 0$ and B is a linear automorphism of \mathbb{Q}^n , it is easy to verify that the image $B^{-1}(\mathcal{C}_{\mathcal{U}})$ is precisely the negative orthant $\{\eta < 0\} \subset \mathbb{R}^n$. But the negative orthant clearly has faces $\{\eta_j = 0\}, j = 1, \dots, n$, which corresponds to the faces H_j of $\mathcal{C}_{\mathcal{U}}$.

Now, if $C \in M_n(\mathbb{Q})$ is another matrix (satisfying (3.3)) such that

$$\mathcal{U} = \{z \in \mathbb{C}^n : \rho(z)^C \text{ is defined and } \rho(z)^C < \mathbb{1}^T\},$$

then the faces of $\mathcal{C}_{\mathcal{U}}$ are also given by $\{c^j \xi = 0\}, 1 \leq j \leq n$, so that there exists a permutation $\sigma \in S_n$ such that

$$\{c^j \xi = 0\} = H_{\sigma(j)} = \{b^{\sigma(j)} \xi = 0\}$$

for each $1 \leq j \leq n$. Then the transposes $(c^j)^T$ and $(b^{\sigma(j)})^T$ are both normal to $H_{\sigma(j)}$, so that there is a $\delta_j \in \mathbb{Q} \setminus \{0\}$ such that $c^j = \delta_j b^{\sigma(j)}$. Therefore, $C = DPB$, where D is the diagonal matrix $\text{diag}(\delta_1, \dots, \delta_n)$ and P is the permutation matrix whose j th column is $e_{\sigma(j)}$. Note that P^{-1} is also a permutation matrix, and hence $P^{-1} > 0$. Therefore, since $C^{-1} = B^{-1}P^{-1}D^{-1}$ and we have $C^{-1} > 0, B^{-1} > 0$ by Proposition 3.2, we have $D > 0$ since $B^{-1}P^{-1} > 0$. Therefore,

$$\begin{aligned} \max_{1 \leq j \leq n} h(C^{-1}e_j) &= \max_{1 \leq j \leq n} h(B^{-1}P^{-1}D^{-1}e_j) = \max_{1 \leq j \leq n} h\left(\frac{1}{\delta_j} B^{-1}e_{\sigma(j)}\right) \\ &= \max_{1 \leq j \leq n} h(B^{-1}e_{\sigma(j)}) = \max_{1 \leq j \leq n} h(B^{-1}e_j). \end{aligned}$$

This shows that $\kappa(\mathcal{U})$ as in 1.5 is well-defined, being independent of the matrix B . ■

3.2 The Jacobian determinant of Φ_A

For a subset $E \subseteq \{1, \dots, n\}$, define

$$(3.6) \quad \mathbb{C}_E^n := \{z \in \mathbb{C}^n : z_k \neq 0 \text{ for all } k \in E\} = \prod_{k=1}^n \Omega_k,$$

where $\Omega_k = \mathbb{C}$ if $k \notin E$ and $\Omega_k = \mathbb{C}^*$ if $k \in E$. Then for $P \in M_n(\mathbb{Q})$ define

$$(3.7) \quad K(P) = \{1 \leq k \leq n : p_k^j < 0 \text{ for some } 1 \leq j \leq n\},$$

that is, $K(P)$ is the set of indices of those columns of P which have at least one negative entry. Notice that if $P \in M_n(\mathbb{Z})$ the matrix power z^P is defined for $z \in \mathbb{C}^n$ if and only if $z \in \mathbb{C}_{K(P)}^n$, i.e., in computing z^P we do not raise zero to a negative power. Similarly, for $P \in M_n(\mathbb{Q})$ and a vector $r \in \mathbb{R}^n$ with $r_j \geq 0$, the rational power r^P is defined provided $r \in \mathbb{R}_{K(P)}^n$ for exactly the same reason.

The following analog of the calculus formula $\frac{d}{dx}x^n = nx^{n-1}$ may be found in [NP09, Lemma 4.2], and will be needed frequently in the sequel.

Lemma 3.8 For any $A \in M_n(\mathbb{Z})$ and any $z \in \mathbb{C}_{K(A)}^n$ we have

$$(3.9) \quad \det \Phi'_A(z) = \det A \cdot z^{\mathbb{1}A-1},$$

where $\Phi_A(z) = z^A$ and $\Phi'_A(z) : \mathbb{C}^n \rightarrow \mathbb{C}^n$ is the complex derivative of Φ_A at the point $z \in \mathbb{C}_{K(A)}^n$.

Proof First assume that z is such that $z_k \neq 0$ for each k . For $1 \leq j, k \leq n$, the entry in the j th row and k th column of the complex derivative matrix $\Phi'_A(z)$ is $\frac{\partial z^{a^j}}{\partial z_k}$, where recall that a^j denotes the multi-index in $(\mathbb{Z}^n)^\dagger$ whose entries constitute the j th row of A . If $a^j_k \neq 0$ then

$$\frac{\partial z^{a^j}}{\partial z_k} = a^j_k z_k^{a^j_k-1} \prod_{\substack{\ell=1 \\ \ell \neq k}}^n z_\ell^{a^j_\ell} = a^j_k \cdot \frac{z^{a^j}}{z_k}.$$

If $a^j_k = 0$, then $\frac{\partial z^{a^j}}{\partial z_k} = 0$, so in fact the above formula holds for all j, k . Now, by the representation of a determinant as an explicit polynomial in the matrix entries, we find

$$\begin{aligned} \det \Phi'_A(z) &= \sum_{\sigma \in \mathcal{S}_n} \operatorname{sgn}(\sigma) \prod_{j=1}^n \frac{\partial z^{a^j}}{\partial z_{\sigma(j)}} = \sum_{\sigma \in \mathcal{S}_n} \operatorname{sgn}(\sigma) \prod_{j=1}^n \left(a^j_{\sigma(j)} \frac{z^{a^j}}{z_{\sigma(j)}} \right) \\ &= \sum_{\sigma \in \mathcal{S}_n} \operatorname{sgn}(\sigma) \prod_{j=1}^n a^j_{\sigma(j)} \cdot \frac{\prod_{j=1}^n z^{a^j}}{\prod_{j=1}^n z_{\sigma(j)}} \\ &= \det A \cdot \frac{\prod_{j=1}^n z^{a^j}}{z^{\mathbb{1}}}, \end{aligned}$$

where we have used the fact that $\prod_{j=1}^n z_{\sigma(j)} = \prod_{j=1}^n z_j = z^{\mathbb{1}}$. Also,

$$\prod_{j=1}^n z^{a^j} = \prod_{j=1}^n \prod_{k=1}^n z_k^{a^j_k} = \prod_{k=1}^n z_k^{\sum_{j=1}^n a^j_k} = z^{\mathbb{1}A}.$$

Therefore,

$$\det \Phi'_A(z) = \det A \cdot \frac{z^{\mathbb{1}A}}{z^{\mathbb{1}}} = \det A \cdot z^{\mathbb{1}A-1},$$

for z such that $z_k \neq 0$ for each k . The result follows by analytic continuation. ■

3.3 A branched covering of the domain \mathcal{U}

In this section, we will construct a quotient map under a group action from a domain with simple geometry (a product of some copies of discs with some copies

of punctured discs) to the domain \mathcal{U} , and this construction will be fundamental in the proof of Theorem 1.2. We first specify what we mean by a quotient map.

Definition 3.10 Let $\Omega_1, \Omega_2 \subset \mathbb{C}^n$ be domains, let $\Phi : \Omega_1 \rightarrow \Omega_2$ be a proper holomorphic mapping. Let $\Gamma \subset \text{Aut}(\Omega_1)$ be a group of biholomorphic automorphisms of Ω_1 . We will say that Φ is of *quotient type with group* Γ if there exist closed lower-dimensional complex-analytic subvarieties $Z_j \subset \Omega_j, j = 1, 2$ such that Φ restricts to a covering map

$$\Phi : \Omega_1 \setminus Z_1 \rightarrow \Omega_2 \setminus Z_2$$

and for each $z \in \Omega_2 \setminus Z_2$, the action of Γ on Ω_1 restricts to a transitive action on the fiber $\Phi^{-1}(z)$. Other names in the literature for such proper maps include *regular* and *Galois* proper maps. The group Γ will be referred to as the group of deck transformations of Φ (sometimes called the Galois group).

Notice that the restricted map $\Phi : \Omega_1 \setminus Z_1 \rightarrow \Omega_2 \setminus Z_2$ becomes a so called *regular* covering map (see [Mas91, p. 135 ff.]), i.e., the covering map gives rise to a biholomorphism between $\Omega_2 \setminus Z_2$ and the quotient $(\Omega_1 \setminus Z_1)/\Gamma$, where it can be shown that Γ acts properly and discontinuously on $\Omega_1 \setminus Z_1$. Further, it follows that Γ is in fact the full group of deck transformations of the covering map $\Phi : \Omega_1 \setminus Z_1 \rightarrow \Omega_2 \setminus Z_2$, and that this covering map has exactly $|\Gamma|$ sheets. Notice that by analytic continuation, the relation $\Phi \circ \sigma = \Phi$ holds for each σ in Γ on all of Ω_1 .

Now let B be a matrix, let the set $K(B) \subset \{1, \dots, n\}$ be as in (3.7), the set of indices of those columns of B which have at least one negative entry. Define a subset $L(B) \subset \{1, \dots, n\}$ by setting

$$(3.11) \quad L(B) = \{1 \leq \ell \leq n : a_\ell^k \neq 0 \text{ for some } k \in K(B)\}.$$

We define the domain $\mathbb{D}_{L(B)}^n = \mathbb{D}^n \cap \mathbb{C}_{L(B)}^n$, i.e.,

$$\mathbb{D}_{L(B)}^n = \{z \in \mathbb{D}^n : z_\ell \neq 0 \text{ for all } \ell \in L(B)\}.$$

Notice that $\mathbb{D}_{L(B)}^n$ is the product of some copies of the unit disc with some copies of the punctured unit disc.

Theorem 3.12 Let \mathcal{U} be the domain of Theorem 1.2, and assume (without loss of generality) that \mathcal{U} is represented as in (3.1), where the matrix $B = (b_k^j)$ satisfies the conditions (3.3), and let $A = \text{adj } B$. Then the monomial map Φ_A of (2.6) maps $\mathbb{D}_{L(B)}^n$ onto \mathcal{U} , and the map so defined

$$(3.13) \quad \Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$$

is a proper holomorphic map of quotient type with group Γ consisting of the automorphisms $\sigma_v : \mathbb{D}_{L(B)}^n \rightarrow \mathbb{D}_{L(B)}^n$ given by

$$(3.14) \quad \sigma_v(z) = \exp(2\pi i A^{-1}v) \odot z$$

for $v \in \mathbb{Z}^n$. Further, the group Γ has exactly $\det A$ elements.

Some related results may be found in [NP20, Zwo99].

Proof By the conditions (3.3), we have $A = (\det B)B^{-1} \geq 0$, so Φ_A is defined on all of $\mathbb{D}_{L(B)}^n$. We first claim that the image $\Phi_A(\mathbb{D}_{L(B)}^n)$ is contained in $\mathbb{C}_{K(B)}^n$. Indeed, if $z \in \Phi_A(\mathbb{D}_{L(B)}^n)$, then $z_\ell \neq 0$ if $a_\ell^k \neq 0$ for some $k \in K(B)$. Therefore the k th element of the vector z^A , i.e.,

$$(3.15) \quad z^{a^k} = z_1^{a_1^k} \dots z_n^{a_n^k}$$

is nonzero, so $z^A \in \mathbb{C}_{K(B)}^n$. Now notice that

$$\mathcal{U} = \{z \in \mathbb{C}^n : \rho(z)^B \text{ is defined and } \rho(z)^B < \mathbb{1}^T\} \subset \mathbb{C}_{K(B)}^n,$$

since $\rho(z)^B$ (and z^B) are defined if and only if $z \in \mathbb{C}_{K(B)}^n$. Therefore, as in the proof of Proposition 3.2, we have

$$(\rho(z)^A)^B = \rho(z)^{BA} = \rho(z)^{(\det B)I} < \mathbb{1},$$

for each point $z \in \mathbb{D}_{L(B)}^n$ since $\det B > 0$. This shows that $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$ is a well defined holomorphic map.

We now show that, if we think of Φ_A as a map from \mathbb{C}^n to itself (recall that $A \geq 0$), then we have

$$(3.16) \quad \Phi_A^{-1}(\mathcal{U}) \subset \mathbb{D}_{L(B)}^n.$$

If $z \in \mathbb{C}^n$ is such that $\Phi_A(z) = z^A \in \mathcal{U}$, then $\rho(z^A)^B$ is well defined and $\rho(z^A)^B < \mathbb{1}^T$. The fact that $\rho(z^A)^B$ is well defined is equivalent to $z^A \in \mathbb{C}_{K(B)}^n$, i.e., for each $k \in K(B)$ we have the entry $z^{a^k} \neq 0$. Now from (3.15), we see that whenever $a_\ell^k \neq 0$ (so $a_\ell^k \geq 1$ since $A \geq 0$), we must have $z_\ell \neq 0$, i.e., $z \in \mathbb{C}_{L(B)}^n$. The other condition $\rho(z^A)^B < \mathbb{1}^T$ on the point z shows that

$$\rho(z)^{(\det B)I} = \rho(z)^{BA} = \rho(z^A)^B < \mathbb{1}^T.$$

Since $\det B > 0$, this shows that $\rho(z) < \mathbb{1}^T$, i.e., $z \in \mathbb{D}^n$. Together these conditions show that $z \in \mathbb{D}_{L(B)}^n$, proving the desired inclusion of (3.16).

We can now show that $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$ is a proper holomorphic map. Let $K \subset \mathcal{U}$ be compact. Since the topology induced on K from \mathcal{U} is the same as that induced from \mathbb{C}^n , it follows that K is closed in \mathbb{C}^n (and bounded). As Φ_A is continuous, $\Phi_A^{-1}(K)$ is closed in \mathbb{C}^n and in fact $\Phi_A^{-1}(K) \subseteq \mathbb{D}_{L(B)}^n$ by (3.16), so that it is bounded as well. Thus, $\Phi_A^{-1}(K)$ is compact for every compact $K \subset \mathcal{U}$, which is to say Φ_A is proper.

Now we want to verify that the proper holomorphic map $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$ is of quotient type in the sense of Definition 3.10 with group $\Gamma = \{\sigma_\nu : \nu \in \mathbb{Z}^n\}$, with σ_ν as in (3.14) above. Let

$$Z_1 = Z_2 = \{z^1 = 0\} = \bigcup_{j=1}^n \{z \in \mathbb{C}^n : z_j = 0\}$$

be the union of the coordinate hyperplanes, which is an analytic variety in \mathbb{C}^n of codimension one. Notice that the formula (3.9) shows that the set of regular points of the map Φ_A (i.e., the set of points $\{z \in \mathbb{D}_{L(B)}^n : \det \Phi'_A(z) \neq 0\}$) contains $\mathbb{D}_{L(B)}^n \setminus Z_1$, and from the fact that $\Phi_A(z) = z^A$, we see that the regular values of Φ_A contains the open set $\mathcal{U} \setminus Z_2$. Notice that both $\mathbb{D}_{L(B)}^n \setminus Z_1$ and $\mathcal{U} \setminus Z_2$ are open subsets of the “algebraic torus” $(\mathbb{C}^*)^n$, and it is clear that Φ_A maps $(\mathbb{C}^*)^n$ into itself (and is even a group homomorphism, if $(\mathbb{C}^*)^n$ is given the group structure from the elementwise multiplication operation \odot). Since a proper holomorphic map is a covering map restricted to its regular points, it now follows that $\Phi_A : \mathbb{D}_{L(B)}^n \setminus Z_1 \rightarrow \mathcal{U} \setminus Z_2$ is a holomorphic covering map. We need to show that it is regular.

Let $w \in \mathcal{U} \setminus Z_2$ have polar representation $w = \rho(w) \odot \exp(i\theta)$, where $\theta \in \mathbb{R}^n$. Let $z \in \mathbb{D}_{L(B)}^n$ be such that $\Phi_A(z) = z^A = w$. Comparing the radial and angular parts, one such preimage is

$$z = \rho(w)^{A^{-1}} \odot \exp(iA^{-1}\theta).$$

Now notice that the angular vector θ is known only up to an additive ambiguity of $2\pi\mathbb{Z}^n$, i.e., two values of θ corresponding to the same w differ by $2\pi v$ for some integer vector $v \in \mathbb{Z}^n$. Therefore two different preimages z, z' of the point w (corresponding to the choices θ and $\theta + 2\pi v$ of the angular vector in the polar representation of w) are related by

$$\begin{aligned} z' &= \rho(w)^{A^{-1}} \odot \exp(iA^{-1}(\theta + 2\pi v)) = \rho(w)^{A^{-1}} \odot \exp(iA^{-1}\theta) \odot \exp(2\pi iA^{-1}v) \\ &= \exp(2\pi iA^{-1}v) \odot z = \sigma_v(z), \end{aligned}$$

which shows that the group Γ acts transitively on the fiber $\Phi^{-1}(w)$.

Now we want to show that the group Γ has exactly $\det A$ elements. Consider the map $\psi : \mathbb{Z}^n \rightarrow \Gamma$ given by $\psi(v) = \sigma_v$. For any $v, \mu \in \mathbb{Z}^n$ and any $z \in \mathbb{D}_{L(B)}^n$ we find that

$$\begin{aligned} \psi(v + \mu)(z) &= \sigma_{v+\mu}(z) = \exp(2\pi iA^{-1}(v + \mu)) \odot z \\ &= \exp(2\pi iA^{-1}v) \odot (\exp(2\pi iA^{-1}\mu) \odot z) \\ &= \sigma_v \circ \sigma_\mu(z) \\ &= \psi(v) \circ \psi(\mu)(z). \end{aligned}$$

Hence, $\psi : \mathbb{Z}^n \rightarrow \Gamma$ is a surjective group homomorphism, and so we have an isomorphism

$$\bar{\psi} : \mathbb{Z}^n / \ker \psi \rightarrow \Gamma.$$

given by $\psi([v]) = \sigma_v$ where $[v] \in \mathbb{Z}^n / \ker \psi$ is the class of $v \in \mathbb{Z}^n$. Notice that $v \in \ker \psi$ if and only if $\exp(2\pi iA^{-1}v) = \mathbb{1}^T$. This means that $A^{-1}v \in \mathbb{Z}^n$, so it follows that $\ker \psi = A(\mathbb{Z}^n)$. Therefore, the group Γ is isomorphic to the quotient $\mathbb{Z}^n / A(\mathbb{Z}^n)$ by the isomorphism $\bar{\psi} : \mathbb{Z}^n / A(\mathbb{Z}^n) \rightarrow \Gamma$

To complete the proof, it is sufficient to recall that for any matrix $A \in M_n(\mathbb{Z})$ with $\det A \neq 0$ we have

$$(3.17) \quad |\mathbb{Z}^n / A(\mathbb{Z}^n)| = |\det A|.$$

For completeness, we give two proofs of (3.17), neither of which unfortunately avoids high technology. The first (see [NP20] and [Zwo99, Theorem 2.1]) is based on the Smith Canonical Form of an integer matrix (see [Mun84, p. 53 ff.]): for any $A \in M_n(\mathbb{Z})$ there exist $P, Q \in GL_n(\mathbb{Z})$ and a diagonal matrix $D \in M_n(\mathbb{Z})$, say $D = \text{diag}(\delta_1, \dots, \delta_n)$, such that $A = PDQ$. Consider the automorphism of \mathbb{Z}^n given by $x \mapsto P^{-1}x$. This maps $A(\mathbb{Z}^n)$ to $D(\mathbb{Z}^n)$ isomorphically, and therefore leads to an isomorphism $\mathbb{Z}^n/A(\mathbb{Z}^n)$ with $\mathbb{Z}^n/D(\mathbb{Z}^n)$, but $\mathbb{Z}^n/D(\mathbb{Z}^n)$ is isomorphic to the product abelian group $\prod_{j=1}^n \mathbb{Z}/\delta_j\mathbb{Z}$, which has exactly $\prod_{j=1}^n |\delta_j|$ elements. On the other hand note that

$$|\det A| = |\det(PDQ)| = |\det P \det D \det Q| = \left| \pm 1 \cdot \prod_{j=1}^n \delta_j \cdot \pm 1 \right|,$$

which completes the first proof.

The second proof of (3.17) is geometric, and based on noticing that the inclusion of abelian groups $A(\mathbb{Z}^n) \hookrightarrow \mathbb{Z}^n$ gives rise to a covering map of tori

$$(3.18) \quad p : \mathbb{R}^n/A(\mathbb{Z}^n) \rightarrow \mathbb{R}^n/\mathbb{Z}^n$$

given by $x + A(\mathbb{Z}^n) \mapsto x + \mathbb{Z}^n$. If we endow \mathbb{R}^n with the Euclidean metric (thought of as a Riemannian metric), we obtain quotient (flat) Riemannian structures on both $\mathbb{R}^n/A(\mathbb{Z}^n)$ and $\mathbb{R}^n/\mathbb{Z}^n$, so that the covering map (3.18) is actually a local isometry. Therefore, the number of sheets of this covering map coincides with the ratio $\text{vol}(\mathbb{R}^n/A(\mathbb{Z}^n))/\text{vol}(\mathbb{R}^n/\mathbb{Z}^n)$. But we know that $\text{vol}(\mathbb{R}^n/\mathbb{Z}^n)$ is 1, being also the volume of the unit parallelepiped, and similarly $\text{vol}(\mathbb{R}^n/A(\mathbb{Z}^n))$ is the volume of the image of the unit parallelepiped under the action of A , which is therefore $|\det A|$. On the other hand, by covering space theory, we see that the number of sheets is equal to the order of the group of deck transformations of the covering p , which is in turn isomorphic to $\pi_1(\mathbb{R}^n/\mathbb{Z}^n)/p_*\pi_1(\mathbb{R}^n/A(\mathbb{Z}^n)) = \mathbb{Z}^n/A(\mathbb{Z}^n)$. ■

3.4 Rationality of the Bergman kernel of \mathcal{U}

For a domain $\Omega \subset \mathbb{C}^n$, denote by $\text{Rat}(\Omega)$ the rational functions of \mathbb{C}^n restricted to Ω . An element f of Ω is defined on $\Omega \setminus Z_f$, where Z_f is an affine algebraic variety in \mathbb{C}^n . Algebraically, $\text{Rat}(\Omega)$ is a field, and is isomorphic to the field of rational functions $\mathbb{C}(z_1, \dots, z_n)$ in n indeterminates. The map $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$ of (3.13) induces a mapping of fields

$$\Phi_A^* : \text{Rat}(\mathcal{U}) \rightarrow \text{Rat}\left(\mathbb{D}_{L(B)}^n\right)$$

given by $\Phi_A^*(f) = f \circ \Phi_A$. Denote by k the image

$$(3.19) \quad k = \Phi_A^*(\text{Rat}(\mathcal{U})) \subset \text{Rat}\left(\mathbb{D}_{L(B)}^n\right).$$

Then k is clearly a subfield of $\text{Rat}(\mathbb{D}_{L(B)}^n)$. The following lemma records a special case of a very general phenomenon:

Lemma 3.20 *The field extension $\text{Rat}(\mathbb{D}_{L(B)}^n)/k$ is Galois, and the map*

$$(3.21) \quad \Theta : \Gamma \rightarrow \text{Gal} \left(\text{Rat} \left(\mathbb{D}_{L(B)}^n \right) / k \right), \quad \Theta(\sigma)(f) = f \circ \sigma^{-1},$$

where $f \in \text{Rat}(\mathbb{D}_{L(B)}^n)$, $\sigma \in \Gamma$, is an isomorphism of the group of deck transformations of the proper map Φ_A with the Galois group of the extension.

Proof Denote by w_j the j th component of the map Φ_A , where $1 \leq j \leq n$. Notice that w_j can be identified with the function $\Phi_A^*(\pi_j) = \pi_j \circ \Phi_A$ where π_j is the j th coordinate function on \mathcal{U} . Since $\text{Rat}(\mathcal{U}) = \mathbb{C}(\pi_1, \dots, \pi_n)$, the field k is generated over \mathbb{C} by the functions w_1, \dots, w_n , i.e., $k = \mathbb{C}(w_1, \dots, w_n)$. Denoting the coordinate functions on $\mathbb{D}_{L(B)}^n$ by z_1, \dots, z_n , we see that $\text{Rat}(\mathbb{D}_{L(B)}^n) = k(z_1, \dots, z_n)$. To show that $\text{Rat}(\mathbb{D}_{L(B)}^n)/k$ is Galois, we need to show that each z_j is algebraic over k , and all the conjugates of z_j over k are already present in $\text{Rat}(\mathbb{D}_{L(B)}^n)$.

Notice that by definition for each $1 \leq j \leq n$, we have $w_j = z_1^{a_1^j} \dots z_n^{a_n^j} = z^{a^j}$, which follows from the fact that $w = \Phi_A(z) = z^A$. Therefore we have

$$z^{\det A \cdot I} = z^{\text{adj } A \cdot A} = (z^A)^{\text{adj } A} = w^{\text{adj } A}.$$

Denoting the j th row of the integer matrix $\text{adj } A$ by c^j we see that $z_j^{\det A} = w^{c^j} \in k$, so that z_j is a root of the polynomial $t^{\det A} - w^{c^j} \in k[t]$. Further, all the roots of this polynomial (which are of the form $\omega \cdot z_j$ for a $(\det A)$ th root of unity, $\omega \in \mathbb{C}$) are clearly in $\text{Rat}(\mathbb{D}_{L(B)}^n)$. It follows that $\text{Rat}(\mathbb{D}_{L(B)}^n)$ is Galois over the subfield k .

For each $\sigma \in \Gamma$, it is clear that $\Theta(\sigma)$ is an automorphism of the field $\text{Rat}(\mathbb{D}_{L(B)}^n)$. We need to show that it fixes k . But for $h = \Phi_A^* g \in k$, where $g \in \text{Rat}(\mathcal{U})$, we have

$$\Theta(\sigma)h = h \circ \sigma^{-1} = g \circ \Phi_A \circ \sigma^{-1} = g \circ \Phi_A = h,$$

since σ is a deck transformation of Φ_A . It follows that $\Theta(\sigma) \in \text{Gal}(\text{Rat}(\mathbb{D}_{L(B)}^n)/k)$. It is easily verified by direct computation that Θ is an injective group homomorphism. To show that Θ is surjective, for $\gamma \in \text{Gal}(\text{Rat}(\mathbb{D}_{L(B)}^n)/k)$, we find a deck transformation $\theta \in \Gamma$ such that $\gamma(f) = f \circ \theta$, so that $\gamma = \Theta(\theta^{-1})$. Let $\theta_j \in \text{Rat}(\mathbb{D}_{L(B)}^n)$ be the function $\theta_j = \gamma(z_j)$, where $z_j \in \text{Rat}(\mathbb{D}_{L(B)}^n)$ is the j th coordinate function. As we saw above, we have $\theta_j = \omega_j \cdot z_j$, where ω_j is a $(\det A)$ th root of unity. Then $\theta = (\theta_1, \dots, \theta_n)$ defines a diagonal unitary mapping of \mathbb{C}^n and therefore maps $\mathbb{D}_{L(B)}^n$ (a product of discs and punctured discs) into itself. Also, the j th component of $\Phi_A \circ \theta$ is given by

$$\theta^{a^j} = \theta_1^{a_1^j} \dots \theta_n^{a_n^j} = \gamma(z_1)^{a_1^j} \dots \gamma(z_n)^{a_n^j} = \gamma(z^{a^j}) = \gamma(w_j) = w_j,$$

which is precisely the j th component of Φ_A . Therefore, $\Phi_A \circ \theta = \Phi_A$, and $\theta \in \Gamma$. Finally, for a rational function $f = f(z_1, \dots, z_n) \in \text{Rat}(\mathbb{D}_{L(B)}^n)$, we have

$$\gamma(f) = f(\gamma(z_1), \dots, \gamma(z_n)) = f(\theta_1, \dots, \theta_n) = f \circ \theta.$$

■

Proposition 3.22 *The Bergman kernel of \mathcal{U} is a rational function.*

Proof Choosing a local continuous branch θ_j of $\arg w_j$ for each coordinate of $w \in \mathcal{U}$, we can write $w = \rho(w) \odot \exp(i\theta)$ for $\theta \in \mathbb{R}^n$. Then $\psi(w) = w^{A^{-1}} = \rho(w)^{A^{-1}} \odot \exp(iA^{-1}\theta)$ is a local inverse to the proper holomorphic map $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$. Since Φ_A is of the quotient type, it follows that the local branches of Φ_A^{-1} are $\{\sigma \circ \psi : \sigma \in \Gamma\}$. By Theorem 3.12, the map $\sigma : \mathbb{D}_{L(B)}^n \rightarrow \mathbb{D}_{L(B)}^n$ is the restriction of a unitary linear map on \mathbb{C}^n , so, noting that the argument behind the proof of (3.9) also applies for (locally defined) rational matrix powers, we have

$$\begin{aligned} \det(\sigma \circ \psi)'(w) &= \det \sigma' \cdot \det \psi'(w) = \det \sigma \cdot \det A^{-1} \cdot w^{\mathbb{1}A^{-1}-\mathbb{1}} \\ &= \det \sigma \cdot \det A^{-1} \cdot (w^{A^{-1}})^{\mathbb{1}} \cdot \frac{1}{w^{\mathbb{1}}} = \frac{\det \sigma \cdot \psi(w)^{\mathbb{1}}}{\det A \cdot w^{\mathbb{1}}}. \end{aligned}$$

Therefore, by the Bell transformation formula, for $z \in \mathbb{D}_{L(B)}^n$ and $w \in \mathcal{U}$, the Bergman kernels $K_{\mathcal{U}}$ and $K_{\mathbb{D}_{L(B)}^n} = K_{\mathbb{D}^n}$ are related by

$$\begin{aligned} \det \Phi_A'(z) \cdot K_{\mathcal{U}}(\Phi_A(z), w) &= \sum_{\sigma \in \Gamma} K_{\mathbb{D}^n}(z, \sigma \circ \psi(w)) \cdot \overline{\det(\sigma \circ \psi)'(w)} \\ &= \frac{1}{\det A \cdot w^{\mathbb{1}}} \cdot \sum_{\sigma \in \Gamma} K_{\mathbb{D}^n}(z, \sigma \circ \psi(w)) \cdot \overline{\det \sigma \cdot \psi(w)^{\mathbb{1}}}. \end{aligned}$$

For a fixed $z \in \mathbb{D}_{L(B)}^n$, consider the function L on $\mathbb{D}_{L(B)}^n$ given by

$$L(\zeta) = \sum_{\sigma \in \Gamma} K_{\mathbb{D}^n}(\bar{z}, \sigma(\zeta)) \cdot \det \sigma \cdot \zeta^{\mathbb{1}},$$

where $\bar{z} = (\bar{z}_1, \dots, \bar{z}_n) \in \mathbb{D}_{L(B)}^n$. Recalling that

$$(3.23) \quad K_{\mathbb{D}^n}(z, w) = \frac{1}{\pi^n} \prod_{j=1}^n \frac{1}{(1 - z_j \bar{w}_j)^2},$$

we see that $L \in \text{Rat}(\mathbb{D}_{L(B)}^n)$, and

$$K_{\mathcal{U}}(z^A, w) = \frac{1}{(\det A)^2 \cdot z^{\mathbb{1}A^{-1}} \cdot w^{\mathbb{1}}} \cdot \overline{L(\psi(w))}.$$

We now claim that $L \in k$, where $k \subset \text{Rat}(\mathbb{D}_{L(B)}^n)$ is as in (3.19). Since the extension $\text{Rat}(\mathbb{D}_{L(B)}^n)/k$ is Galois, it suffices to show that for each field-automorphism $\tau \in \text{Gal}(\text{Rat}(\mathbb{D}_{L(B)}^n)/k)$, we have $\tau(L) = L$. But by Lemma 3.20, for each $\tau \in \text{Gal}(\text{Rat}(\mathbb{D}_{L(B)}^n)/k)$ there is a $\theta \in \Gamma$ such that $\tau(L) = L \circ \theta$. Notice that thanks to (3.14),

we can think of the map $\theta : \mathbb{D}_{L(B)}^n \rightarrow \mathbb{D}_{L(B)}^n$ as a linear map of \mathbb{C}^n represented by a diagonal matrix. Therefore, we have $(\theta(\zeta))^{\mathbb{1}} = \det \theta \cdot \zeta^{\mathbb{1}}$. Consequently

$$\begin{aligned} \tau(L)(\zeta) &= L(\theta(\zeta)) = \sum_{\sigma \in \Gamma} K_{\mathbb{D}^n}(\bar{z}, \sigma(\theta(\zeta))) \cdot \det \sigma \cdot (\theta(\zeta))^{\mathbb{1}} \\ &= \sum_{\sigma \in \Gamma} K_{\mathbb{D}^n}(\bar{z}, (\sigma \circ \theta)(\zeta)) \cdot \det \sigma \cdot \det \theta \cdot \zeta^{\mathbb{1}} \\ &= \sum_{\sigma \in \Gamma} K_{\mathbb{D}^n}(\bar{z}, (\sigma \circ \theta)(\zeta)) \cdot \det(\sigma \circ \theta) \cdot \zeta^{\mathbb{1}} \\ &= L(\zeta), \end{aligned}$$

establishing the claim.

Now since $L \in k$, there is a function $R(z, \cdot) \in \text{Rat}(\mathcal{U})$ such that $L = \Phi_A^* R(z, \cdot)$, i.e.,

$$L(\zeta) = R(z, \Phi_A(\zeta)) = R(z, \zeta^A).$$

Therefore, we have

$$\begin{aligned} K_{\mathcal{U}}(z^A, w) &= \frac{1}{(\det A)^2 \cdot z^{\mathbb{1}A-1} \cdot \bar{w}^{\mathbb{1}}} \cdot \overline{R(z, \Phi_A(\psi(w)))} \\ &= \frac{1}{(\det A)^2 \cdot z^{\mathbb{1}A-1} \cdot \bar{w}^{\mathbb{1}}} \cdot \overline{R(z, w)}, \end{aligned}$$

which shows that for each fixed $z \in \mathbb{D}_{L(B)}^n$, the function $K_{\mathcal{U}}(z, \cdot)$ is rational. By the Reinhardt symmetry of $K_{\mathcal{U}}$, there is a function \tilde{K} such that

$$K_{\mathcal{U}}(z, w) = \tilde{K}(z_1 \bar{w}_1, \dots, z_n \bar{w}_n) = \tilde{K}(z \odot \bar{w}).$$

Since for a fixed z , the function $w \mapsto \tilde{K}(z \odot \bar{w})$ is rational, it follows that the function $t \mapsto \tilde{K}(t) = \tilde{K}(t_1, \dots, t_n)$ is also rational. Therefore, the function $K_{\mathcal{U}}$ is rational on $\mathcal{U} \times \mathcal{U}$. ■

4 Transformation of L^p -Bergman spaces under quotient maps

4.1 Definitions

In order to state our results, we introduce some terminology:

Definition 4.1 We say that a linear map T between Banach spaces $(E_1, \|\cdot\|_1)$ and $(E_2, \|\cdot\|_2)$ is a *homothetic isomorphism* if it is a continuous bijection (and therefore has a continuous inverse) and there is a constant $C > 0$ such that for each $x \in E_1$ we have

$$\|Tx\|_2 = C \|x\|_1.$$

Remark If T is a homothetic isomorphism between Banach spaces, then $\frac{1}{\|T\|} \cdot T$ is an isometric isomorphism of Banach spaces, so a homothetic isomorphism is simply the product of an isometric isomorphism and a scalar operator. In particular, a homothetic isomorphism between Hilbert spaces preserves angles, and in particular orthogonality of vectors.

The following definition and facts are standard:

Definition 4.2 Let $\Omega \subset \mathbb{C}^n$ be a domain, let $\lambda > 0$ be a continuous function (the weight), and let $0 < p < \infty$. Then we define

$$L^p(\Omega, \lambda) = \left\{ f : \Omega \rightarrow \mathbb{C} \text{ measurable} : \int_{\Omega} |f|^p \lambda \, dV < \infty \right\}$$

and

$$A^p(\Omega, \lambda) = \left\{ f : \Omega \rightarrow \mathbb{C} \text{ holomorphic} : \int_{\Omega} |f|^p \lambda \, dV < \infty \right\},$$

where the latter is called a *weighted Bergman space*. If $p \geq 1$, then each of $L^p(\Omega, \lambda)$ and $A^p(\Omega, \lambda)$ is a Banach space with the natural *weighted norm*, and $A^p(\Omega, \lambda)$ is a closed subspace of $L^p(\Omega, \lambda)$.

We will make extensive use of the following notion:

Definition 4.3 Given a group G of biholomorphic automorphisms of a domain $\Omega \subset \mathbb{C}^n$, and a space \mathfrak{F} of functions on Ω , we denote by $[\mathfrak{F}]^G$ the subspace of \mathfrak{F} consisting of functions which are G -invariant in the following sense

$$(4.4) \quad [\mathfrak{F}]^G = \{ f \in \mathfrak{F} : f = \sigma^\sharp(f) \text{ for all } \sigma \in G \},$$

where σ^\sharp is the pullback induced by σ as in (1.8).

Remark Interpreting \mathfrak{F} as a space of holomorphic forms on Ω by associating $f \in \mathfrak{F}$ with the form $f dz_1 \wedge \cdots \wedge dz_n$, this simply says that the forms in $[\mathfrak{F}]^G$ are invariant under pullback by elements of G .

4.2 Transformation of Bergman spaces

With the above definitions, we are ready to state and prove the following elementary facts. For completeness, we give details of the proofs.

Proposition 4.5 Let Ω_1, Ω_2 be domains in \mathbb{C}^n , and let $\Phi : \Omega_1 \rightarrow \Omega_2$ be a proper holomorphic map of quotient type with group $\Gamma \subset \text{Aut}(\Omega_1)$. Then for $1 < p < \infty$, the pullback map Φ^\sharp gives rise to a homothetic isomorphism

$$(4.6) \quad \Phi^\sharp : L^p(\Omega_2) \rightarrow [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma.$$

This restricts to a homothetic isomorphism

$$(4.7) \quad \Phi^\sharp : A^p(\Omega_2) \rightarrow [A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma.$$

Proof Let f be a function on Ω_2 , and let $g = \Phi^\sharp f$ be its pullback to Ω_1 , then we have for each $\sigma \in \Gamma$ that

$$\sigma^\sharp(g) = \sigma^\sharp(\Phi^\sharp f) = (\Phi \circ \sigma)^\sharp f = \Phi^\sharp f = g,$$

where we have used the contravariance of the pullback $(\Phi \circ \sigma)^\sharp = \sigma^\sharp \circ \Phi^\sharp$ and the fact that $\Phi \circ \sigma = \Phi$ which follows since the action of Γ on Ω_1 restricts to actions on each of the fibers. This shows that the range of Φ^\sharp consists of Γ -invariant functions. Special cases of this invariance were already noticed [MSRZ13, CKY20].

To complete the proof of (4.6) we must show that

(i) for each $f \in L^p(\Omega_2)$,

$$(4.8) \quad \|\Phi^\sharp f\|_{L^p(\Omega_1, |\det \Phi'|^{2-p})}^p = |\Gamma| \cdot \|f\|_{L^p(\Omega_2)}^p,$$

(ii) The image $\Phi^\sharp(L^p(\Omega_2))$ is precisely $[L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$.

Let Z_1, Z_2 be as in the Definition 3.10 of proper holomorphic maps of quotient type, i.e., Φ is a regular covering map from $\Omega_1 \setminus Z_1$ to $\Omega_2 \setminus Z_2$. Let U be an open set in $\Omega_2 \setminus Z_2$ which is evenly covered by Φ , and let V be an open set of $\Omega_1 \setminus Z_1$ which is mapped biholomorphically by Φ onto U . Then the inverse image $\Phi^{-1}(U)$ is the finite disjoint union $\bigcup_{\sigma \in \Gamma} \sigma V$. Therefore, if $f \in L^p(\Omega_2)$ is supported in U , then $\Phi^\sharp f$ is supported in $\bigcup_{\sigma \in \Gamma} \sigma V$, and we have

$$\begin{aligned} & \|\Phi^\sharp f\|_{L^p(\Omega_1, |\det \Phi'|^{2-p})}^p \\ &= \int_{\Omega_1} |f \circ \Phi \cdot \det \Phi'|^p |\det \Phi'|^{2-p} dV \\ &= \sum_{\sigma \in \Gamma} \int_{\sigma V} |f \circ \Phi|^p |\det \Phi'|^2 dV = \sum_{\sigma \in \Gamma} \int_U |f|^p dV = |\Gamma| \cdot \|f\|_{L^p(\Omega_2)}^p, \end{aligned}$$

where we have used the change of variables formula applied to the biholomorphic map $\Phi|_{\sigma V}$ along with the fact that the real Jacobian determinant of the map Φ is equal to $|\det \Phi'|^2$.

For a general $f \in L^p(\Omega_2)$, modify the proof as follows. There is clearly a collection of pairwise disjoint open sets $\{U_j\}_{j \in J}$ in $\Omega_2 \setminus Z_2$ such that each U_j is evenly covered by Φ and $\Omega_2 \setminus \bigcup_{j \in J} U_j$ has measure zero. Set $f_j = f \cdot \chi_j$, where χ_j is the indicator function of U_j , so that each $f_j \in L^p(\Omega_2)$ and $f = \sum_j f_j$. Also, the functions $\Phi^\sharp f_j$ have pairwise disjoint supports in Ω_1 . Therefore we have

$$\begin{aligned} \|\Phi^\sharp f\|_{L^p(\Omega_1, |\det \Phi'|^{2-p})}^p &= \sum_{j \in J} \|\Phi^\sharp f_j\|_{L^p(\Omega_1, |\det \Phi'|^{2-p})}^p \\ &= |\Gamma| \sum_{j \in J} \|f_j\|_{L^p(\Omega_2)}^p = |\Gamma| \|f\|_{L^p(\Omega_2)}^p. \end{aligned}$$

To complete the proof, we need to show that Φ^\sharp is surjective in both (4.6) and (4.7). Let $g \in [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$. Let $\{U_j\}_{j \in J}$ be as in the previous paragraph, and set $g_j = g \cdot \chi_{\Phi^{-1}(U_j)}$, where $\chi_{\Phi^{-1}(U_j)}$ is the indicator function of $\Phi^{-1}(U_j)$. Notice that $g_j \in [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$. Let $V_j \subset \Phi^{-1}(U_j)$ be such that Φ maps V_j biholomorphically to U_j , and let $\Psi : U_j \rightarrow V_j$ be the local inverse of Φ onto V_j . Define

$$(4.9) \quad f_j = \Psi^\sharp(g_j).$$

We claim that f_j is defined independently of the choice of V_j . Indeed, any other choice is of the form σV_j for some $\sigma \in \Gamma$ and the corresponding local inverse is $\sigma \circ \Psi$. But we have

$$(\sigma \circ \Psi)^\sharp g_j = \Psi^\sharp \circ \sigma^\sharp g_j = \Psi^\sharp g_j = f_j,$$

where we have used the fact that $\sigma^\sharp g_j = g_j$ since $g_j \in [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$.

Now we define $f = \sum_j f_j$. Notice that the f_j have pairwise disjoint support, and it is easily checked that $\Phi^\sharp f = g$. This establishes that (4.6) is a homothetic isomorphism.

It is clear that if f is holomorphic on Ω_2 , then $\Phi^\sharp f$ is holomorphic on Ω_1 , therefore, Φ^\sharp maps $A^p(\Omega_2)$ into $[A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$. Now, in the argument in the previous paragraph showing that the image $\Phi^\sharp(L^p(\Omega_2))$ is $[L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$, local definition of the inverse map (4.9) shows that if $g \in [A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$, then the f constructed by this procedure is holomorphic, and therefore lies in $A^p(\Omega_2)$. This completes the proof of the proposition. ■

4.3 Bell transformation law for quotient maps

The following is a refinement (for the class of proper holomorphic maps of quotient type) of a classic result of Bell (see [Bel81, Theorem 1], [Bel82, Equation 2.2]).

Proposition 4.10 *Let Ω_1, Ω_2 be domains in \mathbb{C}^n and let $\Phi : \Omega_1 \rightarrow \Omega_2$ be a proper holomorphic map of quotient type with group $\Gamma \subset \text{Aut}(\Omega_1)$. Then the following diagram commutes:*

$$(4.11) \quad \begin{array}{ccc} L^2(\Omega_2) & \xrightarrow[\cong]{\Phi^\sharp} & [L^2(\Omega_1)]^\Gamma \\ \downarrow \mathbf{B}_{\Omega_2} & & \downarrow \mathbf{B}_{\Omega_1} \\ A^2(\Omega_2) & \xrightarrow[\cong]{\Phi^\sharp} & [A^2(\Omega_1)]^\Gamma. \end{array}$$

In order to prove the proposition, we need the following simple lemma, which shows that the Bergman projection interacts well with the action of automorphisms:

Lemma 4.12 *Let $\Omega \subset \mathbb{C}^n$ be a domain, and let \mathbf{B}_Ω be its Bergman projection operator.*

(1) *If $\sigma \in \text{Aut}(\Omega)$ is a biholomorphic automorphism, then*

$$(4.13) \quad \mathbf{B}_\Omega \circ \sigma^\sharp = \sigma^\sharp \circ \mathbf{B}_\Omega.$$

(2) *If $G \subset \text{Aut}(\Omega)$ is a group of biholomorphic automorphisms, then \mathbf{B}_Ω restricts to the orthogonal projection operator from $[L^2(\Omega)]^G$ onto $[A^2(\Omega)]^G$.*

Proof For (1), note that σ^\sharp is a unitary operator on $L^2(\Omega)$ and \mathbf{B}_Ω is an orthogonal projection on $L^2(\Omega)$, therefore the unitarily similar operator $Q = \sigma^\sharp \circ \mathbf{B}_\Omega \circ (\sigma^\sharp)^{-1}$ is also an orthogonal projection. Since σ^\sharp (and therefore its inverse) leaves $A^2(\Omega)$ invariant, it follows that the range of Q is $A^2(\Omega)$. Therefore, $Q = \mathbf{B}_\Omega$.

For (2), let $f \in [L^2(\Omega)]^G$. Then using (4.13), we have for $\sigma \in G$

$$\sigma^\sharp(\mathbf{B}_\Omega f) = \mathbf{B}_\Omega(\sigma^\sharp(f)) = \mathbf{B}_\Omega f,$$

which shows that $\mathbf{B}_\Omega f \in [A^2(\Omega)]^G$, so that \mathbf{B}_Ω maps the G -invariant functions $[L^2(\Omega)]^G$ into the G -invariant holomorphic functions $[A^2(\Omega)]^G$. Since \mathbf{B}_Ω restricts to the identity on $[A^2(\Omega)]^G$, it follows that the range of \mathbf{B}_Ω is $[A^2(\Omega)]^G$. Observe that

$$\ker(\mathbf{B}_\Omega|_{[L^2(\Omega)]^G}) \subseteq \ker \mathbf{B}_\Omega = A^2(\Omega)^\perp \subseteq ([A^2(\Omega)]^G)^\perp,$$

which shows that kernel of the restriction of \mathbf{B}_Ω to $[L^2(\Omega)]^G$ is orthogonal to its range, and therefore an orthogonal projection. ■

Proof of Proposition 4.10 By Proposition 4.5, the Φ^\sharp represented by the top (resp. bottom) horizontal arrow is a homothetic isomorphism from the Hilbert space $L^2(\Omega_2)$ (resp. $A^2(\Omega_2)$) onto the Hilbert space $[L^2(\Omega_1)]^\Gamma$ (resp. $[A^2(\Omega_1)]^\Gamma$). Therefore, Φ^\sharp preserves angles and in particular orthogonality. Now consider the map $P : [L^2(\Omega_1)]^\Gamma \rightarrow [A^2(\Omega_1)]^\Gamma$ defined by

$$(4.14) \quad P = \Phi^\sharp \circ \mathbf{B}_{\Omega_2} \circ (\Phi^\sharp)^{-1},$$

which, being a composition of continuous linear maps, is a continuous linear mapping of Hilbert spaces. Notice that

$$P^2 = \Phi^\sharp \circ \mathbf{B}_{\Omega_2} \circ (\Phi^\sharp)^{-1} \circ \Phi^\sharp \circ \mathbf{B}_{\Omega_2} \circ (\Phi^\sharp)^{-1} = \Phi^\sharp \circ \mathbf{B}_{\Omega_2} \circ (\Phi^\sharp)^{-1} = P,$$

so P is a projection in $[L^2(\Omega_1)]^\Gamma$, with range contained in $[A^2(\Omega_1)]^\Gamma$. Since $(\Phi^\sharp)^{-1}$ and $\Phi^\sharp|_{A^2(\Omega_2)}$ are isomorphisms, and \mathbf{B}_{Ω_2} is surjective, it follows that P is a projection onto $[A^2(\Omega_1)]^\Gamma$. We claim that P is in fact the orthogonal projection onto $[A^2(\Omega_1)]^\Gamma$, i.e., the kernel of P is $([A^2(\Omega_1)]^\Gamma)^\perp$, the orthogonal complement of $[A^2(\Omega_1)]^\Gamma$ in $[L^2(\Omega_1)]^\Gamma$. Since in formula (4.14), the maps $(\Phi^\sharp)^{-1}$ and Φ^\sharp are isomorphisms, it follows that $f \in \ker P$ if and only if $(\Phi^\sharp)^{-1}f \in \ker \mathbf{B}_{\Omega_2}$. But $\ker \mathbf{B}_{\Omega_2} = A^2(\Omega_2)^\perp$, since the Bergman projection is orthogonal. It follows that $\ker P = \Phi^\sharp(A^2(\Omega_2)^\perp)$. Notice that Φ^\sharp , being a homothetic isomorphism of Hilbert spaces, preserves orthogonality, and maps $A^2(\Omega_2)$ to $[A^2(\Omega_1)]^\Gamma$ isomorphically, therefore, $\Phi^\sharp((A^2(\Omega_2)^\perp)^\perp) = ([A^2(\Omega_1)]^\Gamma)^\perp$, which establishes the claim.

Therefore, we have shown that $P = \Phi^\sharp \circ \mathbf{B}_{\Omega_2} \circ (\Phi^\sharp)^{-1}$ is the orthogonal projection from $[L^2(\Omega_1)]^\Gamma$ to the subspace $[A^2(\Omega_1)]^\Gamma$. To complete the proof, we only need to show that the restriction of the Bergman projection \mathbf{B}_{Ω_1} to the Γ -invariant subspace $[L^2(\Omega_1)]^\Gamma$ is also the orthogonal projection from $[L^2(\Omega_1)]^\Gamma$ onto $[A^2(\Omega_1)]^\Gamma$. But this follows from Lemma 4.12 above.

Thus $P = \mathbf{B}_{\Omega_1}|_{[L^2(\Omega_1)]^\Gamma}$, and the commutativity of (4.11) follows. ■

4.4 Transformation of the Bergman projection in L^p -spaces

The following result will be our main tool on studying L^p -regularity of the Bergman projection in the domain \mathcal{U} :

Theorem 4.15 *Let Ω_1, Ω_2 be bounded domains in \mathbb{C}^n , let $\Phi : \Omega_1 \rightarrow \Omega_2$ be a proper holomorphic map of quotient type with group $\Gamma \subset \text{Aut}(\Omega_1)$. Let $p \geq 1$. The following two assertions are equivalent:*

(1) The Bergman projection \mathbf{B}_{Ω_2} gives rise to a bounded operator mapping

$$L^p(\Omega_2) \rightarrow A^p(\Omega_2).$$

(2) The Bergman projection \mathbf{B}_{Ω_1} gives rise to a bounded operator mapping

$$[L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma \rightarrow [A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma.$$

If one of the conditions (1) or (2) holds (and therefore both hold), then the following diagram commutes, where \mathbf{B}_{Ω_j} , $j = 1, 2$ denote the extension by continuity of the Bergman projections:

$$(4.16) \quad \begin{array}{ccc} L^p(\Omega_2) & \xrightarrow[\cong]{\Phi^\sharp} & [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma \\ \downarrow \mathbf{B}_{\Omega_2} & & \downarrow \mathbf{B}_{\Omega_1} \\ A^p(\Omega_2) & \xrightarrow[\cong]{\Phi^\sharp} & [A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma \end{array}$$

Remark Statement (1) in Theorem 4.15 means the following: the restriction of the Bergman projection to a dense subspace of $L^p(\Omega_2)$ given by

$$\mathbf{B}_{\Omega_2} : L^2(\Omega_2) \cap L^p(\Omega_2) \rightarrow A^2(\Omega_2)$$

is bounded in the L^p -norm, i.e., there is a $C > 0$ such that for all $f \in L^2(\Omega_2) \cap L^p(\Omega_2)$,

$$\|\mathbf{B}_{\Omega_2} f\|_{L^p(\Omega_2)} \leq C \|f\|_{L^p(\Omega_2)}.$$

By continuity \mathbf{B}_{Ω_2} extends to a bounded linear operator from $L^p(\Omega_2)$ to $A^p(\Omega_2)$.

Similarly, Statement (2) means the following: the restriction of the Bergman projection to the dense subspace of $[L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$ given by

$$\mathbf{B}_{\Omega_1} : [L^2(\Omega_1)]^\Gamma \cap [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma \rightarrow A^2(\Omega_1)$$

is bounded in the $L^p(\Omega_1, |\det \Phi'|^{2-p})$ -norm, i.e., there is a $C > 0$ such that for all $f \in [L^2(\Omega_1)]^\Gamma \cap [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma$,

$$\|\mathbf{B}_{\Omega_1} f\|_{L^p(\Omega_1, |\det \Phi'|^{2-p})} \leq C \|f\|_{L^p(\Omega_1, |\det \Phi'|^{2-p})}.$$

We now see by Lemma 4.12 that

$$\mathbf{B}_{\Omega_1} \left([L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma \right) \subseteq [A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma,$$

where we have used continuity to extend the operator.

Proof Proposition 4.5 says that Φ^\sharp is a homothetic isomorphism, mapping

$$L^p(\Omega_2) \rightarrow [L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma,$$

and that it restricts to a homothetic isomorphism on the holomorphic subspaces. Similarly, $(\Phi^\sharp)^{-1}$ has the same properties with the domains and ranges switched.

First assume Statement (2). From the diagram (4.11), we write

$$(4.17) \quad \mathbf{B}_{\Omega_2} = (\Phi^\sharp)^{-1} \circ \mathbf{B}_{\Omega_1} \circ \Phi^\sharp.$$

By hypothesis, \mathbf{B}_{Ω_1} is a bounded linear operator mapping

$$[L^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma \rightarrow [A^p(\Omega_1, |\det \Phi'|^{2-p})]^\Gamma.$$

Consequently, this composition maps $L^p(\Omega_2)$ boundedly into $A^p(\Omega_2)$, giving statement (1). A similar argument shows that (1) implies (2).

For the commutativity of the diagram, rewrite (4.17) and see that on the subspace $L^p(\Omega_2) \cap L^2(\Omega_2)$ of $L^p(\Omega_2)$ we have the relation

$$(4.18) \quad \Phi^\sharp \circ \mathbf{B}_{\Omega_2} = \mathbf{B}_{\Omega_1} \circ \Phi^\sharp.$$

Using the hypothesis (for the \mathbf{B}_{Ω_j}) and Proposition 4.5 (for Φ^\sharp), we see that each of the four maps in the diagram (4.16) extends to the respective domain in that diagram and is continuous. By continuity, (4.18) continues to hold for the extended maps. This shows that the diagram (4.16) is commutative. ■

Remark Diagram (4.11) is a special case of diagram (4.16) for $p = 2$.

5 The unboundedness of the Bergman projection on \mathcal{U}

Using Proposition 5.1 and the results of Section 4, we will prove in this section the following partial form of Theorem 1.2:

Proposition 5.1 *The Bergman projection is not bounded in $L^p(\mathcal{U})$, provided*

$$(5.2) \quad p \geq \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) - 1},$$

where $\kappa(\mathcal{U}) \in \mathbb{N}$ is the complexity of the domain \mathcal{U} , as defined in Section 1.1.

5.1 Reinhardt domains

Recall some elementary facts about holomorphic function theory on Reinhardt domains (which are always assumed to be centered at the origin). Let $\Omega \subset \mathbb{C}^n$ be a Reinhardt domain. Every holomorphic function $f \in \mathcal{O}(\Omega)$ admits a unique *Laurent expansion*

$$(5.3) \quad f = \sum_{\alpha \in (\mathbb{Z}^n)^\dagger} a_\alpha(f) \varphi_\alpha,$$

where for $\alpha \in (\mathbb{Z}^n)^\dagger$, $\varphi_\alpha(z)$ is the Laurent monomial z^α as in (2.4), and where $a_\alpha(f) \in \mathbb{C}$ is the α th Laurent coefficient. The Laurent series of f converges absolutely and uniformly to f on every compact subset of Ω .

When f lies in the Bergman space $A^2(\Omega)$, we can say more about the series (5.3): it is actually an orthogonal series converging in the Hilbert space $A^2(\Omega)$, and the family of monomials

$$(5.4) \quad \left\{ \frac{\varphi_\alpha}{\|\varphi_\alpha\|_{L^2}} : \varphi_\alpha \in L^2(\Omega) \right\}$$

forms an orthonormal basis of $A^2(\Omega)$. In particular, if $f \in A^2(\Omega)$ then the Laurent series (5.3) can have $a_\alpha(f) \neq 0$ only when $\varphi_\alpha \in L^2(\Omega)$. It is possible to generalize some of these results to the spaces $A^p(\Omega)$; see [CEM19].

5.2 A criterion for unboundedness of the Bergman projection

We now give an easily checkable condition which shows L^p -Bergman unboundedness on any Reinhardt domain.

Lemma 5.5 *Let Ω be a bounded Reinhardt domain in \mathbb{C}^n , and let $p \geq 2$. Suppose that there is a multi-index $\beta \in (\mathbb{Z}^n)^\dagger$ such that*

$$(5.6) \quad \varphi_\beta \in L^2(\Omega) \setminus L^p(\Omega).$$

Then the Bergman projection \mathbf{B}_Ω fails to map $L^p(\Omega) \rightarrow L^p(\Omega)$.

Proof Define subsets $\mathcal{J}_\beta, \mathcal{K}_\beta \subset \{1, 2, \dots, n\}$ with

$$\mathcal{J}_\beta = \{j : \beta_j \geq 0\}, \quad \mathcal{K}_\beta = \{k : \beta_k < 0\},$$

and let

$$f(w) = \prod_{j \in \mathcal{J}_\beta} w_j^{\beta_j} \times \prod_{k \in \mathcal{K}_\beta} (\overline{w_k})^{-\beta_k}.$$

Then f is a bounded function on Ω , and therefore $f \in L^p(\Omega)$. We now show that $\mathbf{B}_\Omega f = C\varphi_\beta$ for some $C \neq 0$. Since $\varphi_\beta \notin L^p(\Omega)$, this will show that \mathbf{B}_Ω fails to map the element $f \in L^p(\Omega)$ to a function in $L^p(\Omega)$. This will imply that \mathbf{B}_Ω is not bounded in the L^p -norm, since if it were so, it would extend to a map from the dense subspace $L^p(\Omega) \cap L^2(\Omega)$ to the whole of $L^p(\Omega)$.

Let $\gamma = (|\beta_1|, \dots, |\beta_n|) \in \mathbb{N}^n$ be the multi-index obtained by replacing each entry of β by its absolute value. Write the polar form of w as $w = \rho(w) \odot \exp(i\theta)$ for a $\theta \in \mathbb{R}^n$. Then $f(w) = \rho(w)^\gamma e^{i\beta\theta}$, and for $\alpha \in (\mathbb{Z}^n)^\dagger$, we have

$$\varphi_\alpha(w) = w^\alpha = (\rho(w) \odot \exp(i\theta))^\alpha = \rho(w)^\alpha e^{i\alpha\theta}.$$

Further, denote by $|\Omega| = \{\rho(z) : z \in \Omega\} \subset \mathbb{R}^n$ the Reinhardt shadow of Ω , and let \mathbb{T}^n be the unit torus of n dimensions. Then, for each $\alpha \in \mathbb{Z}^n$, we have

$$\begin{aligned} \langle \mathbf{B}_\Omega f, \varphi_\alpha \rangle_{A^2(\Omega)} &= \langle f, \varphi_\alpha \rangle_{L^2(\Omega)} = \int_\Omega f \overline{\varphi_\alpha} dV \\ &= \int_\Omega r^\gamma e^{i\beta\theta} \cdot r^\alpha e^{-i\alpha\theta} dV = \int_{|\Omega|} r^{\gamma+\alpha} r^{\mathbb{1}} dr \times \int_{\mathbb{T}^n} e^{i((\beta-\alpha)\cdot\theta)} d\theta \\ &\quad \times \begin{cases} = 0 & \text{if } \alpha \neq \beta \\ > 0 & \text{if } \alpha = \beta. \end{cases} \end{aligned}$$

Since (5.4) is an orthonormal basis of $A^2(\Omega)$ it follows that all the Fourier coefficients of $\mathbf{B}_\Omega f$ with respect this basis vanish, except the β th coefficient, which is nonzero. Therefore, $\mathbf{B}_\Omega f = C\varphi_\beta \notin L^p(\Omega)$, for some constant $C \neq 0$. ■

5.3 Preliminaries

In this section (and the following Section 6), we will use the following default notation and conventions:

- (1) $B \in M_n(\mathbb{Z})$ is a matrix such that the domain \mathcal{U} is represented as in (3.1),
- (2) B satisfies the properties (3.3), which is not a loss of generality by Proposition 3.2.
- (3) We have $A = \text{adj } B$.

Observe that then by Theorem 3.12, the monomial map Φ_A is a proper holomorphic map of quotient type.

We note the following computation:

Proposition 5.7 *The upper bound in (1.3) is given by*

$$(5.8) \quad \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) - 1} = \min_j \frac{2 \cdot \mathbb{1} \cdot a_j}{\mathbb{1} \cdot a_j - \text{gcd}(a_j)},$$

where, as usual, a_j is the j th column of the matrix A , where notation is as above.

Proof Recall the definition of the projective height function as in (1.4). Then we have, by projective invariance, for each $1 \leq j \leq n$:

$$h(B^{-1}e_j) = h(\det B \cdot B^{-1}e_j) = h_j(\text{adj } B \cdot e_j) = h(Ae_j) = h(a_j).$$

Since $A > 0$, it follows that a_j is a vector of non-negative integers, and the vector $\frac{1}{\text{gcd}(a_j)}a_j$ is such that its entries are coprime non-negative integers. Therefore,

$$h(B^{-1}e_j) = h(a_j) = h\left(\frac{1}{\text{gcd}(a_j)}a_j\right) = \sum_{k=1}^n \frac{|a_j^k|}{\text{gcd}(a_j)} = \frac{\mathbb{1} \cdot a_j}{\text{gcd}(a_j)}.$$

So

$$\kappa(\mathcal{U}) = \max_j h(B^{-1}e_j) = \max_j \frac{\mathbb{1} \cdot a_j}{\text{gcd}(a_j)}.$$

Since the function $x \mapsto \frac{2x}{x-1}$ is strictly decreasing for $x > 1$, we have

$$\frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) - 1} = \frac{2 \cdot \max_j \frac{\mathbb{1} \cdot a_j}{\text{gcd}(a_j)}}{\max_j \frac{\mathbb{1} \cdot a_j}{\text{gcd}(a_j)} - 1} = \min_j \frac{\frac{2 \cdot \mathbb{1} \cdot a_j}{\text{gcd}(a_j)}}{\frac{\mathbb{1} \cdot a_j}{\text{gcd}(a_j)} - 1} = \min_j \frac{2 \cdot \mathbb{1} \cdot a_j}{\mathbb{1} \cdot a_j - \text{gcd}(a_j)}.$$

■

5.4 p -allowable multi-indices

Let $\beta \in (\mathbb{Z}^n)^\dagger$ be a multi-index. We say that β is p -allowable on a Reinhardt domain Ω if we have that the monomial $\varphi_\beta \in L^p(\Omega)$, i.e.,

$$\int_\Omega |\varphi_\beta|^p dV < \infty.$$

We denote the collection of p -allowable multi-indices on Ω by $\mathcal{S}_p(\Omega)$. We first compute the p -allowable multi-indices on the domain \mathcal{U} . Recall that the conventions introduced in Section 5.3 are in force.

Proposition 5.9 *Let $\beta \in (\mathbb{Z}^n)^\dagger$ and $p > 0$. Then $\beta \in \mathcal{S}_p(\mathcal{U})$ if and only if*

$$(5.10) \quad (p \cdot \beta + 2 \cdot \mathbb{1})A > 0.$$

Proof By Theorem 3.12, $\Phi_A : \mathbb{D}_{L(B)}^n \rightarrow \mathcal{U}$ is a proper holomorphic map of quotient type with $\det A$ sheets. Therefore, by Proposition 4.5 (in particular (4.8)) we see that

$$\begin{aligned} \int_{\mathcal{U}} |\varphi_\beta|^p dV &= \frac{1}{|\Gamma|} \int_{\mathbb{D}_{L(B)}^n} |\Phi_A^\#(\varphi_\beta)| dV \\ &= \frac{1}{\det A} \int_{\mathbb{D}_{L(B)}^n} |\varphi_\beta \circ \Phi_A|^p |\det \Phi_A'|^2 dV \\ &= \frac{1}{\det A} \int_{\mathbb{D}_{L(B)}^n} |z^{\beta \cdot A}|^p |\det A \cdot z^{\mathbb{1} \cdot A - \mathbb{1}}|^2 dV(z) \end{aligned}$$

using (2.7f) and (3.9)

$$\begin{aligned} &= (2\pi)^n \det A \int_{(0,1)^n} r^{p \cdot \beta A} \cdot r^{2 \cdot \mathbb{1} \cdot A - 2 \cdot \mathbb{1}} \cdot r^{\mathbb{1}} dr \\ &= (2\pi)^n \det A \int_{(0,1)^n} r^{p \cdot \beta A + 2 \cdot \mathbb{1} \cdot A - \mathbb{1}} dr \\ &= (2\pi)^n \det A \prod_{j=1}^n \int_0^1 r_j^{p \cdot \beta a_j + 2 \cdot \mathbb{1} \cdot a_j - 1} dr_j, \end{aligned}$$

where $a_j \in \mathbb{Z}^n$ is the j th column of A . It is clear that $\int_{\mathcal{U}} |\varphi_\beta|^p dV < \infty$ if and only if $p \cdot \beta a_j + 2 \cdot \mathbb{1} a_j > 0$ for each $1 \leq j \leq n$. This completes the proof. ■

Remark The above proposition can be seen as a special case of [Zwo00, Lemma 2.2.1].

We now consider the important case $p = 2$, so that $\mathcal{S}_2(\Omega)$ corresponds to the monomials in the Bergman space. For a matrix $A \in M_n(\mathbb{Z})$, none of whose columns are zero, denote by $\gcd(a_j)$ the greatest common divisor of the entries in the j th column of A . We then let

$$(5.11) \quad g(A) = (\gcd(a_1), \dots, \gcd(a_n)) \in (\mathbb{Z}^n)^\dagger$$

be the integer row vector whose j th entry is the greatest common divisor of the j th column of A .

Proposition 5.12

(1) *Let $\beta \in (\mathbb{Z}^n)^\dagger$. Then $\beta \in \mathcal{S}_2(\mathcal{U})$ if and only if*

$$(5.13) \quad (\beta + \mathbb{1})A \geq g(A).$$

(2) *For $1 \leq j \leq n$ let Π_j be the integer hypersurface determined by equality in the j th entry of (5.13), that is*

$$(5.14) \quad \Pi_j = \left\{ \beta \in (\mathbb{Z}^n)^\dagger : (\beta + \mathbb{1})a_j = \sum_{k=1}^n (\beta_k + 1)a_j^k = \gcd(a_j) \right\}.$$

Then we have

$$\Pi_j \cap \mathcal{S}_2(\mathcal{U}) \neq \emptyset.$$

The following Lemma will be needed in the proof of part (2) of the proposition.

Lemma 5.15 *Let P be an $m \times n$ integer matrix of rank m for positive integers $n \geq m$, and let $q \in (\mathbb{Z}^n)^\dagger$. Then there is an $x \in (\mathbb{Z}^m)^\dagger$ such that $xP \geq q$.*

Proof Let $\mathcal{O} = \{x \in (\mathbb{R}^m)^\dagger : xP \geq q\}$. Then \mathcal{O} is an unbounded convex set, so at least one of the coordinates x_1, \dots, x_m is unbounded on \mathcal{O} . Rename the coordinates so that x_1 is unbounded. It follows that the projection

$$\{x_1 \in \mathbb{R} : (x_1, \dots, x_m) \in \mathcal{O}\}$$

of \mathcal{O} on the coordinate axis x_1 is an unbounded convex set, and therefore a ray or the whole of \mathbb{R} . For an integer N let $C_N = \{x_1 = N\} \cap \mathcal{O}$, which is naturally thought of as a subset of $(\mathbb{R}^{m-1})^\dagger$. Therefore, either there is an N_1 such that C_N is nonempty if $N \geq N_1$ or there is an N_2 such that C_N is nonempty if $N \leq N_2$. Assuming the former, we see that the sets C_N are convex subsets of $(\mathbb{R}^{m-1})^\dagger$ and similar to each other, i.e., they are dilations of the same set. As the size of each C_N becomes infinite as $N \rightarrow \infty$, for large N , the set C_N contains cubes of arbitrarily large size, where a cube is a product of intervals of the same size in each coordinate. As soon as C_N contains a cube of side $(1 + \varepsilon)$ for some $\varepsilon > 0$, we see that there is a point $M \in (\mathbb{Z}^{m-1})^\dagger$ that belongs to C_N . It follows that the point $(N, M) \in \mathbb{Z} \times (\mathbb{Z}^{m-1})^\dagger$ belongs to \mathcal{O} . ■

Proof of Proposition 5.12 (1) If $p = 2$, the condition (5.10) becomes $2(\beta + \mathbb{1})A > 0$, which is equivalent to

$$(5.16) \quad (\beta + \mathbb{1})A > 0.$$

Now the j th entry of the row vector on the left of the above equation is given by $(\beta + \mathbb{1})a_j = \sum_{k=1}^n (\beta_k + 1)a_j^k$, which is a positive integer divisible by $\gcd(a_j) = \gcd(a_j^1, \dots, a_j^n)$. It follows that (5.16) holds if and only if

$$(\beta + \mathbb{1})a_j \geq \gcd(a_j),$$

which is precisely the content of (5.13).

(2) Fix $1 \leq j \leq n$. By the Euclidean algorithm, $\Pi_j \neq \emptyset$. Choose $y \in \Pi_j$. Define a \mathbb{Z} -module homomorphism $\phi : (\mathbb{Z}^n)^\dagger \rightarrow \mathbb{Z}$ by setting $\phi(x) = xa_j$, i.e.,

$$\phi(x_1, \dots, x_n) = \sum_{k=1}^n x_k a_j^k.$$

We then see that $\Pi_j = y + \ker \phi$. Since \mathbb{Z} is a principal ideal domain, $\ker \phi$ is a free \mathbb{Z} -submodule of $(\mathbb{Z}^n)^\dagger$ of rank $\leq n$ (see [DF04, Theorem 4, Chapter 12 (p. 460)]). Moreover as ϕ is surjective, the quotient \mathbb{Z} -module $(\mathbb{Z}^n)^\dagger / \ker \phi$ is isomorphic to \mathbb{Z} . It can be seen, by tensoring with \mathbb{Q} for example, that the rank of $\ker \phi$ is $n - 1$. Let D be an $(n - 1) \times n$ integer matrix whose rows are a \mathbb{Z} -basis of $\ker \phi$. Then the map $f : (\mathbb{Z}^{n-1})^\dagger \rightarrow (\mathbb{Z}^n)^\dagger$ given by $f(t) = y + tD$ is a parametrization of Π_j , i.e., it is one-to-one and its range is precisely Π_j . To complete the proof of the result, it is sufficient

to show that $f^{-1}(\mathcal{S}_2(\mathcal{U}))$ is a nonempty subset of $(\mathbb{Z}^{n-1})^\dagger$. Notice now that an integer vector $t \in f^{-1}(\mathcal{S}_2(\mathcal{U}))$, i.e., $f(t) \in \mathcal{S}_2(\mathcal{U}) \cap \Pi_j$ if and only if

$$(y + tD + \mathbb{1})A \geq g(A), \quad \text{i.e.,} \quad tDA \geq g(A) - (y + \mathbb{1})A.$$

By Lemma 5.15, there is an integer vector $t \in (\mathbb{Z}^{n-1})^\dagger$ that satisfies the above system of inequalities. This concludes the proof of part (2) ■

5.5 Proof of Proposition 5.1

Recall that p satisfies (5.2). Now by (5.8),

$$p \geq \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) - 1} = \min_j \frac{2 \cdot \mathbb{1} \cdot a_j}{\mathbb{1} \cdot a_j - \gcd(a_j)},$$

so that there is a J with $1 \leq J \leq n$ such that

$$(5.17) \quad p \geq \frac{2 \cdot \mathbb{1} \cdot a_J}{\mathbb{1} \cdot a_J - \gcd(a_J)}.$$

By part 2 of Proposition 5.12, there is an $\beta \in (\mathbb{Z}^n)^\dagger$ which lies in $\mathcal{S}_2(\mathcal{U}) \cap \Pi_J$. Such a β satisfies:

$$(5.18) \quad (\beta + \mathbb{1})a_J = \gcd(a_J),$$

and also

$$(5.19) \quad (\beta + \mathbb{1})A \geq g(A),$$

with $g(A)$ as in (5.11). By construction, $\beta \in \mathcal{S}_2(\mathcal{U})$. We now claim that $\beta \notin \mathcal{S}_p(\mathcal{U})$. By Lemma 5.5 this shows that the Bergman projection is not bounded in $L^p(\mathcal{U})$. To establish the claim we note that the J th entry of the row vector $(p \cdot \beta + 2 \cdot \mathbb{1})A$ is

$$(5.20) \quad \begin{aligned} (p \cdot \beta + 2 \cdot \mathbb{1})a_J &= p \cdot (\beta_k + \mathbb{1}) \cdot a_J + (2 - p) \cdot \mathbb{1} \cdot a_J \\ &= p \cdot \gcd(a_J) + (2 - p) \cdot \mathbb{1} \cdot a_J \\ &= p \cdot (\gcd(a_J) - \mathbb{1} \cdot a_J) + 2 \cdot \mathbb{1} \cdot a_J, \end{aligned}$$

where in the second line we have used (5.18). Thanks to the inequality (5.17) it follows that the quantity in (5.20) is not positive. It follows by Proposition 5.9 that β is not in $L^p(\mathcal{U})$, which establishes the claim and completes the proof.

6 Boundedness of the Bergman projection

In this section, we obtain the following part of Theorem 1.2:

Proposition 6.1 *Let*

$$(6.2) \quad 2 \leq p < \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U}) - 1},$$

then the Bergman projection is bounded on $L^p(\mathcal{U})$.

We begin by recalling the following fact, which will be the main “hard analysis” ingredient of the proof:

Proposition 6.3 *The Bergman projection on the polydisc \mathbb{D}^n gives rise to a bounded operator $B_{\mathbb{D}^n} : L^p(\mathbb{D}^n) \rightarrow A^p(\mathbb{D}^n)$ for all $1 < p < \infty$.*

Proof For the polydisc \mathbb{D}^n , the Bergman projection has the well-known integral representation

$$B_{\mathbb{D}^n} f(z) = \int_{\mathbb{D}^n} K(z, w) f(w) dV(w), \quad f \in L^2(\mathbb{D}^n),$$

where K is the Bergman kernel of the polydisc, which is easily shown to be given by the well-known formula (3.23).

The case $n = 1$ of Proposition 6.3 is by now a staple result in Bergman theory, going back to [ZJ64], where it was proved using the L^p -boundedness of a Calderón–Zygmund singular integral operator. Another approach, based on Schur’s test for L^p -boundedness of an integral operator, was used in [FR75]. An alternative proof of the main estimate needed in this method can be found in [Ax188] and in the monograph [DS04].

Since \mathbb{D}^n is a product domain, the theorem in higher dimensions follows from a textbook application of Fubini’s theorem to the case $n = 1$. ■

6.1 Two lemmas

The following two simple lemmas will be used to deduce monomially weighted estimates starting from Proposition 6.3:

Lemma 6.4 *Let $1 \leq p < \infty$, let $n \geq 1$, and let $\gamma \in (\mathbb{R}^n)^+$ be such that $\gamma > -2 \cdot \mathbb{1}$. Then there is a $C > 0$ such that for any $f \in A^p(\mathbb{D}^n)$ we have*

$$(6.5) \quad \int_{\mathbb{D}^n} |f|^p \rho^\gamma dV \leq C \int_{\mathbb{D}^n} |f|^p dV,$$

where as usual, $\rho(z)^\gamma = \prod_{j=1}^n |z_j|^{\gamma_j}$.

Proof Throughout this proof, C will denote a constant that depends only on p and γ . The actual value of C may change from line to line.

Proceed by induction on the dimension n . First consider the base case $n = 1$. We have, by the Bergman inequality (cf. [DS04, Theorem 1]) that there is a $C > 0$ such that

$$\sup_{|z| < \frac{1}{2}} |f(z)| \leq C \|f\|_{L^p(\mathbb{D})}$$

for all $f \in A^p(\mathbb{D})$. Therefore, for $f \in A^p(\mathbb{D})$ we have the estimate

$$(6.6) \quad \int_{|z| < \frac{1}{2}} |f(z)|^p |z|^\gamma dV(z) \leq \sup_{|z| < \frac{1}{2}} |f(z)|^p \cdot \int_{|z| < \frac{1}{2}} |z|^\gamma dV(z) < C \cdot \|f\|_{L^p(\mathbb{D})}^p,$$

where we have used the fact that since $\gamma > -2$ we have

$$\int_{|z| < \frac{1}{2}} |z|^\gamma dV(z) = 2\pi \int_0^{\frac{1}{2}} r^{\gamma+1} dr < \infty.$$

On the other hand,

$$(6.7) \quad \int_{\frac{1}{2} \leq |z| < 1} |f(z)|^p |z|^\gamma dV(z) \leq C \|f\|_{L^p(\mathbb{D})}^p,$$

where we have used the fact that $\sup_{\frac{1}{2} \leq |z| < 1} |z|^\gamma < \infty$.

Adding (6.6) and (6.7), the estimate (6.5) follows in the case $n = 1$.

For the general case, assume the result established in $n - 1$ dimensions. Write the coordinates of \mathbb{C}^n as $z = (z', z_n) \in \mathbb{C}^{n-1} \times \mathbb{C}$, and $\gamma = (\gamma', \gamma_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. Then, using Fubini's theorem

$$\begin{aligned} \int_{\mathbb{D}^n} |f|^p \rho^\gamma dV &= \int_{\mathbb{D}^{n-1}} \rho(z')^{\gamma'} \left(\int_{\mathbb{D}} |f(z', z_n)|^p |z_n|^{\gamma_n} dV(z_n) \right) dV(z') \\ &\leq C \int_{\mathbb{D}^{n-1}} \rho(z')^{\gamma'} \left(\int_{\mathbb{D}} |f(z', z_n)|^p dV(z_n) \right) dV(z') \\ &\leq C \int_{\mathbb{D}} \left(\int_{\mathbb{D}^{n-1}} |f(z', z_n)|^p \rho(z')^{\gamma'} dV(z') \right) dV(z_n) \\ &\leq C \int_{\mathbb{D}} \left(\int_{\mathbb{D}^{n-1}} |f(z', z_n)|^p dV(z') \right) dV(z_n) \\ &= C \int_{\mathbb{D}^n} |f(z', z_n)|^p dV(z', z_n), \end{aligned}$$

which proves the result. ■

Lemma 6.8 *Let $1 \leq p < \infty$, let $n \geq 1$ and let $\lambda \in \mathbb{N}^n$ be a multi-index of non-negative integers. Then there is a $C > 0$ such that for all $f \in A^p(\mathbb{D}^n)$ we have the estimate:*

$$(6.9) \quad \int_{\mathbb{D}^n} |f|^p dV \leq C \int_{\mathbb{D}^n} |\varphi_\lambda f|^p dV,$$

where $\varphi_\lambda(z) = z^\lambda$ is as in (2.4).

Proof We need only to prove the case in which $\lambda = (1, 0, \dots, 0)$, so that $\varphi_\lambda(z) = z_1$. Once this special case is established, the general result follows by repeatedly applying it and permuting the coordinates.

In what follows, C will denote some positive constant that depends only on p and λ , where the actual value of C may change from line to line. First, consider the one dimensional case, so that we have to show that for a holomorphic function f on the disc we have

$$\int_{\mathbb{D}} |f(z)|^p dV(z) \leq C \int_{\mathbb{D}} |zf(z)|^p dV(z),$$

where the left hand side is assumed to be finite (and therefore the right hand side is finite.) First note that we obviously have

$$(6.10) \quad \int_{\frac{1}{2} \leq |z| < 1} |f(z)|^p dV(z) \leq 2^p \int_{\frac{1}{2} \leq |z| < 1} |zf(z)|^p dV(z).$$

On the other hand, if $|z| = \frac{1}{2}$, we have

$$\begin{aligned} |f(z)|^p &= 2^p |zf(z)|^p \\ &\leq 2^p \sup_{|w| \leq \frac{1}{2}} |wf(w)|^p && \text{(maximum principle)} \\ &\leq C \int_{\mathbb{D}} |wf(w)|^p dV(w) && \text{(Bergman's inequality)} \end{aligned}$$

The maximum principle now implies

$$\sup_{|z| \leq \frac{1}{2}} |f(z)|^p \leq C \int_{\mathbb{D}} |zf(z)|^p dV(z),$$

so that we have

$$(6.11) \quad \int_{|z| \leq \frac{1}{2}} |f(z)|^p dV(z) \leq C \int_{\mathbb{D}} |zf(z)|^p dV(z).$$

Combining (6.10) and (6.11) the result follows for $n = 1$.

For the higher-dimensional case, denote the coordinates of \mathbb{C}^n as (z_1, z') where $z' = (z_2, \dots, z_n)$. Then for $f \in A^p(\mathbb{D}^n)$ we have

$$\begin{aligned} \int_{\mathbb{D}^n} |f(z_1, z')|^p dV(z_1, z') &= \int_{\mathbb{D}^{n-1}} \left(\int_{\mathbb{D}} |f(z_1, z')|^p dV(z_1) \right) dV(z') \\ &\leq C \int_{\mathbb{D}^{n-1}} \left(\int_{\mathbb{D}} |z_1 f(z_1, z')|^p dV(z_1) \right) dV(z') \\ &= C \int_{\mathbb{D}^n} |z_1 f(z_1, z')|^p dV(z_1, z'). \end{aligned}$$

■

6.2 Determination of Γ -invariant subspaces

In this section and Section 6.3, we use the notation established in Section 5.3 above, so that B and A have the same meaning as there. The group Γ is as in Theorem 3.12: the deck transformation group associated to the map Φ_A . We will now determine the Γ -invariant subspaces of holomorphic functions, in the sense of (4.4):

Proposition 6.12 *Let $\alpha \in (\mathbb{Z}^n)^\dagger$. Then the monomial $\varphi_\alpha(z) = z^\alpha$ belongs to the space $[\mathcal{O}((\mathbb{D}^*)^n)]^\Gamma$ of Γ -invariant holomorphic functions on $(\mathbb{D}^*)^n$ if and only if there is a $\beta \in (\mathbb{Z}^n)^\dagger$ such that*

$$\alpha = \beta A - \mathbb{1}.$$

Proof Recall that, by definition, the monomial φ_α is invariant under the group action of Γ if and only if $\sigma_v^\sharp(\varphi_\alpha) = (\varphi_\alpha \circ \sigma_v) \det \sigma'_v = \varphi_\alpha$ for all $\sigma_v \in \Gamma$, where for $v \in \mathbb{Z}^n$, the automorphism $\sigma_v \in \Gamma$ is as in (3.14). Denote the rows of A^{-1} by c^1, \dots, c^n . By the linearity of σ_v

$$\det \sigma'_v = \det \sigma_v = \prod_{j=1}^n e^{2\pi i c^j v} = e^{2\pi i \sum_{j=1}^n c^j v} = e^{2\pi i (\mathbb{1} A^{-1}) v},$$

for $v \in \mathbb{Z}^n$. Also, for $z \in (\mathbb{D}^*)^n$ and $v \in \mathbb{Z}^n$,

$$\varphi_\alpha \circ \sigma_v(z) = \varphi_\alpha(\exp(2\pi i A^{-1}v) \odot z) = (\exp(2\pi i A^{-1}v) \odot z)^\alpha = e^{2\pi i \alpha A^{-1}v} z^\alpha.$$

Therefore,

(6.13)

$$\sigma_v^\sharp(\varphi_\alpha)(z) = e^{2\pi i \alpha A^{-1}v} \cdot z^\alpha \cdot e^{2\pi i \mathbb{1} A^{-1}v} = e^{2\pi i (\alpha + \mathbb{1}) A^{-1}v} \cdot z^\alpha = e^{2\pi i (\alpha + \mathbb{1}) A^{-1}v} \cdot \varphi_\alpha(z).$$

Hence, $\varphi_\alpha \in [\mathcal{O}(\mathbb{D}_{\mathbb{L}(B)}^n)]^\Gamma$ if and only if $\sigma_v^\sharp(\varphi_\alpha) = \varphi_\alpha$ for all $v \in \mathbb{Z}^n$, i.e.,

$$e^{2\pi i (\alpha + \mathbb{1}) A^{-1}v} \cdot z^\alpha = z^\alpha$$

for all $z \in (\mathbb{D}^*)^n$ and $v \in \mathbb{Z}^n$, i.e., if and only if $(\alpha + \mathbb{1}) A^{-1}v \in \mathbb{Z}$ for all $v \in \mathbb{Z}^n$.

Now if there is $\beta \in (\mathbb{Z}^n)^\dagger$ such that $\alpha = \beta A - \mathbb{1}$, then clearly, $(\alpha + \mathbb{1}) A^{-1}v = \beta v \in \mathbb{Z}$ for all $v \in \mathbb{Z}^n$. Conversely, assume that $(\alpha + \mathbb{1}) A^{-1}v \in \mathbb{Z}$ for all $v \in \mathbb{Z}^n$ and let e_1, \dots, e_n be the standard basis of \mathbb{Z}^n . Then the j th column of $(\alpha + \mathbb{1}) A^{-1} = (\alpha + \mathbb{1}) A^{-1} I$ (where I is the $n \times n$ identity matrix) is $(\alpha + \mathbb{1}) A^{-1} e_j$ which is therefore in \mathbb{Z} . Therefore, we have $(\alpha + \mathbb{1}) A^{-1} = \beta \in (\mathbb{Z}^n)^\dagger$. It follows that $\alpha = \beta A - \mathbb{1}$ as desired. ■

Corollary 6.14 *Let $f \in [\mathcal{O}(\mathbb{D}^n)]^\Gamma$. Then there is an $h \in \mathcal{O}(\mathbb{D}^n)$ such that*

$$f(z) = z^{g(A)-\mathbb{1}} \cdot h(z),$$

with $g(A)$ as in (5.11).

Proof Let $f(z) = \sum_{\alpha \geq 0} a_\alpha z^\alpha$, be the Taylor expansion of f . If f is Γ invariant, then we claim that for each α such that $a_\alpha \neq 0$, we have that z^α is Γ -invariant. Indeed using (6.13) we have

$$f = \sigma_v^\sharp f = \sigma_v^\sharp \left(\sum_{\alpha \geq 0} a_\alpha \varphi_\alpha \right) = \sum_{\alpha \geq 0} a_\alpha e^{2\pi i (\alpha A^{-1} + \mathbb{1})v} \cdot \varphi_\alpha,$$

comparing this with the Taylor expansion of f and equating coefficients the claim follows. Therefore, the Taylor expansion of f is of the form

$$(6.15) \quad f(z) = \sum_{\substack{\beta \in (\mathbb{Z}^n)^\dagger \\ \beta A \geq \mathbb{1}}} a_{\beta A - \mathbb{1}} z^{\beta A - \mathbb{1}}.$$

Notice that $z^{\beta A - \mathbb{1}} = \prod_{j=1}^n z_j^{\beta a_j - 1}$. The integer $\beta a_j = \sum_{k=1}^n \beta_k a_j^k$ which occurs in the exponent is divisible by $\gcd(a_j)$. Further since $\sum_{k=1}^n \beta_k a_j^k \geq 1$ it follows that $\beta a_j \geq \gcd(a_j)$. It follows that the monomial $z^{\beta A - \mathbb{1}}$ is of the form $z^{g(A)-\mathbb{1}} \cdot z^\gamma$, where $\gamma \geq 0$. The corollary follows from (6.15). ■

6.3 Proof of Proposition 6.1

By Theorem 3.12, $\Phi_A : \mathbb{D}_{\mathbb{L}(B)}^n \rightarrow \mathcal{U}$ is a proper holomorphic map of quotient type with group Γ . Therefore, thanks to Theorem 4.15, the Bergman projection

$$B_{\mathcal{U}} : L^p(\mathcal{U}) \rightarrow L^p(\mathcal{U})$$

is bounded if and only if

$$(6.16) \quad \mathbf{B}_{\mathbb{D}_{L(B)}^n} : [L^p(\mathbb{D}_{L(B)}^n, |\det \Phi'_A|^{2-p})]^\Gamma \rightarrow [A^p(\mathbb{D}_{L(B)}^n, |\det \Phi'_A|^{2-p})]^\Gamma$$

is bounded. We therefore show that the map (6.16) is bounded if $2 \leq p < \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})-1}$.

To do this we express the map of (6.16) as a composition of three maps, and show that each of these three maps is continuous:

- (1) We first show that we have an inclusion of Banach spaces

$$[L^p(\mathbb{D}_{L(B)}^n, |\det \Phi'_A|^{2-p})]^\Gamma \subseteq [L^p(\mathbb{D}_{L(B)}^n)]^\Gamma,$$

and that the inclusion map

$$(6.17) \quad \iota : [L^p(\mathbb{D}^n, |\det \Phi'_A|^{2-p})]^\Gamma \hookrightarrow [L^p(\mathbb{D}^n)]^\Gamma$$

is bounded.

It is clearly sufficient to show that there is a continuous inclusion of the full space $L^p(\mathbb{D}_{L(B)}^n, |\det \Phi'_A|^{2-p})$ into $L^p(\mathbb{D}_{L(B)}^n)$. For $z \in \mathbb{D}^n$ we have for $p \geq 2$

$$|\det \Phi'_A(z)|^{2-p} = |\det Az^{1-A-1}|^{2-p} = |\det A|^{2-p} \rho(z)^{(1 \cdot A-1)(2-p)} \geq |\det A|^{2-p},$$

where we use the fact that the exponent of $\rho(z)$ is nonpositive, since A has non-negative integer entries and $p \geq 2$. Thus for any $p \geq 2$ and any $f \in L^p(\mathbb{D}_{L(B)}^n, |\det \Phi'_A|^{2-p})$ we have

$$(6.18) \quad \int_{\mathbb{D}_{L(B)}^n} |f|^p dV \leq \frac{1}{|\det A|^{2-p}} \int_{\mathbb{D}_{L(B)}^n} |f|^p |\det \Phi'_A|^{2-p} dV,$$

which proves the inclusion and its continuity.

- (2) Let $P_\Gamma = \mathbf{B}_{L(B)}^n \Big|_{[L^p(\mathbb{D}_{L(B)}^n)]^\Gamma}$ be the restriction of the Bergman projection operator to the Γ -invariant functions. We claim that the operator of Banach spaces

$$(6.19) \quad P_\Gamma : [L^p(\mathbb{D}_{L(B)}^n)]^\Gamma \rightarrow [A^p(\mathbb{D}_{L(B)}^n)]^\Gamma$$

is bounded. Indeed, we have that $L^p(\mathbb{D}_{L(B)}^n) = L^p(\mathbb{D}^n)$ (since $\mathbb{D}_{L(B)}^n$ is obtained from \mathbb{D}^n by removing an analytic set Z , which is of measure zero) and $A^p(\mathbb{D}_{L(B)}^n) = A^p(\mathbb{D}^n)$, in the sense that the analytic set Z is a removable singularity of functions integrable in the p th power, $p \geq 2$ (see [Bel82, p. 687]). By Proposition 6.3, the Bergman projection maps $L^p(\mathbb{D}^n)$ to $A^p(\mathbb{D}^n)$ boundedly. Finally, by part (2) of Lemma 4.12, the Bergman projection maps Γ -invariant functions to Γ -invariant functions. The boundedness of P_Γ follows.

- (3) Finally, we show that there is an inclusion $[A^p(\mathbb{D}_{L(B)}^n)]^\Gamma \subseteq [A^p(\mathbb{D}_{L(B)}^n, |\det \Phi'_A|^{2-p})]^\Gamma$, and the inclusion map so determined is bounded. As noted above $A^p(\mathbb{D}_{L(B)}^n) = A^p(\mathbb{D}^n)$, so it will suffice to show that there is a continuous inclusion

$$(6.20) \quad j : [A^p(\mathbb{D}^n)]^\Gamma \hookrightarrow [A^p(\mathbb{D}^n, |\det \Phi'_A|^{2-p})]^\Gamma.$$

Let $f \in [A^p(\mathbb{D}^n)]^\Gamma$, so by Corollary 6.14, there exists $h \in \mathcal{O}(\mathbb{D}^n)$ such that

$$(6.21) \quad f(z) = z^{g(A)-1} h(z).$$

In fact, $h \in A^p(\mathbb{D}^n)$, since

$$\int_{\mathbb{D}^n} |h(z)|^p dV = \int_{\rho(z) \leq \frac{1}{2}} |h(z)|^p dV + \int_{\frac{1}{2} < \rho(z) < 1} |h(z)|^p dV,$$

where the first of the two integrals is clearly finite and the second integral is

$$= \int_{\frac{1}{2} < \rho(z) < 1} \left| \frac{f(z)}{z^{g(A)-1}} \right|^p dV \leq \left(\prod_{j=1}^n 2^{\gcd(a_j)-1} \right)^p \cdot \int_{\frac{1}{2} < \rho(z) < 1} |f(z)|^p dV < \infty.$$

Now we have that

$$\begin{aligned} \|f\|_{A^p(\mathbb{D}^n, |\det \Phi'_A|^{2-p})}^p &= \int_{\mathbb{D}^n} |f(z)|^p |\det \Phi'_A(z)|^{2-p} dV(z) \\ &= \int_{\mathbb{D}^n} |h(z)|^p |z^{g(A)-1}|^p |\det A \cdot z^{1 \cdot A-1}|^{2-p} dV(z) \\ &\quad \text{using (6.21) and (3.9)} \\ &= |\det A|^{2-p} \cdot \int_{\mathbb{D}^n} |h(z)|^p |z^{p(g(A)-1)+(2-p)(1 \cdot A-1)}| dV(z) \\ (6.22) \quad &= |\det A|^{2-p} \cdot \int_{\mathbb{D}^n} |h|^p \rho^{p(g(A)-1 \cdot A)+2 \cdot 1 \cdot A-2 \cdot 1} dV \end{aligned}$$

Combining Hypotheses (6.2) and (5.8),

$$p < \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})-1} = \min_j \frac{2 \cdot 1 \cdot a_j}{1 \cdot a_j - \gcd(a_j)},$$

so for each $1 \leq j \leq n$,

$$p < \frac{2 \cdot 1 \cdot a_j}{1 \cdot a_j - \gcd(a_j)}.$$

Therefore, the j th component of the exponent of ρ in (6.22) is

$$\begin{aligned} p(\gcd(a_j) - 1 a_j) + 2 \cdot 1 \cdot a_j - 2 &> \left(\frac{2 \cdot 1 \cdot a_j}{1 \cdot a_j - \gcd(a_j)} \right) \\ &\quad \times (\gcd(a_j) - 1 \cdot a_j) + 2 \cdot 1 a_j - 2 = -2, \end{aligned}$$

where we have used the obvious fact that $\gcd(a_j) - 1 \cdot a_j \leq 0$. Therefore, we have

$$p(g(A) - 1A) + 2 \cdot 1A - 2 \cdot 1 > -2 \cdot 1,$$

and so we have, for constants C_1, C_2 independent of the function f :

$$\begin{aligned} (6.22) &\leq C_1 \int_{\mathbb{D}^n} |h|^p dV && \text{by Lemma (6.4)} \\ &\leq C_2 \int_{\mathbb{D}^n} |z^{g(A)-1} \cdot h(z)|^p dV(z) && \text{by Lemma (6.8)} \\ &= C_2 \int_{\mathbb{D}^n} |f|^p dV. \end{aligned}$$

It follows that the inclusion j of (6.20) is continuous.

Therefore, the map $\mathbf{B}_{\mathbb{D}_{L(B)}^n}$ of Banach spaces in (6.16) can be represented as a composition

$$\mathbf{B}_{\mathbb{D}_{L(B)}^n} = j \circ P_\Gamma \circ \iota,$$

where ι , P_Γ and j are as in (6.17), (6.19) and (6.20), respectively and each of which has already been shown to be continuous. The proof of Proposition 6.1 is complete.

7 Conclusion

7.1 End of proof of Theorem 1.2

Recall that, as a consequence of the self-adjointness of the Bergman projection on L^2 , the set of p for which the Bergman projection on a domain is L^p -bounded is Hölder-symmetric (see [EM16, CZ16]), i.e., p belongs to this set if and only if the conjugate index p' also belongs to it, where $\frac{1}{p'} + \frac{1}{p} = 1$. Notice now that the index conjugate to $\frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})-1}$ is $\frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})+1}$. Now by combining Propositions 6.1 and 5.1, we see that for $p \geq 2$, the Bergman projection on \mathcal{U} is bounded if and only if $p \in \left[2, \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})-1}\right)$. By Hölder symmetry, for $p \leq 2$, the Bergman projection is bounded if and only if $p \in \left(\frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})+1}, 2\right]$. The bounds claimed in (1.3) are proved, and therefore Theorem 1.2 holds.

7.2 Comments and questions

The methods used to prove Theorem 1.2 are more general than the result itself, and apply to the Bergman projection on various quotient domains of simple domains with known Bergman kernels, provided the quotient type proper holomorphic map is a monomial map. For example, we may deduce using a modification of our arguments, the range of p for which the Bergman projection is bounded in L^p on the domain

$$\left\{ \left(|z_1|^2 + \dots + |z_{n-1}|^2 \right)^{\frac{k_1}{2}} < |z_n|^{k_2} < 1 \right\} \subset \mathbb{C}^n,$$

where k_1, k_2 are positive integers.

Thanks to Theorem 1.2, the Bergman projection is no longer bounded in $L^p(\mathcal{U})$ if $p \geq \frac{2\kappa(\mathcal{U})}{\kappa(\mathcal{U})-1}$. It is natural to ask if there is an alternate projection from $L^p(\mathcal{U})$ to $A^p(\mathcal{U})$ for such p . In the special case $\mathcal{U} = H_{m/n}$, the generalized Hartogs triangle of (1.6), it is possible to construct for each $p \geq 2$ a *sub-Bergman projection* which gives rise to a bounded projection on L^p . It would be interesting to see whether a similar statement holds for the monomial polyhedra considered in this paper.

Finally, we would like to understand the precise geometric significance of the arithmetic complexity of \mathcal{U} without reference to the representation in terms of the matrix B . Such a description will pave the way of generalizing the results of this paper to wider classes of domains.

Acknowledgments We would like to thank Yuan Yuan for very interesting discussions with DC and LE about this problem and the results of [CKY20] during the 2019 Midwest Several Complex Variables Conference at Dearborn, MI, and Steven Krantz for comments on the same paper made to LE during a visit to Washington University at St. Louis, MO, in the same year. DC would like to thank the mathematics department of the University of Michigan, Ann Arbor for its hospitality in Fall 2019. We also thank Shuo Zhang for pointing out his work in [Zha20, Zha19], and Włodzimierz Zwonek for pointing out his work in [Zwo99, Zwo00].

References

- [Axl88] S. Axler, *Bergman spaces and their operators*. In: Surveys of some recent results in operator theory, Vol. I, Pitman Res. Notes Math. Ser., 171, Longman Sci. Tech., Harlow, 1988, pp. 1–50.
- [BŞ12] D. Barrett and S. Şahutoğlu, *Irregularity of the Bergman projection on worm domains in \mathbb{C}^n* . Michigan Math. J. 61(2012), no. 1, 187–198.
- [Bar84] D. E. Barrett, *Irregularity of the Bergman projection on a smooth bounded domain in \mathbb{C}^2* . Ann. Math. (2) 119(1984), no. 2, 431–436.
- [Beh33] H. Behnke, *Zur Theorie der Singularitäten der Funktionen mehrerer komplexen Veränderlichen*. Math. Ann. 108(1933), no. 1, 91–104.
- [BC82] S. Bell and D. Catlin, *Boundary regularity of proper holomorphic mappings*. Duke Math. J. 49(1982), no. 2, 385–396.
- [Bel81] S. R. Bell, *Proper holomorphic mappings and the Bergman projection*. Duke Math. J. 48(1981), no. 1, 167–175.
- [Bel82] S. R. Bell, *The Bergman kernel function and proper holomorphic mappings*. Trans. Amer. Math. Soc. 270(1982), no. 2, 685–691.
- [Cat80] D. Catlin, *Boundary behavior of holomorphic functions on pseudoconvex domains*. J. Differential Geom. 15(1980/1981), no. 4, 605–625.
- [Cha19] D. Chakrabarti, *On an observation of Sibony*. Proc. Amer. Math. Soc. 147(2019), no. 8, 3451–3454.
- [CEM19] D. Chakrabarti, L. D. Edholm, and J. D. McNeal, *Duality and approximation of Bergman spaces*. Adv. Math. 341(2019), 616–656.
- [CKMM20] D. Chakrabarti, A. Konkel, M. Mainkar, and E. Miller, *Bergman kernels of elementary Reinhardt domains*. Pacific J. Math. 306(2020), no. 1, 67–93.
- [CZ16] D. Chakrabarti and Y. E. Zeytuncu, *L^p mapping properties of the Bergman projection on the Hartogs triangle*. Proc. Amer. Math. Soc. 144(2016), no. 4, 1643–1653.
- [Che17] L. Chen, *The L^p boundedness of the Bergman projection for a class of bounded Hartogs domains*. J. Math. Anal. Appl. 448(2017), no. 1, 598–610.
- [CJY20] L. Chen, M. Jin, and Y. Yuan, *Bergman projection on the symmetrized bidisk*. Preprint, 2020. arXiv:2004.02785.
- [CKY20] L. Chen, S. G. Krantz, and Y. Yuan, *L^p regularity of the Bergman projection on domains covered by the polydisc*. J. Funct. Anal. 279(2020), no. 2, 108522.
- [Col66] L. Collatz, *Functional analysis and numerical mathematics*. Academic Press, New York, London, 1966. Translated from the German by Hansjörg Oser.
- [DF82] K. Diederich and J. E. Fornæss, *Boundary regularity of proper holomorphic mappings*. Invent. Math. 67(1982), no. 3, 363–384.
- [DF04] D. S. Dummit and R. M. Foote, *Abstract algebra*. 3rd ed., John Wiley & Sons, Inc., Hoboken, NJ, 2004.
- [DS04] P. Duren and A. Schuster, *Bergman spaces*. Mathematical Surveys and Monographs, 100, American Mathematical Society, Providence, RI, 2004.
- [Edh16] L. D. Edholm, *Bergman theory of certain generalized Hartogs triangles*. Pacific J. Math. 284(2016), no. 2, 327–342.
- [EM16] L. D. Edholm and J. D. McNeal, *The Bergman projection on fat Hartogs triangles: L^p boundedness*. Proc. Amer. Math. Soc. 144(2016), no. 5, 2185–2196.
- [EM17] L. D. Edholm and J. D. McNeal, *Bergman subspaces and subkernels: degenerate L^p mapping and zeroes*. J. Geom. Anal. 27(2017), no. 4, 2658–2683.

- [EM20] L. D. Edholm and J. D. McNeal, *Sobolev mapping of some holomorphic projections*. *J. Geom. Anal.* 30(2020), no. 2, 1293–1311.
- [FR75] F. Forelli and W. Rudin, *Projections on spaces of holomorphic functions in balls*. *Indiana Univ. Math. J.*, 24(1974/1975), 593–602.
- [Grü03] B. Grünbaum, *Convex polytopes*. 2nd ed., Graduate Texts in Mathematics, 221, Springer-Verlag, New York, 2003. Prepared and with a preface by Volker Kaibel, Victor Klee and Günter M. Ziegler.
- [HS80] M. Hakim and N. Sibony, *Spectre de $A(\bar{\Omega})$ pour des domaines bornés faiblement pseudoconvexes réguliers*. *J. Funct. Anal.* 37(1980), no. 2, 127–135.
- [Hed02] H. Hedenmalm, *The dual of a Bergman space on simply connected domains*. *J. Anal. Math.* 88(2002), 311–335.
- [HKZ00] H. Hedenmalm, B. Korenblum, and K. Zhu, *Theory of Bergman spaces*. Graduate Texts in Mathematics, 199, Springer-Verlag, New York, 2000.
- [HS00] M. Hindry and J. H. Silverman, *Diophantine geometry*. Graduate Texts in Mathematics, 201, Springer-Verlag, New York, 2000. An introduction.
- [Huo18] Z. Huo, *L^p estimates for the Bergman projection on some Reinhardt domains*. *Proc. Amer. Math. Soc.* 146(2018), no. 6, 2541–2553.
- [HW19] Z. Huo and B. D. Wick, *Weak-type estimates for the Bergman projection on the polydisc and the Hartogs triangle*. Preprint, 2019. arXiv:1909.05902.
- [JP08] M. Jarnicki and P. Pflug, *First steps in several complex variables: Reinhardt domains*. EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2008.
- [Kob59] S. Kobayashi, *Geometry of bounded domains*. *Trans. Amer. Math. Soc.* 92(1959), 267–290.
- [Kra13] S. G. Krantz, *Geometric analysis of the Bergman kernel and metric*. Graduate Texts in Mathematics, 268, Springer, New York, 2013.
- [KP08] S. G. Krantz and M. M. Peloso, *Analysis and geometry on worm domains*. *J. Geom. Anal.* 18(2008), no. 2, 478–510.
- [Mas91] W. S. Massey, *A basic course in algebraic topology*. Graduate Texts in Mathematics, 127, Springer-Verlag, New York, 1991.
- [MS94] J. D. McNeal and E. M. Stein, *Mapping properties of the Bergman projection on convex domains of finite type*. *Duke Math. J.* 73(1994), no. 1, 177–199.
- [MSRZ13] G. Misra, S. S. Roy, and G. Zhang, *Reproducing kernel for a class of weighted Bergman spaces on the symmetrized polydisc*. *Proc. Amer. Math. Soc.* 141(2013), no. 7, 2361–2370.
- [Mun84] J. R. Munkres, *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- [NP09] A. Nagel and M. Pramanik, *Maximal averages over linear and monomial polyhedra*. *Duke Math. J.* 149(2009), no. 2, 209–277.
- [NP20] A. Nagel and M. Pramanik, *Bergman spaces under maps of monomial type*. *J. Geometr. Anal.* 31(2021), no. 5, 4531–4560.
- [Par18] J.-D. Park, *The explicit forms and zeros of the Bergman kernel for 3-dimensional Hartogs triangles*. *J. Math. Anal. Appl.* 460(2018), no. 2, 954–975.
- [PS77] D. H. Phong and E. M. Stein, *Estimates for the Bergman and Szegő projections on strongly pseudo-convex domains*. *Duke Math. J.* 44(1977), no. 3, 695–704.
- [Sib75] N. Sibony, *Prolongement des fonctions holomorphes bornées et métrique de Carathéodory*. *Invent. Math.* 29(1975), no. 3, 205–230.
- [Vla66] V. S. Vladimirov, *Methods of the theory of functions of many complex variables*. The M.I.T. Press, Cambridge, MA, London, 1966. Translated from the Russian by Scripta Technica, Inc. Translation edited by Leon Ehrenpreis.
- [ZJ64] V. P. Zaharjuta and V. I. Judovic, *The general form of a linear functional on H_p^1* . *Uspekhi Mat. Nauk.* 19(1964), no. 2, 139–142.
- [Zha19] S. Zhang, *Mapping properties of the Bergman projections on elementary Reinhardt domains*. Submitted, 2019.
- [Zha20] S. Zhang, *L^p boundedness for the Bergman projections over n -dimensional generalized Hartogs triangles*. *Complex Var. Elliptic Equ.* (2020). <https://doi.org/10.1080/17476933.2020.1769085>.
- [Zie95] G. M. Ziegler, *Lectures on polytopes*. Graduate Texts in Mathematics, 152, Springer-Verlag, New York, 1995.
- [Zwo99] W. Zwonek, *On hyperbolicity of pseudoconvex Reinhardt domains*. *Arch. Math. (Basel)* 72(1999), no. 4, 304–314.
- [Zwo00] W. Zwonek, *Completeness, Reinhardt domains and the method of complex geodesics in the theory of invariant functions*. *Dissertationes Math. (Rozprawy Mat.)* 388(2000), 103.

Department of Mathematics, Central Michigan University, Mt Pleasant, MI 48859, USA

e-mail: bende1cc@cmich.edu chakr2d@cmich.edu maink1m@cmich.edu

URL: <http://people.cst.cmich.edu/chakr2d>

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA

e-mail: edholm@umich.edu