

## Existence, uniqueness and stability of positive solutions to sublinear elliptic systems

**Jann-Long Chern and Yong-Li Tang**

Department of Mathematics, National Central University,  
Chung-Li 32001, Taiwan  
(chern@math.ncu.edu.tw; tangyl@math.ncu.edu.tw)

**Chang-Shou Lin**

Department of Mathematics, Taida Institute for Mathematical Sciences,  
National Taiwan University, Taipei 10617, Taiwan  
(cslin@math.ntu.edu.tw)

**Junping Shi**

Department of Mathematics, College of William and Mary,  
Williamsburg, VA 23187-8795, USA (shij@math.wm.edu)

(MS received 20 July 2009; accepted 19 March 2010)

The existence, stability and uniqueness of positive solutions to a semilinear elliptic system with sublinear nonlinearities are proved. It is shown that the precise global bifurcation diagram of the positive solutions is a monotone curve with different asymptotical behaviour according to the form of the nonlinearities. Equations with Hölder continuous nonlinearities are also considered.

### 1. Introduction

Reaction–diffusion systems are used to model many chemical and biological phenomena in the natural world [25, 26], and systems of coupled partial differential equations are also used in other physical models such as nonlinear Schrödinger systems in multi-component Bose–Einstein condensates and nonlinear optics [19, 23]. The steady-state solutions or standing-wave solutions of such systems of nonlinear partial differential equations satisfy a nonlinear elliptic system with more than one equation. Much effort has been devoted to the existence of solutions of such systems (see, for example, [4, 8, 10, 11, 13, 15–17, 24, 30, 33, 34]), but it is usually difficult to determine whether or not the solution is unique.

We consider the positive solutions of a semilinear elliptic system of the form

$$\left. \begin{aligned} \Delta u + \lambda f(u, v) &= 0, & x \in \Omega, \\ \Delta v + \lambda g(u, v) &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \quad (1.1)$$

where  $\lambda > 0$  and  $\Omega$  is a bounded smooth domain. Here  $f$  and  $g$  are real-valued functions defined on  $\mathbb{R}_+^2 := (0, \infty) \times (0, \infty)$  which satisfy

(A1)  $f, g \in C^\alpha(\overline{\mathbb{R}_+^2}) \cap C^1(\mathbb{R}_+^2)$  for  $\alpha \in (0, 1)$ ,

(A2) (cooperativeness) define the Jacobian of the vector field  $(f, g)$  as

$$J(u, v) = \begin{pmatrix} \frac{\partial f}{\partial u}(u, v) & \frac{\partial f}{\partial v}(u, v) \\ \frac{\partial g}{\partial u}(u, v) & \frac{\partial g}{\partial v}(u, v) \end{pmatrix} \equiv \begin{pmatrix} f_u(u, v) & f_v(u, v) \\ g_u(u, v) & g_v(u, v) \end{pmatrix}. \quad (1.2)$$

Then  $f_v(u, v) \geq 0$  and  $g_u(u, v) \geq 0$  for  $(u, v) \in \mathbb{R}_+^2$ .

We consider the existence, uniqueness and stability of positive solutions to (1.1). From assumption (A1), we look for positive solutions  $(u, v) \in C_0^{2,\alpha}(\bar{\Omega})$ . The stability of a solution is determined by the following eigenvalue problem:

$$\left. \begin{aligned} \Delta \xi + \lambda f_u(u, v)\xi + \lambda f_v(u, v)\eta &= -\mu\xi, & x \in \Omega, \\ \Delta \eta + \lambda g_u(u, v)\xi + \lambda g_v(u, v)\eta &= -\mu\eta, & x \in \Omega, \\ \xi(x) = \eta(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \quad (1.3)$$

From the maximum principle of cooperative elliptic systems (see lemma 2.1 and [40]), equation (1.3) has a real principal eigenvalue  $\mu_1(u, v)$ , which has the smallest real part among all spectrum points. A solution  $(u, v)$  is *stable* if  $\mu_1(u, v) > 0$  and it is *unstable* otherwise.

For a scalar semilinear elliptic equation

$$\Delta u + \lambda f(u) = 0, \quad x \in \Omega, \quad u(x) = 0, \quad x \in \partial\Omega, \quad (1.4)$$

the stability of a solution can be defined in a similar fashion. If the nonlinear function is sublinear, that is, if  $f(u)/u$  is a non-increasing function, then it is well known that a positive solution of (1.4) is stable and one can prove that the solution must be unique with various methods. Brezis and Kamin [2] presented several different proofs for the uniqueness (see also [20, 27, 28, 35, 36]).

Here we use the stability defined above to prove that under some general sublinear conditions similar to the one for scalar equations, the positive solution of (1.1) is stable. Then one can use bifurcation theory to prove the existence and uniqueness of the positive solution. We also obtain the precise global bifurcation diagrams of the system in  $(\lambda, u, v)$  space under these conditions. In all cases that we consider, the bifurcation diagram is a single monotone solution curve (see §§ 3 and 4 for more precise statements). The notation of sublinearity and superlinearity of the nonlinear vector field  $(f(u, v), g(u, v))$  or of those in higher dimensions were considered in [39]. But our definition is quite different and our purpose is to prove uniqueness under a sublinear assumption. Our definition of sublinear nonlinearity is similar to the one in [28] for the scalar case.

Dalmasso [11, 12] obtained an existence and uniqueness result for a more special sublinear system and it was extended by Shi and Shivaji [37]. The uniqueness of positive solutions for large  $\lambda$  was proved in [15–18]. For superlinear-type systems, Clément *et al.* [8] obtained *a priori* estimates of positive solutions and used topological methods to prove the existence of solutions. Rellich–Pohozaev-type identities can be established for systems with variational structure [24], which is very useful

for the non-existence result, among others. We note that many of these results are for the special Hamiltonian system case,

$$\left. \begin{aligned} \Delta u + \lambda f(v) &= 0, & x \in \Omega, \\ \Delta v + \lambda g(u) &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{1.5}$$

while we consider equations in a more general form.

For (1.1) satisfying (A2) with  $\Omega$  being a finite ball or the whole space, it is known that a positive solution is radially symmetric [3,41]. Hence, the system can be converted into a system of ordinary differential equations. Several authors have taken that approach for the existence of the solutions [30,33,34] and much success has been achieved for Lane–Emden systems. Using the scaling invariant in the Lane–Emden system, the uniqueness of the radial positive solution has been shown in [10,11,22] for a system with three equations. Korman [21] obtained a uniqueness and exact multiplicity result for the one-dimensional case. A more general approach, using shooting method and linearized equations for the radial case, has been taken by the present authors [6,7].

We recall the maximum principle and prove the main stability result in §2. In §3 we use the stability result and bifurcation theory to prove the existence and uniqueness of solution under several different assumptions on the differentiable nonlinearities; in §4 we consider a similar question for nonlinearities that are merely Hölder continuous, and we use the monotonicity method for the existence and a method in [2,11] for the uniqueness.

**2. The maximum principle and stability**

Let  $(u, v)$  be a solution of (1.1). The stability of  $(u, v)$  is determined by the linearized equation (1.3), which can be written as

$$L\mathbf{u} = J\mathbf{u} + \mu\mathbf{u}, \tag{2.1}$$

where

$$\mathbf{u} = \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \quad L\mathbf{u} = \begin{pmatrix} -\Delta\xi \\ -\Delta\eta \end{pmatrix} \quad \text{and} \quad J = \lambda \begin{pmatrix} f_u & f_v \\ g_u & g_v \end{pmatrix}. \tag{2.2}$$

If we assume that  $(f, g)$  is cooperative (satisfying (A2)), then system (2.1), (2.2) is a linear elliptic system of cooperative type. If we also assume that  $f_v(u(x), v(x)) \not\equiv 0$  and  $g_u(u(x), v(x)) \not\equiv 0$ , then  $J$  is irreducible, and the maximum principles hold for irreducible cooperative systems. We now recall some known results.

LEMMA 2.1. *Suppose that  $\Omega$  is a bounded, open, connected subset of  $\mathbb{R}^n$  satisfying a uniform exterior cone condition, and  $u, v \in C^0(\bar{\Omega})$ , with  $L$  and  $J$  as given in (2.2). Suppose that the entries of  $J$  are in  $L^q(\Omega)$  with  $q \geq pn/(p - n)$  and that  $(f, g)$  satisfies (A2),  $f_v(u(x), v(x)) \not\equiv 0$  and  $g_u(u(x), v(x)) \not\equiv 0$ . Let  $X \equiv [W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)]^2$  and  $Y \equiv [L^p(\Omega)]^2$ , where  $p > n$ . Then we have the following.*

- (i)  $\mu_1 = \inf\{\text{Re}(\mu) : \mu \in \text{spt}(L - J)\}$  is a real eigenvalue of  $L - J$ , where  $\text{spt}(L - J)$  is the spectrum of  $L - J$ .

- (ii) For  $\mu = \mu_1$ , there exists a unique (up to a constant multiple) eigenfunction  $\mathbf{u}_1 \in X$ , and  $\mathbf{u}_1 > 0$  in  $\Omega$ .
- (iii) For  $\mu < \mu_1$ , the equation  $L\mathbf{u} = J\mathbf{u} + \mu\mathbf{u} + \mathbf{f}$  is uniquely solvable for any  $\mathbf{f} \in Y$ , and the solution  $\mathbf{u}(\in X) > 0$  as long as  $\mathbf{f} \geq 0$ .
- (iv) (Maximum principle.) For  $\mu \leq \mu_1$ , suppose that  $\mathbf{u} \in [W^{2,p}(\Omega)]^2$  satisfies  $L\mathbf{u} \geq J\mathbf{u} + \mu\mathbf{u}$  in  $\Omega$ , and  $\mathbf{u} \geq 0$  on  $\partial\Omega$ . Then  $\mathbf{u} \geq 0$  in  $\Omega$ .
- (v) If there exists a  $\mathbf{u} \in [W^{2,p}(\Omega)]^2$  that satisfies  $L\mathbf{u} \geq J\mathbf{u}$  in  $\Omega$ ,  $\mathbf{u} \geq 0$  on  $\partial\Omega$  and either  $\mathbf{u} \not\equiv 0$  on  $\partial\Omega$  or  $L\mathbf{u} \not\equiv J\mathbf{u}$  in  $\Omega$ , then  $\mu_1 > 0$ .

For a more general result and proofs see proposition 3.1, theorem 1.1 and remark 1.4 of [40]. Moreover, from a standard compactness argument, the eigenvalues  $\{\mu_i\}$  of  $L - J$  are countable, and  $|\mu_i - \mu_1| \rightarrow \infty$  as  $i \rightarrow \infty$ . We note that  $\mu_i$  is not necessarily real valued.

We prove the following basic result on the stability of a positive solution.

**THEOREM 2.2.** *Suppose that  $f, g$  satisfies (A1) and (A2) and that  $(u, v)$  is a positive solution of (1.1). We assume that there exists  $q > n$  such that*

$$\text{each entry of } J(u, v) \text{ is in } L^q(\Omega). \tag{2.3}$$

If  $f$  and  $g$  also satisfy the condition

(A3) for any  $(u, v) \in \mathbb{R}_+^2$ ,

$$f(u, v) > f_v(u, v)v + g_v(u, v)u, \quad g(u, v) > g_u(u, v)u + f_u(u, v)v, \tag{2.4}$$

then  $(u, v)$  is stable.

*Proof.* We choose  $p > n$  such that  $q \geq pn/(p - n)$ . Then lemma 2.1 can be applied to  $L - J$  defined in (2.1) and (2.2). Let  $(u, v)$  be a positive solution of (1.1), and let  $(\mu_1, \xi, \eta)$  be the corresponding principal eigenpair such that  $\xi > 0$  and  $\eta > 0$  in  $\Omega$ . Multiplying the equation for  $u$  in (1.1) by  $\eta$ , multiplying the equation for  $\eta$  in (1.3) by  $u$ , integrating over  $\Omega$  and subtracting, we obtain

$$\lambda \int_{\Omega} f\eta \, dx = \lambda \int_{\Omega} g_u u \xi \, dx + \lambda \int_{\Omega} g_v u \eta \, dx + \mu_1 \int_{\Omega} u \eta \, dx. \tag{2.5}$$

Similarly, from the equations for  $v$  and  $\xi$ , we find

$$\lambda \int_{\Omega} g\xi \, dx = \lambda \int_{\Omega} f_u v \xi \, dx + \lambda \int_{\Omega} f_v v \eta \, dx + \mu_1 \int_{\Omega} v \xi \, dx. \tag{2.6}$$

Suppose that  $\mu_1 \leq 0$ . Then we have

$$\lambda \int_{\Omega} f\eta \, dx \leq \lambda \int_{\Omega} g_u u \xi \, dx + \lambda \int_{\Omega} g_v u \eta \, dx \tag{2.7}$$

and

$$\lambda \int_{\Omega} g\xi \, dx \leq \lambda \int_{\Omega} f_u v \xi \, dx + \lambda \int_{\Omega} f_v v \eta \, dx. \tag{2.8}$$

If (A3) is satisfied, then, adding (2.7) and (2.8), we obtain

$$\int_{\Omega} f\eta \, dx + \int_{\Omega} g\xi \, dx \leq \int_{\Omega} (g_v u + f_v v)\eta \, dx + \int_{\Omega} (f_u v + g_u u)\xi \, dx, \tag{2.9}$$

which is in contradiction with (A3) and  $\xi > 0, \eta > 0$  in  $\Omega$ . Hence,  $\mu_1 > 0$  if (A3) is satisfied.  $\square$

The maximum principle can also be used to construct solutions of the elliptic system (1.1) which satisfy the cooperative condition (A2). Let  $X = [C_0^{2,\alpha}(\bar{\Omega})]^2$ . A pair of functions  $(\bar{u}, \bar{v}) \in X$  is an *upper solution* for (1.1) if

$$\left. \begin{aligned} \Delta \bar{u} + \lambda f(\bar{u}, \bar{v}) &\leq 0, & x \in \Omega, \\ \Delta \bar{v} + \lambda g(\bar{u}, \bar{v}) &\leq 0, & x \in \Omega, \\ \bar{u}(x) \geq 0, \quad \bar{v}(x) &\geq 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{2.10}$$

and  $(\underline{u}, \underline{v}) \in X$  is a *lower solution* for (1.1) if

$$\left. \begin{aligned} \Delta \underline{u} + \lambda f(\underline{u}, \underline{v}) &\geq 0, & x \in \Omega, \\ \Delta \underline{v} + \lambda g(\underline{u}, \underline{v}) &\geq 0, & x \in \Omega, \\ \underline{u}(x) \leq 0, \quad \underline{v}(x) &\leq 0, & x \in \partial\Omega. \end{aligned} \right\} \tag{2.11}$$

The following existence result based on the monotone iteration method is well known [29, 32].

**THEOREM 2.3.** *Suppose that  $f$  and  $g$  satisfy (A1) and (A2), and suppose that  $(\bar{u}, \bar{v})$  and  $(\underline{u}, \underline{v})$  are pairs of upper and lower solutions which satisfy  $\bar{u}(x) \geq \underline{u}(x)$  and  $\bar{v}(x) \geq \underline{v}(x)$  for  $x \in \Omega$ . Define  $X_1 \subset X$  by*

$$X_1 = \{(u, v) \in X : \underline{u}(x) \leq u(x) \leq \bar{u}(x), \underline{v}(x) \leq v(x) \leq \bar{v}(x)\}.$$

*Then (1.1) possesses a minimal solution  $(u_m, v_m)$  and a maximal solution  $(u_M, v_M)$  in  $X_1$ ; that is, for any solution  $(u_a, v_a) \in X_1$ , then*

$$u_m \leq u_a \leq u_M \quad \text{and} \quad v_m \leq v_a \leq v_M.$$

### 3. Existence and uniqueness: smooth case

In this section we always assume that  $f$  and  $g$  are smooth, i.e. they satisfy

$$(A1') \quad f, g \in C^1(\overline{\mathbb{R}_+^2}).$$

We note that if (A1') is satisfied, then  $J(u, v) \in C^0(\bar{\Omega})$  and (2.3) is automatically true. Let  $(\lambda_1, \varphi_1)$  be the principal eigenpair of

$$-\Delta\varphi = \lambda\varphi, \quad x \in \Omega, \quad \varphi(x) = 0, \quad x \in \partial\Omega, \tag{3.1}$$

such that  $\varphi_1(x) > 0$  in  $\Omega$  and  $\|\varphi_1\|_{\infty} = 1$ .

First we recall an existence and uniqueness result proved in [37].

THEOREM 3.1. Consider

$$\left. \begin{aligned} \Delta u + \lambda f_1(v) &= 0, & x \in \Omega, \\ \Delta v + \lambda g_1(u) &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega. \end{aligned} \right\} \quad (3.2)$$

Suppose that  $f_1$  and  $g_1$  satisfy, for any  $u \geq 0$  and  $v \geq 0$ ,

$$f_1'(v) > 0, \quad g_1'(u) > 0, \quad (3.3)$$

$$\frac{d}{dv} \left( \frac{f_1(v)}{v} \right) < 0, \quad \frac{d}{du} \left( \frac{g_1(u)}{u} \right) < 0 \quad (3.4)$$

and

$$\lim_{v \rightarrow \infty} \frac{f_1(v)}{v} = \lim_{u \rightarrow \infty} \frac{g_1(u)}{u} = 0. \quad (3.5)$$

Then

- (i) if at least one of  $f_1(0)$  and  $g_1(0)$  is positive, then (3.2) has a unique positive solution  $(u(\lambda), v(\lambda))$  for all  $\lambda > 0$ ,
- (ii) if  $f_1(0) = g_1(0) = 0$ , and  $f_1'(0) > 0$  and  $g_1'(0) > 0$ , then, for some  $\lambda_* = \lambda_1 / \sqrt{f_1'(0)g_1'(0)} > 0$ , (3.2) has no positive solution when  $\lambda \leq \lambda_*$ , and (3.2) has a unique positive solution  $(u(\lambda), v(\lambda))$  for  $\lambda > \lambda_*$ .

Moreover,  $\{(\lambda, u(\lambda), v(\lambda)) : \lambda > \lambda_*\}$  (in the first case, we assume that  $\lambda_* = 0$ ) is a smooth curve, so  $u(\lambda)$  and  $v(\lambda)$  are strictly increasing in  $\lambda$ , and  $(u(\lambda), v(\lambda)) \rightarrow (0, 0)$  as  $\lambda \rightarrow \lambda_*^+$ .

Theorem 3.1 is identical to theorem 6.1 of [37] and we omit the proof here. Note that if  $f_1$  and  $g_1$  satisfy (3.4), then it is necessary that  $f_1(0) \geq 0$  and  $g_1(0) \geq 0$ . Hence,  $f_1$  and  $g_1$  are positive for  $u, v > 0$  here. If  $f_1(0) = 0$ , then we must have  $f_1'(0) > 0$  since  $f_1'(0) > f_1(v)/v$  for  $v > 0$ . Some examples of smooth sublinear functions satisfying conditions (3.3)–(3.5) are

$$\begin{aligned} f(u) &= \ln(u+1) + k, \\ f(u) &= 1 - e^{-u} + k, \\ f(u) &= (1+u)^p - 1 + k, \quad 0 < p < 1, \end{aligned}$$

and

$$f(u) = \frac{u}{(m+u)} + k, \quad k \geq 0.$$

In the following we consider the case when  $f$  and  $g$  depend on both  $u$  and  $v$ . In all results in this section we assume that  $f$  and  $g$  are additionally separated in the form

$$f(u, v) = f_1(v) + f_2(u) \quad \text{and} \quad g(u, v) = g_1(u) + g_2(v). \quad (3.6)$$

All of our results generalize theorem 3.1, but the structure and bifurcation of the solution sets are different. First we have the following.

THEOREM 3.2. Consider

$$\left. \begin{aligned} \Delta u + \lambda[f_1(v) + f_2(u)] &= 0, & x \in \Omega, \\ \Delta v + \lambda[g_1(u) + g_2(v)] &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega. \end{aligned} \right\} \tag{3.7}$$

Suppose that  $f_1(v)$  and  $g_1(u)$  satisfy (3.3)–(3.5), and suppose that  $f_2(u)$  and  $g_2(v)$  satisfy, for any  $u \geq 0$  and  $v \geq 0$ ,

$$f_2(u) \geq 0, \quad f_2'(u) \leq 0, \quad g_2(v) \geq 0, \quad g_2'(v) \leq 0 \tag{3.8}$$

and

$$f_1(0) + f_2(0) > 0 \quad \text{or} \quad g_1(0) + g_2(0) > 0. \tag{3.9}$$

Then (3.7) has a unique positive solution  $(u(\lambda), v(\lambda))$  for all  $\lambda > 0$ . Moreover,  $\{(\lambda, u(\lambda), v(\lambda)) : \lambda > 0\}$  is a smooth curve so that  $u(\lambda)$  and  $v(\lambda)$  are strictly increasing in  $\lambda$ , and  $(u(\lambda), v(\lambda)) \rightarrow (0, 0)$  as  $\lambda \rightarrow 0^+$ .

*Proof.* Our proof follows that of theorem 6.1 in [37]. First we extend  $f_i$  and  $g_i$  to be defined on  $\mathbb{R}$  for  $u, v < 0$  in the following way: if  $f_i(0) = 0$  or  $g_i(0) = 0$ , then we define  $f_i(x)$  or  $g_i(x) \equiv 0$  for  $x < 0$ ; if  $f_i(0) > 0$  or  $g_i(0) > 0$ , then we define  $f_i(x)$  or  $g_i(x) \equiv 0$  for  $x < -\theta$  for some  $\theta > 0$  and we define the function properly so it is continuous on  $\mathbb{R}$ . From the assumptions,  $f(u, v) = f_1(v) + f_2(u)$  and  $g(u, v) = g_1(u) + g_2(v)$  satisfy (A3). Hence, from theorem 2.2, any positive solution of (3.7) is stable. We define

$$F(\lambda, u, v) = \begin{pmatrix} \Delta u + \lambda[f_1(v) + f_2(u)] \\ \Delta v + \lambda[g_1(u) + g_2(v)] \end{pmatrix}, \tag{3.10}$$

where  $\lambda \in \mathbb{R}$  and  $u, v \in C_0^{2,\alpha}(\bar{\Omega})$ . Here, since  $f_i$  and  $g_i$  are  $C^1$ ,  $F: \mathbb{R} \times X \rightarrow Y$  is continuously differentiable, where  $X = [C_0^{2,\alpha}(\bar{\Omega})]^2$  and  $Y = [C^\alpha(\bar{\Omega})]^2$ .

Apparently  $(\lambda, u, v) = (0, 0, 0)$  is a solution of (3.7). We apply the implicit function theorem at  $(\lambda, u, v) = (0, 0, 0)$ . Note that the Fréchet derivative of  $F$  is given here by

$$F_{(u,v)}(\lambda, u, v) \begin{pmatrix} \phi \\ \psi \end{pmatrix} = \begin{pmatrix} \Delta\phi + \lambda[f_1'(v)\psi + f_2'(u)\phi] \\ \Delta\psi + \lambda[g_1'(u)\phi + g_2'(v)\psi] \end{pmatrix}, \tag{3.11}$$

Thus,  $F_{(u,v)}(0, 0, 0)(\phi, \psi)^T = (\Delta\phi, \Delta\psi)^T$ , and it is an isomorphism from  $X$  to  $Y$ . The implicit function theorem implies that  $F(\lambda, u, v) = 0$  has a unique solution  $(\lambda, u(\lambda), v(\lambda))$  for  $\lambda \in (0, \delta)$  for some small  $\delta > 0$ , and that  $(u'(0), v'(0))$  is the unique solution of

$$\begin{aligned} \Delta\phi + f_1(0) + f_2(0) &= 0, & \Delta\psi + g_1(0) + g_2(0) &= 0, & x \in \Omega, \\ \phi(x) = \psi(x) &= 0, & x \in \partial\Omega. \end{aligned} \tag{3.12}$$

Then  $(u'(0), v'(0)) = ([f_1(0)+f_2(0)]e, [g_1(0)+g_2(0)]e)$ , where  $e$  is the unique positive solution of

$$\Delta e + 1 = 0, \quad x \in \Omega, \quad e(x) = 0, \quad x \in \partial\Omega. \tag{3.13}$$




Figure 1. Possible graphs of  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$  in (3.15). Here  $f_1(v) = \ln(v + 1)$  and  $g_1(u) = \ln(2u + 1)$ . (a)  $a = b = 0.9$ , unique intersection; (b)  $a = b = 1.5$ , no intersection.

If  $f_1(0) + f_2(0) > 0$  and  $g_1(0) + g_2(0) > 0$ , then  $(u(\lambda), v(\lambda))$  is positive for  $\lambda \in (0, \delta)$ . If  $f_1(0) + f_2(0) > 0$  and  $g_1(0) + g_2(0) = 0$ , then  $u(\lambda) > 0$  for  $\lambda \in (0, \delta)$  and  $g_2(0) = 0$ . Thus,  $g_2(v) = 0$  for all  $v \in \mathbb{R}$  by (3.8) and the extension of  $g_2$  to  $\mathbb{R}$ . However,

$$\Delta v(\lambda) = -\lambda[g_1(u(\lambda)) + g_2(v(\lambda))] = -\lambda g_1(u(\lambda)) < 0.$$

Hence,  $v(\lambda) > 0$  as well. A similar conclusion holds when  $f_1(0) + f_2(0) = 0$  and  $g_1(0) + g_2(0) > 0$ . Therefore, (3.7) has a positive solution  $(u(\lambda), v(\lambda))$  for  $\lambda \in (0, \delta)$ .

Now we can follow the proof of theorem 6.1 in [37] to obtain the remaining part of the proof. In particular, we can show that the solution  $(\lambda, u(\lambda), v(\lambda))$  is strictly increasing in  $\lambda$  as  $(\partial u(\lambda)/\partial \lambda, \partial v(\lambda)/\partial \lambda)$  satisfies the following equation:

$$F_{(u,v)}(\lambda, u, v) \begin{pmatrix} \frac{\partial u(\lambda)}{\partial \lambda} \\ \frac{\partial v(\lambda)}{\partial \lambda} \end{pmatrix} = - \begin{pmatrix} f_1(v) + f_2(u) \\ g_1(u) + g_2(v) \end{pmatrix}. \tag{3.14}$$

Then  $(\partial u(\lambda)/\partial \lambda, \partial v(\lambda)/\partial \lambda) > 0$  from the maximum principle (lemma 2.1(iii)) and the fact that  $\mu_1((u(\lambda), v(\lambda))) > 0$  from theorem 2.2.  $\square$

Next we consider the case when  $f(u, v)$  and  $g(u, v)$  are not necessarily positive for all  $(u, v) \in \mathbb{R}_+^2$ . Consider

$$\left. \begin{aligned} \Delta u + \lambda[f_1(v) - au] &= 0, & x \in \Omega, \\ \Delta v + \lambda[g_1(u) - bv] &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{3.15}$$

where  $a > 0$  and  $b > 0$ .

First we observe the following geometric properties of the curves  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$  (see figure 1).

LEMMA 3.3. *Suppose that  $f_1(v)$  and  $g_1(u)$  satisfy (3.3)–(3.5) and that  $a, b > 0$ .*

- (i) *If  $f_1(0) = g_1(0) = 0$  and  $ab \geq f_1'(0)g_1'(0)$ , then the curves  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$  have no intersection points in the first quadrant. Furthermore, for any  $(u, v) \in \mathbb{R}_+^2$ ,  $f_1(v) - au \leq 0$  or  $g_1(u) - bv \leq 0$ .*



- (ii) If  $f_1(0) = g_1(0) = 0$  and  $ab < f_1'(0)g_1'(0)$ , then the curves  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$  have a unique intersection point  $(u_*, v_*)$  in the first quadrant.
- (iii) If at least one of  $f_1(0)$  and  $g_1(0)$  is positive, then the curves  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$  have a unique intersection point  $(u_*, v_*)$  in the first quadrant.

*Proof.* First we assume that  $f_1(0) = g_1(0) = 0$ . If  $(u_*, v_*)$  is an intersection point of  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$ , then, from (3.4),

$$f_1'(0) \geq \frac{f_1(v_*)}{v_*} = \frac{au_*}{v_*}, \quad g_1'(0) \geq \frac{g_1(u_*)}{u_*} = \frac{bv_*}{u_*}.$$

Thus, by multiplying the two inequalities, we obtain  $f_1'(0)g_1'(0) \geq ab$ . If  $ab = f_1'(0)g_1'(0)$ , then we must have  $(u_*, v_*) = (0, 0)$  from the strict decreasing property in (3.4). Hence, when  $ab \geq f_1'(0)g_1'(0)$ ,  $f_1(v) - au = 0$  and  $g_1(u) - bv = 0$  have no positive intersection points.

If  $ab < f_1'(0)g_1'(0)$ , then near  $(u, v) = (0, 0)$ , the curve  $f_1(v) - au = 0$  is below  $g_1(u) - bv = 0$ , and hence the existence of an intersection is clear from (3.5). Suppose that  $(u_*, v_*)$  and  $(u_{**}, v_{**})$  are two distinct intersection points. Without loss of generality we assume that  $u_{**} > u_*$ ; then, from the monotonicity of  $f_1$  or  $g_1$ , it is necessary that  $v_{**} > v_*$ . But we have

$$\frac{f_1(v_*)}{av_*} > \frac{f_1(v_{**})}{av_{**}} = \frac{bu_{**}}{g_1(u_{**})} > \frac{bu_*}{g_1(u_*)} = \frac{f_1(v_*)}{av_*}, \tag{3.16}$$

which is a contradiction. Hence, the intersection point  $(u_*, v_*)$  is unique. The case that at least one of  $f_1(0)$  and  $g_1(0)$  is positive is similar and we therefore omit the details. □

The following result classifies the structure of the solution set of (3.15) under different conditions on  $f_1, g_1$  at  $u, v = 0$  and the parameters  $a, b$ .

**THEOREM 3.4.** *Suppose that  $f_1(v)$  and  $g_1(u)$  satisfy (3.3)–(3.5) and that  $a = b > 0$ .*

- (i) If  $f_1(0) = g_1(0) = 0$ ,  $f_1'(0) > 0$ ,  $g_1'(0) > 0$  and  $a^2 \geq f_1'(0)g_1'(0)$ , then (3.15) has no positive solution for any  $\lambda > 0$ .
- (ii) If  $f_1(0) = g_1(0) = 0$ ,  $f_1'(0) > 0$ ,  $g_1'(0) > 0$  and  $a^2 < f_1'(0)g_1'(0)$ , then, for some

$$\lambda_* = \frac{\lambda_1}{\sqrt{f_1'(0)g_1'(0)} - a} \geq \frac{2\lambda_1}{f_1'(0) + g_1'(0) - 2a}, \tag{3.17}$$

equation (3.15) has no positive solution when  $\lambda \leq \lambda_*$  and has a unique positive solution  $(u(\lambda), v(\lambda))$  for  $\lambda > \lambda_*$ .

- (iii) If at least one of  $f_1(0)$  and  $g_1(0)$  is positive, then (3.15) has a unique positive solution  $(u(\lambda), v(\lambda))$  for all  $\lambda > 0$ .

Moreover, in the last two cases,  $\{(\lambda, u(\lambda), v(\lambda)) : \lambda > \lambda_*\}$  (in the second case, we assume  $\lambda_* = 0$ ) is a smooth curve,  $(u(\lambda), v(\lambda)) \rightarrow (0, 0)$  as  $\lambda \rightarrow \lambda_*^+$ .

*Proof.* First we prove that if  $(u(x), v(x))$  is a positive solution of (3.15),  $u(x_1) = \max_{x \in \bar{\Omega}} u(x)$  and  $v(x_2) = \max_{x \in \bar{\Omega}} v(x)$ , then, for  $i = 1, 2$ ,

$$f_1(v(x_i)) - au(x_i) > 0 \quad \text{and} \quad g_1(u(x_i)) - av(x_i) > 0. \quad (3.18)$$

In fact, from the maximum principle and the monotonicity of  $f_1$  and  $g_1$ ,

$$0 \geq \Delta u(x_1) = \lambda[au(x_1) - f_1(v(x_1))] \geq \lambda[au(x_2) - f_1(v(x_2))]. \quad (3.19)$$

Similarly, we also have  $g_1(u(x_i)) - av(x_i) \geq 0$ , and the strict inequalities hold because of the strong maximum principle.

If  $f_1(0) = g_1(0) = 0$ ,  $f_1'(0) > 0$ ,  $g_1'(0) > 0$  and  $a^2 \geq f_1'(0)g_1'(0)$ , then, from lemma 3.3, for any  $(u, v) \in \mathbb{R}_+^2$  either  $f_1(v) - au \leq 0$  or  $g_1(u) - av \leq 0$ . Hence, (3.15) has no positive solution from (3.18). In other cases, from (3.18), we must have  $0 < u(x) < u_*$  and  $0 < v(x) < v_*$  for  $x \in \Omega$  if  $(u(x), v(x))$  is a positive solution of (3.15).

Now we assume that  $f_1(0) = g_1(0) = 0$ ,  $f_1'(0) > 0$ ,  $g_1'(0) > 0$  and  $a^2 < f_1'(0)g_1'(0)$ . We claim that, when

$$\lambda < \frac{2\lambda_1}{f_1'(0) + g_1'(0) - 2a}, \quad (3.20)$$

(3.15) has no positive solution. First of all, the denominator in (3.20) is positive, since

$$f_1'(0) + g_1'(0) \geq 2\sqrt{f_1'(0)g_1'(0)} > 2a.$$

Now, from integration of the equations, we obtain

$$\int_{\Omega} (|\nabla u|^2 + |\nabla v|^2 + a\lambda u^2 + a\lambda v^2) dx = \int_{\Omega} \lambda(f_1(v)u + g_1(u)v) dx. \quad (3.21)$$

From the left-hand side of (3.21), we see that

$$\begin{aligned} \int_{\Omega} (|\nabla u|^2 + |\nabla v|^2 + a\lambda u^2 + a\lambda v^2) dx &\geq \int_{\Omega} (\lambda_1 u^2 + \lambda_1 v^2 + a\lambda u^2 + a\lambda v^2) dx \\ &\geq (\lambda_1 + a\lambda) \int_{\Omega} (u^2 + v^2) dx, \end{aligned} \quad (3.22)$$

while the right-hand side of (3.21) can be estimated as

$$\begin{aligned} \int_{\Omega} \lambda(f_1(v)u + g_1(u)v) dx &\leq \lambda \int_{\Omega} f_1'(0)uv dx + \lambda \int_{\Omega} g_1'(0)uv dx \\ &\leq \frac{1}{2}\lambda(f_1'(0) + g_1'(0)) \int_{\Omega} (u^2 + v^2) dx. \end{aligned} \quad (3.23)$$

Now (3.22) and (3.23) imply that

$$\lambda \geq \frac{2\lambda_1}{f_1'(0) + g_1'(0) - 2a} \quad (3.24)$$

is necessary for the existence of a positive solution  $(u, v)$ .

Next, similarly to the proof of theorem 3.2, we define

$$F(\lambda, u, v) = \begin{pmatrix} \Delta u + \lambda[f_1(v) - au] \\ \Delta v + \lambda[g_1(u) - av] \end{pmatrix}, \tag{3.25}$$

where  $\lambda \in \mathbb{R}$  and  $u, v \in C_0^{2,\alpha}(\bar{\Omega})$ . Then  $(\lambda, 0, 0)$  is a trivial solution of (3.15) for any  $\lambda > 0$ . The linearization of  $F$  at  $(\lambda, 0, 0)$  is

$$F_{(u,v)}(\lambda, 0, 0) \begin{pmatrix} \phi \\ \psi \end{pmatrix} = \begin{pmatrix} \Delta\phi + \lambda[f_1'(0)\psi - a\phi] \\ \Delta\psi + \lambda[g_1'(0)\phi - a\psi] \end{pmatrix}. \tag{3.26}$$

Since  $f_1, g_1$  satisfy (3.3), then, from lemma 2.1,  $F_{(u,v)}(\lambda, 0, 0)$  has a principal eigenvalue  $\mu_1(\lambda)$ , which is the only eigenvalue with positive eigenfunction. Then one can verify that the principal eigenpair of  $F_{(u,v)}(\lambda, 0, 0)$  is  $(\mu_1(\lambda), \varphi_1, c\varphi_1)$ , where  $\mu = \mu_1(\lambda)$  and  $c$  satisfy

$$(\mu + \lambda a + \lambda_1)^2 = f_1'(0)g_1'(0)\lambda^2, \quad c = \frac{\mu + \lambda a + \lambda_1}{\lambda f_1'(0)}. \tag{3.27}$$

In particular, when  $\mu_1(\lambda) = 0$ , we can show that the corresponding  $\lambda = \lambda_*$  is the positive root of

$$[f_1'(0)g_1'(0) - a^2]\lambda^2 - 2a\lambda_1\lambda - \lambda_1^2 = 0, \tag{3.28}$$

and  $c = c_*$  is the positive root of

$$f_1'(0)c^2 - g_1'(0) = 0. \tag{3.29}$$

Hence, when  $\lambda = \lambda_*$ ,  $F_{(u,v)}(\lambda_*, 0, 0)$  is not invertible and  $\lambda = \lambda_*$  is a potential bifurcation point. More precisely, the null space  $N(F_{(u,v)}(\lambda_*, 0, 0)) = \text{span}\{(\varphi_1, c_*\varphi_1)\}$ , the range space

$$R(F_{(u,v)}(\lambda_*, 0, 0)) = \left\{ (\phi, \psi) \in Y : \int_{\Omega} [c_*\phi + \psi]\varphi_1 \, dx = 0 \right\}.$$

Next, note that  $(c_*\varphi_1, \varphi_1)$  is the principal eigenvector of the conjugate operator  $F_{(u,v)}^*(\lambda_*, 0, 0)$ . Finally, we can verify that

$$F_{\lambda(u,v)}(\lambda_*, 0, 0)[\varphi_1, c_*\varphi_1]^T \notin R(F_{(u,v)}(\lambda_*, 0, 0)).$$

If this is not true, then

$$F_{\lambda(u,v)}(\lambda_*, 0, 0) \begin{pmatrix} \varphi_1 \\ c_*\varphi_1 \end{pmatrix} = \begin{pmatrix} (c_*f_1'(0) - a)\varphi_1 \\ (g_1'(0) - ac_*)\varphi_1 \end{pmatrix} \in R(F_{(u,v)}(\lambda_*, 0, 0)),$$

and the definition of  $R(F_{(u,v)}(\lambda_*, 0, 0))$  implies that

$$f_1'(0)c_*^2 - 2ac_* + g_1'(0) = 0. \tag{3.30}$$

But, (3.29) and (3.30) would, taken together, imply that  $c_* = g_1'(0)/a = a/f_1'(0)$ , which is in contradiction with  $a^2 < f_1'(0)g_1'(0)$ . Hence,  $F_{\lambda(u,v)}(\lambda_*, 0, 0)[\varphi_1, c_*\varphi_1]^T \notin R(F_{(u,v)}(\lambda_*, 0, 0))$ .

Now we are in a position to apply a bifurcation from a simple eigenvalue theorem of Crandall and Rabinowitz [9]. The non-trivial solutions of  $F(\lambda, u, v) = (0, 0)$  near  $(\lambda_*, 0, 0)$  are in the form of  $(\lambda(s), u(s), v(s))$  for  $s \in (-\delta, \delta)$ , where  $u(s) = s\varphi_1 + o(s)$  and  $v(s) = c_*s\varphi_1 + o(s)$ . In particular, the solution  $(u(s), v(s))$  is positive when  $s \in (0, \delta)$  from the positivity of  $\varphi_1$ . Moreover, one can use a global bifurcation theorem of Rabinowitz [31] (see also [38]) to conclude that an unbounded continuum  $\Sigma$  of positive solutions of  $F(\lambda, u, v) = (0, 0)$  emanates from  $(\lambda_*, 0, 0)$ . Since (3.15) has no positive solution when

$$\lambda < \frac{2\lambda_1}{f'_1(0) + g'_1(0) - 2a},$$

and all solutions  $(u, v)$  are bounded by  $(u_*, v_*)$ ,  $\Sigma$  must be unbounded in the positive  $\lambda$  direction. This proves that (3.15) has at least one positive solution for each  $\lambda > \lambda_*$ .

Since (A3) is satisfied for  $f(u, v) = f_1(v) - au$  and  $g(u, v) = g_1(u) - av$ , any positive solution  $(u, v)$  of (3.15) is stable from theorem 2.2. Hence, the implicit function theorem can be applied to any positive solution  $(\lambda, u, v)$  so that it belongs to a smooth curve of positive solutions  $\Gamma = \{(\lambda, u(\lambda), v(\lambda))\}$ . Let

$$\lambda_{**} = \inf\{\lambda : (\lambda, u(\lambda), v(\lambda)) \in \Gamma\}.$$

Then

$$\lambda_{**} \geq \frac{2\lambda_1}{f'_1(0) + g'_1(0) - 2a} > 0.$$

From the boundedness of  $(\lambda, u(\lambda), v(\lambda)) \in \Gamma$ , there exists a sequence  $(\lambda^n)$  such that  $\lambda^n > \lambda_{**}$ ,  $\lambda^n \rightarrow \lambda_{**}$ , and  $(u(\lambda^n), v(\lambda^n)) \rightarrow (u_{**}, v_{**})$  in  $[C^2(\bar{\Omega})]^2$  as  $n \rightarrow \infty$ . Hence,  $(u_{**}, v_{**})$  satisfies  $F(\lambda_{**}, u_{**}, v_{**}) = 0$  and  $u_{**}(x) \geq 0$ ,  $v_{**}(x) \geq 0$ . From the maximum principle, either  $u_{**}(x) > 0$  and  $v_{**}(x) > 0$ , or  $u_{**}(x) \equiv 0$  and  $v_{**}(x) \equiv 0$  for  $x \in \Omega$ . But if  $(u_{**}, v_{**})$  is positive, then it is stable, so one can apply implicit function theorem to extend  $\Gamma$ , which contradicts with the definition of  $\lambda_{**}$ . Thus  $(u_{**}, v_{**}) = (0, 0)$ , and we must have  $\lambda_{**} = \lambda_*$  since it is the only bifurcation point for the line of trivial solutions. Since we can use the same argument for each smooth curve of positive solutions, there is only one such curve, which is  $\Sigma$  from the global bifurcation theorem, and  $\Sigma = \{(\lambda, u(\lambda), v(\lambda)) : \lambda > \lambda_*\}$ . This also implies that (3.15) has no positive solution when  $\lambda \leq \lambda_*$ . Here we note that the argument that stability implies uniqueness described above was used in [5].

The case where at least one of  $f_1(0)$  and  $g_1(0)$  is positive is similar, and the existence of a curve of positive solutions can be proved via the implicit function theorem at  $(\lambda, u, v) = (0, 0, 0)$  in a similar way to the proof of theorem 3.2.  $\square$

We note that the condition  $a = b$  in theorem 3.4 is needed for the stability condition (A3). When  $a \neq b$ , the bifurcation arguments can still be applied if  $ab < f'_1(0)g'_1(0)$ , but it is not known whether the stability result in theorem 2.2 holds.

Our last result is about sublinear nonlinearities with positive linear terms. Consider

$$\left. \begin{aligned} \Delta u + \lambda[f_1(v) + au] &= 0, & x \in \Omega, \\ \Delta v + \lambda[g_1(u) + bv] &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{3.31}$$

where  $a, b > 0$ .

**THEOREM 3.5.** *Suppose that  $f_1(v)$  and  $g_1(u)$  satisfy (3.3)–(3.5) and  $a = b > 0$ . Let  $\lambda^* = \lambda_1/a$ , and we define  $\lambda_*$  as follows:*

(i) *if  $f_1(0) = g_1(0) = 0$ ,  $f'_1(0) > 0$  and  $g'_1(0) > 0$ , then*

$$\lambda_* = \frac{\lambda_1}{\sqrt{f'_1(0)g'_1(0) + a}} > 0; \tag{3.32}$$

(ii) *if at least one of  $f_1(0)$  and  $g_1(0)$  is positive, then  $\lambda_* = 0$ .*

*Then (3.31) has no positive solution when  $\lambda \leq \lambda_*$  or  $\lambda \geq \lambda^*$ , and (3.31) has a unique positive solution  $(u(\lambda), v(\lambda))$  for  $\lambda \in (\lambda_*, \lambda^*)$ . Moreover,*

$$\{(\lambda, u(\lambda), v(\lambda)) : \lambda \in (\lambda_*, \lambda^*)\}$$

*is a smooth curve so that  $u(\lambda)$  and  $v(\lambda)$  are strictly increasing in  $\lambda$ ,  $(u(\lambda), v(\lambda)) \rightarrow (0, 0)$  as  $\lambda \rightarrow \lambda_*^+$  and  $\|u(\lambda)\|_\infty + \|v(\lambda)\|_\infty \rightarrow \infty$  as  $\lambda \rightarrow (\lambda^*)^-$ .*

*Proof.* First we assume that  $f_1(0) = g_1(0) = 0$ ,  $f'_1(0) > 0$  and  $g'_1(0) > 0$ . We multiply the equation of  $u$  in (3.31) by  $\varphi_1$  and integrate on  $\Omega$ , obtaining

$$\lambda_1 \int_{\Omega} u\varphi_1 \, dx = \lambda \int_{\Omega} [f_1(v)\varphi_1 + au\varphi_1] \, dx > \lambda a \int_{\Omega} u\varphi_1 \, dx. \tag{3.33}$$

Hence,  $\lambda < \lambda_1/a$ . On the other hand, from (3.33), we have

$$(\lambda_1 - \lambda a) \int_{\Omega} u\varphi_1 \, dx = \lambda \int_{\Omega} f_1(v)\varphi_1 \leq \lambda f'_1(0) \int_{\Omega} v\varphi_1 \, dx. \tag{3.34}$$

Similarly,

$$(\lambda_1 - \lambda a) \int_{\Omega} v\varphi_1 \, dx = \lambda \int_{\Omega} g_1(u)\varphi_1 \leq \lambda g'_1(0) \int_{\Omega} u\varphi_1 \, dx. \tag{3.35}$$

Multiplying (3.34) and (3.35), we obtain that

$$(\lambda_1 - \lambda a)^2 \leq \lambda^2 f'_1(0)g'_1(0). \tag{3.36}$$

Hence, if  $(\lambda, u, v)$  is a positive solution of (3.31), then  $\lambda$  must satisfy (3.36) and

$$\lambda < \frac{\lambda_1}{a}. \tag{3.37}$$

Define

$$h(\lambda) = (\lambda_1 - \lambda a)^2 - \lambda^2 f'_1(0)g'_1(0). \tag{3.38}$$

Then there exists a unique  $\lambda_* \in (0, \lambda_1/a)$  (defined as in (3.32)) such that  $h(\lambda) > 0$  for  $0 \leq \lambda < \lambda_*$  and  $h(\lambda) < 0$  when  $\lambda_* < \lambda < \lambda_1/a$ . Hence, (3.31) can have positive solutions only when  $\lambda \in (\lambda_*, \lambda^*)$ .

A local bifurcation analysis using bifurcation from a simple eigenvalue theorem similar to the one in the proof of theorem 3.4 can be carried out at  $\lambda = \lambda_*$ , and we omit the details. The positive solution  $(\lambda, u(\lambda), v(\lambda))$  is strictly increasing in  $\lambda$  as  $(\partial u(\lambda)/\partial \lambda, \partial v(\lambda)/\partial \lambda)$  satisfies

$$F_{(u,v)}(\lambda, u, v) \begin{pmatrix} \frac{\partial u(\lambda)}{\partial \lambda} \\ \frac{\partial v(\lambda)}{\partial \lambda} \end{pmatrix} = - \begin{pmatrix} f_1(v) + au \\ g_1(u) + av \end{pmatrix}, \tag{3.39}$$

and hence  $(\partial u(\lambda)/\partial \lambda, \partial v(\lambda)/\partial \lambda) > 0$  from the same arguments in the proof of theorem 3.2. The proof of existence of solutions when at least one of  $f_1(0)$  and  $g_1(0)$  is positive is also similar.

Note that (A3) is satisfied for  $f(u, v) = f_1(v) + au$  and  $g(u, v) = g_1(u) + av$ , so any positive solution  $(u, v)$  of (3.31) is stable from theorem 2.2. Hence, in all cases, we can show as before that any positive solution is on a smooth curve, that there is only one such solution curve  $\Sigma$ , and that the left endpoint of  $\Sigma$  is the bifurcation point (depending on the case studied) from the trivial solutions. It only remains to prove that the  $\Sigma$  can be extended to  $\lambda^*$ .

Let  $\lambda^{**} = \sup\{\lambda : (\lambda, u(\lambda), v(\lambda)) \in \Sigma\}$ . From (3.37), we know that  $\Sigma$  can be extended at most to  $\lambda^*$ . Hence,  $\lambda^{**} \leq \lambda^*$ . As  $\lambda \rightarrow (\lambda^{**})^-$ , we must have

$$M_\lambda = \|u(\lambda)\|_\infty + \|v(\lambda)\|_\infty \rightarrow \infty,$$

otherwise a limiting process will yield a positive solution at  $\lambda^{**}$ ; then  $\Sigma$  can be extended further beyond  $\lambda = \lambda^{**}$ , which contradicts with the definition of  $\lambda^{**}$ . Now we define  $U(\lambda) = u(\lambda)/M_\lambda$  and  $V(\lambda) = v(\lambda)/M_\lambda$ . Then  $(U(\lambda), V(\lambda))$  satisfies

$$\left. \begin{aligned} \Delta U + \lambda \left[ \frac{f_1(v)}{M_\lambda} + aU \right] &= 0, & x \in \Omega, \\ \Delta V + \lambda \left[ \frac{g_1(u)}{M_\lambda} + aV \right] &= 0, & x \in \Omega, \\ U(x) = V(x) &= 0, & x \in \partial\Omega. \end{aligned} \right\} \tag{3.40}$$

From the boundedness of  $(U(\lambda), V(\lambda))$  we can obtain a sequence  $(\lambda^n)$  such that  $\lambda^n < \lambda^{**}$ ,  $\lambda^n \rightarrow \lambda^{**}$  and  $(U(\lambda^n), V(\lambda^n)) \rightarrow (U^{**}, V^{**})$  in  $[C^2(\bar{\Omega})]^2$  as  $n \rightarrow \infty$ . From (3.5),  $(\lambda^{**}, U^{**}, V^{**})$  satisfies

$$\left. \begin{aligned} \Delta U^{**} + \lambda^{**} aU^{**} &= 0, & x \in \Omega, \\ \Delta V^{**} + \lambda^{**} aV^{**} &= 0, & x \in \Omega, \\ U^{**}(x) = V^{**}(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \tag{3.41}$$

$\|U^{**}\|_\infty + \|V^{**}\|_\infty = 1$ , and  $U^{**} \geq 0, V^{**} \geq 0$ . Therefore, we must have  $\lambda^{**} = \lambda_1/a$ ,  $U^{**} = s\varphi_1$  and  $V^{**} = (1-s)\varphi_1$  for some  $s \in [0, 1]$ . This proves that  $\lambda^{**} = \lambda^*$ , which completes the proof.  $\square$

**4. Existence and uniqueness: the Hölder continuous case**

In this section we always assume that  $f$  and  $g$  satisfy (A1) and (A2), so  $f$  and  $g$  may not be differentiable. Then the nonlinear operator  $F(\lambda, u, v)$  defined in §3 is not necessarily differentiable, and hence the implicit function theorem and bifurcation theorems cannot be used easily. We shall prove results similar to theorems 3.4 and 3.5 by using different methods.

**THEOREM 4.1.** *Consider (3.15). Suppose that  $f_1(0) = g_1(0) = 0$ ,  $f_1(v)$  and  $g_1(u)$  satisfy (3.5) for any  $u > 0$  and any  $v > 0$ ,*

$$f_1'(v) > 0, \quad g_1'(u) > 0, \tag{4.1}$$

and there exists  $0 < p, q < 1$  such that, for any  $u > 0$  and any  $v > 0$ ,

$$\frac{d}{dv} \left( \frac{f_1(v)}{v^q} \right) \leq 0, \quad \frac{d}{du} \left( \frac{g_1(u)}{u^p} \right) \leq 0. \tag{4.2}$$

Then, for  $\lambda \in (0, \infty)$ , (3.15) has a unique positive solution  $(u(\lambda), v(\lambda))$ . Moreover,

$$\{(\lambda, u(\lambda), v(\lambda)) : \lambda > 0\}$$

is a continuous curve, so  $(u(\lambda), v(\lambda)) \rightarrow (0, 0)$  as  $\lambda \rightarrow 0^+$ .

*Proof.* First we show that the curves  $-au + f_1(v) = 0$  and  $-bv + g_1(u) = 0$  intersect at a unique point in  $\mathbb{R}_+^2$ . From (3.5), for large  $u$ ,  $v = g_1(u)/b$  is below the curve  $u = f_1(v)/a$ . Suppose that the two curves do not intersect. Then, for any  $u > 0$ , we have

$$f_1^{-1}(au) > \frac{g_1(u)}{b}. \tag{4.3}$$

We note that (4.2) implies that, for  $\tau \in (0, 1)$ ,

$$f_1(\tau v) \geq \tau^q f_1(v) \quad \text{and} \quad g_1(\tau u) \geq \tau^p g_1(u). \tag{4.4}$$

For  $u > 0$  small,  $g_1(u) < 1$  and  $u < 1$ , and then (4.3) and (4.4) imply that

$$au > f_1\left(\frac{g_1(u)}{b}\right) \geq [g_1(u)]^q f_1\left(\frac{1}{b}\right) \geq u^{pq} [g_1(1)]^q f_1\left(\frac{1}{b}\right), \tag{4.5}$$

which contradicts  $pq < 1$ . Hence, the two curves must intersect. Suppose that  $(u_*, v_*)$  is an intersection point, and that  $(u_{**}, v_{**})$  is another. Without loss of generality we assume  $u_{**} > u_*$ ; then, from the monotonicity of  $f_1$  or  $g_1$ , it is necessary that  $v_{**} > v_*$ . From

$$a = \frac{f_1(v_*)}{u_*} = \frac{f_1(v_*) v_*^q}{v_*^q u_*} \geq \frac{f_1(v_{**}) v_*^q}{v_{**}^q u_*} = a \left(\frac{v_*}{v_{**}}\right)^q \frac{u_{**}}{u_*}.$$

Thus, we have

$$\left(\frac{v_{**}}{v_*}\right)^q \geq \frac{u_{**}}{u_*}, \tag{4.6}$$

and similarly

$$\left(\frac{u_{**}}{u_*}\right)^p \geq \frac{v_{**}}{v_*}. \tag{4.7}$$

But (4.6) and (4.7) yield

$$\left(\frac{u_{**}}{u_*}\right)^{pq} \geq \frac{u_{**}}{u_*},$$

which contradicts  $u_{**} > u_*$  and  $pq < 1$ . Therefore, the curves  $-au + f_1(v) = 0$  and  $-bv + g_1(u) = 0$  intersect at only one point.

Now we use the monotonicity method to prove the existence of a solution. Let  $(u_*, v_*)$  be the unique intersection point of  $-au + f_1(v) = 0$  and  $-bv + g_1(u) = 0$ . Then, from the proof of theorem 3.4, any positive solution of (3.15) satisfies  $0 < u(x) < u_*$  and  $0 < v(x) < v_*$ . Let  $(\bar{u}, \bar{v}) = (u_*, v_*)$ . Then it is clear that  $(\bar{u}, \bar{v})$  is an upper solution of (3.15).

Let

$$L_f = \frac{f_1(v_*)}{v_*^q}, \quad L_g = \frac{g_1(u_*)}{u_*^p}. \tag{4.8}$$

Since  $0 \leq u \leq u_*$  and  $0 \leq v \leq v_*$ , in this range  $f_1(v) \geq L_f v^q$  and  $g_1(u) \geq L_g u^p$ . We construct a lower solution in the form of  $(\underline{u}, \underline{v}) = (\varepsilon_1 \varphi_1, \varepsilon_2 \varphi_1)$ , where  $\varepsilon_1, \varepsilon_2$  will be specified later. Recall that  $\varphi_1$  is the positive principal eigenfunction with  $\|\varphi_1\|_\infty = 1$ . Now

$$\begin{aligned} \Delta(\varepsilon_1 \varphi_1) + \lambda[-a\varepsilon_1 \varphi_1 + f_1(\varepsilon_2 \varphi_1)] &\geq -\varepsilon_1 \lambda_1 \varphi_1 - \lambda a \varepsilon_1 \varphi_1 + \lambda L_f \varepsilon_2^q \varphi_1^q \\ &= \varepsilon_1 \varphi_1^q [-(\lambda a + \lambda_1) \varphi_1^{1-q} + \lambda L_f \varepsilon_2^q \varepsilon_1^{-1}] \\ &\geq \varepsilon_1 \varphi_1^q [-(\lambda a + \lambda_1) + \lambda L_f \varepsilon_2^q \varepsilon_1^{-1}]. \end{aligned}$$

Similarly,

$$\Delta(\varepsilon_2 \varphi_1) + \lambda[-b\varepsilon_2 \varphi_1 + g_1(\varepsilon_1 \varphi_1)] \geq \varepsilon_2 \varphi_1^p [-(\lambda b + \lambda_1) + \lambda L_g \varepsilon_1^p \varepsilon_2^{-1}].$$

Hence, if we choose

$$\left. \begin{aligned} 0 < \varepsilon_1 &\leq \left(\frac{\lambda L_g}{\lambda b + \lambda_1}\right)^{q/(1-pq)} \left(\frac{\lambda L_f}{\lambda a + \lambda_1}\right)^{1/(1-pq)}, \\ \left(\frac{\varepsilon_1(\lambda a + \lambda_1)}{\lambda L_f}\right)^{1/q} &\leq \varepsilon_2 \leq \frac{\lambda L_g}{\lambda b + \lambda_1} \varepsilon_1^p, \end{aligned} \right\} \tag{4.9}$$

then  $(\underline{u}, \underline{v}) = (\varepsilon_1 \varphi_1, \varepsilon_2 \varphi_1)$  is a lower solution of (3.15). We can choose smaller  $\varepsilon_1$  and  $\varepsilon_2$  but still satisfy (4.9), so that  $(\underline{u}, \underline{v}) \leq (\bar{u}, \bar{v}) = (u_*, v_*)$ . Then from theorem 2.3, there exists a positive solution  $(u, v)$  of (3.15) between the lower and upper solutions.

Next we prove the uniqueness of the solution for any  $\lambda > 0$ . Here we follow an argument of [2, 11]. We fix  $\lambda > 0$ . We define  $G_c(x, y)$  as the Green function of the linear elliptic operator  $-\Delta + cI$  with zero boundary condition where  $c \geq 0$ . Then it is well known that  $G_c(x, y) > 0$  for  $x, y \in \Omega$ . Let  $(u_j, v_j)$ ,  $j = 1, 2$ , be two positive solutions of (3.15). Define

$$S = \{s \in (0, 1]: u_1 - tu_2 \geq 0, v_1 - tv_2 \geq 0 \text{ for } t \in [0, s]\}.$$



Since  $u_j, v_j \in C^{2,\alpha}(\bar{\Omega})$  with  $\alpha = \min\{p, q\}$ , there exists small  $\varepsilon > 0$  such that  $\varepsilon \in S$ , and thus  $S \neq \emptyset$ . Let  $\theta = \sup S$  and assume that  $\theta < 1$ . Then  $u_1 - \theta u_2 \geq 0$  and  $v_1 - \theta v_2 \geq 0$  in  $\bar{\Omega}$ . Then, from (4.1) and (4.4), we have

$$\begin{aligned} u_1(x) &= \lambda \int_{\Omega} G_{\lambda a}(x, y) f_1(v_1(y)) \, dy \\ &\geq \lambda \int_{\Omega} G_{\lambda a}(x, y) f_1(\theta v_2(y)) \, dy \\ &\geq \lambda \theta^q \int_{\Omega} G_{\lambda a}(x, y) f_1(v_2(y)) \, dy \\ &= \theta^q u_2(x). \end{aligned} \tag{4.10}$$

Similarly, we have  $v_1(x) \geq \theta^p v_2(x)$ . From these relations, we obtain

$$\begin{aligned} (-\Delta + \lambda a)(u_1 - \theta u_2) &= \lambda [f_1(v_1) - \theta f_1(v_2)] \\ &\geq \lambda [f_1(\theta^p v_2) - \theta f_1(v_2)] \\ &\geq \lambda(\theta^{pq} - \theta) f_1(v_2) \end{aligned} \tag{4.11}$$

and, similarly,

$$(-\Delta + \lambda b)(v_1 - \theta v_2) \geq \lambda(\theta^{pq} - \theta) g_1(u_2). \tag{4.12}$$

Since  $f_1(v_2(x)) > 0$  and  $g_1(u_2(x)) > 0$  for  $x \in \Omega$ ,  $\theta^{pq} - \theta > 0$ , from the strong maximum principle, (4.11) and (4.12), we have

$$u_1(x) - \theta u_2(x) > 0, \quad v_1(x) - \theta v_2(x) > 0 \quad \text{for } x \in \Omega, \tag{4.13}$$

and the Hopf lemma implies that the normal derivatives on the boundary satisfy

$$\frac{u_1(x) - \theta u_2(x)}{\partial \nu} < 0, \quad \frac{v_1(x) - \theta v_2(x)}{\partial \nu} < 0 \quad \text{for } x \in \partial \Omega. \tag{4.14}$$

This would allow that  $u_1(x) - (\theta + \varepsilon)u_2(x) \geq 0$  and  $v_1(x) - (\theta + \varepsilon)v_2(x) \geq 0$  for a small  $\varepsilon > 0$ , which contradicts the definition of  $\theta$ . Hence, we must have  $\theta = 1$ ,  $u_1 \geq u_2$  and  $v_1 \geq v_2$ . On the other hand, substituting  $u_1, v_1$  with  $u_2, v_2$ , we have  $u_1 \leq u_2$  and  $v_1 \leq v_2$ . Therefore,  $u_1 \equiv u_2$  and  $v_1 \equiv v_2$ , and this proves the uniqueness of the solution.

Finally, we prove that the set of solutions  $\{(\lambda, u(\lambda), v(\lambda)) : \lambda > 0\}$  is a continuous curve. Indeed, for any  $\lambda_a > 0$ , choose a sequence  $\lambda^n \rightarrow \lambda_a$ . Then, from the uniqueness of solutions and the form of the upper/lower solution, one can see that  $(u(\lambda^n), v(\lambda^n)) \rightarrow (u(\lambda_a), v(\lambda_a))$  as  $n \rightarrow \infty$ . Hence, the set of solutions is a continuous curve.  $\square$

**THEOREM 4.2.** *Consider (3.31) with  $a, b > 0$ . Suppose that  $f_1(0) = g_1(0) = 0$ ,  $f_1(v)$  and  $g_1(u)$  satisfy (3.5) and, for any  $u > 0$  and  $v > 0$ , (4.1) and (4.2) hold. Then (3.31) has a unique positive solution  $(u(\lambda), v(\lambda))$  for  $\lambda \in (0, \lambda^*)$ , where  $\lambda^* = \min\{\lambda_1/a, \lambda_1/b\}$ . Moreover,  $\{(\lambda, u(\lambda), v(\lambda)) : \lambda \in (0, \lambda^*)\}$  is a continuous curve, so  $u(\lambda)$  and  $v(\lambda)$  are strictly increasing in  $\lambda$ ,  $(u(\lambda), v(\lambda)) \rightarrow (0, 0)$  as  $\lambda \rightarrow 0^+$ , and*

$$\|u(\lambda)\|_{\infty} + \|v(\lambda)\|_{\infty} \rightarrow \infty \quad \text{as } \lambda \rightarrow (\lambda^*)^-.$$

*Proof.* The proof is similar to that of theorem 4.1, so we only point out the difference. For the existence part we still use  $(\underline{u}, \underline{v}) = (\varepsilon_1 \varphi_1, \varepsilon_2 \varphi_1)$  as a lower solution of (3.31), but now  $\varepsilon_1, \varepsilon_2 \in (0, 1)$  satisfy

$$\left. \begin{aligned} 0 < \varepsilon_1 &\leq \left( \frac{\lambda D_g}{\lambda_1 - \lambda b} \right)^{q/(1-pq)} \left( \frac{\lambda D_f}{\lambda_1 - \lambda a} \right)^{1/(1-pq)} \\ \left( \frac{\varepsilon_1(\lambda_1 - \lambda a)}{\lambda D_f} \right)^{1/q} &\leq \varepsilon_2 \leq \frac{\lambda D_g}{\lambda_1 - \lambda b} \varepsilon_1^p, \end{aligned} \right\} \quad (4.15)$$

where

$$D_f = \left( \inf_{x \in \Omega} \varphi_1^{q-1}(x) \right) f_1(1), \quad D_g = \left( \inf_{x \in \Omega} \varphi_1^{p-1}(x) \right) g_1(1),$$

and here we assume that  $\lambda < \min\{\lambda_1/a, \lambda_1/b\}$ . Note that, from the proof of theorem 3.5, we can only have the solution for  $\lambda < \min\{\lambda_1/a, \lambda_1/b\}$ .

For any  $c < \lambda_1$  we define  $e_c$  to be the unique positive solution of

$$\Delta e + ce + 1 = 0, \quad x \in \Omega, \quad e(x) = 0, \quad x \in \partial\Omega. \quad (4.16)$$

For the upper solution we choose  $(\bar{u}, \bar{v}) = (M_1 e_{\lambda a}, M_2 e_{\lambda b})$ , where  $M_1 \geq 1$  and  $M_2 \geq 1$ . Then

$$\begin{aligned} \Delta(M_1 e_{\lambda a}) + \lambda[aM_1 e_{\lambda a} + f_1(M_2 e_{\lambda b})] \\ &= -M_1 + \lambda f_1(M_2 e_{\lambda b}) \\ &\leq -M_1 + \lambda M_2^q f_1(e_{\lambda b}) \\ &\leq -M_1 + \lambda M_2^q C_f, \end{aligned}$$

where  $C_f = \max_{x \in \bar{\Omega}} f_1(e_{\lambda b}(x))$ . Similarly, if  $C_g = \max_{x \in \bar{\Omega}} g_1(e_{\lambda a}(x))$ , then

$$\Delta(M_2 e_{\lambda b}) + \lambda[bM_2 e_{\lambda b} + g_1(M_1 e_{\lambda a})] \leq -M_2 + \lambda M_1^p C_g.$$

Thus, if we choose  $M_1$  and  $M_2$  by

$$M_1 \geq (\lambda^{1+q} C_g^q C_f)^{1/(1-pq)} \quad \text{and} \quad \left( \frac{M_1}{\lambda C_f} \right)^{1/q} \geq M_2 \geq \lambda M_1^p C_g, \quad (4.17)$$

then  $(\bar{u}, \bar{v}) = (M_1 e_{\lambda a}, M_2 e_{\lambda b})$  is an upper solution of (3.31). By choosing  $M_1$  and  $M_2$  larger (but still satisfying (4.17)) so that  $(\underline{u}, \underline{v}) \leq (\bar{u}, \bar{v})$ , we have a pair of upper and lower solutions needed in theorem 2.3. Hence, the existence of a positive solution is proved.

For the uniqueness proof we only change the operator  $-\Delta + \lambda c$  to  $-\Delta - \lambda c$  for  $c = a, b$ . The elliptic operator is still positive and the maximum principle still holds, since  $\lambda < \min\{\lambda_1/a, \lambda_1/b\}$ . The other parts of the proof are same as those in the proof of theorem 3.5.  $\square$

We remark that the stability defined in § 1 can still be established for  $f$  and  $g$  in theorems 4.1 and 4.2, although  $f_v$  and  $g_u$  become  $\infty$  near  $\partial\Omega$ . By using remark 3.1 of [1] we can extend the spectral theory to the case when the Jacobian has a singularity of the form  $u^{p-1}$  or  $v^{q-1}$  with  $0 < p, q < 1$ .

EXAMPLE 4.3. We can apply the existence and uniqueness results of theorems 4.1 and 4.2 to

$$\left. \begin{aligned} \Delta u + \lambda[\pm au + v^q] &= 0, & x \in \Omega, \\ \Delta v + \lambda[\pm bv + u^p] &= 0, & x \in \Omega, \\ u(x) = v(x) &= 0, & x \in \partial\Omega, \end{aligned} \right\} \quad (4.18)$$

where  $0 < p, q < 1$ . When  $p, q > 1$  and  $pq$  are subcritical, (4.18) (with  $-$  sign) was considered in [14]. Here we prove the existence and uniqueness of the positive solution for the sublinear case.

### Acknowledgements

We thank the anonymous referee for helpful suggestions and comments. J.L.C. was partly supported by the National Science Council of Taiwan Grant no. 97-2115-M008-011. J.S. was partly supported by the National Natural Science Foundation of China Grant no. 10671049, the Longjiang Professorship and the US National Science Foundation. Part of this work was done when J.S. visited the National Central University in May 2009, and he thanks the NCU for their warm hospitality.

### References

- 1 A. Ambrosetti, H. Brézis and G. Cerami. Combined effects of concave and convex nonlinearities in some elliptic problems. *J. Funct. Analysis* **122** (1994), 519–543.
- 2 H. Brézis and S. Kamin. Sublinear elliptic equations in  $\mathbb{R}^n$ . *Manuscr. Math.* **74** (1992), 87–106.
- 3 J. Busca and B. Sirakov. Symmetry results for semilinear elliptic systems in the whole space. *J. Diff. Eqns* **163** (2000), 41–56.
- 4 R. S. Cantrell and C. Cosner. On the positive problem for elliptic systems. *Indiana Univ. Math. J.* **34** (1985), 517–532.
- 5 R. S. Cantrell and C. Cosner. Diffusive logistic equations with indefinite weights: population models in disrupted environments. II. *SIAM J. Math. Analysis* **22** (1991), 1043–1064.
- 6 Z.-Y. Chen, J.-L. Chern, J. Shi and Y.-L. Tang. On the uniqueness and structure of solution to a coupled elliptic system. Preprint.
- 7 J.-L. Chern, C. Lin and J. Shi. Uniqueness of solution to a coupled cooperative system. Preprint.
- 8 Ph. Clément, D. G. de Figueiredo and E. Mitidieri. Positive solutions of semilinear elliptic systems. *Commun. PDEs* **17** (1992), 923–940.
- 9 M. G. Crandall and P. H. Rabinowitz. Bifurcation from simple eigenvalues. *J. Funct. Analysis* **8** (1971), 321–340.
- 10 R. Cui, Y. Wang and J. Shi. Uniqueness of the positive solution for a class of semilinear elliptic systems. *Nonlin. Analysis* **67** (2007), 1710–1714.
- 11 R. Dalmaso. Existence and uniqueness of positive solutions of semilinear elliptic systems. *Nonlin. Analysis* **39** (2000), 559–568.
- 12 R. Dalmaso. Existence and uniqueness of positive radial solutions for the Lane–Emden system. *Nonlin. Analysis* **57** (2004), 341–348.
- 13 D. G. de Figueiredo. Semilinear elliptic systems. In *Nonlinear functional analysis and applications to differential equations*, pp. 122–152 (World Scientific, 1998).
- 14 D. G. de Figueiredo and J. Yang. Decay, symmetry and existence of solutions of semilinear elliptic systems. *Nonlin. Analysis* **33** (1998), 211–234.
- 15 D. D. Hai. Existence and uniqueness of solutions for quasilinear elliptic systems. *Proc. Am. Math. Soc.* **133** (2005), 223–228.
- 16 D. D. Hai. Uniqueness of positive solutions for semilinear elliptic systems. *J. Math. Analysis Applic.* **313** (2006), 761–767.

- 17 D. D. Hai and R. Shivaji. An existence result on positive solutions for a class of semilinear elliptic systems. *Proc. R. Soc. Edinb. A* **134** (2004), 137–141.
- 18 D. D. Hai and R. Shivaji. Uniqueness of positive solutions for a class of semipositone elliptic systems. *Nonlin. Analysis* **66** (2007), 396–402.
- 19 D. S. Hall, M. R. Matthews, J. R. Ensher, C. E. Wieman and E. A. Cornell. Dynamics of component separation in a binary mixture of Bose–Einstein condensates. *Phys. Rev. Lett.* **81** (1998), 1539–1542.
- 20 P. Hess. On uniqueness of positive solutions of nonlinear elliptic boundary value problems. *Math. Z.* **154** (1977), 17–18.
- 21 P. Korman. Global solution curves for semilinear systems. *Math. Meth. Appl. Sci.* **25** (2002), 3–20.
- 22 P. Korman and J. Shi. On Lane–Emden type systems. *Discrete Contin. Dynam. Syst. suppl.* (2005), 510–517.
- 23 T.-C. Lin and J. Wei. Ground state of  $N$  coupled nonlinear Schrödinger equations in  $\mathbb{R}^n$ ,  $n \leq 3$ . *Commun. Math. Phys.* **255** (2005), 629–653.
- 24 E. Mitidieri. A Rellich type identity and applications. *Commun. PDEs* **18** (1993), 125–151.
- 25 J. D. Murray. *Mathematical biology. I. An introduction*, 3rd edn. Interdisciplinary Applied Mathematics, vol. 17 (Springer, 2002).
- 26 J. D. Murray. *Mathematical biology. II. Spatial models and biomedical applications*, Interdisciplinary Applied Mathematics, vol. 18 (Springer, 2003).
- 27 T. Ouyang and J. Shi. Exact multiplicity of positive solutions for a class of semilinear problem. *J. Diff. Eqns* **146** (1998), 121–156.
- 28 T. Ouyang and J. Shi. Exact multiplicity of positive solutions for a class of semilinear problem. II. *J. Diff. Eqns* **158** (1999), 94–151.
- 29 C. V. Pao. On nonlinear reaction–diffusion systems. *J. Math. Analysis Applic.* **87** (1982), 165–198.
- 30 L. A. Peletier and R. C. A. M. Van der Vorst. Existence and nonexistence of positive solutions of nonlinear elliptic systems and the biharmonic equation. *Diff. Integ. Eqns* **5** (1992), 747–767.
- 31 P. H. Rabinowitz. Some global results for nonlinear eigenvalue problems. *J. Funct. Analysis* **7** (1971), 487–513.
- 32 D. H. Sattinger. Monotone methods in nonlinear elliptic and parabolic boundary value problems. *Indiana Univ. Math. J.* **21** (1972), 979–1000.
- 33 J. Serrin and H. Zou. Existence of positive solutions of the Lane–Emden system. *Atti Sem. Mat. Fis. Univ. Modena* **46** (1998), 369–380.
- 34 J. Serrin and H. Zou. Existence of positive entire solutions of elliptic Hamiltonian systems. *Commun. PDEs* **23** (1998), 577–599.
- 35 J. Shi. *Solution set of semilinear elliptic equations: global bifurcation and exact multiplicity* (World Scientific, 2009).
- 36 J. Shi and R. Shivaji. Global bifurcation for concave semipositon problems. In *Advances in evolution equations: proceedings in honor of J. A. Goldstein's 60th birthday* (ed. G. R. Goldstein, R. Nagel and S. Romanelli), pp. 385–398 (New York: Marcel Dekker, 2003).
- 37 J. Shi and R. Shivaji. Exact multiplicity of positive solutions to cooperative elliptic systems. Preprint.
- 38 J. Shi and X. Wang. On global bifurcation for quasilinear elliptic systems on bounded domains. *J. Diff. Eqns* **246** (2009), 2788–2812.
- 39 B. Sirakov. Notions of sublinearity and superlinearity for nonvariational elliptic systems. *Discrete Contin. Dynam. Syst.* **13** (2005), 163–174.
- 40 G. Sweers. Strong positivity in  $C(\bar{\Omega})$  for elliptic systems. *Math. Z.* **209** (1992), 251–271.
- 41 W. C. Troy. Symmetry properties in systems of semilinear elliptic equations. *J. Diff. Eqns* **42** (1981), 400–413.

(Issued 25 February 2011)