

# Neighborhood poverty and suicidal thoughts and attempts in late adolescence

V. Dupéré<sup>1,2\*</sup>, T. Leventhal<sup>2</sup> and É. Lacourse<sup>3</sup>

<sup>1</sup> Department of Psychology, Université de Montréal, Canada

<sup>2</sup> Eliot-Pearson Department of Child Development, Tufts University, Medford, MA, USA

<sup>3</sup> Research Unit on Children's Psychosocial Maladjustment, Biopsychosocial Research Unit, Hôpital Ste-Justine and Department of Sociology, Université de Montréal, Canada

**Background.** Suicide tends to concentrate in disadvantaged neighborhoods, and neighborhood disadvantage is associated with many important risk factors for youth suicide. However, no study has directly investigated the link between neighborhood poverty and youth suicidal behaviors, while controlling for pre-existing vulnerabilities. The objective of this study was to determine whether living in a poor neighborhood is associated with suicidal thoughts and attempts in late adolescence over and above background vulnerabilities, and whether this association can be explained by late-adolescence psychosocial risks: depression, social support, negative life events (NLEs), delinquent activities, substance abuse and exposure to suicide. The potential moderating role of neighborhood poverty was also examined.

**Method.** A subset of 2776 participants was selected from the Canadian National Longitudinal Survey of Children and Youth (NLSCY). Late-adolescence suicidal behaviors and risk factors were self-reported. The 2001 Canadian Census was used to characterize neighborhoods during early and middle adolescence. Late-childhood family and individual controls were assessed through parent-report.

**Results.** At the bivariate level, the odds of reporting suicidal thoughts were about twice as high in poor than non-poor neighborhoods, and the odds of attempting suicide were about four times higher. After controlling for background vulnerabilities, neighborhood poverty remained significantly associated with both suicidal thoughts and attempts. However, these associations were not explained by late-adolescence psychosocial risks. Rather, youth living in poor neighborhoods may be at greater risk through the amplification of other risk factors in disadvantaged neighborhoods.

**Conclusions.** Potential explanations for the increased vulnerability of youth living in poor neighborhoods are discussed.

Received 31 March 2008; Revised 13 August 2008; Accepted 3 September 2008; First published online 10 October 2008

**Key words:** Adolescence, neighborhood poverty, suicidal thoughts, suicide attempts.

## Introduction

Understanding the antecedents of youth suicidal thoughts and attempts is important because these behaviors indicate severe distress that may persist for years, sometimes leading to mental illness well into adulthood (Fergusson *et al.* 2005; Reinherz *et al.* 2006). Suicidal behaviors also represent the most potent risk factor for subsequent completed suicides (Gould *et al.* 2003; Spirito & Overholser, 2003; Bridge *et al.* 2006). In addition to these severe consequences for youth,

repercussions of suicide and associated behaviors are colossal for significant others, including of course parents and siblings (Pfeffer *et al.* 2002; Murphy *et al.* 2003), but also treating clinicians (Hendin *et al.* 2004). Despite the significance of suicide and associated behaviors, our understanding of these behaviors remains limited (Joiner *et al.* 2005). Comprehensive models including a broad range of risk factors, from the biological to the social, are thought to represent the best avenue for improving our understanding of these behaviors (Joiner *et al.* 2005; Spirito & Esposito-Smythers, 2006). Neighborhood disadvantage is a potentially important social risk factor for youth suicide that has received little research attention. The aim of the present study was to examine the association between neighborhood poverty and suicidal thoughts and attempts among

---

\* Address for correspondence: V. Dupéré, Ph.D., Eliot-Pearson Department of Child Development, Medford, MA 02155, USA.  
(Email: verodupere@yahoo.com)

This paper was presented at the biennial meeting of the Society for Research on Adolescence, Chicago, March 2008.

older adolescents, a group particularly at risk for these outcomes (Kessler *et al.* 1999).

Indirect evidence suggests that neighborhood disadvantage could improve explanatory models of youth suicide. First, epidemiological studies show that adult suicide deaths tend to concentrate in disadvantaged neighborhoods (Rehkopf & Buka, 2005; Middleton *et al.* 2006), perhaps especially so among younger residents (Middleton *et al.* 2004; Miller *et al.* 2005; Exeter & Boyle, 2007). Second, results from a study that included both individual and community data revealed that neighborhood disadvantage was independently associated with adult suicide deaths even after taking into account individual socioeconomic characteristics (Cubbin *et al.* 2000). Third, although neighborhood effects on adolescent suicidal behaviors have yet to be demonstrated, neighborhood disadvantage is associated with an array of major risk factors for youth suicide. Importantly, youth living in disadvantaged neighborhoods exhibit higher rates of internalizing symptoms such as depression and hopelessness (Perez-Smith *et al.* 2002; Xue *et al.* 2005). They also present more externalizing problems, such as delinquency and substance abuse (Leventhal & Brooks-Gunn, 2000; Duncan *et al.* 2002), which are also associated with youth suicide (Gould *et al.* 2003; Bridge *et al.* 2006; Spirito & Esposito-Smythers, 2006). Thus, neighborhood disadvantage could increase youth suicidal behaviors through its impact on more proximal risk factors in the internalizing and externalizing spheres.

Why is neighborhood disadvantage associated with these major risk factors for youth suicide? Its association with internalizing symptoms is thought to be rooted in exposure to stressful events in the community, in the form of social and physical disorder (Aneshensel & Sucoff, 1996; Ross, 2000; Wheaton & Clarke, 2003; Hill *et al.* 2005). Examples of social disorder prevalent in disadvantaged communities that could be perceived as threatening and affect youths' mental health include exposure to violence and intimidation, substance abuse and conspicuous illegal activities. In an effort to avoid such stressful situations, youth and their parents often choose to avoid public spaces in disadvantaged neighborhoods (Jarrett, 1999; Molnar *et al.* 2004), resulting in increased isolation and reduced social support. Physical disorder, such as dilapidated housing or poorly kept infra-structures, including schools and parks, might be interpreted as additional evidence of a lack of investment in building a safe environment that provides positive opportunities for youth (Molnar *et al.* 2004). It is easy to envisage how repeated exposure to stressors and reduced access to support in disadvantaged communities might induce feelings of hopelessness and

depression. For these reasons, stressful life events, reduced social support and depression are considered in this study as potential risk factors linking neighborhood disadvantage and youth suicidal behaviors.

Neighborhood disadvantage not only increases exposure to social disorder but also increases the likelihood that youth engage in problematic behaviors themselves, including delinquent activities and substance abuse (Leventhal & Brooks-Gunn, 2000; Duncan *et al.* 2002). Youth living in disadvantaged neighborhoods are thought to be more likely to be involved in delinquency because of the difficulty residents experience exerting effective collective control over youth groups within this environment (Sampson *et al.* 1997; Leventhal & Brooks-Gunn, 2000). Residents living in disadvantaged areas often withdraw from a community perceived as dangerous, thus impeding the development of social cohesion and mutual trust, two collective resources considered essential for efficient supervision of youth groups (Sampson *et al.* 1997). Drugs are also more accessible in poor than non-poor neighborhoods because drug markets are more prolific when social cohesion is low (Saxe *et al.* 2001). In this study, involvement in delinquent activities and substance abuse, two problem behaviors associated with both neighborhood disadvantage and youth suicidal behaviors, were examined as potential explanatory mechanisms.

Exposure to suicide is another aspect that could increase suicide risk in disadvantaged neighborhoods. As mentioned earlier, epidemiological studies show that adult suicide tends to concentrate in disadvantaged areas (e.g. Rehkopf & Buka, 2005). As such, adolescents raised in disadvantaged neighborhoods could be more likely to have known people who have committed suicide. Exposure to suicide is a documented environmental risk factor for youth suicide, possibly because of contagion or imitation effects (Gould *et al.* 2003; Spirito & Esposito-Smythers, 2006). Thus, it was also explored as a potential explanatory mechanism.

So far, it has been argued that neighborhood disadvantage could increase suicidal thoughts and attempts by facilitating the emergence of important individual risk factors for suicide. However, neighborhood disadvantage could not only be associated with a higher level of these risks but also amplify their impact. Recent results indicate that neighborhood disadvantage, perhaps because it limits access to support and coping resources, could indeed modify the impact of other risk factors. For example, one study found that negative life events (NLEs) were more likely to lead to depression among adult residents of disadvantaged neighborhoods, as compared to those of better-off neighborhoods (Cutrona *et al.* 2005).

Among adolescents, neighborhood disadvantage has been shown to amplify the impact of individual vulnerabilities on youth externalizing outcomes (Lynam *et al.* 2000; Obeidallah *et al.* 2004; Dupéré *et al.* 2008). In keeping with these results, the moderating impact of neighborhood disadvantage on other risk factors was explored.

The primary objective of this study was to examine whether youth living in poor neighborhoods are at an increased risk for suicidal thoughts and attempts in late adolescence, after controlling for pre-existing vulnerabilities. Introducing control variables is necessary in neighborhood research to reduce potential selection biases, given that otherwise disadvantaged families are more likely to live in poor areas (Leventhal & Brooks-Gunn, 2000). Family socio-economic status (SES), family disruption and maternal depression, and also late-childhood internalizing symptoms and hyperactivity/impulsivity problems, were included as controls. Including individual-level controls in addition to family socio-economic characteristics allows a more thorough control of potentially confounding factors (Xue *et al.* 2005). A second aim was to examine whether neighborhood effects could be explained by increased levels, in late adolescence, of important risk factors for youth suicide and associated behaviors, including depression, NLEs, low social support, substance abuse, delinquent activities, and exposure to suicide. Finally, the potential moderating role of neighborhood poverty on other risk factors was investigated.

## Method

### Sample

The National Longitudinal Survey of Children and Youth (NLSCY) is a nationally representative, prospective study of Canadian children and adolescents launched in 1994–1995, with ongoing follow-ups conducted biennially (Statistics Canada and Human Resources Development Canada, 1995; Statistics Canada, 2007). The NLSCY is based on a clustered probability sample of private households within the 10 Canadian provinces, excluding children living in remote areas, institutional settings, and on First Nations reserves. Initially, 13 439 households out of 15 579 identified as having at least one dependent child newborn to age 12 years agreed to participate. In each of the survey households, the ‘person most knowledgeable’ about the child was interviewed. In almost all cases, this person was the mother, and is referred to hereafter as such. From 10 years old on, children were also asked to complete self-reported questionnaires. Within each household, up to a maximum of four children were

randomly selected, resulting in an initial total of 22 831, of whom 16 903 were selected to be followed biennially.

In this study, we selected participants who had reached 18–19 years old by cycle 6, the last currently available cycle of data collection. That is, we selected those who were at least 8 years old in cycle 1 ( $n = 4951$ ). Of that number, we selected those who still participated in the survey when they reached 18–19 years old ( $n = 3088$ ). Thus, the retention rate after a decade was  $>60\%$ . Following previous NLSCY neighborhood studies (Kohen *et al.* 2002; Dupéré *et al.* 2008), one child per household was selected at random to avoid within-family clustering, resulting in a final sample size of 2776. This sample was approximately evenly distributed across two cohorts: for the older cohort, aged 10–11 years in cycle 1, we used data collected from cycle 1 to cycle 5 (1994–1995 to 2002–2003); for the younger cohort, aged 10–11 years in cycle 2, we used data collected from cycle 2 to cycle 6 (1996–1997 to 2004–2005).

To take into account attrition as well as the sampling design, all analyses were conducted using normalized longitudinal survey weights. The use of longitudinal weights helps to preserve the representativeness of the sample despite survey drop-out, but it does not eliminate potential bias associated with partial non-response. Partial non-response arises when participants omit answering some of the survey questions. In this study, the rate of partial non-response varied between 0% and 4% for mother-reported variables, and between 6% and 9% for self-reported variables. Partial non-response on the predictors was treated through multiple imputation (Allison, 2001; von Hippel, 2007), using the SAS procedures MI and MIANALYZE (SAS Institute Inc., Cary, NC, USA). It is noteworthy that had simple list-wise deletion been used to manage non-response, the main conclusions of the present study would remain unchanged.

### Measures

Suicidal behaviors and risk factors were self-reported in late adolescence (18–19 years old). Neighborhood poverty was assessed at two time-points during adolescence (12–13 and 14–15 years old), using the 2001 Canadian census. Family and individual controls were measured through parent-report in late childhood (10–11 years old).

#### *Suicidal behavior (18–19 years old)*

Suicidal thoughts were assessed by asking participants: ‘During the past 12 months, did you seriously

consider attempting suicide?’ For those who answered ‘yes’ (7.7%), suicide attempts were measured by asking: ‘During the past 12 months, how many times did you attempt suicide?’ For this sample, 3.2% reported having attempted suicide at least once. Similar items have been used to measure suicidal behavior in other studies using community-based and national samples (Kessler *et al.* 1999; Johnson *et al.* 2002). The pattern of prevalence of suicide ideations and attempts is consistent with those reported in other Canadian (Weissman *et al.* 1999; Langlois & Morrison, 2002; Fotti *et al.* 2006) and American (Kessler *et al.* 1999; Johnson *et al.* 2002) samples.

#### *Late-adolescence risk factors (18–19 years old)*

Current depression was assessed with a 12-item shortened version of the Center for Epidemiological Studies Depression Scale (CES-D; Radloff, 1977). The items tapped the presence of depressive symptoms in the past 7 days, including depressed mood, hopelessness, poor appetite, trouble concentrating and restless sleep (mean = 6.9, s.d. = 6.6). Answers were coded on a four-point scale, ranging from score 0 (rarely; <1 day) to 3 (most or all of the time; 5–7 days). The shortened scale showed good psychometric properties in the NLSCY as well as in other Canadian samples (Statistics Canada and Human Resources Development Canada, 1995; Poulin *et al.* 2005). Social support (mean = 19.76, s.d. = 3.55) was measured by eight items (e.g. there are people I can count on in an emergency; if something went wrong, no one would help me) derived from the Social Provision Scale (Cutrona, 1984). The response scale included four choices (0 = strongly disagree; to 3 = strongly agree). Five items assessed the occurrence of NLEs (mean = 1.18, s.d. = 1.13) during the past 2 years (i.e. painful break-up with boyfriend/girlfriend, serious problem at school or at work, death of someone close, divorce or separation of parents and other difficult events). The major domains in which NLEs may occur for adolescents were tapped through these items (Williamson *et al.* 1998).

Substance abuse was measured through two self-reported items asking about the frequency of alcohol abuse (frequency of being drunk or having five drinks or more on one occasion in the past 12 months) and marijuana consumption (frequency of marijuana/hashish/cannabis use in the past 12 months). Those who abused alcohol (17%) or used marijuana (17%) more than once a week were considered at risk. Involvement in delinquent activities was assessed through one question asking participants if they were part, in the past 12 months, of a gang that broke the law by stealing, hurting people or damaging property (4%). Because of item inconsistencies between cohorts,

we had to rely on a single item to define this variable. However, it is important to note that the relationship between involvement in delinquent groups and delinquency is strong (Lacourse *et al.* 2003) and that similar single-item measures have shown adequate validity (Thornberry *et al.* 2003; Lacourse *et al.* 2006). Finally, participants were asked if someone they had known personally had committed suicide as a measure of exposure to suicide. About a quarter reported that they did.

#### *Early and middle adolescence neighborhood poverty (12–13 and 14–15 years old)*

Neighborhood poverty was assessed at two time-points, in early (12–13 years old) and middle (14–15 years old) adolescence, using the 2001 Canadian Census. The dissemination area (DA) was the geographic unit used to approximate the neighborhood environment (Puderer, 2001). The DA has a population range of 400 to 700 individuals and represents a convenient geographical unit that has been used in other NLSCY neighborhood studies (Kohen *et al.* 2002; Dupéré *et al.* 2008). All DAs with 20% or more residents under Statistics Canada’s Low-Income Cut-off qualified as poor neighborhoods. In early adolescence, middle adolescence, or both, 28% of the sample resided in a poor neighborhood.

#### *Late-childhood family and individual controls (10–11 years old)*

Family SES was measured through an index computed by Statistics Canada (Willms & Shields, 1996). Five standardized variables were combined in this index, including mothers’ and fathers’ educational attainment, mothers’ and fathers’ occupational prestige, and household income (mean = 0.0, s.d. = 0.8). Non-intact family status represented those (29%) who did not live with two biological or adoptive parents. Maternal depression was assessed with the 12-item shortened version of the CES-D described above (mean = 4.8, s.d. = 6.0). Two parent-reported scales measuring internalizing problems and hyperactivity/impulsivity were used to measure individual control variables. The items included in the scales were previously validated in other Canadian samples (Boyle *et al.* 1987; Tremblay *et al.* 1994). Both scales showed good psychometric properties (Statistics Canada and Human Resources Development Canada, 1995). Response scales with three categories were used (0 = never to 2 = often). Internalizing problems (mean = 2.9, s.d. = 2.8) were assessed with eight items (e.g. seems unhappy, sad or depressed; is nervous, high-strung or tense). Hyperactivity/impulsivity

(mean = 4.2, s.d. = 3.6) was measured with eight items (e.g. is restless or hyperactive; is impulsive, acts without thinking).

## Results

### Attrition

Measures obtained at 10–11 years of age were used to assess differential attrition. Males [ $\chi^2(1, n = 4951) = 7.3, p = 0.007$ ], non-Whites [ $\chi^2(1, n = 4484) = 4.9, p = 0.027$ ] and participants from lower-SES families [ $F(1, 4674) = 32.4, p < 0.001$ ] were more likely to have dropped out by late adolescence than girls, Whites and higher SES families respectively. However, no significant differences in attrition were found by cohort, age, family status, parental depression, hyperactivity, internalizing symptoms and neighborhood poverty.

### Descriptives statistics

Table 1 presents the intercorrelations among suicidal thoughts and attempts and individual and family predictors. The results indicate that suicidal thoughts and attempts are significantly ( $p < 0.001$ ) associated with most late-childhood controls and late-adolescence risks. The associations tended to be stronger for variables measured in late adolescence, as compared with demographics and controls measured in late childhood. Not surprisingly, late-adolescence depression showed the strongest correlation with both suicidal thoughts and attempts. Other risk factors strongly associated with suicidal thoughts or attempts ( $p < 0.001$ ) included maternal depression, hyperactivity, neighborhood poverty, alcohol and marijuana use, NLEs, lack of social support and exposure to suicide.

As hypothesized, living in a poor neighborhood was associated with both suicidal thoughts (6.1% in non-poor neighborhoods; 11.6% in poor neighborhoods) and suicide attempts (1.8% in non-poor neighborhoods; 6.6% in poor neighborhoods). Based on the results of univariate logistic regressions, the odds of reporting suicidal thoughts were about two times higher among those who lived in poor neighborhoods during adolescence as compared to those who did not [odds ratio (OR) 2.0, 95% confidence interval (CI) 1.5–2.7], whereas the odds of attempting suicide were about four times higher (OR 4.0, 95% CI 2.5–6.3). Table 1 also shows that growing up in a poor neighborhood was negatively associated, at  $p < 0.001$ , with White race/ethnicity, residence in a rural area, family SES and social support and positively associated with non-intact family status and maternal and youth depression. At  $p < 0.05$ , neighborhood poverty was also positively associated with hyperactivity and NLEs.

### Multivariate models

Hierarchical logistic regressions were used to predict suicidal thoughts and attempts. The first model included demographics and late-childhood family and individual controls, as well as neighborhood poverty, while also adjusting for cohort and region of residence. This model tested whether there was an independent association between neighborhood poverty and suicidal thoughts and attempts. The second model examined potential mediation mechanisms of neighborhood effects, by incorporating late-adolescence risk factors, including depression, low social support, NLEs, alcohol and marijuana abuse, delinquent activities and exposure to suicide into the previous model. The last model assessed potential moderating effects by incorporating interactions found between neighborhood poverty and family and individual controls and risk factors measured in early childhood and late adolescence. Only significant interactions were incorporated.

#### Suicidal thoughts (Table 2)

Model 1 (Table 2) demonstrates that when demographics and late-childhood controls were included simultaneously along with neighborhood poverty, neighborhood poverty had a significant independent effect on youth suicidal thoughts. The effect of neighborhood poverty was essentially unchanged when compared with univariate results. This pattern of findings suggests that its association with suicidal thoughts is probably not attributable to individual and family vulnerabilities.

Model 2 shows that the effect of neighborhood poverty again remained essentially unchanged when late-adolescence risks (depression, low social support, NLEs, delinquent activities, substance abuse and exposure to suicide) were incorporated. Thus, a mediation hypothesis was not supported.

Two interaction effects, between neighborhood poverty and hyperactivity ( $p = 0.06$ ) and exposure to suicide ( $p = 0.05$ ), emerged as marginally significant over and above the effect of other family and individual risk factors. These interaction effects were incorporated into Models 3 and 4 respectively. In these models, the main effects of neighborhood poverty was appreciably reduced, suggesting that youth living in poor neighborhoods could be partly more at risk through the amplified effect of other risk factors. Fig. 1(a, b) illustrates these findings.

#### Suicide attempts (Table 3)

In general, the results for suicide attempts followed a similar pattern to those for suicidal thoughts

**Table 1.** Intercorrelations of suicide thoughts and attempts and predictor variables

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1. Suicide thoughts	–																	
2. Suicide attempts	<b>0.63</b>	–																
3. Age	–0.04	–0.02	–															
4. White	0.02	0.01	0.01	–														
5. Rural	0.01	0.00	0.00	<b>0.09</b>	–													
6. Male	–0.01	–0.03	–0.02	–0.01	–0.02	–												
7. SES	–0.05	–0.07	–0.02	<b>0.10</b>	<b>–0.16</b>	0.04	–											
8. Non-intact family	0.02	0.04	0.01	–0.04	–0.05	–0.01	<b>–0.25</b>	–										
9. Maternal depression	0.08	<b>0.09</b>	–0.05	–0.04	0.01	0.00	<b>–0.27</b>	<b>0.25</b>	–									
10. Hyperactivity	<b>0.09</b>	<b>0.14</b>	0.00	0.03	0.02	<b>0.16</b>	<b>–0.15</b>	<b>0.13</b>	<b>0.19</b>	–								
11. Internalizing symptoms	0.07	0.06	0.05	<b>0.07</b>	–0.01	0.02	<b>–0.10</b>	<b>0.08</b>	<b>0.28</b>	<b>0.46</b>	–							
12. Poor neighborhood	<b>0.09</b>	<b>0.13</b>	0.01	<b>–0.14</b>	<b>–0.07</b>	0.03	<b>–0.26</b>	<b>0.21</b>	<b>0.11</b>	0.04	0.02	–						
13. Depression	<b>0.34</b>	<b>0.24</b>	0.01	–0.03	–0.06	<b>–0.11</b>	–0.05	<b>0.12</b>	<b>0.12</b>	<b>0.14</b>	<b>0.16</b>	<b>0.11</b>	–					
14. Delinquent activities	0.06	0.05	–0.02	–0.06	–0.02	0.00	–0.05	0.03	0.02	0.01	0.02	0.03	0.06	–				
15. Alcohol	<b>0.11</b>	0.04	–0.01	–0.02	0.02	<b>0.14</b>	–0.02	0.02	<b>0.08</b>	0.04	–0.01	0.04	0.04	0.03	–			
16. Marijuana	<b>0.14</b>	<b>0.11</b>	0.03	–0.03	–0.01	<b>0.16</b>	–0.03	<b>0.08</b>	0.03	<b>0.10</b>	0.01	0.01	<b>0.11</b>	0.05	<b>0.31</b>	–		
17. NLEs	<b>0.19</b>	<b>0.16</b>	0.04	0.03	–0.03	<b>–0.12</b>	–0.07	<b>0.12</b>	<b>0.09</b>	<b>0.11</b>	<b>0.11</b>	0.07	<b>0.33</b>	0.03	<b>0.09</b>	<b>0.09</b>	–	
18. Social support	<b>–0.17</b>	<b>–0.15</b>	–0.06	<b>0.12</b>	0.04	<b>–0.15</b>	<b>0.14</b>	<b>–0.12</b>	<b>–0.09</b>	<b>–0.13</b>	<b>–0.07</b>	<b>–0.11</b>	<b>–0.29</b>	–0.01	–0.06	<b>–0.13</b>	–0.06	–
19. Exposure	<b>0.09</b>	<b>0.11</b>	0.06	0.06	0.05	<b>–0.11</b>	<b>–0.07</b>	<b>0.09</b>	0.03	0.05	0.03	–0.02	0.06	0.06	<b>0.07</b>	0.06	<b>0.22</b>	–0.01

SES, Socio-economic status; NLE, negative life event.

Boldface indicates coefficient significant at  $p < 0.001$ .


**Table 2.** Logistic regressions predicting young adults' suicidal thoughts<sup>a</sup>

Measure	Model 1		Model 2		Model 3		Model 4	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
<b>Demographics</b>								
Male	0.77	0.57–1.05	0.89	0.61–1.28	0.91	0.63–1.32	0.91	0.63–1.33
Age	0.69*	0.51–0.94	0.53***	0.38–0.75	0.52***	0.37–0.74	0.52***	0.37–0.73
White	1.78	0.89–3.55	2.59*	1.13–5.96	2.39*	1.05–5.43	2.58*	1.13–5.90
Rural	1.17	0.72–1.89	1.57†	0.92–2.69	1.53	0.89–2.62	1.58†	0.92–2.71
<b>Late-childhood controls</b>								
SES	1.04	0.84–1.29	0.99	0.76–1.29	0.99	0.76–1.29	0.98	0.75–1.28
Non-intact family	0.86	0.60–1.22	0.47***	0.31–0.72	0.46***	0.30–0.70	0.47***	0.31–0.71
Maternal depression	1.02	0.99–1.05	1.01	0.98–1.05	1.01	0.98–1.05	1.01	0.98–1.05
Internalizing symptoms	1.01	0.95–1.07	1.02	0.95–1.09	1.02	0.95–1.09	1.02	0.95–1.09
Hyperactivity/impulsivity	1.08***	1.03–1.13	1.01	0.96–1.07	0.97	0.91–1.04	1.01	0.95–1.06
<b>Early- and middle-adolescence neighborhood context</b>								
Poor neighborhood	2.04***	1.47–2.85	2.03***	1.39–2.95	1.31	0.72–2.39	1.58†	1.00–2.49
<b>Late-adolescence risks</b>								
Depression			1.15***	1.12–1.18	1.15***	1.12–1.18	1.15***	1.12–1.19
NLEs			1.30**	1.11–1.51	1.30**	1.11–1.52	1.31***	1.12–1.53
Social support			0.90***	0.86–0.95	0.90***	0.85–0.94	0.90***	0.86–0.95
Delinquent activities			1.46	0.70–3.06	1.45	0.69–3.05	1.40	0.67–2.94
Substance abuse								
Alcohol			2.01***	1.34–3.01	2.02***	1.34–3.04	2.01***	1.34–3.03
Marijuana			2.13***	1.43–3.17	2.17***	1.45–3.24	2.15***	1.44–3.21
Exposure to suicide			1.74**	1.17–2.60	1.75**	1.17–2.62	1.34	0.82–2.18
<b>Interactions</b>								
Poor neighborhood × hyperactivity					1.09†	0.99–1.20		
Poor neighborhood × exposure							2.09†	1.00–4.40

SES, Socio-economic status; NLE, negative life event; OR, odds ratio; CI, confidence interval.

<sup>a</sup> All models control for cohort and region of residence.

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ , †  $p < 0.10$ .



**Fig. 1.** Predicted odds of (a, b) suicidal thoughts and (c) suicidal attempts as a function of exposure to suicide (non-exposed *versus* exposed), hyperactivity (average level of hyperactivity *versus* 2 s.d. above the mean), and negative life events (NLEs) (average level of NLEs *versus* 2 s.d. above the mean), by neighborhood poverty status (other risk factors were held constant at their average level). ◆---◆, Non-poor neighborhood; ■—■, poor neighborhood.

(Table 3). Neighborhood poverty had an independent effect over and above demographics and late-childhood controls (see Model 1), and the size of the

neighborhood poverty effect was not reduced when late-adolescence risks were added to the model (see Model 2).

**Table 3.** Logistic regressions predicting young adults' suicidal attempts<sup>a</sup>

Measure	Model 1		Model 2		Model 3	
	OR	95% CI	OR	95% CI	OR	95% CI
<b>Demographics</b>						
Male	0.46**	0.28–0.75	0.53*	0.30–0.95	0.57†	0.32–1.02
Age	0.74	0.46–1.19	0.48**	0.28–0.83	0.47**	0.27–0.81
White	1.67	0.59–4.77	2.76	0.80–9.58	3.23†	0.90–11.65
Rural	0.92	0.41–2.03	1.03	0.41–2.54	1.07	0.43–2.66
<b>Late-childhood controls</b>						
SES	1.00	0.71–1.40	0.89	0.56–1.39	0.90	0.57–1.42
Non-intact family	0.78	0.45–1.34	0.35**	0.18–0.68	0.34**	0.17–0.66
Maternal depression	1.04*	1.00–1.07	1.04†	1.00–1.08	1.04†	0.99–1.08
Internalizing symptoms	0.94	0.86–1.02	0.96	0.87–1.06	0.96	0.86–1.06
Hyperactivity/impulsivity	1.23***	1.15–1.31	1.16***	1.07–1.26	1.16***	1.07–1.26
<b>Early- and middle-adolescence neighborhood context</b>						
Poor neighborhood	4.31***	2.60–7.17	5.33***	2.90–9.79	2.41†	0.85–6.81
<b>Late-adolescence risks</b>						
Depression			1.12***	1.08–1.17	1.12***	1.08–1.17
NLEs			1.32*	1.03–1.68	1.05	0.75–1.48
Social support			0.87***	0.80–0.93	0.88***	0.81–0.95
Delinquent activities			1.72	0.61–4.88	1.74	0.60–5.01
Substance abuse						
Alcohol			0.88	0.44–1.76	0.85	0.42–1.73
Marijuana			4.20***	2.28–7.75	4.27***	2.32–7.86
Exposure to suicide			3.21***	1.68–6.12	3.46***	1.83–6.57
<b>Interactions</b>						
Poor neighborhood × NLEs					1.57†	0.97–2.52

SES, Socio-economic status; NLE, negative life event; OR, odds ratio; CI, confidence interval.

<sup>a</sup> All models control for cohort and region of residence.

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ , †  $p < 0.10$ .

For suicide attempts, only one interaction effect, between neighborhood poverty and NLEs, emerged as marginally significant ( $p = 0.06$ ). The main effects of neighborhood poverty and NLEs were appreciably reduced when the interaction effect was included, suggesting that NLEs had a pronounced association with youths' risk of suicide attempts in poor neighborhoods, and that living in a poor neighborhood was especially problematic *vis-à-vis* suicidal behaviors for those experiencing NLEs (see Model 3). Fig. 1c illustrates this finding.

## Discussion

The aim of this study was to examine the association between neighborhood poverty and suicidal thoughts and attempts among older adolescents. Even though neighborhood disadvantage is associated with an array of risk factors for youth suicide (e.g. Leventhal & Brooks-Gunn, 2000), its independent role as a social

risk factor for adolescent suicidal behaviors has not been examined in previous studies. As hypothesized, the results showed that neighborhood poverty was independently associated with suicidal thoughts and attempts, after controlling for pre-existing family and individual vulnerabilities, including family SES, family disruption, maternal depression, internalizing symptoms and hyperactivity/impulsivity problems. These results represent an important addition to the literature, given that previous epidemiological studies looking at the link between neighborhood characteristics and suicidal behaviors did not focus on youth, a group particularly at risk for suicide attempts, and did not include family- or individual-level controls, except for one study conducted among adults that controlled for individual socio-economic characteristics (Cubbin *et al.* 2000).

At the descriptive level, the odds of suicidal ideations were about two times higher in poor than non-poor neighborhoods, and the odds of attempting



suicide were about four times higher in disadvantaged neighborhoods. This increased vulnerability is comparable to that of other groups considered at high risk for suicide, such as Native youth or gay, lesbian and bisexual adolescents, for whom the rates of suicide have been found to be two to six times higher than in the general population (Spirito & Esposito-Smythers, 2006; Kirmayer *et al.* 2007). Even if the processes underlying vulnerability are likely to vary among these groups, the results of this study suggest that youth living in poor neighborhoods are also highly vulnerable and should be a target of youth suicide prevention and intervention efforts.

Another unique contribution of this study is the examination of potential mechanisms for explaining neighborhood effects on suicide. Contrary to expectations, psychosocial risks, including depression, low social support, NLEs, delinquent activities, substance abuse and exposure to suicide, did not explain neighborhood effects. In other words, although adolescents raised in poor neighborhoods tended, as expected, to exhibit higher levels of depression, experience more NLEs and perceive lower levels of social support in late adolescence, these factors did not explain their higher rates of suicidal thoughts and attempts. Other potential mechanisms proposed in the neighborhood literature, such as exposure to violence and abuse, lack of social cohesion in the neighborhood, lower quality relationships with family members or peers, parental criminality, low-quality institutional resources such as school or lack of positive opportunities, need to be examined in future research (Leventhal & Brooks-Gunn, 2000). In addition, considering differential biological responses as a function of neighborhood characteristics could also enhance our understanding of 'neighborhood effects' on youth behavior (Hill *et al.* 2005; Manuck *et al.* 2005).

The results hint at an alternative pathway that might place adolescents raised in poor neighborhood at risk. Marginally significant interaction effects suggested that known risk factors for suicide, namely hyperactivity, exposure to suicide and NLEs, could exert stronger effects among youth living in disadvantaged neighborhoods. Youth who are otherwise at risk for suicide could be more likely to exhibit suicidal thoughts and to attempt suicide when they are exposed daily to a stressful environment that is less likely to provide strong emotional, social and institutional resources in the face of a crisis (Cutrona *et al.* 2005). Thus, a combination of known risk factors for suicide, along with exposure to a challenging environment, might compound the risk of considering suicide as a solution when confronted with a difficult situation. These effects parallel those reported in an emerging literature looking at youth externalizing

problems, illustrating that the effects of important individual risk factors are magnified in disadvantaged neighborhoods (Lynam *et al.* 2000; Obeidallah *et al.* 2004; Dupéré *et al.* 2008). The results suggest that similar amplifying effects might also apply to internalizing problems, although this finding needs replication and should be regarded with caution given the marginal levels of significance of the interactions and the risks inherent in examining person  $\times$  environment interaction effects with a dichotomous outcome (Eaves, 2006).

Other study limitations should be mentioned. First, non-experimental neighborhood studies are subject to selection bias. Indeed, even if major potential confounders are controlled, there is always a possibility that apparent neighborhood effects are attributable to unmeasured characteristics associated with both neighborhood disadvantage and the outcome (Leventhal & Brooks-Gunn, 2000). Some limitations also arise from the use of a large-scale national data set. Large-scale data sets provide statistical power and representativeness but, perhaps inevitably, broadness is often obtained at the cost of depth. For instance, only two questions focused on suicidal thoughts and attempts, so that the specific timing and circumstances surrounding these events were unknown. Furthermore, because of the unequal time-frames (e.g. depressive symptoms were assessed in reference to the past week whereas suicide outcomes were assessed with regard to the past 12 months), temporal precedence and directionality are uncertain. In addition, because the NLSCY was not primarily designed to assess neighborhood effects, hierarchical multi-level modeling strategies could not be used because of insufficient within-neighborhood clustering of participants. For the same reason, potential mechanisms underlying youth suicide attempts were limited to individual-level factors and did not include any potentially relevant neighborhood-level mechanisms, such as exposure to community violence.

Despite these limitations, this study makes a valuable contribution to the literature by examining the link between neighborhood poverty and suicide attempts among youths. The results have potentially important implications for prevention and treatment of vulnerable youth. They suggest that it could prove highly beneficial to implement prevention programs designed to reduce adolescents' suicidal behaviors in disadvantaged communities with high rates of youth suicide. Educational programs for general practitioners working in disadvantaged areas represent one promising avenue (Szanto *et al.* 2007). Another strategy that could reduce the gap between adolescents in poor and non-poor neighborhoods is to improve access to school-based health services. Indeed,

high schools located in lower SES communities are less likely to offer school-based health services (Billy *et al.* 2000), even though the presence of such services are beneficial for youth in high-risk communities (Borowsky *et al.* 1999). School-based prevention programs tested among high-risk groups, such as life skill-training programs, could also prove effective. These programs have been demonstrated to enhance problem-solving strategies and to foster a sense of personal ability to cope with life difficulties (Thompson *et al.* 2001). In short, the prevention and intervention literature suggests many avenues that could significantly reduce the likelihood that adolescents in poor neighborhoods select suicidal behavior as a coping strategy in the face of important life difficulties.

### Acknowledgments

V.D. was supported by fellowships from the Canadian Institutes of Health Research (CIHR) and the Social Sciences and Humanities Research Council of Canada (SSHRC).

The research and analysis in this paper are based on data from Statistics Canada, and the opinions expressed do not represent the views of Statistics Canada.

### Declaration of Interest

None.

### References

- Allison PD (2001). *Missing Data*. Sage: Thousand Oaks, CA.
- Aneshensel CS, Sucoff CA (1996). The neighborhood context of adolescent mental health. *Journal of Health and Social Behavior* **37**, 293–310.
- Billy JOG, Grady WR, Wenzlow AT, Brener ND, Collins JL, Kann L (2000). Contextual influences on school provision of health services. *Journal of Adolescent Health* **27**, 12–24.
- Borowsky IW, Resnick MD, Ireland M, Blum RW (1999). Suicide attempts among American Indian and Alaska Native youth: risk and protective factors. *Archives of Pediatrics and Adolescent Medicine* **153**, 573–580.
- Boyle MH, Offord DR, Hofmann HG, Catlin GP, Byles JA, Cadman DT (1987). Ontario Child Health Study. I. Methodology. *Archives of General Psychiatry* **44**, 826–831.
- Bridge JA, Goldstein TR, Brent DA (2006). Adolescent suicide and suicidal behavior. *Journal of Child Psychology and Psychiatry* **47**, 372–394.
- Cubbin C, LeClere FB, Smith GS (2000). Socioeconomic status and injury mortality: individual and neighborhood determinants. *Journal of Epidemiology and Community Health* **54**, 517–524.
- Cutrona CE (1984). Social support and stress in the transition to parenthood. *Journal of Abnormal Psychology* **93**, 378–390.
- Cutrona CE, Russell DW, Brown PA, Clark LA, Hessling RM, Gardner KA (2005). Neighborhood context, personality, and stressful life events as predictors of depression among African American women. *Journal of Abnormal Psychology* **114**, 3–15.
- Duncan SC, Duncan TE, Strycker LA (2002). A multilevel analysis of neighborhood context and youth alcohol and drug problems. *Prevention Science* **3**, 125–133.
- Dupéré V, Lacourse E, Willms JD, Leventhal T, Tremblay RE (2008). Neighborhood poverty and early transition to sexual activity in young adolescents: a developmental ecological approach. *Child Development* **79**, 805–806.
- Eaves LJ (2006). Genotype × environment interaction in psychopathology: fact or artifact? *Twin Research and Human Genetics* **9**, 1–8.
- Exeter DJ, Boyle PJ (2007). Does young adult suicide cluster geographically in Scotland? *Journal of Epidemiology and Community Health* **61**, 731–736.
- Fergusson DM, Horwood LJ, Ridder EM, Beautrais A (2005). Suicidal behavior in adolescence and subsequent mental health outcomes in young adulthood. *Psychological Medicine* **35**, 983–993.
- Fotti SA, Katz LY, Afifi TO, Cox BJ (2006). The associations between peer and parental relationships and suicidal behaviours in early adolescents. *Canadian Journal of Psychiatry* **51**, 698–703.
- Gould MS, Greenberg T, Velting DM, Shaffer D (2003). Youth suicide risk and preventive interventions: a review of the past 10 years. *Journal of the American Academy of Child and Adolescent Psychiatry* **42**, 386–405.
- Hendin H, Haas AP, Maltzberger JT, Szanto K, Rabinowicz H (2004). Factors contributing to therapists' distress after the suicide of a patient. *American Journal of Psychiatry* **161**, 1442–1446.
- Hill TD, Ross CE, Angel RJ (2005). Neighborhood disorder, psychophysiological distress, and health. *Journal of Health and Social Behavior* **46**, 170–186.
- Jarrett RL (1999). Successful parenting in high risk neighborhoods. *Future of Children* **9**, 45–50.
- Johnson JG, Cohen P, Gould MS, Kasen S, Brown JS, Brook JS (2002). Childhood adversities, interpersonal difficulties, and risk for suicide attempts during late adolescence and early adulthood. *Archives of General Psychiatry* **59**, 741–749.
- Joiner TE, Brown JS, Wingate LR (2005). The psychology and neurobiology of suicidal behavior. *Annual Review of Psychology* **56**, 287–314.
- Kessler RC, Borges G, Walters EE (1999). Prevalence and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Archives of General Psychiatry* **56**, 617–626.
- Kirmayer LJ, Brass GM, Holton T, Paul K, Simpson C, Tait C (2007). Suicide among Aboriginal people in Canada. The Aboriginal Healing Foundation: Ottawa, ON.
- Kohen DE, Brooks-Gunn J, Leventhal T, Hertzman C (2002). Neighborhood income and physical and social disorder in Canada: associations with young children's competencies. *Child Development* **73**, 1844–1860.
- Lacourse E, Nagin D, Tremblay RE, Vitaro F, Claes M (2003). Developmental trajectories of boys' delinquent group membership and facilitation of violent behaviors

- during adolescence. *Development and Psychopathology* **15**, 183–197.
- Lacourse E, Nagin DS, Vitaro F, Côté S, Arseneault L, Tremblay RE** (2006). Prediction of early onset deviant peer group affiliation: a 12-year longitudinal study. *Archives of General Psychiatry* **63**, 562–568.
- Langlois S, Morrison P** (2002). Suicide deaths and suicide attempts. *Health Reports* **13**, 9–22.
- Leventhal T, Brooks-Gunn J** (2000). The neighborhoods they live in: the effects of neighborhood residence on child and adolescent outcomes. *Psychological Bulletin* **126**, 309–337.
- Lynam DR, Caspi A, Moffitt TE, Wikström PH, Loeber R, Novak SP** (2000). The interaction between impulsivity and neighborhood context on offending: the effects of impulsivity are stronger in poorer neighborhoods. *Journal of Abnormal Psychology* **109**, 563–574.
- Manuck SB, Bleil M, Petersen KL, Flory JD, Mann JJ, Ferrell RE, Muldoon MF** (2005). The socio-economic status of communities predicts variation in brain serotonergic responsivity. *Psychological Medicine* **35**, 519–528.
- Middleton N, Sterne JAC, Gunnell D** (2006). The geography of despair among 15–44 year-old men in England and Wales: putting suicide on the map. *Journal of Epidemiology and Community Health* **60**, 1040–1047.
- Middleton N, Whitley E, Frankel S, Dorling D, Sterne J, Gunnell D** (2004). Suicide risk in small areas in England and Wales, 1991–1993. *Social Psychiatry and Psychiatric Epidemiology* **39**, 45–52.
- Miller JR, Piper TM, Ahern J, Tracy M, Tardiff KJ, Vlahov D, Galea S** (2005). Income inequality and risk of suicide in New York city neighborhoods: a multilevel case-control study. *Suicide and Life-Threatening Behavior* **35**, 448–459.
- Molnar BE, Gortmaker SL, Bull FC, Buka SL** (2004). Unsafe to play? Neighborhood disorder and lack of safety predict reduced physical activity among urban children and adolescents. *American Journal of Health Promotion* **18**, 378–386.
- Murphy SA, Johnson LC, Wu L, Fan JJ, Lohan J** (2003). Bereaved parents' outcomes 4 to 60 months after their children's deaths by accident, suicide, or homicide: a comparative study demonstrating differences. *Death Studies* **27**, 39–61.
- Obeidallah D, Brennan RT, Brooks-Gunn J, Earls F** (2004). Links between pubertal timing and neighborhood contexts: implications for girls' violent behavior. *Journal of the American Academy of Child and Adolescent Psychiatry* **43**, 1460–1468.
- Perez-Smith A, Spirito A, Boergers J** (2002). Neighborhood predictors of hopelessness among adolescent suicide attempters: preliminary investigation. *Suicide and Life-Threatening Behavior* **32**, 139–145.
- Pfeffer CR, Jiang H, Kakuma T, Hwang J, Metsch M** (2002). Group intervention for children bereaved by the suicide of a relative. *Journal of the American Academy of Child and Adolescent Psychiatry* **41**, 505–513.
- Poulin C, Hand D, Boudreau B** (2005). Validity of a 12-item version of the CES-D used in the National Longitudinal Study of Children and Youth. *Chronic Diseases in Canada* **26**, 65–72.
- Puderer H** (2001). *Introducing the Dissemination Area for the 2001 Census: An Update*. Geography Working Paper Series. Statistics Canada: Ottawa.
- Radloff LS** (1977). The CES-D scale: a self-report of depression scale for research in the general population. *Applied Psychological Measurement* **1**, 385–401.
- Rehkopf DH, Buka SL** (2005). The association between suicide and the socio-economic characteristics of geographical areas: a systematic review. *Psychological Medicine* **36**, 145–157.
- Reinherz HZ, Tanner JL, Berger SR, Beardslee WR, Fitzmaurice GM** (2006). Adolescent suicidal ideation as predictive of psychopathology, suicidal behavior, and compromised functioning at age 30. *American Journal of Psychiatry* **163**, 1226–1232.
- Ross CE** (2000). Neighborhood disadvantage and adult depression. *Journal of Health and Social Behavior* **41**, 177–187.
- Sampson RJ, Raudenbush SW, Earls F** (1997). Neighborhoods and violent crime: a multilevel study of collective efficacy. *Science* **277**, 918–924.
- Saxe L, Kadushin C, Beveridge A, Livert D, Tighe E, Rindskopf D, Ford J, Brodsky A** (2001). The visibility of illicit drugs: implications for community-based drug control strategies. *American Journal of Public Health* **91**, 1987–1994.
- Spirito A, Esposito-Smythers C** (2006). Attempted and completed suicide in adolescence. *Annual Review of Clinical Psychology* **2**, 237–266.
- Spirito A, Overholser JC (eds)** (2003). *Evaluating and Treating Adolescent Suicide Attempters: From Research to Practice*. Academic Press: San Diego, CA.
- Statistics Canada** (2007). *Microdata User Guide National Longitudinal Survey of Children and Youth Cycle 6, September 2004 to June 2005*. Statistics Canada: Ottawa, ON, Canada.
- Statistics Canada and Human Resources Development Canada** (1995). *National Longitudinal Survey of Children: Overview of Survey Instruments for 1994–95 Data Collection, Cycle 1*. Catalogue no. 89F0078XIE. Statistics Canada and Human Resources Development Canada: Ottawa, ON, Canada.
- Szanto K, Kalmar S, Hendin H, Rihmer Z, Mann JJ** (2007). A suicide prevention program in a region with a very high suicide rate. *Archives of General Psychiatry* **64**, 914–920.
- Thompson EA, Eggert LL, Randell BP, Pike KC** (2001). Evaluation of indicated suicide risk prevention approaches for potential high school dropouts. *American Journal of Public Health* **91**, 742–752.
- Thornberry TP, Krohn MD, Lizotte AJ, Smith CA, Tobin K** (2003). *Gangs and Delinquency in Developmental Perspective*. Cambridge University Press: Cambridge.
- Tremblay RE, Pihl RO, Vitaro F, Dobkin PL** (1994). Predicting early onset of male antisocial behavior from preschool behavior: a test of two personality theories. *Archives of General Psychiatry* **51**, 732–739.
- von Hippel PT** (2007). Regression with missing Ys: an improved strategy for analyzing multiply-imputed data. *Sociological Methodology* **37**, 83–117.
- Weissman MM, Bland RC, Canino GJ, Greenwald S, Hwu HG, Joyce PR, Karam EG, Lee CK, Lellouch J, Lepine JP, Newman SC, Rubio-Stipec M, Wells JE,**

- Wickramaratne PJ, Wittchen HU, Yeh EK** (1999). Prevalence of suicide ideation and suicide attempts in nine countries. *Psychological Medicine* **29**, 9–17.
- Wheaton B, Clarke P** (2003). Space meets time: integrating temporal and contextual influences on mental health in early adulthood. *American Sociological Review* **68**, 680–706.
- Williamson DE, Birmaher B, Frank E, Anderson B, Matty MK, Kupfer DJ** (1998). Nature of life events and difficulties in depressed adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, **37**, 1049–1057.
- Willms JD, Shields M** (1996). *A Measure of Socioeconomic Status for the National Longitudinal Survey of Children and Youth*. Canadian Research Institute for Social Policy: Fredericton, NB.
- Xue Y, Leventhal T, Brooks-Gunn J, Earls F** (2005). Neighborhood residence and mental health problems of 5- to 11-year-olds. *Archives of General Psychiatry* **62**, 554–563.