

# Mesozoic non-marine petroleum source rocks determined by palynomorphs in the Tarim Basin, Xinjiang, northwestern China

DE-XIN JIANG\*, YONG-DONG WANG†‡, ELEANORA I. ROBBINS§,  
JIANG WEI¶ & NING TIAN†

\*Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029,  
People's Republic of China

†Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008,  
People's Republic of China

§U.S. Geological Survey, Reston VA 20192, USA

¶Norfolk State University, Norfolk VA 23504, USA

(Received 30 July 2007; accepted 26 March 2008; First published online 30 July 2008)

**Abstract** – The Tarim Basin in Northwest China hosts petroleum reservoirs of Cambrian, Ordovician, Carboniferous, Triassic, Jurassic, Cretaceous and Tertiary ages. The sedimentary thickness in the basin reaches about 15 km and with an area of 560 000 km<sup>2</sup>, the basin is expected to contain giant oil and gas fields. It is therefore important to determine the ages and depositional environments of the petroleum source rocks. For prospective evaluation and exploration of petroleum, palynological investigations were carried out on 38 crude oil samples collected from 22 petroleum reservoirs in the Tarim Basin and on additionally 56 potential source rock samples from the same basin. In total, 173 species of spores and pollen referred to 80 genera, and 27 species of algae and fungi referred to 16 genera were identified from the non-marine Mesozoic sources. By correlating the palynomorph assemblages in the crude oil samples with those in the potential source rocks, the Triassic and Jurassic petroleum source rocks were identified. Furthermore, the palynofloras in the petroleum provide evidence for interpretation of the depositional environments of the petroleum source rocks. The affinity of the miospores indicates that the petroleum source rocks were formed in swamps in brackish to lacustrine depositional environments under warm and humid climatic conditions. The palynomorphs in the crude oils provide further information about passage and route of petroleum migration, which is significant for interpreting petroleum migration mechanisms. Additionally, the thermal alternation index (TAI) based on miospores indicates that the Triassic and Jurassic deposits in the Tarim Basin are mature petroleum source rocks.

Keywords: spores and pollen, petroleum source rocks, petroleum migration, oil field, Triassic–Jurassic, Tarim Basin.

## 1. Introduction

Petroleum and source rock correlation is a classic tool for source rock identification. Geochemists use chemical biomarkers as indicators to correlate petroleum and source rocks. Palynomorphs can also be used as indicators for petroleum and source rock correlation, because the walls of spores and pollen are resistant to the thermal alteration in the process of petroleum genesis, as well as to the effects of petroleum migration. Moreover, palynomorphs can indicate the geological age and sedimentary environments of source rocks. Consequently, palynology is a useful scientific method in petroleum source research, especially in non-marine sediments.

There are several publications dealing with the identification of oil source rocks. The first was by Sanders (1937), who extracted spores, algae and fungi from Cretaceous and Tertiary crude oil samples from Mexico, and Tertiary crude oil samples from Romania. Waldschmidt (1941) extracted diatom and

plant fragments from Permian crude oils of Colorado, USA. Timofeev & Karimov (1953) made palynological investigations on crude oils of Russia. De Jersey (1965) reported plant microfossils in crude oils of the Moonie oil field in Queensland, Australia. On the other hand, Hunt (1979) documented that spores and pollen are too large to migrate along with liquid hydrocarbons, and he considered palynomorphs in petroleum to be *in situ*, deriving from the reservoir rocks themselves. However, Jiang & Yang (1980) described Cretaceous spores and pollen from crude oils in Tertiary reservoirs and in Silurian metamorphic rocks of the Yumen oil field in the Jiuxi Basin, China. The presence of spores and pollen in the crude oils indicates that these can migrate along with petroleum. Hua & Lin (1989) suggested that microfissures resulting from abnormal pressures occurring in source rocks should constitute the important pathways of petroleum primary migration in the Jiuxi Basin. In addition, Jiang (1990, 1996) found Carboniferous and Permian miospores in crude oils from an igneous rock reservoir of the Junggar Basin, China. The discovery of these spores and pollen in the igneous reservoir may serve as a direct

‡Author for correspondence: ydwang@nigpas.ac.cn, ydwang-67@163.com

indication that they were expelled from source rocks and migrated along with petroleum into the reservoir. McGregor (1996) reviewed studies of palynomorphs in petroleum and considered that these studies merit wider attention, because the results and interpretations of researchers working on this subject have achieved credibility.

The Tarim Basin of Xinjiang, northwest China, is a continental petroliferous basin where some prospective large oil–gas fields have been found. Graham *et al.* (1990) reported analyses of potential petroleum source rocks of the Xinjiang basins, and suggested that the Upper Triassic to Middle Jurassic sequences which were sufficiently buried comprise a potentially significant oil source in the northern Tarim Basin. Hendrix *et al.* (1995) provided a detailed organic geochemical database for organic-rich Lower and Middle Jurassic strata throughout central Xinjiang. They presented field and laboratory evidence demonstrating that organic-rich Lower and Middle Jurassic strata are dominated by terrestrial-derived type III kerogens, and concluded that Jurassic coaly strata have significant potential as petroleum source rocks in the northern Tarim, southern Junggar and Turpan basins. Hanson *et al.* (2000) conducted organic geochemical analyses on a large suite of oils and source rocks extracted from the Tarim Basin. On the basis of statistical cluster analysis, they suggested that most of the oils originated from source rocks deposited in either the Middle–Upper Ordovician or the Upper Triassic to Lower–Middle Jurassic. Based on previous preliminary studies of spores and pollen in crude oils from several petroliferous provinces in the Tarim Basin, Jiang & Yang (1983, 1986, 1992, 1996, 1999) suggested that the Triassic and Jurassic systems should contain favourable petroleum source rocks. This paper addresses a further method of correlation between petroleum and source rocks with palynomorphs from the non-marine Mesozoic deposits, presents analysis of additional material, and discusses the depositional environments of the petroleum source

rocks as well as mechanisms of petroleum migration in the Tarim basin.

## 2. Geological background

The Tarim Basin in the Xinjiang Uygur Autonomous Region of Northwestern China lies between 36 and 42° N latitude, and 74 and 90° E longitude (Fig. 1). The basin is bounded to the north by the Tianshan Mountain Range, to the southwest by the Kunlun Mountain Range, and to the southeast by the Altun Mountain Range. It is a large cratonic basin with a superimposed sedimentary thickness of 13 to 15 km. The Hercynian orogeny of Carboniferous to Permian age resulted in uplift of the Tianshan fold belt and the Kunlun fold belt, as well as evolution of the non-marine sedimentary basin (Zhou & Zheng, 1990).

Mesozoic strata of the Tarim Basin are non-marine, with the exception of Upper Cretaceous shallow-marine strata in the western basin (Zhou & Chen, 1990; Zhou, 2001) (Fig. 2). During Triassic times, lacustrine sedimentary sequences developed in the Kuqa and the North Tarim depressions. The Lower Triassic Ehuobulake Formation consists primarily of greyish brown sandstones and conglomerates intercalated with greyish green and dark grey mudstones, with a total thickness reaching 548 to 592 m. The Middle Triassic Karamay Formation is mainly composed of dark grey, greenish grey and black mudstones and grey siltstones intercalated with greyish brown sandstones with a thickness of 424 to 572 m. The lower Upper Triassic Huangshanjie Formation consists of dark grey mudstones and black carbonaceous mudstones intercalated with greyish green fine-grained sandstones, grey argillo-calcareous rocks and coals; the thickness is 135 to 413 m. The uppermost Triassic Taliqike Formation includes grey sandstones, dark grey and greenish grey mudstones, grey argillo-calcareous rocks, black carbonaceous mudstones and coal beds with a thickness of 545 m to 836 m.




Figure 1. Sketch map showing tectonic subdivisions, and locations of oil fields and boreholes in the Tarim Basin, northwestern China.

Age	Epoch Symbol	Thickness (m)	Reservoirs	Oil fields	Formations (Fm.)/ Groups (Gr.)	Sedimentary facies
Tertiary	N		▲	Kelatu Kekeya		
	E		▲	Yingmaili		
Cretaceous	K <sub>2</sub>	450	▲		Yengisar Gr.	Shallow-marine and lagoon facies
	K <sub>1</sub>	300-1500			Kapushaliang (Kizilsu) Gr.	
Jurassic	J <sub>3</sub>	256-278	▲	Yiqikelike Yakela	Qigu (Kuzigongsu) Fm.	Terrestrial clastic deposits
	J <sub>2</sub>	872-1245	▲		Qiakemake (Taerga) Fm. ◆ Kezilenuer (Yangye) Fm. ◆	Lacustrine and swampy facies Meandering fluvial facies
	J <sub>1</sub>	523-1160	▲		Yangxia (Kangsu) Fm. ◆ Ahe(Shalitashi) Fm.	(Kuqa, North Tarim, Southwest Tarim depressions)
Triassic	T <sub>3</sub>	545-836	▲	Yakela Lunnan	Taliqike Fm. ◆	Braided fluvial facies Lacustrine facies (Kuqa and North Tarim depressions)
		135-413	▲		Huangshanjie Fm. ◆	
	T <sub>2</sub>	424-572	▲		Karamay Fm. ◆	
	T <sub>1</sub>	548-592	▲		Ehuobulake Fm. ◆	
Ordovician	O <sub>2+3</sub>	792	▲	Lunnan		
	O <sub>1</sub>	808	▲	Yakela		

Abbreviations: O<sub>1</sub>-Lower Ordovician; O<sub>2+3</sub>-Middle and Upper Ordovician; T<sub>1</sub>-Lower Triassic; T<sub>2</sub>-Middle Triassic; T<sub>3</sub>-Upper Triassic; J<sub>1</sub>-Lower Jurassic; J<sub>2</sub>-Middle Jurassic; J<sub>3</sub>-Upper Jurassic; K<sub>1</sub>-Lower Cretaceous; K<sub>2</sub>-Upper Cretaceous; E-Eocene; N-Neogene; ▲-Oil reservoir; ◆-Principal petroleum source.

Figure 2. A brief stratigraphic framework showing principal reservoirs, oil fields, major petroleum source rocks and their sedimentary facies in the Tarim Basin, Xinjiang, China.

The Ehuobulake, Karamay, Huangshanjie and Taliqike formations contain various kinds of fossils, including plants, miospores, megaspores, acritarchs, charophytes and conchostracans (Zhou & Chen, 1990; Liu, 2003). The Upper Triassic dark grey and black organic-rich mudstones probably represent lacustrine deposits (Ma & Wen, 1991). In addition, an Upper Triassic braided fluvial facies was reported in the north Tarim basin (Hendrix *et al.* 1992) (Fig. 2).

During the Jurassic period, the lacustrine area further expanded, and both lacustrine and swampy sequences are well developed in the Kuqa Depression, the North Tarim Depression and the Southwest Tarim Depression. The Lower Jurassic Ahe (Shalitashi) and Yangxia (Kangsu) formations consist mainly of grey sandstones and dark grey mudstones intercalated with black carbonaceous mudstones and coals containing plants, miospores, megaspores and acritarchs (Liu, 2003); the thickness varies from 523 to 1160 m. The Middle Jurassic Kezilenuer (Yangye) and Qiakemake (Taerga) formations include grey sandstones, dark grey and greenish grey mudstones, black carbonaceous mud-

stones, coal beds and oil shales containing miospores, megaspores, estherids, ostracods and bivalves; the thickness ranges from 872 to 1245 m (Liu, 2003). Most Lower and Middle Jurassic strata in the basin consist of interbedded sandstone, siltstones, shales and coals, and were interpreted as meandering fluvial facies by Hendrix *et al.* (1995). The Upper Jurassic Qigu Formation consists of brown and brownish red mudstones intercalated with sandstones, and contains charophytes and bivalves, and the thickness is 256 to 278 m (Ma & Wen, 1991) (Fig. 2).

The Lower Cretaceous Kapushaliang (Kizilsu) Group in the Tarim Basin consists of brownish red sandstone and conglomerate intercalated with greyish green siltstones and mudstones, containing ostracods, estherids, charophytes and miospores; the thickness is about 300 to 1500 m (Jiang, He & Dong, 1988; Ma & Wen, 1991; Li, 2000; Jiang *et al.* 2006, 2007). The Upper Cretaceous Yengisar Group in the western basin, dominated by shallow-marine, littoral and lagoonal deposits, and with a thickness of 450 m, carries a rich marine fauna of Tethyan forms (Huang & Chen, 1987;

Ma & Wen, 1991). In addition, dinoflagellate cysts and acritarchs from the Yengisar Group were reported in Xinjiang (Yu & Zhang 1980) (Fig. 2).

It is noteworthy that tectonism in the Tarim Basin created four depressions, three uplifts, several stratigraphic angular unconformities, and many structural traps and faults (Zhou & Zheng, 1990). The depressions are favourable for preservation of organic material and formation of petroleum; the uplifts provide favourable traps for accumulation of petroleum. Unconformable contacts and faults can act as available passages for migration of petroleum. Structural traps within a depression or within an uplift between two depressions are usually the targets of petroleum migration. In fact, commercial oil and gas fields have been found in the Kuqa Depression, the Southwest Tarim Depression, and the North Tarim Uplift located between the Kuqa Depression and the North Tarim Depression (Fig. 1). These depressions can provide sufficient petroleum sources for large oil and gas fields.

### 3. Material and methods

Thirty-eight crude oil samples collected from 22 petroleum reservoirs in seven oil fields in the Tarim Basin were investigated in our study. These oilfields include the Yiqikelike, Yakela, Lunnan, Yingmaili, Kelatu, Maigaiti and Kekeya oil fields (Fig. 1). In addition, 32 rock samples collected from Triassic and Jurassic strata that crop out near Kuqa, Aksu and Kashi, and 24 core samples collected from the boreholes SC-1, S-9, S-14, K-2 and K-6 in the basin (Fig. 1), were used for correlation between petroleum and source rocks.

The method described by Jiang (1990) for extraction of spores and pollen from crude oil samples was adopted in this study to extract palynomorphs, including spores, pollen, algae and fungi from the petroleum. More than five litres of crude oil were used for each sample. The procedure of this method includes oil sample dilution with benzene or gasoline, oil sample filtration in a heater (70–75 °C), insoluble organic matter (kerogen) extraction in a Soxhlet apparatus using benzene ether, ketone and alcohol, and fossil concentration by heavy liquid flotation. The rock samples, including those from outcrops and cores, were prepared by standard methods using 10 % hydrochloric acid, 40 % hydrofluoric acid and 5 % potassium hydroxide. Gravity separation with a cadmium iodide–potassium iodide solution ( $\text{CdI}:\text{KI}:\text{H}_2\text{O} = 10:9:9$ ) was used to concentrate palynomorphs from rocks. All the microspore fossils were mounted in glycerin jelly for study by light microscopy.

Assuming the palynomorphs found in the crude oils and in the rocks have been correctly identified, the palynomorphs recovered from the oils are used to determine the geological age of the rock that provided the sources of the oil. Correlations between palynomorphs in oils and those in rocks have been applied to determine geological ages and stratohorizons of petroleum source rocks. Thermal alteration

index (TAI) based on spore/pollen colour was used to judge the maturity of petroleum source rocks (Traverse, 1988).

### 4. Palynomorphs in the Tarim Basin crude oils

In the Tarim Basin, a total of 173 species of spores and pollen referred to 80 genera, and 27 species of algae and fungi referred to 16 genera, were identified. Most of the palynomorphs in crude oils are Triassic and Jurassic species (Figs 3, 4), and the rest are time-transitional palynomorphs.

#### 4.a. Triassic palynomorphs

Our investigation demonstrates that Triassic palynomorphs are found in crude oils from every petroleum reservoir in the North Tarim Uplift. Sixty species of Triassic spores and pollen are identified in crude oils from the Ordovician, the Triassic and the Jurassic reservoirs of the Yakela oil field and the Lunnan oil field, as well as in the Cretaceous reservoir of the Yakela oil field in the North Tarim Uplift. It is noted that these microspores have previously been reported from the Keuper stage, or upper Triassic Rhaetian stage, or the Triassic in Europe, Australia and in Xinjiang, Shanxi and Yunnan provinces of China (Table 1).

#### 4.b. Jurassic palynomorphs

Jurassic palynomorphs are found in crude oils from the different petroleum reservoirs in the North Tarim Uplift, the Kuqa Depression and the Southwest Tarim Depression. Sixty-two species of Jurassic spores and pollen are found in crude oils from the Ordovician, Triassic, Jurassic, Cretaceous and Palaeogene reservoirs in the north Tarim, and Neogene reservoirs in the Southwest Tarim. These microspores are widely distributed in the Jurassic strata of Eurasia, North America and Australia (Table 2). Some of them were initially documented from the Lower to Middle Jurassic deposits covering different regions; others have been recorded ranging through the Jurassic sequences.

In addition, some fossil fungi, algae and acritarchs are found in crude oils from the Yakela oil field in the North Tarim, and from the Kekeya oil field and the Maigaiti oil field in the Southwest Tarim (Table 3). They are significant for indicating sedimentary environments of the petroleum source rocks.

### 5. Identification of petroleum source rocks in the Tarim Basin

The microspores extracted from crude oils usually form a three-part assemblage that represents the source bed, carrier bed and reservoir bed, each of which is different in geological age. The reservoir rocks of an oil field are always known, so spores and pollen deriving from the reservoir bed itself can be easily separated






Figure 3. Triassic and Jurassic miospores in crude oils from boreholes L-1, S-9, S-14, and K-6 in the Tarim Basin (see Fig. 1 for borehole location). (a) *Cyathidites australis* Couper (no. L1-146); (b) *Cyathidites minor* Couper (no. L1-16); (c) *Dictyophyllidites harrisii* Couper (no. K6-26); (d) *Undulatisporites pflugii* Pocock (no. S14-124); (e) *Concavisporites toralis* (Leschik) Nilsson (no. L1-4); (f) *Granulatisporites jurassicus* Pocock (no. L1-26); (g) *Apiculatisporis parvispinosus* (Leschik) Qu (no. S14-168); (h, i) *Apiculatisporis spiniger* (Leschik) Qu (h, no. L1-6; i, no. S14-80); (j) *Duplexisporites amplexiformis* (Kara-Murza) Playford & Dettmann (no. L1-23); (k) *Duplexisporites anagrammensis* (Kara-Murza) Playford & Dettmann (no. L1-169); (l) *Duplexisporites scanicus* (Nilsson) Playford & Dettmann (no. L1-1); (m) *Lycopodiumsporites subrotundum* (Kara-Murza) Pocock (no. L1-92); (n) *Lycopodiumsporites paniculatooides* Tralau (no. L1-105); (o) *Lycopodiacidites rhaeticus* Schulz (no. S9-21); (p, q) *Lycopodiacidites kuepperi* Klaus (p, no. S14-50; q, no. S14-40); (r, s) *Lophotriteles corrugatus* Ouyang & Li (r, no. L1-29; s, no. L1-1); (t) *Retusotriteles mesozoicus* Klaus (no. S9-32); (u) *Lundbladisporea subornata* Ouyang & Li (no. S9-22); (v) *Limatulasporites parvus* Qu & Wang (no. S14-103); (w) *Limatulasporites dalongkouensis* Qu & Wang (no. S14-64); (x) *Tigrisporites halleinis* Klaus (no. S14-101); (y) *Zebbrasporites kahleri* Klaus (no. S14-53); (z) *Verrucosporites contactus* Clarke (no. S14-40); (a1) *Verrucosporites remyanus* Madler (no. L1-111); (b1) *Aratrisporites fischeri* (Klaus) Playford & Dettmann (no. L1-1).






Figure 4. Triassic and Jurassic miospores in crude oils from Boreholes L-1, S-9, S-14, K-2, K-6 and Y-1 in the Tarim Basin (see Fig. 1 for borehole location). (a) *Aratrisporites scabratus* Klaus (no. S14-61); (b) *Aratrisporites paenulatus* Playford & Dettmann (no. L1-1); (c, d) *Aratrisporites granulatus* (Klaus) Playford & Dettmann (c, no. S14-119; d, no. S14-46); (e) *Aratrisporites strigosus* Playford (no. S14-144); (f) *Cycadopites typicus* (Mal.) Pocock (no. S14-134); (g) *Cerebropollenites carlylensis* Pocock (no. Y1-11); (h) *Aratrisporites fischeri* (Klaus) Playford & Dettman (no. L1-165); (i) *Aratrisporites tenuispinosus* Playford (no. L1-41); (j, k) *Aratrisporites parvispinosus* Leschik (j, no. L1-92; k, no. L1-191); (l) *Cycadopites nitidus* (Balme) Pocock (no. S14-3); (m) *Chordasporites orientalis* Ouyang & Li (no. L1-50); (n) *Taeniaesporites pellucidus* (Goubin) Balme (no. S9-9); (o) *Bennettitaeapollenites lucifer* (Thierg.) Potonié (no. L1-20); (p) *Chordasporites singulichorda* Klaus (no. S14-58); (q) *Piceites pseudorotundiformis* (Mal.) Pocock (no. L1-15); (r) *Quadraeculina limbata* Maljavkina (no. Y1-9); (s) *Protopinus scanicus* Nilsson (no. Y1-34); (t) *Cedripites minor* Pocock (no. K6-10); (u) *Cycadopites subgranulosus* (Couper) Clarke (no. S14-131); (v) *Podocarpidites florinii* Pocock (no. K2-4).

Table 1. Triassic spores and pollen in crude oils from reservoirs of different ages in the Tarim Basin and their distribution in the Triassic strata of Europe, Asia and Australia

Spores and pollen	Regions				GB	DE	CH	AT	RO	CN	AU
	References				[1]	[2]	[3]	[4]	[5]	[6]	[7]
	Count of specimens in reservoirs										
	O	T	J	K	T <sub>3</sub>	T	T <sub>3</sub>	T <sub>3</sub>	T	T	T
<b>Spores</b>											
<i>Apiculatisporis globosus</i> (Leschik) Playford & Dettmann 1965	4	6	2				*			—	—
<i>A. parvispinosus</i> (Leschik) Qu 1980	5	6	4				*			—	
<i>A. spiniger</i> (Leschik) Qu 1980	6	9	3	1			*			—	
<i>Aratrisporites coryliseminis</i> Klaus 1960	6	4	2					*		—	
<i>A. fischeri</i> (Klaus) Playford & Dettman 1965	5	8						*		—	—
<i>A. granulatus</i> (Klaus) Playford & Dettman 1965	9	7	4	3				*		—	—
<i>A. paenulatus</i> Playford & Dettmann 1965	4	4								—	*
<i>A. parvispinosus</i> Klaus 1960		3						*		—	
<i>A. parvispinosus</i> Leschik 1955	2	5					*			—	
<i>A. scabratus</i> Klaus 1960	7	9	4	2				*		—	
<i>A. strigosus</i> Playford 1965	8	14	3	3						—	*
<i>A. tenuispinosus</i> Playford 1965		5								—	*
<i>Asseretospora gyrata</i> (Playf. & Dettm.) Schuurman 1977	3	8	3	2						—	*
<i>Calamospora nathorstii</i> (Halle) Klaus 1960	7	5	4	2	*			—		—	
<i>C. tener</i> (Leschik) Mädler 1964	3	6	2	1		—		*		—	
<i>Camarozonosporites rudis</i> (Leschik) Klaus 1960		7						*		—	
<i>Conbaculatisporites mesozoicus</i> Klaus 1960	3							*		—	
<i>Limatulasporites dalongkouensis</i> Qu & Wang 1986	6	8	3							*	
<i>L. parvus</i> Qu & Wang 1986	9	11	4	2						*	
<i>Lophotriletes corrugatus</i> Ouyang & Li 1980		5								*	
<i>Lundbladisporea nejburgii</i> Schulz 1964	3	5	2			*			—	—	
<i>L. playfordi</i> Balme 1963		3								—	*
<i>L. plicata</i> Bai 1983	2	4								—	*
<i>L. subornata</i> Ouyang & Li 1980	3	7	2							*	
<i>Lycopodiacidites kuepperi</i> Klaus 1960	7	4	2	1				*		—	
<i>L. rhaeticus</i> Schulz 1967	2	5	2	2		*				—	
<i>Multinodisporites junctus</i> Ouyang & Li 1980		3	1							*	
<i>Osmundacidites alpinus</i> Klaus 1960	4	6	5	2				*		—	
<i>Punctatisporites ambiguus</i> Leschik 1955	3	4						*		—	
<i>P. microtumulosus</i> Playford & Dettmann 1965	4	4								—	*
<i>P. triassicus</i> Schulz 1964	12	15	4	3		*			—	—	
<i>Retusotriletes arcticus</i> Qu & Wang 1986		6	2	1						*	
<i>R. mesozoicus</i> Klaus 1960	5	4	3	3				*		—	
<i>Tigrisporites halleinis</i> Klaus 1960	5							*		—	
<i>Verrucosisporites contactus</i> Clarke 1965	7				*					—	
<i>V. remyanus</i> Mädler 1964		5				*				—	
<i>Zebrasporites kahleri</i> Klaus 1960	5							*		—	
<b>Pollen</b>											
<i>Alisporites aequalis</i> Mädler 1964		4				*				—	
<i>A. australis</i> de Jersey 1962	3	3	4							—	*
<i>A. fusiformis</i> Ouyang & Li 1980		4								*	
<i>A. parvus</i> de Jersey 1962	3	5	5	3						—	*
<i>Cedripites parvisaccus</i> Ouyang & Li 1980	8	5	4	2						*	
<i>Chordasporites impensus</i> Ouyang & Li 1980		3								*	
<i>C. orientalis</i> Ouyang & Li 1980		4	3							*	
<i>C. singulichorda</i> Klaus 1960	4	5	2	1				*		—	
<i>Colpectopollis pseudostratus</i> (Kopytova) Qu & Wang 1986		9	4	4						*	
<i>C. scitulus</i> (Qu & Pu) Qu & Wang 1986		6	2							*	
<i>Enzonalasporites tenuis</i> Leschik 1955	4	2						*		—	
<i>E. vigenis</i> Leschik 1955	5		1					*		—	
<i>Lueckisporites triassicus</i> Clarke 1965	7	8			*					—	
<i>Minutosaccus parvus</i> Qu & Wang 1986	4	6	2							*	
<i>Parcisporites rarus</i> Ouyang & Li 1980	5	5	3	1						*	
<i>P. solutus</i> Leschik 1955	4	3						*		—	
<i>Pinuspollenites normalis</i> Qu & Wang 1986	7	4	3	2						*	
<i>Platysaccus undulatus</i> Ouyang & Li 1980		5	2	1						*	
<i>Podocarpidites queenslandi</i> (deJersey) Qu 1980		7								—	*

Table 1. (Cont.)

	Regions				GB	DE	CH	AT	RO	CN	AU
	References				[1]	[2]	[3]	[4]	[5]	[6]	[7]
	Count of specimens in reservoirs										
Spores and pollen	O	T	J	K	T <sub>3</sub>	T	T <sub>3</sub>	T <sub>3</sub>	T	T	T
<i>P. radialis</i> (Leschik) Qu 1984	5	3					*			–	
<i>Taeniaesporites divisus</i> Qu 1982		5								*	
<i>T. kraeuseli</i> Leschik 1955	5	7					*			–	
<i>T. rhaeticus</i> Schulz 1967	4	4	3			*				–	
<b>Total</b>	217	312	99	42							

Abbreviations: O – Ordovician; T – Triassic; T<sub>3</sub> – Upper Triassic; J – Jurassic; K – Cretaceous; GB – Great Britain; DE – Germany, CH – Switzerland; AT – Austria; RO – Romania; CN – China; AU – Australia.

‘\*’ indicates the original stratigraphic horizon of a taxon that was first described in this locality; ‘–’ indicates other stratigraphic ranges of this taxon that were subsequently described (but not the first record).

References: [1] Clarke, 1965; Batten & Coppelhus, 1996; [2] Schulz, 1964; Mädlar, 1964; [3] Leschik, 1955; [4] Klaus, 1960; [5] Venkatachala, Beju & Kar, 1968; [6] Yang *et al.* 1986; Qu, 1982; Ouyang & Li, 1980; Liu, 2003; [7] De Jersey, 1962; Balme, 1963; Playford, 1965.

Table 2. Jurassic spores and pollen in crude oils from reservoirs of different ages in the Tarim Basin and their distribution in the Jurassic strata of Eurasia, North America, Australia and New Zealand

	Regions						GB	SE	RU	CN	CA	AU	NZ
	References						[1]	[2]	[3]	[4]	[5]	[6]	[7]
	Count of specimens in reservoirs												
Spores and pollen	O	T	J	K	E	N	J <sub>1-2</sub>	J <sub>1-2</sub>	J	J	J	J	J
<b>Spores</b>													
<i>Apiculatisporis ovalis</i> (Nilsson) Norris 1967						11		*		–	–		
<i>A. variabilis</i> Pocock 1970		3	3		2	5				–	*		
<i>Asseretospora amplexiformis</i> Qu et Wang 1990		7	4					*		–		–	
<i>A. anagrammensis</i> Liu 2000		5	3						*			–	
<i>A. scanicus</i> Huang 1993		5	3	2				*				–	
<i>Cibotiumspora paradoxa</i> (Mal.) Chang 1965	4	5	7	4	3	8			*				
<i>Concavisporites toralis</i> (Leschik) Nilsson 1958		4	2					*					
<i>Cyathidites australis</i> Couper 1953	4	6	5	2	2	9	–						*
<i>C. minor</i> Couper 1953	8	9	7	5	7	24	–						*
<i>Deltoidospora gradata</i> (Mal.) Pocock 1970	3		6	3	3	8			*				
<i>D. lineata</i> (Bolch.) Pocock 1970						5			*				
<i>D. perpusilla</i> (Bolch.) Pocock 1970	5	4	7	3	2	14			*				
<i>Dictyophyllidites harrisii</i> Couper 1958	7	5	6	4	5	21							
<i>Gleicheniidites conflexus</i> (Chln.) Xu & Zhang 1980		6	4	3	2	4	*						
<i>G. nilssonii</i> Pocock 1970		5	3									*	
<i>G. rousei</i> Pocock 1970		4	3	3	1	17						*	
<i>Granulatisporites jurassicus</i> Pocock 1970		2	5	2	3	4					*		
<i>G. minor</i> de Jersey 1959		3	4	3	2	4						*	
<i>Klukisporites variegatus</i> Couper 1958						7	*						
<i>Leptolepidites major</i> Couper 1958						4	*						
<i>L. verrucatus</i> Couper 1953						4							*
<i>Lycopodiumsporites paniculatooides</i> Tralau 1968		5	2			3		*					
<i>L. subrotundus</i> (Kara-Murza) Pocock 1970		6	3			4			*		–		
<i>Marattisporites scabratus</i> Couper 1958	7		5		1		*						
<i>Murospora jurassica</i> Pocock 1970						5						*	
<i>M. minor</i> Pocock 1970						7						*	
<i>Osmundacidites wellmanii</i> Couper 1953	8	8	14	6	4	19	–						*
<i>Todisporites major</i> Couper 1958						7	*						
<i>Undulatisporites concavus</i> Kedves 1971 # <sup>1</sup>						7							
<i>U. pflugii</i> Pocock 1970	5	6	4	3								*	
<b>Pollen</b>													
<i>Alisporites lowoodensis</i> de Jersey 1963						4							*
<i>Bennettiteapollenites lucifer</i> (Thierg.) Potonie 1958 # <sup>2</sup>	2	5	5	3		7							
<i>Callialasporites dampieri</i> (Balme) Dev 1961			2										*
<i>C. minus</i> (Tralau) Guy 1971		2	3					*					



Table 2. (Cont.)

	Regions						GB	SE	RU	CN	CA	AU	NZ
	References						[1]	[2]	[3]	[4]	[5]	[6]	[7]
	Count of specimens in reservoirs												
Spores and pollen	O	T	J	K	E	N	J <sub>1-2</sub>	J <sub>1-2</sub>	J	J	J	J	J
<i>Cedripites minor</i> Pocock 1970	5	4	7	3	2	9				–	*		
<i>Cerebropollenites carlylensis</i> Pocock 1970						4				–	*		
<i>Chasmatosporites elegans</i> Nilsson 1958						8		*		–			
<i>C. major</i> Nilsson 1958						7		*		–			
<i>C. minor</i> Nilsson 1958						9		*		–			
<i>Cycadopites minimus</i> (Cookson) Pocock 1970 # <sup>3</sup>			7	8	4	14			–	–	–		
<i>C. nitidus</i> (Balme) Pocock 1970	9	8	14	5	4	17				–	–	*	
<i>C. subgranulosus</i> (Couper) Clarke 1965	7	3	9	3		16	*			–			
<i>C. typicus</i> (Mal.) Pocock 1970	4		8	4	5	12			*	–	–		
<i>Paleoconiferus asaccatus</i> Bolchovitina 1956						6			*	–	–		
<i>Parvisaccites enigmatus</i> Couper 1958			4	1		7	*			–			
<i>Piceites expositus</i> Bolchovitina 1956	2	4	5	2	3	6			*	–			
<i>P. pseudorotundiformis</i> (Mal.) Pocock 1970		5	3	2		4			*	–	–		
<i>Pityosporites parvisaccatus</i> de Jersey 1959		4	2							–			*
<i>Platysaccus lopsinensis</i> (Mal.) Pocock 1970						7			*	–	–		
<i>Podocarpidites florinii</i> Pocock 1970						5				–	*		
<i>P. langii</i> Pocock 1970						5				–	*		
<i>P. multicus</i> (Bolch.) Pocock 1970	3	2	3	2	1	7			*	–	–		
<i>P. rousei</i> Pocock 1970			4			4				–	*		
<i>P. unicus</i> (Bolch.) Pocock 1970	2		3		1	5			*	–	–		
<i>P. wapellensis</i> Pocock 1970						4				–	*		
<i>Protopicea exilioides</i> (Bolch.) Pocock 1970						4			*	–	–		
<i>Protopinus scanicus</i> Nilsson 1958						5		*		–			
<i>Quadraeculina limbata</i> Maljavkina 1949			3	1	2	5			*	–			
<i>Vitreisporites itunensis</i> Pocock 1970						3				–	*		
<i>V. jansonii</i> Pocock 1970						5				–	*		
<i>V. jurassicus</i> Pocock 1970						7				–	*		
<i>V. shouldicei</i> Pocock 1970						4				–	*		
<b>Total</b>	85	135	182	77	59	401							

Abbreviations: O – Ordovician; T – Triassic; J<sub>1-2</sub> – Lower and Middle Jurassic; J – Jurassic; K – Cretaceous; E – Eocene; N – Neogene; GB – Great Britain; SE – Sweden; RU – Russia; CN – China; CA – Canada; AU – Australia; NZ – New Zealand.

References: [1] Couper, 1958; [2] Nilsson 1958; Tralau, 1968; [3] Maljavkina, 1949; Bolchovitina, 1956; [4] Sun, 1989; Huang, 1995; Liu, 1982, 2003; Jiang & Wang, 2002; [5] Pocock, 1970; [6] Balme, 1963; De Jersey, 1959, 1963; [7] Couper, 1953.

\* indicates the original stratigraphic horizon of a taxon that was first described in this locality; ‘–’ indicates other stratigraphic ranges of this taxon that were subsequently described (but not the first record).

#<sup>1</sup> Hungary (Potonié, 1958); #<sup>2</sup> Germany (Potonié, 1958); #<sup>3</sup> Madagascar (Cookson, 1947).

from the three-part assemblage. The remainders of the assemblage are indicative for source and carrier beds. Generally, the oldest miospores in the assemblage indicate the source rocks, those of intermediate age indicate the carrier beds, and the youngest indicate the reservoir rocks (e.g. Jiang, 1988, 1990, 1991). Sometimes the three parts of the assemblage are coeval, indicating that the source rocks, carrier beds and reservoir rocks belong to the same formation (Jiang, 1988). Although the geological circumstances are often complicated, petroleum source rocks can be distinguished from non-source rocks by correlation of the palynological assemblages in crude oils and their supposed source rocks.

Triassic miospores in crude oils from the North Tarim Uplift are common in the Triassic deposit of the Tarim Basin. Twenty-one species of spores and pollen found in oils are also identified in the dark grey and black mudstone of the Lower Triassic Ehuobulake Formation; 37 miospore species identified in petroleum samples are found in the Middle Triassic Karamay

Formation, and 44 species in petroleum samples are found in the dark grey and black mudstone of the Upper Triassic Huangshanjie and Taliqike formations (Table 4).

Jurassic miospores in crude oils from the North Tarim Uplift, the Kuqa Depression and the Southwest Tarim Depression are also common representatives in the Lower Jurassic and Middle Jurassic deposits of the basin. Forty-five species extracted from oils are also found in the dark grey and black mudstone of the Lower Jurassic Ahe and Yangxia formations, the Middle Jurassic Kezilenuer and Qiakemake formations in the North Tarim Uplift and the Kuqa Depression; 60 species found in oils are recorded in the dark grey and black mudstone of the Lower Jurassic Shalitashi and Kangsu formations, as well as the Middle Jurassic Yangye and Taerga formations in the Southwest Tarim Depression (Table 5).

The miospore assemblages identified in the crude oils are similar to those from the rocks of the Triassic and Jurassic successions. This indicates that

Table 3. Fungi, algae and acritarchs in crude oils of the Tarim Basin

Fungi, algae and acritarchs	North Tarim Yakela Oilfield	Southwest Tarim	
		Kekeya Oilfield	Maigaiti Oilfield
<b>Fungi</b>			
<i>Multicellaesporites ovatus</i> Sheffy & Dilcher 1971		+	
<i>M. pachydermus</i> Ke & Shi 1978		+	
<i>M. margaritus</i> Ke & Shi 1978			+
<i>Glomus</i> sp.			+
<b>Algae</b>			
Pyrrhophyta			
<i>Conicoidium</i> cf. <i>granorugosum</i> Jiabo 1978	+		
<i>C.</i> sp.	+		
<i>Dinogymnium granulatum</i> Jiabo 1978		+	
<i>Prominangularia</i> cf. <i>dongyingensis</i> Jiabo 1978		+	
<i>Rhombodella</i> cf. <i>baculata</i> Jiabo 1978		+	
<i>R.</i> cf. <i>variabilis</i> Jiabo 1978		+	
<i>R.</i> sp.		+	
<i>Tenua</i> cf. <i>bellula</i> Jiabo 1978	++	+	
<i>T.</i> sp.	++		
Chlorophyta			
<i>Campenia</i> cf. <i>circellata</i> Jiabo 1978	++		+
<i>Hungarodiscus</i> cf. <i>punctatus</i> Jiabo 1978	++		
<i>H.</i> sp.	++		
<b>Acritarchs</b>			
<i>Dictyotidium</i> cf. <i>asperatum</i> Jiabo 1978	++		
<i>D.</i> sp.	++		
<i>Granodiscus</i> cf. <i>granulatus</i> Mädlar 1963			+
<i>Granoreticella</i> sp.	++		
<i>Heliospermopsis</i> sp.	+		
<i>Porusphaera</i> sp.	++		+
<i>Tectocorpidium</i> cf. <i>rimosum</i> Jiabo 1978	+		
<i>Verrucosphaera</i> cf. <i>verrucosa</i> Jiabo 1978	+		
Acritarcha indet. Type 1	+		
Acritarcha indet. Type 2	+		
Acritarcha indet. Type 3	+		

See Figure 1 for locations. + rare, ++ common.

the palynomorphs in the crude oils were derived from Triassic and Jurassic plants, were deposited and released from these formations during primary migration, and migrated into different reservoirs during secondary migration. The correlation results between oils and rocks with palynomorphs suggest that the Triassic and Lower to Middle Jurassic deposits are important contributors to the petroleum source in the Tarim Basin (Fig. 2).

The colour of the Triassic and Jurassic miospores is dark orange to brown, both in the petroleum and in the rocks. The thermal alteration index (TAI) based on colour of spore/pollen ranges between 2.8 and 3.4. This thermal maturity belongs to the mature main phase of liquid petroleum generation (Traverse, 1988), and consequently the Triassic and Jurassic petroleum source rocks of the Tarim Basin belong to the mature source rock type. These results are supported by the evidence from organic geochemical analyses made on crude oils and their putative source rocks in the Tarim Basin. Based on the bulk geochemical analyses and correlation between crude oils and their supposed source rock extracts, it was concluded that Mesozoic strata, particularly Lower and Middle Jurassic strata deposits, comprise a potentially significant non-marine petroleum source sequence.  $T_{max}$  value of 429 °C to 449 °C and vitrinite reflectance ( $R_o$ ) values of 0.47 % to 0.67 % indicate that these rocks are at or just below

the threshold of oil generation, and samples collected within depocentre sections average higher  $R_o$  values ( $R_o = 0.75$  % for one Middle Jurassic rock sample) (Graham *et al.* 1990; Hendrix *et al.* 1995). Analyses of coals and organic-rich shales show a dominance of terrestrial, higher plant components. Visual kerogen analysis indicates that vitrinite, inertinite and exinite are the dominant minerals, and elemental analysis characterizes most kerogen as type III (Hendrix *et al.* 1995). Pyrolysis–gas chromatography results show the prominence of alkene/alkane doublets, and suggest that the Jurassic strata are capable of liquid hydrocarbon generation. Biomarker correlations show that three-quarters of the petroleum samples are consistent with derivation from the Jurassic strata. The high Pr/Ph ratios for most extracts and oil samples (generally > 2.5) are consistent with a higher plant-dominated non-marine environment.

### 6. Environment of petroleum source rocks

Additional data have been accumulated for improving our understanding of the potential botanical relationships of major dispersed miospore taxa, based upon *in situ* spores of the Mesozoic plants as well as their extant relatives. The closest relatives to the miospores identified in this study include a broad range of plants (Table 6). The extant relatives to most

Table 4. Distribution of important Triassic miospore taxa in crude oils of the Triassic strata in the Tarim Basin

Spores and pollen	Ehuobulake Fm. (T <sub>1</sub> )	Karamay Fm. (T <sub>2</sub> )	Huangshanjie Fm. (T <sub>3</sub> <sup>1</sup> )	Taliqike Fm. (T <sub>3</sub> <sup>2</sup> )
<b>Spores</b>				
<i>Apiculatisporis globosus</i> (Leschik) Playford & Dettmann 1965		+	++	++
<i>A. parvispinosus</i> (Leschik) Qu 1980		++	++	++
<i>A. spiniger</i> (Leschik) Qu 1980	+	++	++	++
<i>Aratrisporites coryliseminis</i> Klaus 1960		+	++	++
<i>A. fischeri</i> (Klaus) Playford & Dettman 1965		+	++	+
<i>A. granulatus</i> (Klaus) Playford & Dettman 1965	+	++	++	++
<i>A. paenulatus</i> Playford & Dettmann 1965		+	++	+
<i>A. parvispinosus</i> Leschik 1955			+	++
<i>A. paraspinosus</i> Klaus 1960			+	++
<i>A. scabratus</i> Klaus 1960	+	++	++	++
<i>A. strigosus</i> Playford 1965		++	++	++
<i>A. tenuispinosus</i> Playford 1965		+	+	++
<i>Asseretospora gyrata</i> (Playf. & Dettm.) Schuurman 1977			++	++
<i>Calamospora nathorstii</i> (Halle) Klaus 1960	+	++	++	++
<i>C. tener</i> (Leschik) Mädlar 1964			++	++
<i>Camarozonosporites rudis</i> (Leschik) Klaus 1960			+	++
<i>Conbaculatisporites mesozoicus</i> Klaus 1960		+	++	++
<i>Limatulasporites dalongkouensis</i> Qu & Wang 1986	++			
<i>L. parvus</i> Qu & Wang 1986	++	+		
<i>Lophotriletes corrugatus</i> Ouyang & Li 1980	++			
<i>Lundbladispora neiburgii</i> Schulz 1964	+	++		
<i>L. playfordi</i> Balme 1963	++			
<i>L. plicata</i> Bai 1983	+	++		
<i>L. subornata</i> Ouyang & Li 1980	++			
<i>Lycopodiacidites kuepperi</i> Klaus 1960			++	++
<i>L. rhaeticus</i> Schulz 1967			++	++
<i>Multinodisporites junctus</i> Ouyang & Li 1980	++	+		
<i>Osmundacidites alpinus</i> Klaus 1960			++	++
<i>Punctatisporites ambiguus</i> Leschik 1955			++	+
<i>P. microtumulosus</i> Playford & Dettmann 1965		+		++
<i>P. triassicus</i> Schulz 1964	+	++	+	+
<i>Retusotriletes arcticus</i> Qu & Wang 1986	++	+		
<i>R. mesozoicus</i> Klaus 1960		+	++	++
<i>Tigrisporites halleinis</i> Klaus 1960		+	++	+
<i>Verrucosisporites contactus</i> Clarke 1965		+	++	++
<i>V. remyanus</i> Mädlar 1964		+	+	++
<i>Zebbrasporites kahleri</i> Klaus 1960			+	++
<b>Pollen</b>				
<i>Alisporites aequalis</i> Mädlar 1964			++	++
<i>A. australis</i> de Jersey 1962		++	++	++
<i>A. fusiformis</i> Ouyang & Li 1980	++			
<i>A. parvus</i> de Jersey 1962		++	++	++
<i>Cedripites parvisaccus</i> Ouyang & Li 1980	++	+		
<i>Chordasporites impensus</i> Ouyang & Li 1980	++			
<i>C. orientalis</i> Ouyang & Li 1980	++			
<i>C. singulichorda</i> Klaus 1960		+	++	+
<i>Colpectopollis pseudostriatum</i> (Kopytova) Qu & Wang 1986		++	++	++
<i>C. scitulus</i> (Qu & Pu) Qu & Wang 1986		++	++	++
<i>Enzonalsporites tenuis</i> Leschik 1955			++	++
<i>E. vigenis</i> Leschik 1955			+	++
<i>Lueckisporites triassicus</i> Clarke 1965		+	++	++
<i>Minutosaccus parvus</i> Qu & Wang 1986		++	+	+
<i>Parcisporites rarus</i> Ouyang & Li 1980	++	+		
<i>P. solutus</i> Leschik 1955			+	++
<i>Pinuspollenites normalis</i> Qu & Wang 1986		++	++	+
<i>Platysaccus undulatus</i> Ouyang & Li 1980	++			
<i>Podocarpidites queenslandi</i> (deJersey) Qu 1980		++	++	++
<i>P. radialis</i> (Leschik) Qu 1984		+	++	+
<i>Taeniaesporites divisus</i> Qu 1982	++	+		
<i>T. kraeuseli</i> Leschik 1955		+	+	++
<i>T. rhaeticus</i> Schulz 1967			++	++

+ rare, ++ common.

Abbreviations: Fm – Formation; T<sub>1</sub> – Lower Triassic; T<sub>2</sub> – Middle Triassic; T<sub>3</sub><sup>1</sup> – lower Upper Triassic; T<sub>3</sub><sup>2</sup> – middle Upper Triassic  
For references to authors of taxon, see Tables 1 and 2 and reference list.



Table 5. Distribution of important Jurassic miospore taxa in crude oils of the Jurassic strata in the Tarim Basin

Spores and pollen	North Tarim				Southwest Tarim			
	Ahe Fm. (J <sub>1</sub> <sup>1</sup> )	Yangxia Fm. (J <sub>1</sub> <sup>2</sup> )	Kezilenuer Fm. (J <sub>2</sub> <sup>1</sup> )	Qiakemake Fm. (J <sub>2</sub> <sup>2</sup> )	Shalitashi Fm. (J <sub>1</sub> <sup>1</sup> )	Kangsu Fm. (J <sub>1</sub> <sup>2</sup> )	Yangye Fm. (J <sub>2</sub> <sup>1</sup> )	Taerga Fm. (J <sub>2</sub> <sup>2</sup> )
<b>Spores</b>								
<i>Apiculatisporis ovalis</i> (Nilsson) Norris 1967							++	++
<i>A. variabilis</i> Pocock 1970			++				++	+
<i>Cibotiumspora paradoxa</i> (Mal.) Chang 1965	+	++	++	+	+	++	++	++
<i>Concavisporites toralis</i> (Leschik) Nilsson 1958		++	++			+	+	
<i>Cyathidites australis</i> Couper 1953	+	++	++	++	+	++	++	++
<i>C. minor</i> Couper 1953	++	+++	+++	+++	++	+++	+++	+++
<i>Deltoidospora gradata</i> (Mal.) Pocock 1970			+	+		+	++	++
<i>D. lineata</i> (Bolch.) Pocock 1970							++	+
<i>D. perpusilla</i> (Bolch.) Pocock 1970	+		++	++		+	++	++
<i>Dictyophyllidites harrisii</i> Couper 1958	+	++	++	++		+++	+++	++
<i>Duplexisporites amplexiformis</i> (Kara-Murza) Playford & Dettmann 1965		++	++	+				
<i>D. anagrammensis</i> (Kara-Murza) Playford & Dettmann 1965			++	+			+	
<i>D. scanicus</i> (Nilsson) Playford & Dettmann 1965			++	+				
<i>Gleicheniidites conflexus</i> (Chln.) Xu & Zhang 1980			++	++			++	++
<i>G. nilssonii</i> Pocock 1970			++	++			+	
<i>G. rousei</i> Pocock 1970			++	++			++	+
<i>Granulatisporites jurassicus</i> Pocock 1970		++	++	++		++	++	++
<i>G. minor</i> de Jersey 1959	+	++			+	++		
<i>Klukisporites variegatus</i> Couper 1958		+	++	+		+	++	+
<i>Leptolepidites major</i> Couper 1958							++	++
<i>L. verrucatus</i> Couper 1953						+	++	+
<i>Lycopodiumsporites paniculatoides</i> Tralau 1968			++	++			+	+
<i>L. subrotundus</i> (Kara-Murza) Pocock 1970		++	++	+		+	+	+
<i>Marattisporites scabratus</i> Couper 1958			++	++			++	++
<i>Murospora jurassica</i> Pocock 1970							++	+
<i>M. minor</i> 1970							++	+
<i>Osmundacidites wellmanii</i> Couper 1953	+	++	++	++	+	++	++	++
<i>Todisporites major</i> Couper 1958							++	+
<i>Undulatisporites concavus</i> Kedves 1971							++	+
<i>U. pflugii</i> Pocock 1970			++				+	
<b>Pollen</b>								
<i>Alisporites lowoodensis</i> de Jersey 1963		++				+	+	
<i>Bennettiteapollenites lucifer</i> (Thierg.) Potonie 1958			++	++			++	++
<i>Callialasporites dampieri</i> (Balme) Dev 1961			+	+			++	+
<i>C. minus</i> (Tralau) Guy 1971			++	++			++	+
<i>Cedripites minor</i> Pocock 1970		++	++	++		++	++	++
<i>Cerebropollenites carlylensis</i> Pocock 1970			+	+			++	++
<i>Chasmatosporites elegans</i> Nilsson 1958		+				++	+	
<i>C. major</i> Nilsson 1958		+				++		
<i>C. minor</i> Nilsson 1958						++	+	
<i>Cycadopites minimus</i> (Cookson) Pocock 1970			+	+			++	++
<i>C. nitidus</i> (Balme) Pocock 1970	+	++	++	++	+	++	++	++
<i>C. subgranulosus</i> (Couper) Clarke 1965		++	++			++	++	+

Table 5. (Cont.)

Spores and pollen	North Tarim				Southwest Tarim			
	Ahe Fm. (J <sub>1</sub> <sup>1</sup> )	Yangxia Fm. (J <sub>1</sub> <sup>2</sup> )	Kezilenuer Fm. (J <sub>2</sub> <sup>1</sup> )	Qiakemake Fm. (J <sub>2</sub> <sup>2</sup> )	Shalitashi Fm. (J <sub>1</sub> <sup>1</sup> )	Kangsu Fm. (J <sub>1</sub> <sup>2</sup> )	Yangye Fm. (J <sub>2</sub> <sup>1</sup> )	Taerga Fm. (J <sub>2</sub> <sup>2</sup> )
<i>C. typicus</i> (Mal.) Pocock 1970	+	++	++	++		++	++	++
<i>Paleoconiferus asaccatus</i> Bolchovitina 1956		+	+	+		+	++	+
<i>Parvisaccites enigmatus</i> Couper 1958			+	+			++	+
<i>Piceites expositus</i> Bolchovitina 1956			++	++			++	++
<i>P. pseudorotundiformis</i> (Mal.) Pocock 1970	+	++	++	++		+	+	+
<i>Pityosporites parvisaccatus</i> de Jersey 1959	+	++				+		
<i>Platysaccus lopsinensis</i> (Mal.) Pocock 1970							++	+
<i>Podocarpidites florinii</i> Pocock 1970							++	+
<i>P. langii</i> Pocock 1970							+	++
<i>P. multicus</i> (Bolch.) Pocock 1970	+	++	++			++	++	
<i>P. rousei</i> Pocock 1970				+			++	++
<i>P. unicus</i> (Bolch.) Pocock 1970			+	+			++	+
<i>P. wapellensis</i> Pocock 1970							+	++
<i>Protopinus scanicus</i> Nilsson 1958		++	+			++	++	
<i>Protopicea exilioides</i> (Bolch.) Pocock 1970			+	+			++	++
<i>Quadraeculina limbata</i> Maljavkina 1949		++	++	+		+	++	++
<i>Vitreisporites itunensis</i> Pocock 1970			++	++			++	+
<i>V. jansonii</i> Pocock 1970							++	
<i>V. jurassicus</i> Pocock 1970							++	+
<i>V. shouldicei</i> Pocock 1970							++	

+ rare, ++ common, +++ abundant.

Abbreviations: Fm. – Formation; J<sub>1</sub><sup>1</sup> – lower Lower Jurassic; J<sub>1</sub><sup>2</sup> – middle Lower Jurassic; J<sub>2</sub><sup>1</sup> – lower Middle Jurassic; J<sub>2</sub><sup>2</sup> – middle Middle Jurassic.

Table 6. Botanical relationships of major dispersed spore/pollen genera identified in crude oils of the Tarim Basin

Pteridophyta	Gymnospermae
<b>Lycopodiaceae</b>	<b>Pteridospermopsida</b>
<i>Aratrisporites</i> (Leschik) Klaus	Caytoniales
<i>Camazonosporites</i> (Potonié) Klaus	<i>Vitreisporites</i> (Leschik) Jansonius
<i>Lycopodiumsporites</i> Thiergart	Incertae sedis
<i>Lundbladispore</i> Balme	<i>Alisporites</i> Daugherty
<b>Filicopsida</b>	<b>Cycadopsida</b>
Marattiaceae	Cycadales or Ginkgoales
<i>Marattisporites</i> Couper	<i>Cycadopites</i> (Wodehouse) Wilson & Webster
Osmundaceae	Bennettiales (?)
<i>Osmundacidites</i> Couper	<i>Bennettiteapollenites</i> Thiergart
<i>Todisporites</i> Couper	Incertae sedis
Schizaeaceae	<i>Chasmatosporites</i> Nilsson
<i>Klukisporites</i> Couper	<b>Coniferopsida</b>
Gleicheniaceae	Podocarpaceae
<i>Gleicheniidites</i> (Ross) Delcourt et Sprumont	<i>Parvisaccites</i> Couper
Cyatheaceae	<i>Platysaccus</i> (Naumova) Potonié & Kremp
<i>Cyathidites</i> Couper	<i>Podocarpidites</i> (Cookson) Potonié
<i>Deltoidospora</i> (Miner) Potonié	Pinaceae
Dicksoniaceae	<i>Cedripites</i> Wodehouse
<i>Cibotiumspora</i> Chang	<i>Piceapollenites</i> Potonié
<i>Cyathidites</i> Couper	<i>Piceites</i> Bolchovitina
Dipteridaceae or Cheiroleuriaceae or Matoniaceae	<i>Pinuspollenites</i> Raatz
<i>Concavisporites</i> (Pflug) Delcourt & Sprumont	Araucariaceae
<i>Dictyophyllidites</i> Couper	<i>Callialasporites</i> Dev
<i>Granulatisporites</i> (Ibrahim) Potonié & Kremp	

This summary is based upon comprehensive results of the *in situ* spore studies of the Mesozoic plants and their living relatives based on major references: e.g. Couper, 1958; Nilsson, 1958; Potonié, 1962; Townrow, 1962; Chang, 1965; Helby & Martin, 1965; Grauvogel-Stamm, 1978; Van Konijnenburg-Van Cittert, 1971, 1975, 1978, 1981, 1989, 1993; Litwin, 1985; Traverse, 1988; Balme, 1995; Wang, 1999, 2002; Wang & Mei, 1999; Wang *et al.* 2001; Abbink, 1998 and the studies on extant spores and pollen from living plants: e.g. Zhang *et al.* 1976; Wang *et al.* 1995.

of these plants are humidogene thermophytes. For instance, lycophytes grow on acidic soils in humid climates; the marattialean ferns are large and tall plants, presently growing in the tropical or subtropical forests; the tree ferns (represented by *Cyathidites*, for example, in the spore record) are growing in temperate-tropical humid areas; and several ground fern taxa (Table 6) are distributed in the tropical and subtropical swamp/marsh lands as understory vegetation. The cycads are typical thermophytes, and so are several of the conifer taxa identified in the miospore assemblages. A case analysis on the Jurassic rock palynomorphs, their affinities, vegetation reconstruction and climatic implications was carried out in the Qaidam Basin, a giant petroliferous Mesozoic basin near the Tarim Basin, northwest China (Wang, Mosbrugger & Zhang, 2005). In summary, the ecological characteristics of the parent plants to which the spores and pollen in this study belong suggest that they grow in warm and humid climate conditions.

The algae in crude oils of the Tarim Basin (Table 3) are informative for interpreting the depositional environments of petroleum source rocks. Pyrrhophyta algae are related to the marine environment, and dinoflagellates usually indicate marine conditions. Chlorophyte algae are mostly produced in freshwater bodies, and *Pediastrum* indicates typical freshwater conditions. Neither dinoflagellates nor *Pediastrum* are found in the petroleum or source rocks of the basin, therefore the depositional environments of the petroleum source rocks are supposed to be neither brines of the marine environment nor typical freshwater. The ecological conditions reflected by the palynology show that the source rocks were probably formed in swamps in brackish to lacustrine environments during warm and humid climatic conditions.

The odd-carbon-chain n-paraffin and olefins are synthesized by marine plants primarily in the C<sub>15</sub> to C<sub>21</sub> range and by land plants primarily in the C<sub>27</sub> to C<sub>35</sub> range. Brackish-water plants synthesize in the intermediate range C<sub>19</sub> to C<sub>27</sub> (Hunt, 1979). Hendrix *et al.* (1995) reported that the dominant n-alkane is either n-C<sub>21</sub>, n-C<sub>23</sub>, or n-C<sub>25</sub>, and a slight to pronounced odd-over-even preference (OEP) is present in the Jurassic rock sample from the Tarim Basin. The dominance of n-alkane (n-C<sub>21</sub> to n-C<sub>25</sub>) suggests that the source rocks were deposited in a brackish-water sedimentary environment.

## 7. Mechanisms of petroleum migration as shown by palynomorphs

Spore/pollen grains which were originally buried in the sediments have contributed their waxy, fatty and oily secretions to the formation of petroleum, leaving only their decay-resistant remains. These fossil spores and pollen could migrate along with liquid and gaseous hydrocarbons; some of them enter petroleum accumulations, and provide information about passage,

direction and route of petroleum migration (Jiang, 1991; Jiang & Yang, 1994, 1999).

The primary migration means the migration of original petroleum from the petroleum source bed. Original petroleum could likely not exit through source rock pore networks, because the pore diameters of petroleum source rocks are generally less than 0.01  $\mu\text{m}$ , and the oil droplet diameters are usually bigger than 1  $\mu\text{m}$  (Tissot & Welte, 1978; Li, 2004). Palynomorphs in petroleum are generally larger than 15  $\mu\text{m}$  in diameter and the pores of source rocks are too small for their passage. Therefore, the presence of miospores derived from petroleum source rocks in crude oils suggests that the possible pathways of petroleum primary migration could be via microfissures in the source rocks, not via pore space. Microfissures formed by abnormal high pressure during the process of diagenesis are common, and such microfissures are presumably available for initial migration and expulsion of petroleum (Tissot & Welte, 1978; Roehl, 1981; Hua & Lin, 1989; Li, 2004). The width of the microfissures is generally less than 100  $\mu\text{m}$  (Li, 2004). Hua & Lin (1989) reported that microfissures fields with bitumen can be observed under microscope in the Upper Jurassic to Lower Cretaceous mudstone from the Jiuxi Basin, China. This is evidence for microfissures serving as pathways of petroleum primary migration. Fossil miospores have the ability to pass through various migration pathways, as they are very thin and flexible. When the hydrocarbon fluid power is strong, the fossil spores and pollen can be pressed into wrinkles and pass through narrow pathways, and subsequently recover their original state when the space around them increases; this flexibility can be observed under the microscope (Jiang & Yang, 1992).

Pore networks associated with secondary migration, such as connected porous openings, interstratified openings, joint fissures, fault fissures and unconformity surfaces, all provide an avenue for migration of expelled spores and pollen away from the source rock. Larger structures (such as faults and unconformities) are also well developed in the Tarim Basin and as previously outlined, and crude oil samples collected from the wells near faults or unconformity surfaces contain numerous spores and pollen.

The phase state of petroleum migration depends in essence on the passages of petroleum migration. Because microfissures are wide enough for the passage of miospores, the passageways must be unblocked for the passage of oil droplets. It follows that the migration of petroleum in the liquid phase is fully possible in the course of primary migration. Liquid phase migration is also common in the course of secondary migration, because the passageways generally are much wider than microfissures.

The palynological results indicate that the routes of petroleum migration in the Tarim Basin mainly are from the Triassic or Jurassic petroleum source rocks to the different petroleum reservoirs, such as the Ordovician, Triassic, Jurassic, Cretaceous and Tertiary



Table 7. Directions and routes of petroleum migration in the Tarim Basin

Oil fields/ Reservoirs	Source beds	Reservoir beds	Migration directions	Migration routes
Kelatu	J	N	Vertical	J→N
Kekeya	J	N	Vertical	J→N
Yiqikelike K	J	K	Vertical	J→K
Yiqikelike J	J	J	Lateral	J→J
Yingmaili	J	E	Vertical	J→E
Yakela K	J	K	Vertical	J→K
	T		Vertical	T→K
Yakela J	J	J	Lateral	J→J
(Lunnan J)	T		Vertical	T→J
Yakela T	J	T	Vertical	J→T
(Lunnan T)	T		Lateral	T→T
Yakela O	J	O	Vertical /Lateral	J→O
(Lunnan O)	T		Vertical /Lateral	T→O

Abbreviations: O – Ordovician; T – Triassic; J – Jurassic; K – Cretaceous; E – Eocene; N – Neogene.

reservoirs. Both vertical and lateral migrations are important in the course of petroleum migration in the Tarim Basin. Importantly, structural deformation could complicate the theory that the occurrence of older palynomorphs in younger reservoir strata requires vertical secondary migration. If structural deformation juxtaposes source rocks and reservoir rocks laterally, lateral migration may be more important. The later juxtaposition of source rocks and reservoir rocks as a consequence of structural deformation, such as Triassic or Jurassic source rocks and Ordovician reservoir rocks, is a possible scenario in the Tarim Basin. Based on the study of palynomorphs in petroleum, the directions and routes of petroleum migration are summarized in Table 7. The petroleum migration routes were complicated in the northern Tarim Basin and relatively simple in the southwestern Tarim Basin.

## 8. Conclusions

The following conclusions may be drawn from the present study:

- (1) The palynomorphs identified from crude oils and rocks in the Triassic and Jurassic of the Tarim Basin provide informative evidence for determining the potential petroleum source rocks. The results of this investigation indicate that dark-coloured argillaceous rocks of the Lower Triassic Ehuobulake Formation, the Middle Triassic Karamay Formation, the Upper Triassic Huangshanjie and Taliqike formations, the Lower Jurassic Yangxia Formation, and the Middle Jurassic Kezilenuer and Qiakemake formations are the probable petroleum source rocks in the northern Tarim Basin. The dark-coloured argillaceous rocks of the Lower Jurassic Kangsu Formation and the Middle Jurassic Yangye and Taerga formations are the probable petroleum source rocks in the southwestern Tarim Basin.
- (2) The thermal alteration index (TAI) based on colour of spore/pollen indicates that the Triassic and Jurassic dark-coloured argillaceous rocks in

the Tarim Basin are mature petroleum source rocks. This conclusion is supported by the results of organic geochemical analyses. The vitrinite reflectance ( $R_o$ ) of the Jurassic rocks from the northern Tarim depocentre reaches 0.75 %, within the oil window.

- (3) The botanical affinities of the spores and pollen identified in crude oils of the Tarim Basin include mosses, ferns, cycads and conifers, and most of these plants prefer warm and humid climates. The ecological characteristics of the palaeoflora indicate that the Triassic and Jurassic petroleum source rocks were formed in brackish to lacustrine swamps during warm and humid climate conditions.
- (4) Judging from the palynomorphs, it may be concluded that microfissures formed by abnormal high pressure during the diagenesis and catagenesis of petroleum source rocks could provide pathways for the primary migration of petroleum. Faults, unconformity surfaces, joints and other fissures could provide passages for the secondary migration of petroleum. The main direction of petroleum migration could be represented by either vertical migration or lateral migration for different reservoir types, and the routes of petroleum migration would be determined by different source beds and different reservoir beds.

**Acknowledgements.** This study was supported by the National Basic Research Program of China (grant no. 2006CB701401), National Science Foundation of China (nos 40472004, 40632010). The authors would like to thank Prof. Yang H. Q. for technical assistance, Prof. He Z. S. and Mr Dong K. L. for providing geological information and some samples, and Mr Sun F., Ms Du J. E., and Ms Lai C. Y. for collecting and preparing samples. Special thanks are due to Prof. Marc Hendrix (University of Montana, USA), Prof. Vivi Vajda (University of Lund, Sweden) and an anonymous referee for providing valuable suggestions and references, which greatly improved the manuscript. This is a contribution to International Geoscience Program – IGCP 506.

## References

- ABBINK, O. A. 1998. *Palynological investigations in the Jurassic of the North Sea region*. LPP Contributions Series no. 8. Utrecht: LPP Foundation, 192 pp.
- BALME, B. E. 1963. Plant microfossils from the Lower Triassic of western Australia. *Palaeontology* **6**, 12–40.
- BALME, B. E. 1995. Fossil in situ spores and pollen grains, an annotated catalogue. *Review of Palaeobotany and Palynology* **87**, 81–323.
- BATTEN, D. J. & COPPELHUS, E. B. 1996. Biostratigraphic significance of uppermost Triassic and Jurassic miospores in northwest Europe. In *Palynology, Principles and Applications*, vol. 2 (eds J. Jansonius & D. C. McGregor), pp. 795–806. Dallas, Texas: American Association of Stratigraphic Palynologists Foundation.
- BOLCHOVITINA, N. A. 1956. Atlas of spores and pollen from the Jurassic and Lower Cretaceous deposits of the Vilyui depression. *Transactions of the Institute of Geology, Academy of Sciences, USSR* **2**, 1–188, pls 1–25 (in Russian).
- CHANG, L. J. 1965. Sporo-pollen assemblage from Yima coal-bearing formation of Mianchi County of Henan Province and its significance. *Acta Palaeontologica Sinica* **13**, 160–96, pls 1–7 (in Chinese with English abstract).
- CLARKE, R. F. A. 1965. Keuper miospores from Worcestershire, England. *Palaeontology* **8**, 294–321, pls 35–9.
- COOKSON, I. C. 1947. Plant microfossils from the Lignites of Kerguelen Archipelago. *B.A.N.Z. Antarctic Research Expedition 1929–1931, Reports Series A 2*, 127–42.
- COUPER, R. A. 1953. Upper Mesozoic and Cenozoic spores and pollen grains from New Zealand. *New Zealand Geological Survey Palaeontological Bulletin* **22**, 1–77.
- COUPER, R. A. 1958. British Mesozoic microspores and pollen grains. A systematic and stratigraphic study. *Palaeontographica B* **103**, 75–179, pls 15–31.
- DE JERSEY, N. J. 1959. Jurassic spores and pollen grains from the Rosewood coalfield. *Queensland Government Mining Journal* **60**, 344–66, pls 1–3.
- DE JERSEY, N. J. 1962. Triassic spores and pollen grains from the Ipswich coalfield. *Geological Survey of Queensland Publication* **307**, 1–18, pls 1–6.
- DE JERSEY, N. J. 1963. Jurassic spores and pollen grains from the Marburg sandstone. *Geological Survey of Queensland Publication* **313**, 1–13, pls 1–3.
- DE JERSEY, N. J. 1965. Plant microfossils in some Queensland crude oil samples. *Geological Survey of Queensland Publication* **329**, 1–9.
- GRAHAM, S. A., BRASSELL, S., CARROLL, A. R., XIAO, X., DEMAISON, G., MCKNIGHT, C. L., LIANG, Y., CHU, J. & HENDRIX, M. S. 1990. Characteristics of selected petroleum source rocks, Xinjiang Uygur Autonomous Region, northwest China. *American Association of Petroleum Geologists Bulletin* **74**, 493–512.
- GRAUVOGEL-STAMM, L. 1978. La flore du Grès a Voltzia (Buntsandstein Supérieur) des Vosges du Nord (France). Morphologie, anatomie, interprétations phylogénique et paléogéographique. *Univ. Louis Pasteur Strasbourg Institute Géologie Science Géologie Mémorie* **50**, 1–225.
- HANSON, A. D., ZHANG, S. C., MOLDOWAN, J. M., LIANG, D. G. & ZHANG, B. M. 2000. Molecular organic geochemistry of the Tarim basin, northwest China. *American Association of Petroleum Geologists Bulletin* **84**, 1109–28.
- HELBY, R. & MARTIN, A. R. H. 1965. *Cylostrobos* gen. nov., cones of lycopsidean plants from the Narrabeen Group (Triassic) of New South Wales. *Australian Journal of Botany* **13**, 389–404.
- HENDRIX, M. S., BRASSELL, S. C., CARROLL, A. R. & GRAHAM, S. A. 1995. Sedimentology, organic geochemistry, and petroleum potential of Jurassic coal measures, Tarim, Junggar, and Turpan basins, northwest China. *American Association of Petroleum Geologists Bulletin* **79**, 929–59.
- HENDRIX, M. S., GRAHAM, S. A., CARROLL, A. R., MCKNIGHT, C. L., SCHULEIN, B. J. & WANG, Z. 1992. Sedimentary record and climatic implications of recurrent deformation in the Tian Shan: Evidence from Mesozoic strata of the north Tarim, south Junggar, and Turpan basins, northwest China. *Geological Society of America Bulletin* **104**, 53–79.
- HUA, B. Q. & LIN, X. X. 1989. Discussion on some problems of oil migration in Jiuxi Basin. *Acta Sedimentologica Sinica* **7**, 39–47 (in Chinese with English abstract).
- HUANG, P. 1995. Early–Middle Jurassic sporopollen assemblages from Dananhu coalfield of Tuha Basin, Xinjiang and their stratigraphical significance. *Acta Palaeontologica Sinica* **34**, 171–93, pls 1–4 (in Chinese with English abstract).
- HUANG, J. Q. & CHEN, B. W. 1987. *The Evolution of Tethys in China and Adjacent Regions*. Beijing: Geological Publishing House, 109 pp.
- HUNT, J. M. 1979. *Petroleum Geochemistry and Geology*. San Francisco: W. H. Freeman and Company, 617 pp.
- JIABO. 1978. On the Paleogene dinoflagellates and acritarches from the coastal region of Bohai. *Nanjing Institute of Geology and Palaeontology, Academia Sinica, Beijing: Science Press*, 1–190 (in Chinese with English abstract).
- JIANG, D. X. 1988. Spores and pollen in oils as indicators of lacustrine source rocks. In *Lacustrine Petroleum Source Rocks* (eds A. J. Fleet, K. Kelts & M. R. Talbot), pp. 159–69. Geological Society of London, Special Publication no. 40.
- JIANG, D. X. 1990. Palynological evidence for identification of nonmarine petroleum source rocks, China. *Ore Geology Reviews* **5**, 553–75, pls 1–6.
- JIANG, D. X. 1991. Fossil spores and pollen in petroleum and their significance. *Chinese Journal of Botany* **3**, 62–76, pls 1–4.
- JIANG, D. X. 1996. Fossil pollen and spores in crude oil from an igneous reservoir. In *Palynology, Principles and Applications*, vol. 3 (eds J. Jansonius & D. C. McGregor), pp. 1123–8. Dallas, Texas: American Association of Stratigraphic Palynologists Foundation.
- JIANG, D. X., HE, Z. S. & DONG, K. L. 1988. Early Cretaceous palynofloras from Tarim Basin, Xinjiang. *Acta Botanica Sinica* **30**, 430–40, pls 1–3 (in Chinese with English abstract).
- JIANG, D. X., WANG, Y. D., HE, Z. S. & DONG, K. L., NI, Q. & TIAN, N. 2006. Early Cretaceous palynofloras from the Kizilsu Group in the Tarim Basin, Xinjiang. *Acta Micropalaeontologica Sinica* **23**, 371–91 (in Chinese with English abstract).
- JIANG, D. X., WANG, Y. D., HE, Z. S. & DONG, K. L. 2007. Early Cretaceous Sporopollen assemblage from the Shuangshanhe Formation in Baicheng Area of the Tarim Basin, Xinjiang. *Acta Micropalaeontologica Sinica* **24**, 247–60 (in Chinese with English abstract).
- JIANG, D. X. & WANG, Y. D. 2002. Middle Jurassic spore-pollen assemblage from the Yanan Formation of Dongsheng, Nei Monggol, China. *Acta Botanica Sinica* **44**, 230–8, pls 1–2.

- JIANG, D. X. & YANG, H. Q. 1980. Petroleum spore-pollen assemblages and oil source rock of Yumen oil-bearing region in Gansu. *Acta Botanica Sinica* **22**, 280–5, pls 1–2 (in Chinese with English abstract).
- JIANG, D. X. & YANG, H. Q. 1983. Petroleum spore-pollen assemblages and oil source rock of Kuche Seg in Xinjiang. *Acta Botanica Sinica* **25**, 179–86, pl. 1 (in Chinese with English abstract).
- JIANG, D. X. & YANG, H. Q. 1986. Petroleum spore-pollen assemblages and oil source rocks of Yecheng Seg in Xinjiang. *Acta Botanica Sinica* **28**, 111–16, pls 1–3 (in Chinese with English abstract).
- JIANG, D. X. & YANG, H. Q. 1992. Spores and pollen in crude oils and petroleum source of Tarim Basin. *Science in China (Series B)* **35**, 1005–12, pls 1–2.
- JIANG, D. X. & YANG, H. Q. 1994. Spores and pollen in oil from igneous rock petroleum pool and petroleum origin of Junggar Basin. *Science in China (Series B)* **37**, 1499–1505, pls 1–2.
- JIANG, D. X. & YANG, H. Q. 1996. Spores and pollen from crude oil of Kashi Depression, Xinjiang. *Acta Botanica Sinica* **38**, 809–13, pls 1–2 (in Chinese with English abstract).
- JIANG, D. X. & YANG, H. Q. 1999. Petroleum spore-pollen assemblages and petroleum source rocks of North Tarim Upheaval in Xinjiang. *Acta Botanica Sinica* **41**, 213–18, pls 1–2 (in Chinese with English abstract).
- KE & SHI. (Scientific Research Institute of Petroleum Exploration and Development and Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences), 1978. *Early Tertiary Spores and Pollen grains from the Coastal Region of Bohai*. Beijing: Science Press, 1–164 (in Chinese with English abstract).
- KLAUS, W. 1960. Sporen der karnischen Stufe der ostalpinen Trias. *Jahrbuch Geologie Bundesanstalt (Austria)* **5**, 107–84, pls 28–38.
- LESCHIK, G. 1955. Die Keuper Flora von Neuwelt bei Basel. II. Die Iso- und Mikrosporen. *Schweizerische Paläontologische Abhandlungen* **72**, 1–70.
- LI, M. C. 2004. *Petroleum and Natural Gas Migration*. Beijing: Petroleum Industry Press, 350 pp. (in Chinese with English abstract).
- LI, W. B. 2000. Early Cretaceous palynoflora from northern Tarim basin. *Acta Palaeontologica Sinica* **39**, 28–45, pls 1–6 (in Chinese with English abstract).
- LITWIN, R. J. 1985. Fertile organs and *in situ* spores of ferns from the Late Triassic Chinle Formation of Arizona and New Mexico, with discussion of the associated dispersed spores. *Review of Palaeobotany and Palynology* **44**, 101–46.
- LIU, Z. S. 1982. Early and Middle Jurassic spore-pollen assemblages from the Shiguai coal-field of Baotou, Nei Monggol. *Acta Palaeontologica Sinica* **21**, 371–9, pls 1–2 (in Chinese with English abstract).
- LIU, Z. S. 2003. Triassic and Jurassic spore-pollen assemblages from the Kuqa Depression, Tarim Basin of Xinjiang, NW China. *Palaeontologia Sinica, New Series A* **190**, 1–244, pls 1–40.
- MA, B. L. & WEN, C. Q. 1991. *Formation and evolution of sedimentary rocks and oil-gas in Tarim Basin*. Beijing: Science Press, 195 pp. (in Chinese with English abstract).
- MÄDLER, K. 1964. Bemerkenswerte Sporenformen aus dem Keuper und unterem Lias. *Fortschritte in der Geologie von Rheinland und Westfalen* **12**, 169–200.
- MALJAVKINA, V. S. 1949. Identification of spores and pollen of the Jurassic and Cretaceous. *Trudy VNIGRI, N.S.* **33**, 1–138, pls 1–51 (in Russian).
- MCGREGOR, D. C. 1996. Palynomorphs in petroleum and formation water, A review. In *Palynology, Principles and Applications, vol. 3* (eds J. Jansonius & D. C. McGregor), pp. 1115–21. Dallas, Texas: American Association of Stratigraphic Palynologists Foundation.
- NILSSON, T. 1958. Über das Vorkommen eines mesozoischen Sapropelgesteins in Schonen. *Lunds Universitets Årsskrift, Avd 2*, **54**, 5–112.
- OUYANG, S. & LI, Z. P. 1980. Microflora from the Kayitou Formation of Fuyan district, E. Yunnan and its bearing on stratigraphy and palaeobotany. In *Stratigraphy and Palaeontology of the Late Permian coal measures of western Guizhou and eastern Yunnan*, pp. 123–83. Beijing: Science Press (in Chinese with English abstract).
- PLAYFORD, G. 1965. Plant microfossils from Triassic sediments near Poatina, Tasmania. *Journal of Geological Society of Australia* **12**, 173–210.
- POCOCK, S. A. J. 1970. Palynology of the Jurassic sediments of western Canada. *Palaeontographica B* **130**, 12–136, pls 5–26.
- POTONIÉ, R. 1958. Synopsis der Gattungen der Sporae dispersae. Teil II. *Geologisches Jahrbuch Beiheft* **31**, 1–114.
- POTONIÉ, R. 1962. Synopsis der Sporae *in situ*. *Beiheft zum Geologischen Jahrbuch* **H 52**, 1–204.
- QU, L. 1982. The palynological assemblage from the Liujiagou Formation of Jiaocheng, Shanxi. *Bulletin of Geological Institute, Chinese Academy of Geological Society* **4**, 83–93, pls 1–2 (in Chinese with English abstract).
- ROEHL, P. O. 1981. Dilation brecciation, proposed mechanism of fracturing, petroleum expulsion, and dolomitization in Monterey Formation, California. *American Association of Petroleum Geologists Bulletin* **65**, 980–1.
- SANDERS, J. M. 1937. The microscopical examination of crude petroleum. *Journal of Institute of Petroleum Technology* **23**, 525–73.
- SCHULZ, E. 1964. Sporen und pollen aus dem mittleren Buntsandstein des germanischen Beckens. *Monatsbericht der Deutscher Akademischer Wissenschaft zu Berlin* **6**, 597–606.
- SCHULZ, E. 1967. Sporenpaläontologische untersuchungen rotliassischer Schichten im Zentralteil des germanischen Beckens. *Palaeontologische Abhandlungen B* **2**, 541–633.
- SHEFFY, M. V. & DILCHER, D. L. 1971. Morphology and taxonomy of fungal spores. *Palaeontographica B* **133**, 34–51.
- SUN, F. 1989. Early and Middle Jurassic spore-pollen assemblages of Qiquanhu coal-field of Turpan, Xinjiang. *Acta Botanica Sinica* **31**, 638–46, pls 1–2.
- TIMOFEEV, B. V. & KARIMOV, A. K. 1953. Spores and pollen in mineral oil. *D. A. N. USSR* **92**, 151–2 (in Russian).
- TISSOT, B. P. & WELTE, D. H. 1978. *Petroleum Formation and Occurrence*. New York, Berlin: Springer-Verlag, 314 pp.
- TOWNROW, J. A. 1962. On some disaccate pollen grains of Permian to middle Jurassic age. *Grana Palynologica* **3**, 13–44.
- TRALAU, H. 1968. Botanical investigation into the fossil flora of Eriksdal in Fyledalen, Scania. II. The middle Jurassic microflora. *Bulletin of Geological Survey of Sweden* **633**, 1–132, pls 1–26.
- TRAVERSE, A. 1988. *Paleopalynology*. Boston: Unwin Hyman, 600 pp.
- VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1971. *In situ* gymnosperm pollen from the Middle Jurassic of Yorkshire. *Acta Botanica Neerlandica* **20**, 1–96.



- VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1975. Some notes on *Marattia anglica* from the Jurassic of Yorkshire. *Review of Palaeobotany and Palynology* **20**, 205–14.
- VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1978. Osmundaceous spores in situ from the Middle Jurassic of Yorkshire, England. *Review of Palaeobotany and Palynology* **26**, 125–41.
- VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1981. Schizaeaceous spores in situ from the Jurassic of Yorkshire, England. *Review of Palaeobotany and Palynology* **33**, 169–81.
- VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1989. Dicksoniaceae spores in situ from the Jurassic of Yorkshire, England. *Review of Palaeobotany and Palynology* **61**, 273–301.
- VAN KONIJNENBURG-VAN CITTERT, J. H. A. 1993. A review of the Matoniaceae based on in situ spores. *Review of Palaeobotany and Palynology* **78**, 235–67.
- VENKATACHALA, B. S., BEJU, D. & KAR, R. K. 1968. Palynological evidence on the presence of Lower Triassic in the Danubean (Moesian) Platform, Rumania. *Palaeobotanist* **16**, 29–37.
- WALDSCHMIDT, W. A. 1941. Progress report on microscopic examination of Permian crude oils. *American Association of Petroleum Geologists Bulletin* **25**, 934.
- WANG, F. H., CHIAN, N. F., ZHANG, Y. L. & YANG, H. Q. 1995. *Pollen Flora of China*. Beijing: Science Press, 461 pp. (in Chinese with English abstract).
- WANG, Y. D. 1999. Fertile organs and in situ spores of *Marattia asiatica* (Kawasaki) Harris (Marattiales) from the Lower Jurassic Hsiangchi Formation in Hubei, China. *Review of Palaeobotany and Palynology* **107**, 125–44.
- WANG, Y. D. 2002. Fern ecological implications from the Lower Jurassic in Western Hubei. *Review of Palaeobotany and Palynology* **119**, 125–41.
- WANG, Y. D., GUIGNARD, G., LUGARDON, B. & BARALE, G. 2001. Ultrastructure of in situ *Marattia asiatica* (Marattiaceae) spores from the Lower Jurassic in Hubei, China. *International Journal of Plant Sciences* **162**, 927–36.
- WANG, Y. D. & MEI, S. W. 1999. Fertile organs and in situ spores of a matoniaceous fern from the Lower Jurassic of West Hubei. *Chinese Science Bulletin* **44**, 1333–7.
- WANG, Y. D., MOSBRUGGER, V. & ZHANG, H. 2005. Early–Middle Jurassic vegetation and climate events in the Qaidam Basin, China. *Palaeogeography, Palaeoclimatology, Palaeoecology* **224**, 200–16.
- YANG, J., QU, L., ZHOU, H., CHENG, Z., ZHOU, T., HOU, J., LI, P., SUN, S., LI, Y., ZHANG, Y., WU, X., ZHANG, Z. & WANG, Z. 1986. *Permian and Triassic strata and fossil assemblages in the Dalongkou area of Jimsar, Xinjiang*. Beijing: Geological Publishing House, 262 pp. (in Chinese with English abstract).
- YU, J. X. & ZHANG, W. P. 1980. Upper Cretaceous Dinoflagellate cysts and Acritarchs of Western Xinjiang. *Bulletin of Chinese Academy of Geological Sciences* **2**, 93–119, pls 1–6 (in Chinese with English abstract).
- ZHANG, Y. L., XI, Y. Z., ZHANG, J. T. & GAO, G. Z. 1976. *Sporae Pteridophytorum Sinicorum*. Beijing: Science Press, 451 pp. (in Chinese with English abstract).
- ZHOU, Z. Y. 2001. *Stratigraphy of the Tarim Basin*. Beijing: Science Press, 359 pp. (in Chinese with English abstract).
- ZHOU, Z. Y. & CHEN, P. J. 1990. *Biostratigraphy and Geological Evolution of the Tarim Basin*. Beijing: Science Press, 366 pp., pls 1–8 (in Chinese with English abstract).
- ZHOU, Q. J. & ZHENG, J. J. 1990. *Tectonics of the Tarim Basin*. Beijing: Science Press, 144 pp. (in Chinese with English abstract).