



Maternal height and double-burden of malnutrition households in Mexico: stunted children with overweight or obese mothers

Lucía Félix-Beltrán^{1,*} , James Macinko^{1,2} and Randall Kuhn³

¹Department of Health Policy and Management, Fielding School of Public Health, University of California at Los Angeles, 650 Charles E Young Dr S, Los Angeles 90095, CA, USA: ²Department of Community Health Sciences, Fielding School of Public Health, University of California at Los Angeles, Los Angeles, CA, USA: ³Department of Community Health Sciences, Fielding School of Public Health, University of California at Los Angeles, Los Angeles, CA, USA

Submitted 25 July 2019: Final revision received 21 June 2020: Accepted 23 July 2020: First published online 1 September 2020

Abstract

Objective: To assess the association between short maternal height and four types of mother–child nutritional status groupings within Mexican households.

Design: We classified mother–child dyads into four groups: stunted child and a non-overweight/non-obese mother (*stunting-only*), non-stunted child and an overweight/obese mother (*overweight-only*), stunted child with an overweight/obese mother (*double-burden*) and households with neither child stunting nor overweight/obese mothers (*neither-condition*). We assessed the association between maternal height and mother–child nutrition status using multinomial logistic regression, controlling for socio-economic covariates.

Setting: Nationally representative cross-section of households from the 2012 Mexican National Health and Nutrition Survey.

Participants: Children <5 years of age were matched to their mothers, resulting in a sample of 4706 mother–child dyads.

Results: We found that among children with stunting, 53.3% have an overweight/obese mother. Double-burden was observed in 8.1% of Mexican households. Maternal short stature increased the probability of stunting-only by 3.5% points (p.p.) and double-burden by 9.7 p.p. ($P < 0.05$). The inverse association was observed for overweight-only and neither-condition households, where the probability of these outcomes decreased by 7.2 and 6 p.p. in households with short-statured mothers ($P < 0.05$), respectively.

Conclusions: Women with short stature are more likely to develop overweight and simultaneously have a stunted child than those who are not short-statured. Our findings underline the challenges faced by public health systems, which have to balance the provision of services for both an undernourished and increasingly overweight/obese population.

Keywords

Double-burden of malnutrition
Mexico
Maternal height
Mother–child dyad

During the past few decades, there has been a worldwide decrease in undernutrition and a concomitant increase in overweight and obesity^(1–3). These shifts occurred as a result of nutrition and epidemiological transitions. As a consequence, the burden of disease has shifted from primarily infectious and vector-transmitted diseases (associated with undernutrition) to non-communicable diseases such as type 2 diabetes and hypertension, which are associated with overweight and obesity⁽⁴⁾. Previous studies have found that as a country's economy increases, so too does an urban lifestyle among its population, leading

to reduced physical activity and increased consumption of energy-dense foods⁽⁵⁾. These factors help explain how populations can experience increases in overweight alongside persistent undernutrition^(3,6–8), a situation that is fairly common for many countries in Latin America^(8–11).

Mexico has experienced a rapid and uneven nutrition transition. With economic growth, stunting prevalence has declined from 26.9% in 1988 to 13.6% in 2012⁽¹²⁾. Over the same time period, there was a 104% increase in the combined prevalence of overweight and obesity among adults, from 34.5% in 1988 to 71.2% in 2012⁽¹³⁾.

*Corresponding author: Email luciafelix@ucla.edu

© The Author(s), 2020. Published by Cambridge University Press on behalf of The Nutrition Society



However, these processes have not occurred in an even manner across Mexican states, regions^(12,14,15) or socio-economic groups⁽¹⁶⁾. Within this context, there has emerged a double-burden of malnutrition, frequently defined as households with both stunted children and overweight or obese mothers^(4,16). This coexistence poses challenges for health and nutrition policies as well as the healthcare system. For instance, widespread public programmes, such as conditional cash transfers that have been effective in reducing some forms of undernutrition, may have inadvertently contributed to obesity in some settings^(17–20).

Prior research in Mexico has identified household characteristics such as living in the southern region and rural residence as well as maternal factors such as lower height, higher birth order, lower schooling and the presence of central maternal adiposity to be positively associated with child stunting^(21,22). The fact that over and undernutrition occur in the same household has given rise to the hypothesis that conditions that were once considered opposite sides of a plateau might actually share common causal factors such as a poor-quality diet⁽²³⁾, physical activity or aspects of the social and cultural environment⁽²⁴⁾. Household food insecurity (HFI) (an indicator of poor household diet)⁽²⁵⁾ has been found to be positively associated with both childhood stunting⁽²⁶⁾ and adult overweight⁽²⁷⁾. Additionally, severe HFI has been found to increase the risk of stunting in children with non-obese mothers, but not among whose mothers are obese⁽²⁸⁾.

Another potential shared causal factor between child stunting and maternal overweight is maternal height. In low- and middle-income countries, maternal height is inversely correlated with child stunting⁽²⁹⁾. Specifically, previous research carried out in Mexico found that as maternal height increases, the probability of having a stunted child decreases⁽²²⁾. Evidence from Bangladesh, Indonesia and Guatemala suggests that short maternal height is associated with a higher risk of double-burden^(30,31). However, to our knowledge, no studies currently document such an association in Mexico.

In the current study, we construct mother–child dyads^(24,32,33) to assess the prevalence of different mutually exclusive groupings of mother and child nutritional status within the same household. These include: (1) households that have a stunted child and a non-overweight/obese mother (*stunting-only*); (2) those that have an overweight or obese mother but a non-stunted child (*overweight-only*); (3) those that have both a stunted child and an overweight or obese mother (*double-burden*) and (4) those with neither a stunted child nor an overweight or obese mother (*neither-condition*). We then test the association between short maternal stature and these different mother–child nutrition status groups, while controlling for important confounders.

The current study adds several insights to the literature. As discussed above, previous studies on double-burden in

Mexico either consider HFI⁽²⁸⁾ or maternal height^(9,22) as potential predictors of double-burden, but to our knowledge, our study is the first to consider both factors simultaneously. Additionally, we construct four mutually exclusive categories of household nutritional status using the mother–child dyad as the unit of analysis⁽⁴⁾ and employ multivariate statistical techniques to control for a number of important potential confounders.

Methods

Data source

We performed secondary data analysis using the 2012 National Health and Nutrition Survey, a two-stage probabilistic, national, rural, urban and state representative survey⁽³⁴⁾. Data were collected from 50 528 households across the 32 Mexican states between October 2011 and May 2012⁽³⁵⁾.

Mother–child dyads are our unit of analysis. These dyads are commonly used when assessing weight discordance because it is assumed that mothers and children are in closer contact and share more resources than do other household members⁽⁴⁾. We excluded 333 households (3.1%) from the study because mothers did not live in the household, had died or were not otherwise identifiable, and 5405 women because they were not randomly selected for anthropometric measures. Dyads were constructed by matching children <5 years old to their non-pregnant mothers aged 12–49 years with anthropometric data (n 4764). The child's age was reported by the mother and no woman surveyed within this age range had more than one surveyed child <5 years old. Finally, 58 dyads (1.1%) were excluded because of missing HFI data. This resulted in an analytic sample of 4706 mother–child dyads (see online supplementary material, Appendix 1).

Outcome variable

Child nutritional status

Stunting was assessed using height and age measurements. Height was assessed with a stadiometer with 1 mm precision⁽³⁶⁾. Height-for-age Z-scores for children 0–59 months old were calculated according to WHO standards⁽³⁷⁾. The range of valid scores included in the dataset was between -6.0 SD and 6.0 SD⁽³⁴⁾. Children were categorised into stunted (height-for-age Z-scores ≤ 2 SD) and non-stunted (height-for-age Z-scores ≥ -2 SD) according to WHO cut-off points⁽³⁸⁾.

Maternal nutritional status

BMI of participants (kg/cm^2) was assessed using height and weight. Weight was assessed using a SECA scale with 100 g precision and height with a stadiometer of 1 mm precision⁽³⁶⁾. For adolescent women (13–19 years old) BMI-for-age Z-scores were calculated using WHO

standards⁽³⁹⁾. BMI values outside the range of 10–58 were excluded from the primary data source. Adolescents with *Z*-scores above +1 SD and below +2 SD were classified as overweight and those with *Z*-scores above +2 SD were classified as obese. Based on their BMI, women 20–49 years of age were classified into the following categories: underweight: BMI 10–18.49, normal weight: 18.5–24.99, overweight: 25–29.99 and obese ≥ 30 ⁽⁴⁰⁾. Mothers classified as overweight or obese were assigned a 1 or 0 otherwise.

Household nutrition status/types of mother–child dyad

Based on the nutritional status of the child and the mother, households were classified into four outcome categories:

- Households with a stunted child and non-overweight/obese mother were classified as *stunting-only*.
- Households with a non-stunted child and overweight/obese mother were classified as *overweight-only*.
- Households that had both a stunted child and an overweight or obese mother were classified as *double-burden*.
- Households that had neither a stunted child nor an overweight/obese mother were classified as *neither-condition*. This category was used as the reference group in multivariate analyses.

Independent variable

Maternal height was available as the mother's height in cm. It was operationalised as a dichotomous variable. Consistent with previous studies, a value of 1 was assigned to mothers with height <150 cm, considered to be 'short stature' in the Mexican population⁽²²⁾. A value of 0 was assigned otherwise.

Control variables

HFI was assessed using The Latin American and Caribbean Food Security Scale⁽⁴¹⁾. This scale uses fifteen questions to measure the perceptions of food access during the past 3 months that result from the availability of household resources. It uses dichotomous questions to inquire on the perceptions of worry, quality and quantity of available foods. Positive response options were assigned a value of 1 and 0 otherwise. The total summation score allowed us to generate a categorical assessment of HFI using pre-established cut-off values⁽⁴¹⁾. Households with no positive answers (score of 0) were considered food secure. Households with members under 18 years of age and 1–5 positive answers or without members under 18 years and a score between 1 and 3 were considered to have mild food insecurity. Households were classified as moderately food insecure when they had scores between 6 and 10 and members under 18 years of age, and 4–6 positive answers without members under 18 years old. Severely food

insecure households were those with members with 18 years and a score between 11 and 15, and those without members under 18 years and 7–8 positive answers⁽⁴¹⁾ (see online supplementary material, Appendix 2). To assess socio-economic status (SES), we used a pre-constructed measure of well-being based on household characteristics including household demographic structure, head of household characteristics, infrastructure conditions, household assets, consumption patterns and marginality level. The National Health and Nutrition Survey 2012 team used principal component analysis on the list of items and divided the resulting principal factor into quintiles⁽⁴²⁾. Quintile 1 (Q1) represents the lowest (poorest) and Quintile 5 (Q5) the highest possible level of SES. A dichotomous variable was used to control for exposure to urban environments. Urban localities were those with ≥ 2500 inhabitants and assigned a 1, and rural localities with <2500 inhabitants were assigned a 0, as provided by the primary data source. In order to control for regional variation, states were classified into four regions: South, North, Centre and Mexico City (classified as a region of its own due to population density)⁽³⁴⁾. Maternal age was operationalised into three categories: 13–19, 20–35 and 36–49 years.

Statistical analysis

Data analysis was performed using Stata version 14.2⁽⁴³⁾ and used the SVY command to incorporate sampling weights and adjust for complex survey design effects. We obtained means for continuous variables, proportions (%) for categorical variables and linearised SE for both. First, we analysed the entire sample and then we stratified results by type of mother–child dyad. Then, we performed bivariate analysis to assess overall statistical differences between categories of household nutrition status using design-corrected *F* tests and adjusted Wald tests to determine statistical significance.

For multivariate analyses, we used a multinomial logistic model (mlogit) to assess the probability of having stunting-only, overweight-only and double-burden and compared with those where neither-condition was observed. We first estimated relative risk ratios (RRR) for maternal short stature (ref = non-short stature) with 95% CI adjusting for SES, locality type, maternal age and region. Additionally, we estimated the predicted probability of each outcome category for households with and without maternal short stature as well as the marginal effect (marginal effect at representative values)⁽⁴⁴⁾. Both of these calculations adjusted for the same covariates in the RRR estimation. To test hypotheses, we used two-sided tests and $P < 0.05$ in all statistical analyses.

Results

Out of the entire sample, 14.5% of children were stunted while 63.4% of mothers were overweight or obese. We



found that 390 (8.1%) households were double-burden, 5.7% were stunting-only, more than half were overweight-only (56.0%) and 30.2% had neither-condition. The mean maternal height was 154 cm (with a SE of 0.30) and 25.9% of the mothers had short stature. Mildly food insecure households were most common, representing 44.0% of households, while severe HFI households were the less frequent (10.5%). Mother-child dyads were evenly distributed across SES categories and concentrated in the Southern and Central regions (33.9 and 31.0%, respectively). Mean maternal age was 28 years and there were < 10% of adolescent mothers (Table 1).

More than half of the children with stunting had an overweight or obese mother (53.3%). Of the remaining children with stunting, 1.6% had underweight mothers and 45.1% had mothers in the normal weight category. Children who were not stunted had a higher proportion of overweight or obese mothers (60.2%) and a lower proportion of normal or underweight ones (38.6 and 1.2%, respectively) (Fig. 1).

Stratified results show statistically significant differences across mother-child dyad types in mean maternal stature, proportion of mothers with short maternal height, locality type, SES, region and maternal age. Mothers in double-burden households had the lowest mean stature (149.9 cm) while those with neither-condition had the highest (155.6 cm). Double-burden households had the highest proportion of mothers with short stature (54.2%), followed by stunting-only households (44.7%), overweight-only (22.7%) and neither-condition households (20.2%). We also observed that double-burden and stunting-only households were mostly located in rural localities (32.4 and 45.9%, respectively), while overweight-only (24.3%) and neither-condition were not (26.0%). Regarding SES, neither-condition and overweight-only were equally distributed across quintiles, while stunting-only households and double-burden ones were mostly poorer and concentrated in Q1 and Q2 (70% of stunting-only were in Q1 and Q2, and about 50% of double-burden households were in these categories). In terms of regional distribution,

Table 1 Nutritional status, socio-economic, demographic characteristics and nutritional status of the sample (*n* 4706)*

Variables	<i>n</i>	%	Linearised SE	95 % CI
Child nutritional status				
Stunted	660	14.5	1.19	12.29, 17.11
Maternal nutrition status				
Underweight	72	1.6	0.29	1.12, 2.33
Normal weight	1585	35.0	1.02	32.94, 37.07
Overweight	1753	35.1	1.19	32.70, 37.51
Obese	1296	28.3	1.14	26.09, 30.71
Household nutrition status				
Child stunting and maternal overweight/obesity (double-burden)	390	8.1	0.69	6.85, 9.66
Child stunting and non-overweight/obese mother (stunting-only)	270	5.7	0.66	4.48, 7.16
Overweight/obese mother and non-stunted child (overweight-only)	2659	56.0	1.19	53.56, 58.36
Non-stunted child and non-overweight/obese mother (neither-condition)	1387	30.2	0.99	28.24, 32.35
Short maternal height (<150 cm)	1367	25.9	1.68	22.61, 29.41
Maternal height (mean cm)	154.4		0.30	153.76, 154.97
Household food insecurity				
Secure	1095	26.0	1.06	23.95, 28.26
Mild	2173	44.0	1.07	41.88, 46.23
Moderate	918	19.5	0.77	17.96, 21.06
Severe	520	10.5	0.55	9.39, 11.61
Location				
Rural (<2500 inhabitants)	1834	26.7	4.72	18.28, 37.27
SES†				
Quintile 1	1242	20.9	2.48	16.28, 26.32
Quintile 2	1097	21.0	1.05	18.97, 23.24
Quintile 3	938	19.2	0.85	17.50, 20.94
Quintile 4	800	19.9	1.37	17.24, 22.80
Quintile 5	629	19.1	1.78	15.73, 22.95
Region				
North	1027	19.8	4.75	11.90, 31.20
Centre	1670	31.0	5.71	20.76, 43.57
Mexico City	185	15.2	7.97	4.89, 38.54
South	1824	33.9	5.72	23.43, 46.25
Maternal age				
13–19 years	512	7.9	0.56	6.81, 9.08
20–35 years	3355	76.8	1.28	73.92, 79.12
36–49 years	839	15.5	1.00	13.58, 17.65
Maternal age (mean years)	28.5		0.15	18.17, 28.78

SES, socio-economic status.

*Used weighted data to adjust for the complex design of the survey.

†Quintile 1 is the lowest and Quintile 5 the highest level of SES.




Fig. 1 Child and maternal nutrition status. ■, Obesity; ■, Overweight; ■, Normal weight; ■, Low weight

double-burden (45.6%) and stunting-only (45.4%) had the highest concentration of pairs in the South, while neither-condition households were concentrated in the Centre (36.2%). Mothers in neither-condition households were younger (26 years) than those in overweight-only, stunting-only or double-burden ones (29 and 30 years) (Table 2).

Our multivariate analyses show that maternal short stature (RRR = 3.5; 95% CI 2.50, 4.09) and severe HFI (RRR = 1.7; 95% CI 1.04, 2.93) were positively associated with double-burden, and that households in the Central region were less likely to develop double-burden than those in the South (RRR = 0.6; 95% CI 0.39, 1.21). Our results also show that the relative risk of having double-burden significantly increases with maternal age (Table 3).

Additionally, the relative risk of stunting-only was positively associated with the presence of maternal short stature (RRR = 2.3; 95% CI 1.51; 3.49) and rural residence (RRR = 1.9; 95% CI 1.29, 2.65). SES was negatively associated with stunting-only (RRR = 0.6; 95% CI 0.34, 1.00). It should be noted that rounding of the CI leads to an upper limit of 1.00, meaning that the association between SES and the presence of a stunted child with a non-overweight/obese mother is fairly small in magnitude. We found that households with mild HFI are more likely to have

Table 2 Nutritional status, socio-economic and demographic characteristics by type of mother-child dyad (*n* 4706)*

	Neither-condition		Overweight-only		Stunting-only		Double-burden		<i>P</i>
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
<i>n</i> (%)	1387	30.2	2659	56.0	270	5.7	390	8.1	
Maternal height (cm)									<0.0000
Mean	155.6		154.7		151.0		149.9		
SE	0.35		0.26		0.26		0.60		
Short maternal height (<150 cm)	309	20.2	695	22.7	141	44.7	222	54.2	<0.0000
Household food insecurity									0.0772
Secure	367	30.0	606	25.3	53	21.5	69	19.9	
Mild	637	42.6	1242	45.3	121	41.9	173	42.7	
Moderate	247	18.4	511	19.1	59	23.4	101	23.3	
Severe	136	9.0	300	10.4	37	13.2	47	14.2	
Location									0.0001
Rural (<2500 inhabitants)	547	26.0	960	24.3	153	45.9	174	32.4	
SES†									<0.0000
Quintile 1	357	20.4	607	17.5	130	40.8	148	31.5	
Quintile 2	330	20.7	602	20.6	53	20.8	112	25.1	
Quintile 3	265	18.8	572	20.1	43	14.3	59	17.1	
Quintile 4	227	17.9	508	22.6	25	13.1	40	13.5	
Quintile 5	208	22.2	371	19.1	19	11.0	31	12.9	
Region									0.0004
North	307	19.4	625	21.8	35	11.2	60	14.2	
Centre	572	36.2	920	30.0	79	27.2	99	21.3	
Mexico City	61	15.5	97	14.5	11	16.2	16	18.9	
South	447	29.0	1017	33.7	145	45.4	215	45.6	
Age									<0.0000
13–19 years	236	13.0	191	4.5	54	15.6	31	5.6	
20–35 years	1004	78.5	1890	75.7	184	73.0	277	76.7	
36–49 years	147	8.4	578	19.5	32	11.4	82	17.7	
Age (years)									<0.0000
Mean	26.3		29.9		29.4		30.0		
SE	0.4		0.2		0.6		0.4		

SES, socio-economic status.

*Used weighted data to adjust for the complex design of the survey. Significance was assessed with two-sided *P* < 0.05.

†Quintile 1 is the lowest and Quintile 5 the highest level of SES.

Table 3 Relative risk ratios (RRR) and 95% CI of the factors associated to households with child stunting and maternal overweight (double-burden), households with child stunting and non-maternal overweight/obesity (stunting-only), households with a non-stunted child and maternal overweight/obesity (overweight-only) compared with households with a non-stunted child and a non-overweight/obese mother (neither-condition) (*n* 4706)†‡

	Double-burden		Stunting-only		Overweight-only	
	RRR	95 % CI	RRR	95 % CI	RRR	95 % CI
Non short-maternal height (≥ 150 cm)	–		–		–	
Short maternal height (<150 cm)	3.5***	2.50, 4.09	2.3***	1.51, 3.49	1.1	0.83, 1.41
Household food insecurity						
Secure	–		–		–	
Mild	1.3	0.85, 2.00	1.1	0.75, 1.66	1.3*	1.00, 1.61
Moderate	1.4	0.88, 2.19	1.3	0.70, 2.31	1.2	0.88, 1.60
Severe	1.7*	1.04, 2.93	1.4	0.68, 3.12	1.4	0.95, 1.90
Location						
Urban (>2500)	–		–		–	
Rural (<2500)	1.1	0.77, 1.50	1.9**	1.29, 2.65	1.0	0.79, 1.13
SES§						
Quintile 1	–		–		–	
Quintile 2	1.1	0.73, 1.55	0.7	0.42, 1.15	1.3*	1.01, 1.62
Quintile 3	0.9	0.50, 1.52	0.6*	0.34, 1.00	1.4*	1.02, 1.82
Quintile 4	0.8	0.48, 1.28	0.7	0.32, 1.32	1.6**	1.12, 1.33
Quintile 5	0.6	0.35, 1.10	0.5	0.21, 1.16	1.0	0.78, 1.33
Region						
South	–		–		–	
North	0.0	0.58, 1.21	0.7	0.40, 1.27	0.9	0.75, 1.19
Centre	0.6**	0.39, 1.21	0.8	0.47, 1.19	0.7***	0.53, 0.82
Mexico City	1.1	0.74, 1.60	1.3	0.63, 2.82	0.7*	0.52, 0.82
Age						
13–19 years	–		–		–	
20–35 years	2.4**	1.37, 4.09	0.9	0.55, 1.37	2.9***	2.14, 3.87
36–49 years	4.6***	2.30, 9.41	1.0	0.50, 2.09	6.5***	4.32, 9.89

RRR, relative risk ratio; SES, socio-economic status.

†Showing adjusted RRR with 95% CI in parentheses from multinomial logistic regression model comparing double-burden, stunting-only and overweight-only households to those with neither-condition.

‡Used weighted data to adjust for the complex design of the survey.

§Quintile 1 is the lowest and Quintile 5 the highest level of SES.

P* < 0.05; ** *P* < 0.01; * *P* < 0.001.

overweight-only (RRR = 1.3; 95% CI 1.00, 1.61) and so are those in higher SES quintiles (RRR = 1.6; 95% CI 1.12, 1.33). Also, dyads in the Central region and Mexico City are less likely to be overweight-only than those in the South (RRR = 0.7; 95% CI 0.53, 0.62 and RRR = 0.7; 95% CI 0.52, 0.82). Results also show that the RRR of overweight-only increases with maternal age (Table 3).

Having maternal short stature significantly increases a household's probability of double-burden from 0.053 to 0.150, a change of 9.7 % points (p.p.) (*P* < 0.001) and from 0.04 to 0.075 (*P* = 0.019), a change of 0.035 in stunting-only households. We observed an inverse association between short maternal height and overweight-only and neither-condition households, where the probability of these outcomes was higher for women who did not have short stature (an absolute change of –7.2 and –6.0 p.p., respectively) (Fig. 2).

Discussion

The current study estimated the prevalence of double-burden – households with child stunting and maternal overweight or obesity – and assessed whether mothers

in double-burden households were more likely to be of short stature. Our results are consistent with previous estimations: the prevalence of double-burden households in Mexico was above 8%⁽¹⁶⁾, and the prevalence of households that have maternal overweight but not child stunting was greater than that of households without either condition.

The prevalence of double-burden households in Mexico was below than the prevalence of this phenomenon in lower middle-income Latin American countries such as Guatemala^(30,45,46), Ecuador⁽⁴⁷⁾ and Honduras⁽⁴⁸⁾, but was higher than in higher middle-income Latin American countries such as Colombia^(24,49), Brazil^(50,51) and Peru⁽⁴⁸⁾. This could be due to the widespread prevalence of adult overweight and obesity in Mexico (63.4% of mothers in our sample), while < 6% of households were stunting-only. Overweight-only households were observed evenly across SES levels, while stunting-only households were found almost exclusively in the lowest SES levels and in rural localities.

Another major finding is that more than half of the children (53%) with stunting had an overweight or obese mother. Of the remaining 47% of stunted children, < 2% had mothers with low weight and 45% were in the




Fig. 2 Predicted probabilities and marginal effect of maternal short stature on mother–child dyad nutrition status. Used weighted data to adjust for the complex design of the survey; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. ■, No short stature; ■, Short stature; ■, Absolute change

normal weight category. A possible explanation for this result is that low weight households tend to have income that is insufficient to increase their BMI⁽⁹⁾, while the normal weight group might be at-risk of becoming overweight. Even though formal forecasting is beyond the scope of this paper, it is possible that if increasing adult overweight trends⁽⁵²⁾ continue, all children with stunting could have overweight or obese mothers and therefore live in double-burden households. This situation merits further exploration and should be monitored.

Short maternal height was observed in 25% of our sample, in more than half of double-burden households and 45% of stunting-only households. Moreover, results from our multinomial logistic regression showed that in households with short maternal height, the likelihood of having stunting-only is twice than that of having neither-condition, consistent with previous studies in Mexico^(9,22). Specifically, there is a 3.5 p.p. (P -value = 0.019) difference of being stunting-only between households with and without maternal short stature. More importantly, the risk of double-burden increases by 9.7 p.p. (P -value < 0.000) when the mother's height is below 150 cm. The opposite association is observed for households that are overweight-only, where maternal short stature decreases the probability of overweight-only by 7.2 p.p. (P -value = 0.007) and of having neither-condition by 6%p.p. (P -value = 0.014). Having short height not only puts the mother at higher risk

of overweight or obesity but also her child at higher risk of stunting. This is consistent with prior studies which have found that maternal short stature increases the risk of double-burden in Indonesia⁽³¹⁾, Guatemala^(30,46) and Brazil⁽⁵¹⁾.

A possible explanation for our findings can be found in a growing body of literature which supports that maternal short stature has adverse short- and long-term consequences on child development. For example, in some low- and middle-income countries, maternal short stature has been negatively associated with small-for-gestational-age, preterm births⁽⁵³⁾, child growth patterns^(29,54,55) and increased risk of infant mortality⁽²⁹⁾. Additionally, prior studies have identified that child undernutrition during the first years has an important influence on adult health outcomes such as obesity and non-communicable diseases^(1,56) and that individuals with poor fetal growth or stunting in the first years of life are more likely to gain excessive weight during adolescence, placing them at higher risk of nutrition-related diseases⁽⁵⁷⁾. Furthermore, poor nutrition in early childhood could be an important driving factor behind obesity in low- and middle-income countries^(10,57). In that vein, within a dyadic context, maternal height becomes a significant predictor of maternal and child nutritional status, possibly reflecting the transmission of disease burden acquired by the mother during her early childhood.

Our results underline the importance of continuing to invest in prenatal care and early childhood development



in order to improve health and well-being across the life course. Since experiences during the first years of life can lead to negative health outcomes and affect future generations, early childhood investments could be cost-effective alternatives to address adult outcomes^(57–59).

In contrast with existing evidence from some Latin American countries⁽³²⁾, our study shows that factors such as exposure to urban lifestyles and SES are not predictors of double-burden in Mexico, but do predict child stunting (without an overweight/obese mother). This could be because Mexico is at an advanced stage of an uneven health transition^(14,16) leading to a situation in which stunting-only households are concentrated in lower socio-economic strata, double-burden households are found in both urban and rural localities, overweight-only households are spread across the country and there is a higher proportion of neither-condition households in the Central region.

According to our results, HFI is a predictor of double-burden and overweight-only but not of stunting-only. We found a weak but statistically significant relationship between severe HFI and double-burden, consistent with other studies that have explored this association⁽⁵⁰⁾. The weakness of this association could be due to the fact that both HFI and maternal overweight are widely spread across the country and that the causal pathways that explain nutrition status are highly complex. Nevertheless, there is evidence to support that HFI may be a shared factor in the causal pathways of child undernutrition and maternal overweight, indicating poor dietary quality^(23,24,60). Our results diverge from those obtained in a previous study⁽²⁸⁾, where severe HFI increased the risk of stunting-only households but not of double-burden, these differences could be due to the fact that we included different variables in our models and a different analytic strategy.

Households with a stunted child but not an overweight or obese mother and those with double-burden share socio-demographic characteristics such as being concentrated in the lowest socio-economic quintiles, in the Southern region and in rural localities. Both conditions are also associated with short maternal height. These types of households are significantly different from maternal overweight-only households and those that have neither-condition, both of which are more evenly distributed across SES quintiles, in urban localities, in the Central region and have fewer mothers with short stature.

One of the limitations of our study is that by using cross-sectional data, we are unable to assess the direction of the association between some covariates such as HFI, SES and nutrition status, leading to possible reverse causality. The need for longitudinal studies to disentangle the complexity of these relationships has previously been acknowledged⁽¹¹⁾. Ideally, information on the mother's early childhood environment would allow for assessing its effect on her adult outcomes and those of her children. However, even with data taken at only one time point, we are able

to measure the significance of maternal height as a predictor of maternal and child nutrition status, potentially reflecting the transmission of nutrition and health conditions from one generation to another. In addition, our operationalisation of household nutrition status combined overweight and obese mothers. This should be taken with caution because there might be specific issues in the extremes of the BMI distribution⁽⁶¹⁾ and overweight and obesity might have different risk factors and effects.

Another limitation is that we do not account for the role played by nutrition and health policies in the development of child stunting and maternal overweight/obesity. As identified in previous studies, health and nutrition programmes originally devised to target undernutrition could be contributing to the development of overweight or obesity. For example, in the case of Mexico's conditional cash transfers, doubling the cash transfer was associated with better motor development and nutrition among children⁽⁶²⁾, but to a higher BMI, diastolic blood pressure and overweight among adults⁽¹⁷⁾. More recent studies found that cash and in-kind transfers not only increase fruit, vegetable and micronutrient consumption but also lead to excess energy consumption⁽¹⁸⁾. In that vein, our findings continue to draw attention to the challenges set by the double-burden of malnutrition for public programmes that seek to improve health and nutrition, which has been pointed out in existing literature⁽⁶³⁾.

It is also important to highlight that studies seeking to understand the drivers of the double-burden of malnutrition may operationalise this outcome in different ways, defining it as the coexistence of maternal adiposity and child stunting⁽²¹⁾, overweight and anaemia⁽⁶⁰⁾, child overweight with concurrent stunting^(9,64) and at least one household member with overweight and one with underweight⁽⁶⁵⁾. Each measurement poses different challenges and can yield different results. Our findings highlight the complexities of the relationships between undernutrition, overweight and their associated factors in a country in an advanced stage of the nutrition transition that has experienced economic growth while widening inequality gaps^(9,14,16).

Conclusions

Maternal overweight and HFI are generalised population issues in Mexico, where six out of ten women between 13 and 49 years of age are overweight or obese and only three in ten households are food secure. Additionally, over 50% of children who are stunted have a mother who is overweight or obese. It is possible that at least 20% of the remaining mothers will develop either of these conditions over the next few years. Women who are short stature are more likely to be overweight and have a stunted child than those who are not short-statured. These findings underline the challenges faced by public healthcare

systems, which have to tend to a population with multiple forms of malnutrition. These difficulties are also faced by public programmes targeting undernutrition, which must develop strategies to prevent unhealthy weight gain among their beneficiary population.

The significance of maternal height as a predictor of maternal overweight and childhood stunting highlights the importance of devising policies that focus on the quality of prenatal care and early childhood development, for they might represent a cost-effective way of preventing child undernutrition and later adverse health outcomes.

Acknowledgements

Acknowledgements: None. *Financial support:* L.F.B. received support from the Graduate Research Summer Mentorship program from the UCLA Graduate Division to conduct part of this research. This research did not receive any other support from funding agencies in the public, commercial or not-for-profit sectors. *Conflict of interest:* None. *Authorship:* L.F.B. conceived the article, developed the conceptual model analysed the data and drafted the manuscript. J.M. drafted and critically revised the manuscript, R.K. interpreted the data and critically revised the manuscript. *Ethics of human subject participation:* We performed secondary analysis using a survey for which respondents provided written consent after being notified of the survey's purpose.

Supplementary material

For supplementary material accompanying this paper visit <https://doi.org/10.1017/S136898002000292X>

References

- Black RE, Victora CG, Walker SP *et al.* (2013) Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* **382**, 427–451.
- Finucane MM, Stevens GA, Cowan MJ *et al.* (2011) National, regional, and global trends in body-mass index since 1980: systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet* **377**, 557–567.
- Popkin BM (1994) The nutrition transition in low-income countries: an emerging crisis. *Nutr Rev* **52**, 285–298. doi: 10.1111/j.1753-4887.1994.tb01460.x.
- Tzioumis E & Adair LS (2014) Childhood dual burden of under- and overnutrition in low- and middle-income countries: a critical review. *Food Nutr Bull* **35**, 230–243.
- Popkin BM, Adair LS & Ng SW (2012) Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev* **70**, 3–21.
- Popkin BM (1999) Urbanization, lifestyle changes and the nutrition transition. *World Dev* **27**, 1905–1916.
- Popkin BM (2011) Contemporary nutritional transition: determinants of diet and its impact on body composition. *Proc Nutr Soc* **70**, 82–91.
- Rivera-Dommarco JA, Barquera S, González-Cossio T *et al.* (2004) Nutrition transition in Mexico and other Latin American countries. *Nutr Rev* **62**, 149–157.
- Fernald LC & Neufeld LM (2007) Overweight with concurrent stunting in very young children from rural Mexico: prevalence and associated factors. *Europ J Clin Nutr* **61**, 623–632.
- Abdullah A (2015) The double burden of undernutrition and overnutrition in developing countries: an update. *Curr Obes Rep* **4**, 337–349.
- Kosaka S & Umezaki M (2017) A systematic review of the prevalence and predictors of the double burden of malnutrition within households. *Brit J Nutr* **117**, 1118–1127.
- Rivera-Dommarco JÁ, Cuevas-Nasu L, González de Cosío T *et al.* (2013) Desnutrición crónica en México en el último cuarto de siglo: análisis de cuatro encuestas nacionales (Stunting in Mexico in the last quarter century: analysis of four national surveys). *Salud Pública de México* **55**, S161–S169.
- Torres F & Rojas A (2018) Obesidad y salud pública en México: transformación del patrón hegemónico de oferta-demanda de alimentos (Obesity and public health in Mexico: transforming the hegemonic food supply and demand pattern). *Problemas del Desarrollo* **49**, 145–169.
- Gomez-Dantes H, Fullman N, Lamadrid-Figueroa H *et al.* (2016) Dissonant health transition in the states of Mexico, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **388**, 2386–2402.
- Rivera JA, Pedraza LS, Aburto TC *et al.* (2016) Overview of the dietary intakes of the Mexican population: results from the National Health and Nutrition Survey 2012. *J Nutr* **146**, 1851s–1855s.
- Kroker-Lobos MF, Pedroza-Tobias A, Pedraza LS *et al.* (2014) The double burden of undernutrition and excess body weight in Mexico. *Am J Clin Nutr* **100**, 1652s–1658s.
- Fernald LC, Gertler PJ & Hou X (2008) Cash component of conditional cash transfer program is associated with higher body mass index and blood pressure in adults. *J Nutr* **138**, 2250–2257.
- Leroy JL, Gadsden P, Rodríguez-Ramírez S *et al.* (2010) Cash and in-kind transfers in poor rural communities in Mexico increase household fruit, vegetable, and micronutrient consumption but also lead to excess energy consumption. *J Nutr* **140**, 612–617.
- Lopez-Arana S, Avendano M, Forde I *et al.* (2016) Conditional cash transfers and the double burden of malnutrition among children in Colombia: a quasi-experimental study. *Brit J Nutr* **115**, 1780–1789.
- Shamah-Levy T, Méndez-Gómez-Humarán I, Gaona-Pineda EB *et al.* (2016) Food assistance programmes are indirectly associated with anaemia status in children <5 years old in Mexico. *Brit J Nutr* **116**, 1095–1102.
- Barquera S, Peterson KE, Must A *et al.* (2007) Coexistence of maternal central adiposity and child stunting in Mexico. *Int J Obes* **31**, 601–607.
- Shamah-Levy T, Nasu LC, Moreno-Macias H *et al.* (2008) Maternal characteristics determine stunting in children of less than five years of age results from a national probabilistic survey. *Clin Med Insights: Pediatr* **1**, 43–52.
- Garrett JL & Ruel MT (2005) Stunted child-overweight mother pairs: prevalence and association with economic development and urbanization. *Food Nutr Bull* **26**, 209–221.
- Parra DC, Gomez LF, Iannotti L *et al.* (2018) Maternal and familial correlates of anthropometric typologies in the nutrition transition of Colombia, 2000–2010. *Public Health Nutr* **21**, 2584–2594.
- Rodríguez LA, Mundo-Rosas V, Mendez-Gomez-Humarán I *et al.* (2016) Dietary quality and household food insecurity among Mexican children and adolescents. *Matern Child Nutr* **13**, e1237.



26. Cuevas-Nasu L, Rivera-Dommarco JA, Shamah-Levy T *et al.* (2014) Inseguridad alimentaria y estado de nutrición en menores de cinco años de edad en México (Food insecurity and nutritional status of preschool children in Mexico). *Salud Pública de México* **56**, s47–s53.
27. Morales-Ruán M, Méndez-Gómez Humarán I, Shamah-Levy T *et al.* (2014) La inseguridad alimentaria está asociada con obesidad en mujeres adultas de México (Food insecurity is associated with obesity in adult women of Mexico). *Salud Pública de México* **56**, s54–s61.
28. Shamah-Levy T, Mundo-Rosas V, Morales-Ruan C *et al.* (2017) Food insecurity and maternal-child nutritional status in Mexico: cross-sectional analysis of the National Health and Nutrition Survey 2012. *BMJ Open* **7**, e014371.
29. Özaltın E, Hill K & Subramanian SV (2010) Association of maternal stature with offspring mortality, underweight, and stunting in low-to middle-income countries. *JAMA* **303**, 1507–1516.
30. Lee J, Houser RF, Must A *et al.* (2010) Disentangling nutritional factors and household characteristics related to child stunting and maternal overweight in Guatemala. *Econ Hum Biol* **8**, 188–196.
31. Oddo V, Rah J, Semba RD *et al.* (2012) Predictors of maternal and child double burden of malnutrition in rural Indonesia and Bangladesh. *Am J Clin Nutr* **95**, 951–958.
32. Parra D, Gomez LF, Iannotti L *et al.* (2018) Multilevel correlates of household anthropometric typologies in Colombian mothers and their infants. *Glob Health Epidemiol Genom* **3**, e6.
33. Parra DC, Iannotti L, Gomez LF *et al.* (2015) The nutrition transition in Colombia over a decade: a novel household classification system of anthropometric measures. *Arch Public Health* **73**, 12.
34. Gutiérrez JP, Rivera-Dommarco J, Shamah-Levy T *et al.* (2012) Encuesta Nacional de Salud y Nutrición 2012 (National Health and Nutrition Survey 2012). *Resultados Nacionales (National Results)*. Cuernavaca, México: Instituto Nacional de Salud Pública (MX).
35. Romero-Martínez M, Shamah-Levy T, Franco-Núñez A *et al.* (2013) Encuesta Nacional de Salud y Nutrición 2012: diseño y cobertura (National Health and Nutrition Survey 2012: design and coverage). *Salud pública de Mexico* **55**, S332–S340.
36. Shamah-Levy T, Rivera-Dommarco JA, Verónica M *et al.* (2012) La doble carga de la mala nutrición: desnutrición y obesidad (The double-burden of malnutrition: undernutrition and obesity). In *Panorama de la Seguridad Alimentaria y Nutricional en México 2012 (Landscape of Food and Nutritional Insecurity in Mexico 2012)* [OdINU]plAylA Secretaría de Desarrollo Social, Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación, Instituto Nacional de Salud Pública, editor]. México: Food and Agriculture Organization of the United Nations. E-ISBN 978-92-5-307623-9.
37. WHO (2010) *WHO Anthro for Personal Computers, Version 3.2. 2: Software for Assessing Growth and Development of the World's Children*. Geneva: WHO.
38. WHO (2006) *WHO Child Growth Standards: Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index-for-Age: Methods and Development* [WMGR Study, editor]. Geneva: World Health Organization.
39. de Onis M, Onyango AW, Borghi E *et al.* (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* **85**, 660–667.
40. WHO (2000) *Obesity: Preventing and Managing the Global Epidemic*. Geneva: World Health Organization.
41. Committee ES (2012) *Escala Latinoamericana y Caribeña de Seguridad Alimentaria (ELCSA): Manual de uso y aplicaciones (Latin American and Caribbean Food Security Scale (ELCSA). Use and Application Manual)*. Rome: Food and Agriculture Organization of the United Nations.
42. Gutiérrez JP (2013) Clasificación socioeconómica de los hogares en la ENSANUT 2012 (Household socioeconomic classification in the National Health and Nutrition Survey 2012). *Salud Pública de México* **55**, S341–S346.
43. StataCorp (2015) *Stata Statistical Software: Release 14*. College Station, TX: StataCorp LP.
44. Long JS & Freese J (2014) *Regression Models for Categorical Dependent Variables Using Stata*. 3rd ed. College Station, TX: Stata Press.
45. Ramirez-Zea M, Kroker-Lobos MF, Close-Fernandez R *et al.* (2014) The double burden of malnutrition in indigenous and nonindigenous Guatemalan populations. *Am J Clin Nutr* **100**, 1644s–1651s.
46. Doak CM, Campos Ponce M, Vossenaar M *et al.* (2016) The stunted child with an overweight mother as a growing public health concern in resource-poor environments: a case study from Guatemala. *Ann Hum Biol* **43**, 122–130.
47. Freire WB, Silva-Jaramillo KM, Ramirez-Luzuriaga MJ *et al.* (2014) The double burden of undernutrition and excess body weight in Ecuador. *Am J Clin Nutr* **100**, 1636s–1643s.
48. Dieffenbach S & Stein AD (2012) Stunted child/overweight mother pairs represent a statistical artifact, not a distinct entity. *J Nutr* **142**, 771–773.
49. Sarmiento OL, Parra DC, Gonzalez SA *et al.* (2014) The dual burden of malnutrition in Colombia. *Am J Clin Nutr* **100**, 1628s–1635s.
50. Gubert MB, Spaniol AM, Segall-Correa AM *et al.* (2016) Understanding the double burden of malnutrition in food insecure households in Brazil. *Matern Child Nutr*. Published online: E-Pub 09 August. doi: 10.1111/mcn.12347.
51. Gea-Horta T, Silva Rde C, Fiaccone RL *et al.* (2016) Factors associated with nutritional outcomes in the mother-child dyad: a population-based cross-sectional study. *Public Health Nutr* **19**, 2725–2733.
52. Pedroza A & Rivera-Dommarco JA (2013) Prevalencia de obesidad en adultos mexicanos, 2000–2012 (Prevalence of obesity in Mexican adults 2000–2012). *Salud Pública de México* **55**, S151–S160.
53. Kozuki N, Katz J, Lee AC *et al.* (2015) Short maternal stature increases risk of small-for-gestational-age and preterm births in low- and middle-income countries: individual participant data meta-analysis and population attributable fraction. *J Nutr*. Published online: E-Pub. doi: 10.3945/jn.115.216374.
54. Addo OY, Stein A, Fall Ch *et al.* (2013) Maternal height and child growth patterns. *J Pediatr* **163**, 549.
55. Ali Z, Saaka M, Adams A-G *et al.* (2017) The effect of maternal and child factors on stunting, wasting and underweight among preschool children in Northern Ghana. *BMC Nutr* **3**, 31.
56. Beltrán-Sánchez H, Crimmins EM, Teruel GM *et al.* (2011) Links between childhood and adult social circumstances and obesity and hypertension in the Mexican population. *J Aging Health* **23**, 1141–1165.
57. Victora CG, Adair L, Fall C *et al.* (2008) Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* **371**, 340–357.
58. Currie J & Vogl T (2013) Early-life health and adult circumstance in developing countries. *Annu Rev Econ* **5**, 1–36.
59. Richter LM, Daelmans B, Lombardi J *et al.* (2017) Investing in the foundation of sustainable development: pathways to scale up for early childhood development. *Lancet* **389**, 103–118.
60. Jones AD, Mundo-Rosas V, Cantoral A *et al.* (2016) Household food insecurity in Mexico is associated with the co-occurrence of overweight and anemia among women of reproductive age, but not female adolescents. *Matern*



- Child Nutr.* Published online: E-Pub 14 December. doi: 10.1111/mcn.12396.
61. Razak F, Corsi DJ, Slutsky AS *et al.* (2015) Prevalence of Body Mass Index Lower Than 16 Among Women in Low- and Middle-Income Countries. *JAMA* **314**, 2164–2171. doi: 10.1001/jama.2015.15666.
 62. Fernald LCH, Gertler PJ & Neufeld LM (2008) Role of cash in conditional cash transfer programmes for child health, growth, and development: an analysis of Mexico's Oportunidades. *Lancet* **371**, 828–837.
 63. Fernald LCH (2013) Promise, and risks, of conditional cash transfer programmes. *Lancet* **382**, 7.
 64. Bates K, Gjonca A & Leone T (2017) Double burden or double counting of child malnutrition? The methodological and theoretical implications of stunting overweight in low and middle income countries. *J Epidemiol Comm Health.* Published online: E-Pub 31 May. doi: 10.1136/jech-2017-209008.
 65. Doak C, LS A, M B *et al.* (2005) The dual burden household and the nutrition transition paradox. *Int J Obes* **29**, 129–136.