

A construction of infinitely many singular weak solutions to the equations of nonlinear elasticity

J. Sivaloganathan

Department of Mathematical Sciences, University of Bath,
Bath BA2 7AY, UK

S. J. Spector

Department of Mathematics, Southern Illinois University,
Carbondale, IL 62901–4408, USA

(MS received 12 January 2001; accepted 17 August 2001)

Radial deformations of a ball composed of a nonlinear elastic material and corresponding to cavitation have been much studied. In this paper we use rescalings to show that each such deformation can be used to construct infinitely many non-symmetric singular weak solutions of the equations of nonlinear elasticity for the same displacement boundary-value problem. Surprisingly, this property appears to have been unnoticed in the literature to date.

1. Introduction

Let $\Omega \subset \mathbb{R}^n$ denote the reference state of a nonlinearly elastic body, let $p \geq 1$ and let

$$\mathbf{u} : \Omega \rightarrow \mathbb{R}^n, \quad \mathbf{u} \in W^{1,p}(\Omega; \mathbb{R}^n), \quad \det \nabla \mathbf{u} > 0 \text{ a.e.},$$

denote a deformation of the body. In hyperelasticity we associate with each deformation the energy

$$E(\mathbf{u}) = \int_{\Omega} W(\nabla \mathbf{u}(\mathbf{x})) \, d\mathbf{x}, \quad (1.1)$$

where $W : M_+^{n \times n} \rightarrow \mathbb{R}^+$ is the stored-energy function and $M_+^{n \times n}$ denotes the $n \times n$ matrices with positive determinant (see, for example, [4]). The equilibrium equations of nonlinear elasticity are the Euler–Lagrange equations for (1.1) and can take a number of forms. For the purposes of this paper we note the following two forms,

$$\frac{\partial}{\partial x^\alpha} \left[W(\nabla \mathbf{u}) \delta_\alpha^\beta - u_{,\beta}^k \frac{\partial W}{\partial F_\alpha^k}(\nabla \mathbf{u}) \right] = 0, \quad \beta = 1, 2, \dots, n, \quad (1.2)$$

and

$$\frac{\partial}{\partial x^\alpha} \left[\frac{\partial W}{\partial F_\alpha^i}(\nabla \mathbf{u}) \right] = 0, \quad i = 1, 2, \dots, n, \quad (1.3)$$

where we have used the summation convention for repeated indices. (Equation (1.2) is sometimes referred to as the energy-momentum form of the equilibrium equations.)

Suppose further that the deformation \mathbf{u} satisfies the affine boundary condition

$$\mathbf{u}(\mathbf{x}) = \mathbf{A}\mathbf{x} \quad \text{for } \mathbf{x} \in \partial\Omega,$$

where $\mathbf{A} \in M_+^{n \times n}$ is given. We denote by \mathbf{u}^h the corresponding homogeneous deformation

$$\mathbf{u}^h(\mathbf{x}) \equiv \mathbf{A}\mathbf{x}.$$

Now let $\Omega = B$, the unit ball in \mathbb{R}^n , and decompose $B \approx \bigcup_{i=1}^\infty \bar{B}_{\varepsilon_i}(\mathbf{x}_i)$ up to (Lebesgue) measure zero as the countable disjoint union of closed balls, i.e.

$$\text{meas}\left\{B \setminus \bigcup_{i=1}^\infty \bar{B}_{\varepsilon_i}(\mathbf{x}_i)\right\} = 0, \quad \bar{B}_{\varepsilon_i}(\mathbf{x}_i) \cap \bar{B}_{\varepsilon_j}(\mathbf{x}_j) = \emptyset \quad \text{for } i \neq j, \tag{1.4}$$

where

$$\bar{B}_{\varepsilon_i}(\mathbf{x}_i) = \{\mathbf{x} : |\mathbf{x} - \mathbf{x}_i| \leq \varepsilon_i\} \subset B$$

and $|\cdot|$ denotes the Euclidean norm on \mathbb{R}^n . This can always be achieved, for example, by Vitali's theorem (see, for example, [6, p. 29]). Note that, by (1.4),

$$\omega_n = \omega_n \sum_{i=1}^\infty \varepsilon_i^n,$$

where ω_n is the volume of the unit ball in \mathbb{R}^n and hence

$$\sum_{i=1}^\infty \varepsilon_i^n = 1. \tag{1.5}$$

Next define $\tilde{\mathbf{u}} : B \rightarrow \mathbb{R}^n$ by

$$\tilde{\mathbf{u}}(\mathbf{x}) = \begin{cases} \varepsilon_i \mathbf{u}\left(\frac{\mathbf{x} - \mathbf{x}_i}{\varepsilon_i}\right) + \mathbf{A}\mathbf{x}_i & \text{if } \mathbf{x} \in B_{\varepsilon_i}(\mathbf{x}_i), \\ \mathbf{A}\mathbf{x} & \text{otherwise.} \end{cases} \tag{1.6}$$

Then $\tilde{\mathbf{u}} \in W^{1,p}(B; \mathbb{R}^n)$ and $(\tilde{\mathbf{u}} - \mathbf{u}^h) \in W_0^{1,p}(B; \mathbb{R}^n)$ (this construction is used in proposition 2.3 of Ball and Murat [2]). Moreover, by (1.6),

$$\begin{aligned} E(\tilde{\mathbf{u}}) &= \int_B W(\nabla \tilde{\mathbf{u}}(\mathbf{x})) \, d\mathbf{x} \\ &= \sum_{i=1}^\infty \int_{B_{\varepsilon_i}(\mathbf{x}_i)} W\left(\nabla \mathbf{u}\left(\frac{\mathbf{x} - \mathbf{x}_i}{\varepsilon_i}\right)\right) \, d\mathbf{x} \\ &= \sum_{i=1}^\infty \varepsilon_i^n \int_B W(\nabla \mathbf{u}(\mathbf{y})) \, d\mathbf{y} \\ &= \int_B W(\nabla \mathbf{u}(\mathbf{y})) \, d\mathbf{y} \end{aligned}$$

(where we have made the change of variables $\mathbf{y} = (\mathbf{x} - \mathbf{x}_i)/\varepsilon_i$ and used (1.5)). Therefore, $\tilde{\mathbf{u}}$ has the same energy as \mathbf{u} . We show in this paper that if \mathbf{u} is a radial cavitation solution, then $\tilde{\mathbf{u}}$ is also a *weak solution* to the equations (1.3) and (1.2) of nonlinear elasticity. We first gather in the next section some basic results on the radial cavitation problem.

2. Hypotheses on W

We assume throughout this paper that $W \in C^2(M_+^{n \times n})$ is isotropic and frame indifferent so that, for any $Q \in SO(n)$ (the $n \times n$ special orthogonal matrices), we have

$$W(FQ) = W(QF) = W(F) \quad \text{for all } F \in M_+^{n \times n}.$$

It is a consequence of the above assumption that

$$W(F) = \Phi(v_1, v_2, \dots, v_n) \quad \text{for all } F \in M_+^{n \times n}, \tag{2.1}$$

where Φ is a symmetric function of its arguments and v_1, v_2, \dots, v_n denote the singular values of F (i.e. the eigenvalues of $\sqrt{F^T F}$).

We assume that W is *strongly elliptic*,

$$\frac{\partial^2 W}{\partial F_\alpha^i \partial F_\beta^j}(F) a^i a^j b^\alpha b^\beta > 0 \tag{2.2}$$

for any $a, b \in \mathbb{R}^n \setminus \{0\}$. We finally assume that there exist $\varepsilon > 0$ and $C > 0$ such that

$$\left\| F^T \frac{\partial W}{\partial F}(AF) \right\| \leq C[W(F) + 1] \quad \text{for all } F \in M_+^{n \times n} \tag{2.3}$$

whenever $\|A - I\| < \varepsilon$. Here, $\|\cdot\|$ denotes the Euclidean norm on $M^{n \times n}$,

$$\|F\|^2 = F : F \quad \text{and} \quad F : G = \text{tr}(F^T G) \quad \text{for } F, G \in M^{n \times n}.$$

3. Results on radial cavitation

Suppose that $A = \lambda I$, $\lambda > 0$, $p \in [1, n)$ and

$$u(x) = \frac{r(R)}{R} x, \quad R = |x|, \quad r : [0, 1] \rightarrow [0, \infty), \tag{3.1}$$

is a radial map of B , where $r \in C^2((0, 1]) \cap C^1([0, 1])$. Then

$$\nabla u = r'(R) \frac{x \otimes x}{R^2} + \frac{r(R)}{R} \left[I - \frac{x \otimes x}{R^2} \right]$$

and, consequently (see, for example, [1]), the singular values, v_1, v_2, \dots, v_n , of $F = \nabla u$ are given by $v_1 = r'(R)$, $v_2 = \dots = v_n = r(R)/R$. Thus, using (2.1),

$$\frac{\partial W}{\partial F}(\nabla u) = \Phi_{,1}(R) \frac{x \otimes x}{R^2} + \Phi_{,2}(R) \left[I - \frac{x \otimes x}{R^2} \right] \tag{3.2}$$

and

$$\nabla u^T \frac{\partial W}{\partial F}(\nabla u) = r'(R) \Phi_{,1}(R) \frac{x \otimes x}{R^2} + \frac{r(R)}{R} \Phi_{,2}(R) \left[I - \frac{x \otimes x}{R^2} \right], \tag{3.3}$$

where $\Phi_{,i}$ denotes differentiation of Φ with respect to its i th argument and $\Phi(R)$ and $\Phi_{,i}(R)$ denote, respectively, Φ and $\Phi_{,i}$ evaluated at the arguments $v_1 = r'(R)$, $v_2 = \dots = v_n = r(R)/R$.

Now suppose that \mathbf{u} , given by (3.1), is a radial cavitation map, i.e. r satisfies

- (a) $r(1) = \lambda$;
- (b) $r'(R) > 0$ for $R \in (0, 1]$;
- (c) the radial equilibrium equation

$$\frac{d}{dR} [R^{n-1} \Phi_{,1}(R)] = (n-1)R^{n-2} \Phi_{,2}(R) \quad \text{for } R \in (0, 1);$$

- (d) $r(0) > 0$ and $\lim_{R \rightarrow 0^+} R^{n-1} \Phi_{,1}(R) = 0$.

REMARK 3.1. See [7] and the references therein for constitutive hypotheses on W under which there exist maps corresponding to radial cavitation.

REMARK 3.2. A straightforward calculation, using (3.2) and (3.3), shows that any solution of the radial equilibrium equation satisfies (1.2) and (1.3) in $B \setminus \{0\}$.

REMARK 3.3. The conditions $r'(R) > 0$ for $R \in (0, 1]$, $r(0) > 0$ and $r(1) = \lambda > 0$ imply that a map \mathbf{u} that corresponds to radial cavitation must be one-to-one on $B \setminus \{0\}$ and satisfy $\mathbf{u}(B) \subset \lambda B$.

REMARK 3.4. The assumption (2.2) of strong ellipticity yields $\Phi_{,11} > 0$ for the corresponding function Φ (given by (2.1)). Thus the radial equilibrium equation is a well-defined second-order differential equation.

REMARK 3.5. By lemma 4.1 of [1], a radial map \mathbf{u} of the form (3.1) lies in $W^{1,p}(B; \mathbb{R}^n)$, $p \geq 1$, if and only if the corresponding scalar function $r(\cdot)$ is absolutely continuous on closed subintervals of $(0, 1)$ and satisfies

$$\int_0^1 R^{n-1} \left[|r'(R)|^p + \left| \frac{r(R)}{R} \right|^p \right] dR < \infty.$$

The next result notes, in particular, that every radial cavitation map \mathbf{u} has finite energy.

LEMMA 3.6. *Let \mathbf{u} be a radial cavitation map (i.e. \mathbf{u} is given by (3.1), where r satisfies conditions (a)–(d) above). Then*

- (i) $E(\mathbf{u}) = \omega_n [\Phi(1) + (r(1) - r'(1)) \Phi_{,1}(1)]$;
- (ii) $\delta^n \left[\Phi(\delta) + \left(\frac{r(\delta)}{\delta} - r'(\delta) \right) \Phi_{,1}(\delta) \right] \rightarrow 0$ as $\delta \rightarrow 0^+$;
- (iii) $\delta^n [\Phi(\delta) - r'(\delta) \Phi_{,1}(\delta)] \rightarrow 0$ as $\delta \rightarrow 0^+$.

Proof. Parts (i) and (ii) of the lemma (stated for $n \geq 2$) follow from a straightforward generalization of proposition 1.13 in [10] (which proves the same results, under hypothesis (2.2), in the case $n = 3$). Part (iii) is then an immediate consequence of part (ii) and condition (d) above. □

LEMMA 3.7. *The radial cavitation map \mathbf{u} satisfies the weak form of the energy-momentum equations,*

$$\int_B \nabla \mathbf{v} : \left[W(\nabla \mathbf{u}) \mathbf{I} - \nabla \mathbf{u}^T \frac{\partial W}{\partial \mathbf{F}}(\nabla \mathbf{u}) \right] d\mathbf{x} = 0 \quad \text{for all } \mathbf{v} \in C_0^1(B; \mathbb{R}^n).$$

REMARK 3.8. The above equations arise as necessary conditions for a minimizer of (1.1) on using inner variations and the condition (2.3) (see [3]).

Proof. We first note that, by (2.3), the fact that \mathbf{u} has finite energy and the dominated convergence theorem,

$$\int_B \nabla \mathbf{v} : \mathbf{M}(\nabla \mathbf{u}) d\mathbf{x} = \lim_{\delta \rightarrow 0^+} \int_{B \setminus B_\delta(0)} \nabla \mathbf{v} : \mathbf{M}(\nabla \mathbf{u}) d\mathbf{x}, \tag{3.4}$$

where

$$\mathbf{M}(\nabla \mathbf{u}) = W(\nabla \mathbf{u}) \mathbf{I} - \nabla \mathbf{u}^T \frac{\partial W}{\partial \mathbf{F}}(\nabla \mathbf{u}). \tag{3.5}$$

Next, since \mathbf{u} satisfies (1.2) in $B \setminus \{0\}$, it follows from (3.3) and the divergence theorem that

$$\begin{aligned} \int_{B \setminus B_\delta(0)} \nabla \mathbf{v} : \mathbf{M}(\nabla \mathbf{u}) d\mathbf{x} &= \int_{\partial B \cup \partial B_\delta(0)} \mathbf{v} \cdot \mathbf{M}(\nabla \mathbf{u}) \mathbf{n} \\ &= \int_{\partial B_\delta(0)} -\mathbf{v} \cdot \mathbf{M}(\nabla \mathbf{u}) \frac{\mathbf{x}}{|\mathbf{x}|} \\ &= \int_{\partial B_\delta(0)} \mathbf{v} \cdot [\Phi(R) - r'(R)\Phi_{,1}(R)] \mathbf{n} \\ &= [\Phi(\delta) - r'(\delta)\Phi_{,1}(\delta)] \int_{B_\delta(0)} \operatorname{div} \mathbf{v} d\mathbf{x}. \end{aligned}$$

However,

$$\left| [\Phi(\delta) - r'(\delta)\Phi_{,1}(\delta)] \int_{B_\delta(0)} \operatorname{div} \mathbf{v} d\mathbf{x} \right| \leq C \delta^n |\Phi(\delta) - r'(\delta)\Phi_{,1}(\delta)| \rightarrow 0$$

as $\delta \rightarrow 0^+$ by lemma 3.6, which, together with (3.4) and the above calculation, yields the desired result. □

A slight modification of the above argument yields the following result, which holds in the case when the test function \mathbf{v} is not necessarily zero on ∂B .

LEMMA 3.9. *The radial cavitation map \mathbf{u} satisfies*

$$\int_B \nabla \mathbf{v} : \left[W(\nabla \mathbf{u}) \mathbf{I} - \nabla \mathbf{u}^T \frac{\partial W}{\partial \mathbf{F}}(\nabla \mathbf{u}) \right] d\mathbf{x} = [\Phi(1) - r'(1)\Phi_{,1}(1)] \int_{\partial B} \mathbf{v} \cdot \mathbf{n}$$

for all $\mathbf{v} \in C^1(\bar{B}; \mathbb{R}^n)$.

REMARK 3.10. It is a consequence of the invariance of the equilibrium equations under the scaling symmetry $(\mathbf{x}, \mathbf{u}) \rightarrow (\epsilon \mathbf{x}, \epsilon \mathbf{u})$ (with $\epsilon > 0$) that any weak solution $\mathbf{u} : B \rightarrow \mathbb{R}^n$ of (1.2) (or (1.3)) generates a corresponding weak solution $\mathbf{u}_\epsilon(\mathbf{x}) = \epsilon \mathbf{u}(\mathbf{x}/\epsilon)$ of (1.2) (or (1.3)) on the domain ϵB .

We note further for later use that, in the particular case that \mathbf{u} is a radial cavitation map, we also have the following consequence of lemma 3.9,

$$\int_{\epsilon B} \nabla \mathbf{v} : \left[W(\nabla \mathbf{u}_\epsilon) \mathbf{I} - \nabla \mathbf{u}_\epsilon^T \frac{\partial W}{\partial \mathbf{F}}(\nabla \mathbf{u}_\epsilon) \right] d\mathbf{x} = [\Phi(1) - r'(1)\Phi_{,1}(1)] \int_{\partial(\epsilon B)} \mathbf{v} \cdot \mathbf{n}$$

for all $\mathbf{v} \in C^1(\epsilon \bar{B}; \mathbb{R}^n)$.

The next two results correspond to lemmas 3.7 and 3.9 for the second form (1.3) of the equilibrium equations. These follow by analogous arguments to the above and so the proofs are omitted.

LEMMA 3.11. *The radial cavitation map satisfies the weak form of the equilibrium equations (1.3), i.e.*

$$\int_B \nabla \mathbf{v} : \frac{\partial W}{\partial \mathbf{F}}(\nabla \mathbf{u}) d\mathbf{x} = 0 \quad \text{for all } \mathbf{v} \in C_0^1(B; \mathbb{R}^n).$$

LEMMA 3.12. *The radial cavitation map satisfies*

$$\int_B \nabla \mathbf{v} : \frac{\partial W}{\partial \mathbf{F}}(\nabla \mathbf{u}) d\mathbf{x} = \Phi_{,1}(1) \int_{\partial B} \mathbf{v} \cdot \mathbf{n} \quad \text{for all } \mathbf{v} \in C^1(\bar{B}; \mathbb{R}^n).$$

REMARK 3.13. Lemmas 3.11 and 3.12 can also be obtained from theorem 4.2 in [1].

4. Construction of infinitely many singular weak solutions

We next apply the construction outlined earlier in (1.4)–(1.6) to a radial cavitation map \mathbf{u} to construct a corresponding map $\tilde{\mathbf{u}} : B \rightarrow \mathbb{R}^n$ with infinitely many discontinuities. The main results of this paper are the following two theorems.

THEOREM 4.1. *The map $\tilde{\mathbf{u}}$ lies in $W^{1,p}(B; \mathbb{R}^n)$ for $1 \leq p < n$, is injective almost everywhere and is a weak solution of the equations*

$$\frac{\partial}{\partial x^\alpha} \left[W(\nabla \mathbf{u}) \delta_\alpha^\beta - u_{,\beta}^k \frac{\partial W}{\partial F_\alpha^k}(\nabla \mathbf{u}) \right] = 0 \quad \text{in } B, \quad \beta = 1, 2, \dots, n,$$

i.e.

$$\int_B \nabla \mathbf{v} : \left[W(\nabla \tilde{\mathbf{u}}) \mathbf{I} - \nabla \tilde{\mathbf{u}}^T \frac{\partial W}{\partial \mathbf{F}}(\nabla \tilde{\mathbf{u}}) \right] d\mathbf{x} = 0 \quad \text{for all } \mathbf{v} \in C_0^1(B; \mathbb{R}^n).$$

Proof. The fact that $\tilde{\mathbf{u}} \in W^{1,p}(B; \mathbb{R}^n)$ and is injective almost everywhere follows directly (see remarks 3.3 and 3.5). Now let $\mathbf{v} \in C_0^1(B; \mathbb{R}^n)$. Then, by the definition

of $\tilde{\mathbf{u}}$, equation (3.5), lemma 3.9 and remark 3.10,

$$\begin{aligned} \int_B \nabla \mathbf{v} : \mathbf{M}(\nabla \tilde{\mathbf{u}}) \, d\mathbf{x} &= \sum_{i=1}^{\infty} \int_{B_{\varepsilon_i}(\mathbf{x}_i)} \nabla \mathbf{v} : \mathbf{M}(\nabla \tilde{\mathbf{u}}) \, d\mathbf{x} \\ &= \sum_{i=1}^{\infty} [\Phi(1) - r'(1)\Phi_{,1}(1)] \int_{\partial B_{\varepsilon_i}(\mathbf{x}_i)} \mathbf{v} \cdot \mathbf{n} \\ &= [\Phi(1) - r'(1)\Phi_{,1}(1)] \sum_{i=1}^{\infty} \int_{B_{\varepsilon_i}(\mathbf{x}_i)} \operatorname{div} \mathbf{v} \, d\mathbf{x} \\ &= [\Phi(1) - r'(1)\Phi_{,1}(1)] \int_B \operatorname{div} \mathbf{v} \, d\mathbf{x} \\ &= [\Phi(1) - r'(1)\Phi_{,1}(1)] \int_{\partial B} \mathbf{v} \cdot \mathbf{n} = 0. \end{aligned}$$

□

THEOREM 4.2. *The map $\tilde{\mathbf{u}}$ is a weak solution of the equations*

$$\frac{\partial}{\partial x^\alpha} \left[\frac{\partial W}{\partial F_i^\alpha}(\nabla \mathbf{u}) \right] = 0 \quad \text{in } B, \quad i = 1, 2, \dots, n,$$

i.e.

$$\int_B \nabla \mathbf{v} : \frac{\partial W}{\partial \mathbf{F}}(\nabla \tilde{\mathbf{u}}) \, d\mathbf{x} = 0 \quad \text{for all } \mathbf{v} \in C_0^1(B; \mathbb{R}^n).$$

Proof. The proof of this result is exactly analogous to that of the last theorem and is therefore omitted. □

5. Concluding remarks

We note that the construction used in the last two theorems is independent of the domain chosen and therefore yields singular weak solutions for any domain (given a domain Ω , write $\Omega \approx \bigcup_{i=1}^{\infty} \bar{B}_{\varepsilon_i}(\mathbf{x}_i)$ up to measure zero, define $\tilde{\mathbf{u}}$ by (1.6) and proceed as in the case $\Omega = B$). Note that, in the case when \mathbf{u} is a radial cavitation map, although the ‘composite deformation’ $\tilde{\mathbf{u}}$ constructed according to (1.6) is continuous away from the singular set $S = \{\mathbf{x}_i : i \in \mathbb{N}\}$, it is not C^1 away from S . In particular, $\tilde{\mathbf{u}}$ is not C^1 on the boundary of the balls $B_{\varepsilon_k}(\mathbf{x}_k)$. This follows from the first equation of (1.6), since $\|\nabla \mathbf{u}\|$ is infinite at the origin and each $\mathbf{x}^* \in \partial B_{\varepsilon_k}(\mathbf{x}_k)$ is an accumulation point of the centres of the balls. (Otherwise, without loss of generality, there exists $\mathbf{x}^* \in \partial \bar{B}_{\varepsilon_1}(\mathbf{x}_1)$ with $\mathbf{x}^* \notin \{\mathbf{x}_i\}$, and hence there is a $\delta > 0$ such that $\mathbf{x}_i \notin B_{3\delta}(\mathbf{x}^*)$ for all i . By (1.5), there exists $M \in \mathbb{N}$ such that $\varepsilon_i < \delta$ and hence $B_\delta(\mathbf{x}^*) \cap \bar{B}_{\varepsilon_i}(\mathbf{x}_i) = \emptyset$ for $i > M$. Also, $\mathbf{x}^* \notin K := \bigcup_{i=2}^M \bar{B}_{\varepsilon_i}(\mathbf{x}_i)$, which is closed, so $d := \operatorname{dist}(\mathbf{x}^*, K) > 0$. Consequently, $D := B_{\min(d,\delta)}(\mathbf{x}^*) \setminus \bar{B}_{\varepsilon_1}(\mathbf{x}_1)$ has positive measure and $D \cap \bar{B}_{\varepsilon_i}(\mathbf{x}_i) = \emptyset$ for every i , a contradiction.)

Though it has been well known to workers in nonlinear elasticity that the construction (1.6) yields deformations $\tilde{\mathbf{u}}$ with equal energy to \mathbf{u} , it seems to have been (surprisingly) overlooked to date that in the case when \mathbf{u} is a radial cavitation map this construction also produces further weak solutions. Hence the examples in

this paper have implications for proving regularity of weak solutions in nonlinear elasticity (see, for example, [5]). In this context, we mention the recent interesting example of Müller and Šverák [9] of a $W^{1,\infty}$ weak solution that is nowhere C^1 . Finally, we note that an analogous approach to that used in this paper can be used to show that the maps $\tilde{\mathbf{u}}$ constructed from rescalings of a radial cavitation map generate weak solutions of the Cauchy form of the equilibrium equations,

$$\frac{\partial}{\partial y^\alpha} T_{i\alpha}(\mathbf{y}) = 0, \quad \mathbf{y} \in \mathbf{u}(\Omega),$$

where

$$T_{i\alpha}(\mathbf{u}(\mathbf{x})) = \frac{1}{\det \nabla \mathbf{u}(\mathbf{x})} \frac{\partial W}{\partial F_\beta^i} (\nabla \mathbf{u}(\mathbf{x})) u_{,\beta}^\alpha(\mathbf{x}).$$

The details are left to the interested reader. (Though, of course, in this case, the deformed body $\tilde{\mathbf{u}}(\Omega)$ will vary according to the particular decomposition of Ω used in constructing $\tilde{\mathbf{u}}$.)

Acknowledgments

J.S. thanks J. M. Ball and R. D. James for helpful discussions related to this work. The authors acknowledge the support of the Isaac Newton Institute (Cambridge) and Southern Illinois University. This material is based upon work supported in part by the National Science Foundation under grant no. 0072414.

References

- 1 J. M. Ball. Discontinuous equilibrium solutions and cavitation in nonlinear elasticity. *Phil. Trans. R. Soc. Lond. A* **306** (1982), 557–611.
- 2 J. M. Ball and F. Murat. $W^{1,p}$ -quasiconvexity and variational problems for multiple integrals. *J. Funct. Analysis* **58** (1984), 225–253.
- 3 P. Baumann, N. C. Owen and D. Phillips. Maximum principles and *a priori* estimates for a class of problems from nonlinear elasticity. *Analyse Nonlineaire* **8** (1991), 119–157.
- 4 P. G. Ciarlet. Three-dimensional elasticity. In *Mathematical elasticity*, vol. 1 (Amsterdam: North-Holland, 1988).
- 5 L. C. Evans. Quasiconvexity and partial regularity in the calculus of variations. *Arch. Ration. Mech. Analysis* **95** (1986), 227–252.
- 6 M. Giaquinta, G. Modica and J. Soucek. *Cartesian currents in the calculus of variations* (Springer, 1998).
- 7 C. O. Horgan and D. A. Polignone. Cavitation in nonlinearly elastic solids: a review. *Appl. Mech. Rev.* **48** (1995), 471–485.
- 8 S. Müller and S. J. Spector. An existence theory for nonlinear elasticity that allows for cavitation. *Arch. Ration. Mech. Analysis* **131** (1995), 1–66.
- 9 S. Müller and V. Šverák. Convex integration for Lipschitz mappings and counterexamples to regularity. Preprint 26/1999, Max-Planck-Institute, Leipzig.
- 10 J. Sivaloganathan. Uniqueness of regular and singular equilibria for spherically symmetric problems of nonlinear elasticity. *Arch. Ration. Mech. Analysis* **96** (1986), 97–136.

(Issued 16 August 2002)