

# *Hammondia* isolated from dogs and foxes are genetically distinct

J. ABEL<sup>1</sup>, G. SCHARS<sup>2</sup>, K. ORZESZKO<sup>3</sup>, R. B. GASSER<sup>3,4</sup> and J. T. ELLIS<sup>1\*</sup>

<sup>1</sup>Department of Cell and Molecular Biology, University of Technology, Sydney, Westbourne Street, Gore Hill, NSW 2065, Australia

<sup>2</sup>Institute of Epidemiology, Friedrich-Loeffler-Institut-Federal Research Institute for Animal Health, Seestrasse 55, 16868 Wusterhausen, Germany

<sup>3</sup>Department of Veterinary Science, The University of Melbourne, 250 Princes Highway, Werribee, Victoria 3030, Australia

<sup>4</sup>Biotechnology Research Institute, Macquarie University, North Ryde, NSW 2109, Australia

(Received 5 May 2005; revised 26 July 2005; accepted 27 July 2005; first published online 27 September 2005)

## SUMMARY

*Hammondia heydorni* is regarded as a protozoan parasite that uses canids, e.g. dogs and foxes, as definitive hosts, but clinical signs of infection are rare. This study therefore took advantage of the opportunity to study an oocyst population from the faeces of a dog suffering from intermittent bouts of diarrhoea. Oocysts from the naturally infected dog were shown to be *H. heydorni* by using the polymerase chain reaction combined with DNA sequencing as a diagnostic tool. † A comparison of the first internal transcribed spacer (ITS1) sequence of ribosomal DNA obtained with those from other dog and fox oocysts, previously regarded as *H. heydorni*, showed these oocysts contained identical ITS1 sequences. However, the oocyst DNA from the fox and dog differed by the presence/absence of a 9 bp insertion/deletion within intron 1 of the alpha tubulin gene, and this difference was conserved across a number of different oocyst populations from the 2 species of host. A PCR assay was established that takes advantage of this insertion/deletion and is able to differentiate between the 2 oocyst populations. This study therefore provides evidence that *H. heydorni* oocysts from dogs and foxes represent 2 distinct genetic lineages that can be differentiated using a PCR, which targets the alpha tubulin locus.

Key words: *Hammondia*, *Neospora*, PCR, DNA sequencing, internal transcribed spacer, alpha tubulin gene, speciation.

## INTRODUCTION

*Hammondia heydorni* is a cyst-forming coccidian parasite with an obligatory 2-host life-cycle (Heydorn, 1973; Fayer, 1974; Dubey and Fayer, 1976). In addition to dogs, also foxes and coyotes are regarded as definitive hosts in the parasite's life-cycle, whereas various herbivores were identified as intermediate hosts (Dubey *et al.* 2002*a*). Experimental data on a fox-derived *Hammondia* isolate suggest that infection of the definitive host occurs by ingestion of raw body tissues (e.g. raw meat) containing tissue cysts and leads to the shedding of unsporulated oocysts (Schars *et al.* 2003). Sporulation of oocysts can be detected within 3 days after their excretion in faeces (Blagburn *et al.* 1988), and intermediate hosts become infected by ingestion of the infective sporulated oocysts (Schars *et al.* 2002; Mohammed *et al.* 2003).

\* Corresponding author: Department of Cell and Molecular Biology, University of Technology, Sydney, Westbourne Street, Gore Hill, NSW 2065, Australia. Tel: +612 9514 4161. Fax: +612 95144003. E-mail: john.ellis@uts.edu.au (J.T. Ellis)

† The nucleotide sequence data reported in this paper are available from GenBank under the following Accession numbers DQ183058, DQ183059 and DQ022687.

*Hammondia heydorni* is closely related to other coccidian parasites, such as *Toxoplasma gondii*, *Neospora caninum* and *Hammondia hammondi*. However, there is still considerable controversy about the taxonomy and phylogenetic relationships of these species and the taxonomic status of *H. heydorni* and *N. caninum* (see Ellis *et al.* 1999*b*; Dubey *et al.* 2002*a*; Heydorn and Mehlhorn, 2002).

In order to learn more about the biology of *H. heydorni*, new sensitive and specific methods such as serological tests and PCR are required (Slapeta *et al.* 2002*a*; Dubey *et al.* 2002*a*). PCR is a useful tool for the detection and identification of the cyst-forming coccidia (Ellis, 1998). In particular, the internal transcribed spacers (ITS) of rDNA or the alpha tubulin sequence have recently been reported as valuable genetic markers for the study of this species (Ellis *et al.* 1999*b*; Siverajah *et al.* 2002; Slapeta *et al.* 2002*a*). These sequences display reasonably high levels of conservation, yet there is sufficient sequence variation among related species to allow discrimination (e.g., Dubey *et al.* 2002*a*; Ellis and Pomroy, 2003).

Recently, *Hammondia* oocysts from foxes were compared with *H. heydorni* obtained from dogs. In contrast to the comparison of the ITS1, a comparison of partial large subunit ribosomal DNA sequences among *Hammondia* isolates from foxes

and dogs showed them to be genetically different from each other (Schaes *et al.* 2002; Mohammed *et al.* 2003). Although the number of nucleotide differences was low, these observations imply that either there may be considerable genetic diversity among *H. heydorni* isolates, as recently shown by Sreekumar *et al.* (2004), or that more than one species may exist within the taxon known as *H. heydorni*.

The present study describes the molecular characterization of oocysts shed by a naturally infected dog. The study, using PCR and sequencing of the ITS1, showed them to be consistent with *H. heydorni*. Partial sequencing of the alpha tubulin locus from these parasites was also performed, and the sequence obtained was compared with that derived from 2 oocyst populations obtained from experimentally infected foxes (designated Fox 2000/1, Schaes *et al.* 2002; and HhO, Mohammed *et al.* 2003); as well as from another naturally infected dog (Siverajah *et al.* 2002; Ellis and Pomroy, 2003). The data show the oocysts from dog and from fox to be genetically distinct within intron 1 of the alpha tubulin gene. A PCR is also described, which takes advantage of an insertion/deletion found at this site that allows for the differentiation of the fox from the dog-derived *Hammondia* oocysts. This test should prove useful in the study of the prevalence and host distribution of these two parasite lineages, which are currently called *H. heydorni*.

## MATERIALS AND METHODS

### Case study

In October 2003, a dog (5-year-old, neutered female Hungarian Vizsla; body weight 26 kg) with signs of haemolytic anaemia was presented in the Department of Veterinary Science of the University of Melbourne. Laboratory findings confirmed the clinical diagnosis and the dog was given immunosuppressive treatment (prednisolone (20 mg), cyclosporine (100 mg) twice per day and azothiaprime (50 mg) once daily) and supportive therapy for 10 days. During this time amoxicillin (500 mg) was also administered twice per day for 7 days. Subsequently, the dog was administered long-term treatment with azothiaprime and prednisolone. After discharge from hospital, the dog developed cystitis and recurrent bouts of diarrhoea. Bacterial culture of urine (obtained by cystocentesis) revealed *Escherichia coli*, and the cystitis was treated with therapeutic doses of enrofloxacin. To investigate causative agents involved in the diarrhoea, coproscopical examinations were conducted. Examination by light microscopy revealed the presence of coccidian oocysts, which were subsequently purified by flotation on saturated zinc sulphate using standard methods.

The oocysts obtained (called HhM) from the dog were allowed to sporulate for 1 week in 2.5% potassium dichromate ( $K_2CrO_4$ ), washed in phosphate-buffered saline (PBS) and then examined for ultraviolet autofluorescence (wavelength: 330–385 nm) employing an Olympus BX51 microscope as described previously (Lindquist *et al.* 2003).

Twenty oocysts were examined by light microscopy and measured by standard procedures. The remaining oocysts were washed and resuspended in PBS, ruptured by vortexing them in an Eppendorf tube with glass beads (710–1180  $\mu$ m, Sigma) followed by multiple cycles of freeze/thaw. Subsequently, the sample was centrifuged and the supernatant transferred to a fresh tube, and an equal amount of lysis buffer (10 mM Tris pH 8, 100 mM EDTA pH 8, 1% SDS) with Proteinase K (100  $\mu$ g/ml) added. The sample was then incubated for 2 h at 56 °C, after which it was extracted twice with a standard mix of phenol/chloroform/isoamyl alcohol (25:24:1), and once with chloroform. The DNA was precipitated with ethanol, washed once with 70% ethanol, and then resuspended in sterile 10 mM Tris-1 mM EDTA and stored at –20 °C for analysis.

### Other DNAs included in this study

Genomic DNA obtained from oocysts shed by experimentally infected foxes (Fox 2000/1, Schaes *et al.* 2002; HhO, Mohammed *et al.* 2003), a naturally infected dog (HhNZ; Ellis and Pomroy, 2003) and an experimentally infected dog (Giessen-1999; Schaes *et al.* 2005) were also used in this study for comparative purposes. Genomic DNA from *N. caninum* (NC-1 strain) and *T. gondii* (RH) were included as controls.

### PCR and DNA sequencing

Purified genomic DNA was used as a template in the PCR. The ITS1 region was amplified as described previously (Ellis *et al.* 1999b; Ellis and Pomroy, 2003), using the primers TIM3 and TIM11.

For the amplification of a small fragment of the alpha tubulin gene spanning intron 1, primers AT9 (5'TGGCTGAGTAAGCACGGATTTTTC) and AT264 (5'GTGTCCTGTTCCCGCGTGTG-AT) were designed from gene sequence data reported by Siverajah *et al.* (2002). The specificity of the primers was evaluated using genomic DNA from *Hammondia*, *N. caninum* and *T. gondii*. A control containing no DNA was also performed, in order to exclude cross-contamination of any reagents. The PCR reaction contained 1  $\times$  PCR buffer (Finzyme), 250  $\mu$ M of each dNTP, 1U Dynazyme II F-501L polymerase (Finzyme), 20  $\mu$ g/ml of bovine serum albumin (Fluka) in a volume of 25  $\mu$ l. The PCR products were subjected




Fig. 1. *Hammondia heydorni* oocysts (HhM) shed by the naturally infected dog. Bar represents 10  $\mu\text{m}$ . (A) Unsporulated oocyst in water. (B) Sporulated oocyst containing 2 sporocysts in water.

to agarose (1.5%) gel electrophoresis in Tris borate-EDTA buffer. For sequencing gel-purified (Qiagen gel extraction kit) amplification products were used. In the case of the Giessen-1999 isolate, unpurified amplification products were used. Amplicons were sequenced at least twice both directions. The PCR cycling conditions were 94 °C for 5 min, followed by 50 cycles of 56 °C for 1 min, 72 °C for 1 min and 94 °C for 1 min. The PCR was finished with 56 °C for 1 min and a final extension at 72 °C for 5 min.

The program AssemblyLign™ (International Biotechnologies) was used to align ITS1 or tubulin sequences, and the consensus sequences obtained were compared against sequences deposited in the NCBI (National Centre of Bioinformatics) database using the program Blast N.

## RESULTS

### Case study

Microscopical examination of the faeces from the dog revealed oocysts consistent with *H. heydorni* or *N. caninum*, based on morphological characteristics. In water the thick-walled oocysts were  $11.4 \pm 0.5 \times 12.3 \pm 0.6 \mu\text{m}$  in size (mean  $\pm$  standard deviation) and contained 2 sporocysts (and 4 sporozoites in each sporocyst) after sporulation (Fig. 1). This morphology was consistent with that described for *N. caninum*, *H. heydorni* and *T. gondii* (see Lindsay, Upton and Dubey, 1999). The oocysts showed autofluorescence under ultraviolet light.

The ITS1 region amplified from HhM oocyst DNA was ~600 bp in size. The PCR product was sequenced and an investigation of present sequence databases with the consensus sequence (475 bp GenBank Accession number DQ183058) revealed 100% identity with/to previous ITS1 sequences from *H. heydorni* from canids (GenBank Accession no. AY189897, Giessen-1999; AF508030, HhNZ).

The species specificity of primers AT9 and AT264 was assessed by PCR using genomic DNA from *Hammondia*, *N. caninum* and *T. gondii*. A PCR product was only obtained from *Hammondia* oocysts under the conditions tested (Fig. 2). The partial




Fig. 2. PCR of genomic DNA from cyst-forming coccidia using 3 different primer pairs. HY, *H. heydorni* Giessen-1999; HV, *Hammondia* sp. Fox-2000/1; TG, *Toxoplasma gondii* (RH strain); NC, *Neospora caninum* (NC-1 strain); BU, No DNA negative control; M, molecular weight marker comprising a 100 bp ladder. All DNAs produce a PCR product with primers TIM3/TIM11 (A), while only *Hammondia* DNA gives a product with primers AT9 and AT264 (B). DNA derived from *Hammondia* oocysts shed by foxes produce a PCR product with primers HhGF and HhGR (C).

alpha tubulin sequence (Accession no. DQ183059) obtained from HhM PCR products (using AT9 and AT264) was the same as that determined previously (Accession no. AY169962). This DNA was extracted from oocysts found in the faeces of a naturally infected dog from New Zealand (HhNZ; Siverajah *et al.* 2002; Ellis and Pomroy, 2003). An identical sequence was obtained for the alpha tubulin gene spanning intron 1 from *H. heydorni* Giessen-1999 (Accession number DQ022687). Based on these PCR and DNA sequencing data results, the evidence shows the oocyst sample HhM is genetically consistent with *H. heydorni*.

### Comparison of tubulin sequences

Comparison of the 3 partial alpha tubulin sequences obtained from dog oocysts (samples HhM, HhNZ and Giessen-1999) with those determined from

<b>FOX2000/1</b>	CGCGTACCATTTA <b><u>CCATGCAC</u></b> <b><u>TCCATGCAC</u></b> ACATATGCAT	
<b>HhO</b>	CGCGTACCATTTA <b><u>CCATGCAC</u></b> <b><u>TCCATGCAC</u></b> ACATATGCAT	
<b>HhM</b>	CGCGTACCATTTA <b><u>CCATGCAC</u></b>	ACATATGCAT
<b>HhNZ</b>	CGCGTACCATTTA <b><u>CCATGCAC</u></b>	ACATATGCAT
<b>Giessen-1999</b>	CGCGTACCATTTA <b><u>CCATGCAC</u></b>	ACATATGCAT

Fig. 3. Comparison of partial  $\alpha$  tubulin gene sequences from 5 isolates of *Hammondia heydorni* (HhM, oocysts from this study; HhNZ from Ellis and Pomroy (2003); Fox 2000/1 described by Schares *et al.* (2002); HhO described by Mohammed *et al.* (2003); Giessen-1999 described by Schares *et al.* (in preparation). The repetitive sequence is in bold and underlined. The fox sequences contain 2 copies of this repeat, whereas the dog sequences contain only 1. Sequence orientation is 5 to 3'.

2 populations of fox oocysts (HhO and FOX 2000/1) showed that both of the latter sequences contained a 9-bp insertion (TCCATGCAC) (Fig. 3). This difference in size was detectable upon agarose gel (1.5%) electrophoresis of the PCR products (not shown). Eight of the bases found in this insertion were also present in the sequence present immediately following the site of insertion. In order to confirm the presence/absence of the insertion/deletion amongst the isolates, PCR primers were designed (HhGF, 5'CGCATTCAAAAAGTGCCACAATTG; HhGR, 5'ACTCCATGCACACATATGCAT) which should exclusively produce a PCR product from DNA of fox oocysts because HhGR bridges the region corresponding to the insertion/deletion. As shown in Fig. 2, a PCR product was obtained exclusively from Fox 2000/1 DNA only. No PCR products were amplified from *H. heydorni* DNA extracted from dog-derived oocysts, *T. gondii*, *N. caninum* or from the no-DNA control.

#### DISCUSSION

Clinical cases of hammondiosis in dogs are rare. This study therefore took advantage of the opportunity to study an oocyst population from the faeces of a dog suffering from intermittent bouts of diarrhoea. This dog had previously been treated for haemolytic anaemia, and it is interesting to note that others have previously suggested that the immunosuppressive treatment for this disease had rendered another dog susceptible to toxoplasmosis (Webb *et al.* 2005).

The purified, sporulated oocysts obtained were  $11.4 \pm 0.5 \times 12.3 \pm 0.6 \mu\text{m}$  in size, when measured in water, contained 2 sporocysts and had a thick, colourless wall. These morphological features are consistent with the morphology of oocysts of *T. gondii* (from felids), *H. heydorni* (from canids) and *N. caninum* (from canids) (Lindsay *et al.* 1999). Similar-sized oocysts from these species ( $12.6 \times 11.9 \mu\text{m}$  and  $10\text{--}14.1 \times 9.2\text{--}13.1 \mu\text{m}$  respectively) respectively are indistinguishable by light microscopy (Heydorn, 1973; Blagburn *et al.* 1988). The

structure of the sporulated oocysts is also similar among all 3 species, containing 2 sporocysts and 4 sporozoites/sporocyst.

A great deal of discussion about the taxonomic status of *H. heydorni* and *N. caninum* arose from the discovery of *N. caninum* oocysts in the faeces of dogs (McAllister *et al.* 1998; Mehlhorn and Heydorn, 2000; Heydorn and Mehlhorn, 2002). However, it is now generally accepted that at least 2 *Hammondia*-/*Neospora*-like species with morphologically similar oocyst stages but with distinct biological, immunological, ultrastructural and molecular features use the dog as definitive host (reviewed by Dubey *et al.* 2002*a,b*). Nevertheless, doubts remain as to whether studies on *H. heydorni*-like oocysts conducted before 1998 were performed using a homogeneous population of parasites. For instance, Müller *et al.* (2001) verified by PCR methodologies that the isolate originally designated as *H. heydorni*-Berlin-1996 contained *N. caninum*.

Molecular tools can assist in making a specific identification of a species of a parasite present in clinical specimens (Ellis, 1998). The ITS1 rDNA region has repeatedly proven useful for the specific identification of coccidian species (Holmdahl and Mattson, 1996; Payne and Ellis, 1996; Buxton *et al.* 1998; Ellis *et al.* 1999*a*; Slapeta *et al.* 2002*a,b*; Ellis and Pomroy, 2003; Mohammed *et al.* 2003). Thus this locus was therefore amplified and sequenced from the oocyst sample HhM. Database searches revealed that the sequence determined herein was identical to that obtained from 2 other *H. heydorni* isolates (AY189897, AF508030). The ITS1 sequence was therefore consistent with *H. heydorni*. These findings were supported by the analysis of partial alpha tubulin sequences, which upon database searches also showed complete identity to a sequence from another *H. heydorni* isolate from a naturally infected dog (AY169962) (Siverajah *et al.* 2002; Ellis and Pomroy, 2003). As shown in the present study, the alpha tubulin gene spanning intron 1 from another dog-derived *H. heydorni* isolate (Giessen-1999) was also identical to both that of HhM and HhNZ.

Genetic analysis of the large subunit of rDNA from oocysts found in the faeces of an Arabian red fox (Mohammed *et al.* 2003) confirmed the identification of sequence divergence in the rDNA of *Hammondia* reported by Schares *et al.* (2002). As the latter sequence is considered to be highly conserved between species, analysis of another conserved locus, the alpha tubulin gene, was also included in this present study. Although the precise structure of the *Hammondia* alpha tubulin gene is unknown, by comparison to the *N. caninum* gene, the region analysed spans what is predicted to be intron 1 (Siverajah *et al.* 2002). The alpha tubulin sequence obtained from the dog oocysts (samples HhM, HhNZ and Giessen-1999) were compared with those obtained from 2 oocyst populations obtained from foxes (Schares *et al.* 2002; Mohammed *et al.* 2003). Analysis by agarose gel electrophoresis revealed that the PCR products were slightly different in size. Subsequent sequencing confirmed this finding by showing that the sequences generated from the two different oocyst populations obtained from foxes both had an insert of 9 bp (TCCATGCAC) when compared with that obtained from the dog oocysts. PCR of oocyst DNA using a primer spanning the region of the insertion/deletion gave a product for the oocysts from fox but not from the oocysts from dog thereby confirming the presence of the deletion. Taken together, the results of the present study suggest that *H. heydorni*-like oocysts from dogs are genetically different from those shed by foxes. A large-scale evaluation including extended sets of isolates from dog and fox need to be examined to test the present proposal.

A recent study of sequence polymorphisms found in a variety of introns (including those of the alpha tubulin locus) among different isolates of *T. gondii* concluded these polymorphisms correlated with distinct clonal lineages (Su *et al.* 2003). This consideration may also apply to these *H. heydorni*-like organisms, which during speciation may have evolved into 2 genetically distinct lineages: one uses dogs as definitive hosts and one employing foxes. Nevertheless, before conclusions regarding the existence of another species of *Hammondia* can be made, further research should focus on establishing whether biological, ecological and epidemiological differences exist between them.

Part of this work was carried out by J.A. as an undergraduate research project for the University of Regensburg, while at the University of Technology, Sydney. We thank Andrea Bärwald for technical assistance. We thank Leonie Richards (Melbourne University) for referring this case to us.

#### REFERENCES

- Blagburn, B. L., Lindsay, D. S., Swango, L. J., Pidgeon, G. L. and Braund, K. G. (1988). Further characterization of the biology of *Hammondia heydorni*. *Veterinary Parasitology* **27**, 193–198.
- Buxton, D., Maley, S. W., Wright, S., Thomson, K. M., Rae, A. G. and Innes, E. A. (1998). The pathogenesis of experimental neosporosis in pregnant sheep. *Journal of Comparative Pathology* **118**, 267–279.
- Dubey, J. P. and Fayer, R. (1976). Development of *Isospora bigemina* in dogs and other mammals. *Parasitology* **73**, 371–380.
- Dubey, J. P., Barr, B. C., Barta, J. R., Bjerkås, I., Björkman, C., Blagburn, B. L., Bowman, D. D., Buxton, D., Ellis, J. T., Gottstein, B., Hemphill, A., Hill, D. E., Howe, D. K., Jenkins, M. C., Kobayashi, Y., Koudela, B., Marsh, A. E., Mattsson, J. G., McAllister, M. M., Modrý, D., Omata, Y., Sibley, L. D., Speer, C. A., Trees, A. J., Uggla, A., Upton, S. J., Williams, D. J. L. and Lindsay, D. S. (2002a). Redescription of *Neospora caninum* and its differentiation from related coccidia. *International Journal for Parasitology* **32**, 929–946.
- Dubey, J. P., Hill, D. E., Lindsay, D. S., Jenkins, M. C., Uggla, A. and Speer, C. A. (2002b). *Neospora caninum* and *Hammondia heydorni* are separate species/organisms. *Trends in Parasitology* **18**, 66–69.
- Ellis, J. (1998). Polymerase chain reaction approaches for the detection of *Neospora caninum* and *Toxoplasma gondii*. *International Journal for Parasitology* **28**, 1053–1060.
- Ellis, J. T., McMillan, D., Ryce, C., Payne, S., Atkinson, R. and Harper, P. A. W. (1999a). Development of a single tube nested polymerase chain reaction assay for the detection of *Neospora caninum* DNA. *International Journal for Parasitology* **29**, 1589–1596.
- Ellis, J. T., Morrison, D. A., Liddell, S., Jenkins, M. C., Mohammed, O. B., Ryce, C. and Dubey, J. P. (1999b). The genus *Hammondia* is paraphyletic. *Parasitology* **118**, 357–362.
- Ellis, J. T. and Pomroy, W. E. (2003). *Hammondia heydorni* oocysts in the faeces of a greyhound in New Zealand. *New Zealand Veterinary Journal* **51**, 38–39.
- Fayer, R. (1974). Development of *Sarcocystis fusiformis* in the small intestines of the dog. *Journal of Parasitology* **60**, 660–665.
- Heydorn, A. O. (1973). Zum Lebenszyklus der kleinen Form von *Isospora bigemina* des Hundes. Rind und Hund als mögliche Zwischenwirte. *Berliner und Münchner Tierärztliche Wochenschrift* **86**, 323–329.
- Heydorn, A. O. and Mehlhorn, H. (2002). *Neospora caninum* is an invalid species name: an evaluation of facts and statements. *Parasitology Research* **88**, 175–184.
- Holmdahl, O. J. M. and Mattsson, J. G. (1996). Rapid and sensitive identification of *Neospora caninum* by *in vitro* amplification of the internal transcribed spacer 1. *Parasitology* **112**, 177–182.
- Lindquist, H. D., Bennett, J. W., Hester, J. D., Ware, M. W., Dubey, J. P. and Everson, W. V. (2003). Autofluorescence of *Toxoplasma gondii* and related coccidian oocysts. *Journal of Parasitology* **89**, 865–867.
- Lindsay, D. S., Upton, S. J. and Dubey, J. P. (1999). A structural study of the *Neospora caninum* oocyst. *International Journal for Parasitology* **29**, 1521–1523.

- McAllister, M. M., Dubey, J. P., Lindsay, D. S., Jolley, W. R., Wills, R. A. and McGuire, A. M.** (1998). Dogs are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* **28**, 473–478.
- Mehlhorn, H. and Heydorn, A. O.** (2000). *Neospora caninum*: is it really different from *Hammondia heydorni* or is it a strain of *Toxoplasma gondii*? An opinion. *Parasitology Research* **86**, 169–178.
- Mohammed, O. B., Davies, A., Hussein, H. S., Daszak, P. and Ellis, J. T.** (2003). *Hammondia heydorni* from the Arabian Mountain Gazelle and Red Fox in Saudi Arabia. *Journal of Parasitology* **89**, 535–539.
- Müller, N., Sager, H., Hemphill, A., Mehlhorn, H., Heydorn, A. O. and Gottstein, B.** (2001). Comparative molecular investigation of Nc5-PCR amplicons from *Neospora caninum* NC-1 and *Hammondia heydorni*-Berlin-1996. *Parasitology Research* **87**, 883–885.
- Payne, S. and Ellis, J. T.** (1996). Detection of *Neospora caninum* DNA by the polymerase chain reaction. *International Journal for Parasitology* **26**, 347–351.
- Schares, G., Heydorn, A. O., Cüppers, A., Mehlhorn, H., Geue, L., Peters, M. and Conraths, F. J.** (2002). In contrast to dogs, red foxes (*Vulpes vulpes*) did not shed *Neospora caninum* upon feeding intermediate host tissues. *Parasitology Research* **88**, 44–52.
- Schares, G., Meyer, J., Bärwald, A., Conraths, F. J., Riebe, R., Bohne, W., Rohn, K. and Peters, M.** (2003). A *Hammondia*-like parasite from the European fox (*Vulpes vulpes*) forms biologically viable tissue cysts in cell culture. *International Journal for Parasitology* **33**, 229–234.
- Schares, G., Pantcher, N., Barutzki, D., Heydorn, A. O., Bauer, C. and Conraths, F. J.** (2005). *Neospora caninum*, *Hammondia heydorni*, *Toxoplasma gondii* and *Hammondia hammondi* oocysts in the faeces of naturally infected dogs in Germany. *International Journal for Parasitology* (in the Press).
- Siverajah, S., Ryce, C., Morrison, D. A. and Ellis, J. T.** (2002). Characterization of an alpha tubulin gene sequence from *Neospora caninum* and *Hammondia heydorni*, and their comparison to homologous genes from Apicomplexa. *Parasitology* **126**, 561–569.
- Slapeta, J. R., Koudela, B., Votypka, J., Modrý, D., Horejs, R. and Lukes, J.** (2002a). Coprodiagnosis of *Hammondia heydorni* in dogs by PCR based amplification of ITS 1 rRNA: differentiation from morphologically indistinguishable oocysts of *Neospora caninum*. *Veterinary Journal* **163**, 147–154.
- Slapeta, J. R., Modrý, D., Kyselova, I., Horejs, R., Lukes, J. and Koudela, B.** (2002b). Dog shedding oocysts of *Neospora caninum*: PCR diagnosis and molecular phylogenetic approach. *Veterinary Parasitology* **109**, 157–167.
- Sreekumar, C., Hill, D. E., Miska, K. B., Rosenthal, B. M., Vianna, M. C. B., Venturini, L., Basso, W., Gennari, S. M., Lindsay, D. S. and Dubey, J. P.** (2004). *Hammondia heydorni*: evidence of genetic diversity among isolates from dogs. *Experimental Parasitology* **107**, 65–71.
- Su, C., Evans, D., Cole, R. H., Kissinger, J. C., Ajioka, J. W. and Sibley, L. D.** (2003). Recent expansion of *Toxoplasma* through enhanced oral transmission. *Science* **299**, 414–416.
- Webb, J. A., Keller, S. L., Southorn, E. P., Armstrong, J., Allen, D. G., Peregrine, A. S. and Dubey, J. P.** (2005). Cutaneous manifestations of disseminated toxoplasmosis in an immunosuppressed dog. *Journal of the American Animal Hospital Association* **41**, 198–202.