

Warwickshire Group (Pennsylvanian) red-beds of the Canonbie Coalfield, England–Scotland border, and their regional palaeogeographical implications

NEIL S. JONES*†‡, DOUGLAS W. HOLLIDAY*§ & JOHN A. McKERVEY*¶

*British Geological Survey, Keyworth, Nottingham NG12 5GG, UK

†Saudi Aramco, PO Box 2001, Dhahran 31311, Saudi Arabia

‡10 Tarn Court, Ilkley, West Yorkshire LS29 8UE, UK

¶16 Harvard House, Rivermead, Wilford Lane, Nottingham NG2 7RB, UK

(Received 23 February 2009; revised version received 19 February 2010; accepted 2 March 2010; first published online 9 June 2010)

Abstract – Late Carboniferous red-beds, < 700 m thick, at outcrop and in the subsurface of the Canonbie Coalfield can be assigned to the Warwickshire Group. They are preserved within the axial part of the Solway Syncline and are divisible into the Eskbank Wood, Canonbie Bridge Sandstone and Becklees Sandstone formations. Sedimentation largely took place on a well-drained alluvial plain, characterized mainly by early, primary oxidation of the strata. Large, northerly-flowing braided river systems were common, with overbank and floodplain fines deposited lateral to the channels; soils formed during intervals of low sediment aggradation. The Canonbie succession includes some of the youngest Carboniferous rocks preserved in the UK. Correlation of the Eskbank Wood Formation is equivocal, but using petrographical, heavy mineral, zircon age dating and palaeocurrent data, the Canonbie Bridge Sandstone Formation can be unambiguously correlated with the Halesowen Formation of Warwickshire, the Pennant Sandstone Formation of South Wales and the offshore Boulton Formation. This suggests that southerly-derived detritus travelled considerable distances from the Variscan highlands of Brittany and/or central Germany across the southern North Sea and UK areas, to a position some hundreds of kilometres north of that previously recognized. The Becklees Sandstone Formation has much in common with the Salop Formation of the English Midlands. It appears to have no preserved equivalent elsewhere in the UK or in the UK sector of the southern North Sea but resembles stratigraphically higher parts of the southern North Sea succession seen in the Dutch sector.

Keywords: Canonbie, Carboniferous, red-beds, correlation, sedimentology.

1. Introduction

The sub-Permian unconformity is a pronounced feature of the geological map of northern England and southern Scotland. It records several tens of millions of years of regional uplift and erosion during and following the Variscan Orogeny (Chadwick *et al.* 1995; Stone *et al.* 2010), and resulted in strata ranging from probable mid-Permian to early Triassic in age resting discordantly on rocks of Early Ordovician to Asturian (Moscowian) age. As a consequence, the youngest, largely red-bed Carboniferous rocks (Bolsovian–Asturian) are only locally preserved in the region and are not well understood.

The largest area in northern England and southern Scotland where Upper Carboniferous (Bolsovian–Asturian) strata are preserved is to the south of the Canonbie Coalfield, along the axis of the NNE-trending Solway Syncline (Figs 1–3). They are mainly concealed by the younger Permo-Triassic rocks of the Carlisle Basin but, in a relatively small area straddling the England–Scotland border near Canonbie, they are locally exposed, notably in the River Esk (Fig. 2),

providing a rare opportunity in this region to examine these rocks in the field.

Previous research in the area has concentrated on the older coal-bearing successions because of their potential economic importance (Peach & Horne, 1903; Barrett & Richey, 1945; Lumsden *et al.* 1967; Ramsbottom *et al.* 1978; Picken, 1988). Apart from establishing a broad, generalized lithological sequence, and the presentation of some faunal and floral data, little has been published on the strata above the *cambriense* Marine Band (Fig. 5).

Stratigraphical subdivision of the younger parts of the Carboniferous is often hampered by the common occurrence of reddening of the strata. Besly & Turner (1983) and Besly, Burley & Turner (1993) have described three phases of reddening of Carboniferous strata: (1) an early syndepositional oxidation of sediments at the time of deposition (termed primary red-beds in this paper), (2) oxidation of initially grey siderite- and coal-bearing sediments by penetrative weathering beneath the Carboniferous land surface soon after deposition, and (3) penetrative or diagenetic reddening of Carboniferous rocks below the sub-Permian unconformity, linked to the ingress of oxidizing groundwaters as a result of periods of subaerial exposure. This penetrative oxidation has been

†Author for correspondence: neil.jones@aramco.com




Figure 1. General location map for the Canonbie Coalfield and the other UK coalfields mentioned in the text (modified from British Geological Survey, 1999). MSNFF – Maryport–Stublick–Ninety Fathom Fault system.

described in detail by Bailey (1926), Trotter (1953, 1954), Mykura (1960), Turner (1980) and Besly (1998). The latter two are grouped here as secondary red-beds.

In this account we present the results of a largely field-based stratigraphical and sedimentological study of the red-beds that occur above the Cambriense Marine Band in the Canonbie area. The paper proposes a new lithostratigraphical nomenclature for these red-beds and comparisons are made with other areas of the UK, both onshore and offshore. At the time of field investigation (March 2004 and July 2005), the strata were particularly well exposed along the River Esk at Canonbie as river levels were low and relatively little river sediment was present (Fig. 2). This has allowed us to present a more detailed description than given previously by Barrett & Richey (1945) and Lumsden *et al.* (1967) of the main River Esk section. Observations from the outcrop sections are supplemented by information from 17 boreholes, drilled into the coal-bearing Langsettian–Duckmantian (Bashkirian) strata (Table 1), and by seismic reflection surveys (Fig. 4). Many of the boreholes were drilled by open-hole rock-bit methods through these mainly red rocks as they were regarded as being of little economic interest. Where cores were taken, they have since been discarded, and descriptions are of variable quality. Fortunately, some of the more recent boreholes, mainly

drilled in the 1980s (Picken, 1988), have wireline geophysical well-log suites (abbreviated hereafter to well-logs) through this part of the succession.

While it was not the primary purpose of this work to carry out a detailed petrographical study, a few samples were analysed by standard optical methods in order to provide further information on the nature of the detrital components within the sandstones (Table 2). These samples were also analysed for their heavy mineral assemblages in order to compare them with the extensive published dataset (Hallsworth, 1992; Hallsworth & Chisholm, 2000; Hallsworth *et al.* 2000; Morton, Clau e-Long & Hallsworth, 2001; Morton, Hallsworth & Moscariello, 2005). Heavy mineral separations from the 63–125 µm fraction were analysed petrographically using the technique of Morton & Hallsworth (1994, 1999), and using the ribbon counting method of Galehouse (1971) for 200 non-opaque heavy mineral grains (Table 3). A portion of the heavy mineral residue for two samples (22 garnets from Canonbie 1 and NJN101; Table 2) were analysed by wavelength-dispersive electron microprobe (Cameca SX-50) operating at 15 kV and 20 nA. In addition, zircon grains were separated from a Canonbie Bridge Sandstone Formation heavy mineral sample and U–Pb isotopic age dates were made using the Sensitive High Resolution Ion Micro Probe Reversed Geometry





Figure 2. (a) Geological map of the Canonbie Coalfield and adjacent areas. The positions of the main boreholes used in the study are also marked. (b) Close-up of the outcrop localities studied. Geological linework from BGS 50 000 scale Digmap data with additions from Picken (1988). The map falls within the National Grid 100 km square NY.

Table 1. General metadata for boreholes used in this study

Borehole name	Quarter sheet	Number	Easting	Northing	Length (m)	Total depth (m)	Date of drilling	Ground level (m)	Rotary table or kelly bushing depth (m)	Datum
BECKHALL	NY37NW	2	333924	575733	420	421	1980	89.34	92.94	Ground level
BECKLEES	NY37SE	3	335166	571578	1370	1370.6	1982	35.25	Unknown	Ground level
BOGRA	NY37NW	3	332864	575529	447	448.3	1983	82.89	Unknown	Ground level
BROADMEADOWS	NY37NE	15	337646	576265	791	791.57	1979	80	83	Kelly bushing
CRANBERRY	NY36NW	3	330724	569485	300	300.1	1982	40.8	Unknown	?Ground level
CROOKHOLM FARM	NY47NW	26	342452	576555	632	632	1956	54.86	Unknown	?Ground level
EVERTOWN	NY37NE	14	336390	575938	777	780.3	1979	92.8	95.8	Kelly bushing
FORGE	NY37NE	7	339456	576720	458	458.47	1893	~30	Unknown	Unknown
GLENZIERFOOT	NY37SE	2	336514	574275	869	869.39	1980	64.93	68.53	Kelly bushing
KNOTTYHOLM	NY37NE	6	339501	577124	649	649.83	1955	42.67	Unknown	Unknown
RIDDINGS	NY47SW	2	340300	574530	143.26	143.26	1914	28.96	Unknown	Ground level
RIDDINGS NO. 1	NY47SW	16	340310	574480	126.79	126.8	1914	~27.43	Unknown	Ground level
ROWANBURN	NY47NW	21	340896	577014	481	481.46	1890	~57	Unknown	Unknown
ROWANBURNFOOT	NY47NW	27	341031	575743	876	876.91	1955	31.7	Unknown	Unknown
STAFFLER	NY37SW	1	332973	572267	708	711.43	1980	52.2	55.2	Ground level
TIMPANHECK	NY37SW	2	332207	574677	558	563.27	1983	83.99	Unknown	?Ground level
WOODHOUSELEES	NY37SE	1	339119	574951	1045	1045.92	1956	57.91	Unknown	?Ground level

Borehole locations are marked on Figure 2.




Figure 3. Isopach map showing the preserved thickness of Pennine Coal Measures and Warwickshire group strata along the Solway Syncline between the Canonbie and West Cumbrian coalfields. Redrawn with modifications from Map 14 of Chadwick *et al.* (1995, p. 88). Contours spaced every 200 metres. GF – Gilnockie Fault; MFS – Maryport Fault system; PCMG – Pennine Coal Measures Group; WG – Warwickshire Group.

(SHRIMP RG); the results of this work are reported elsewhere (Morton, Fanning & Jones, 2010).

2. Tectonic and regional setting

The Pennsylvanian rocks of the Canonbie Coalfield are terminated to the north by the Gilnockie Fault (Barrett & Richey, 1945; Lumsden *et al.* 1967) (Figs 2–4), one of a set of en échelon ENE-trending structures that form the northern margin of the Northumberland–Solway Basin (Chadwick *et al.* 1995). Up to 8000 m

of Carboniferous strata were deposited in this basin which was initiated by rifting in Tournaisian times. The Gilnockie and related southerly-directed faults are interpreted as being antithetic syndepositional structures on the northern margin of the basin. Phases of rifting continued until the end of Visean time, but thereafter, dip-slip faulting was much subdued and basin thermal subsidence dominated. The youngest rocks now preserved are Asturian in age and form the subject of this paper.

Towards the end of Carboniferous time, sedimentation ceased and the rocks of the basin were deformed during the Variscan Orogeny when the rocks were faulted, locally thrust, and gently folded. In several parts of northern England and in the southern North Sea, it has been suggested that deposition of the latest part of the basin-fill was contemporary with the early phases of the Variscan Orogeny and that many of the folds are in part syndepositional (e.g. Leeder & Hardman, 1990; Chadwick *et al.* 1995, p. 36; Corfield *et al.* 1996). Within the study area, Chadwick *et al.* (1995, fig. 27) and Jones & Holliday (2007) have inferred syndepositional thickening of strata into the Solway Syncline beginning in late Visean time. The folding probably occurred as a response to dextral displacement on the Gilnockie Fault and other basin-bounding faults (de Paola *et al.* 2005).

3. Pennine Upper Coal Measures Formation

Following the lithostratigraphical subdivision of Carboniferous rocks proposed by Waters *et al.* (2007) for onshore Great Britain, the oldest rocks described in this account are referred to the Pennine Upper Coal Measures Formation and the overlying red-beds are placed in the Warwickshire Group (Fig. 5). The Pennine Upper Coal Measures Formation comprises mainly grey strata, although secondary red-beds become more common upwards. The formation is about 150 m in




Figure 4. (a) Uninterpreted and (b) interpreted NW to SE seismic profile (ED-86-02) across part of the Canonbie Coalfield. The position of this line is shown on Figure 2. BSF – Becklees Sandstone Formation; Carb – Carboniferous; Hercyn u/c – Hercynian unconformity; PCMG – Pennine Coal Measures Group; PUCM – Pennine Upper Coal Measures Formation; SF – Stainmore Formation.

thickness in the Becklees and Glenzierfoot boreholes (Fig. 7; Table 1). On the NW flank of the Solway Syncline the full thickness of the Pennine Upper Coal Measures Formation is generally not preserved due to erosion beneath the sub-Permian unconformity.

The base of the formation is taken at the Cambriense Marine Band (Figs 5–7), and Picken (1988) suggested that there was a major unconformity at this level, with about 155 m of strata (including the upper part of the Pennine Middle Coal Measures Formation) eroded on the flanks of the coalfield. However, no indication of such a major unconformity has been identified on the seismic reflection sections, and the excellent well-log correlation between boreholes across the coalfield

suggests that in fact no strata are missing. There is some indication that the strata are generally thicker along the axis of the Solway Syncline than on its flanks, and also the gradation to primary red-beds may occur at lower stratigraphical levels on the flanks than in the syncline. The basin flanks do appear to contain much more sandstone than along the axis of the syncline, but the log correlations suggest that these are localized channel sandstones present at several different stratigraphical levels and that they do not constitute evidence for the more widespread unconformity suggested by Picken (1988).

Borehole well-log data indicate that the formation largely comprises mudstones, some coals, *Estheria*-rich mudstone beds and *Spirorbis* limestones

Table 2. General metadata for the Warwickshire Group thin-section and heavy mineral samples used in this study

Specimen number	Stratigraphy (formation)	Easting	Northing	Location information	Comments
NJN-98	BSF	339360	576310	Approx. 300 m downstream of Canonbie Bridge, on SE bank of River Esk	Brick red sandstone; very friable; very fine- to medium-grained, poorly sorted
NJN-99	BSF	339813	575760	Approx. 1.3 km downstream of Canonbie Bridge, on east bank of River Esk; locality known as Mason's Stream	Moderately sorted, brick red sandstone; fine-grained
NJN-100	CBSF	339789	576088	Approx. 600 m ESE from Dead Neuk; close to where track from Park House joins River Esk; outcrop set back from river	Medium-grained lithic sandstone
NJN-101	CBSF	339838	576005	Approx. 600 m ESE from Dead Neuk; close to where track from Park House joins River Esk	Medium- to coarse-grained lithic sandstone
NJN-102	CBSF	339551	576501	Approx. 20 m downstream of Canonbie Bridge, on SE bank of River Esk	Medium-grained lithic sandstone
CAN1	CBSF	339540	576500	Downstream of Canonbie Bridge, on SE bank of River Esk	Medium- to coarse-grained lithic sandstone; sample used for heavy mineral analysis only

BSF – Becklees Sandstone Formation; CBSF – Canonbie Bridge Sandstone Formation.

Table 3. Heavy mineral data for the Canonbie Bridge Sandstone Formation (CBSF) and Becklees Sandstone Formation (BSF): (a) point-counted data, and (b) main ratios of heavy minerals with similar grain sizes and densities that determine the relative abundances of minerals with similar hydraulic and diagenetic behaviour

(a)			Zir	Rut	Tor	Apa	Gnt	Mnz	Cr-S	Ana	Amp	Sta	Count
Sample	Location	Formation	Frequency %										
NJN98	River Esk	BSF	77	17	6	1	0	0	0	0	0	0	200
NJN99	River Esk	BSF	32	9	30	28	1	0	0.5	0.5	0	R	200
NJN100	River Esk	CBSF	60	9	7	22	1	0	2	0	0	0	200
NJN101	River Esk	CBSF	43	5	12	37	3	0.5	1	0	0	R	200
NJN102	River Esk	CBSF	67	16	6	10	1	1	0	0	0	0	168
CAN1	Canonbie	CBSF	44	8	10	36	1	1	0	0	0.5	0	200

(b)			RuZi	ATi	MZi	CZi	GZi	RuZi	ATi	GZi	MZi	CZi
Sample	Location	Formation	Indices					Counts				
NJN98	River Esk	BSF	17.0	<i>41.5</i>	0.0	0.0	0.0	482	41	400	400	400
NJN99	River Esk	BSF	34.5	44.7	0.3	0.7	5.3	438	506	303	288	289
NJN100	River Esk	CBSF	17.5	<i>77.8</i>	0.9	1.3	1.8	269	117	226	224	225
NJN101	River Esk	CBSF	11.0	<i>72.7</i>	1.1	0.0	7.0	300	227	287	270	267
NJN102	River Esk	CBSF	<i>19.4</i>	<i>61.5</i>	<i>1.8</i>	<i>0.0</i>	<i>0.9</i>	139	26	113	114	112
CAN1	Canonbie	CBSF	18.4	70.6	4.1	0.0	3.0	316	218	266	269	258

Italicized numbers indicate heavy mineral index data of relatively poor statistical quality.

Mineral abbreviations: Zir – zircon; Rut – rutile; Tor – tourmaline; Apa – apatite; Gnt – garnet; Mnz – monazite; Cr-S – Cr-spinel; Ana – anatase; Amp – amphibole; Sta – staurolite; R – rare. ATi – apatite–tourmaline; CZi – Chrome-spinel–zircon; GZi – garnet–zircon; MZi – monazite–zircon; RuZi – rutile–zircon.

(Figs 6, 7). Sandstones are generally thin (2–5 m thick), although in the Glenzierfoot and Broadmeadows boreholes a sandstone is present at the base of the formation, forming a single unit up to 20 m in thickness with a distinct blocky gamma-ray character (Fig. 7).

The upper part of the formation can be examined at outcrop along the western bank of the River Esk [between 3905 7715 and 3900 7715] (Fig. 2). Here the succession dominantly comprises pinkish to reddish brown silty claystone, with some sandy siltstone and rare pinkish grey fine- to medium-grained micaceous sandstone. Rooted palaeosols are common, although in places the roots are present only as traces due to the effects of oxidation. Ironstone nodules and thin beds are also commonly present. Oxidized coals also occur, with one example represented by an organic-rich, ferruginous limestone at [NY 3904 7712] (Fig. 8). Mudstones contain a fauna including *Anthraconaia pruvosti* and *Leaia bristolensi*, suggesting this upper part belongs to the *tenuis* Chronozone and thus spans

the interval between late Bolsovian and Asturian (Moscowian) (Lumsden *et al.* 1967; Ramsbottom *et al.* 1978) (Fig. 5).

4. Warwickshire Group

The term Warwickshire Group has been applied to the predominantly red-bed strata that typically form the upper part of the Carboniferous succession in the UK previously referred to by such names as the Barren or Red Measures (see Powell *et al.* 2000). The type area is the Warwickshire Coalfield in central England (Fig. 1), where the succession attains its fullest thickness of 1225 m (Powell *et al.* 2000). In the southern Pennine coalfields the Warwickshire Group is largely unconformable onto older Pennine Coal Measures Group strata, with the ‘Symon Unconformity’ present in places between the two (Powell *et al.* 2000). The group ranges in age from late Bolsovian to Autunian (Mid- and Late Pennsylvanian), although, due to the

CHRONOSTRATIGRAPHY		LITHOSTRATIGRAPHY		NON-MARINE BIVALVE ZONES	MAIN MARINE BANDS	KEY MARKER BEDS IN CANONBIE AREA	
REGIONAL STAGE	REGIONAL SUBSTAGE	GROUP	FORMATION				
?Stephanian	?Cantabrian	Warwickshire	Becklees Sandstone	<i>prolifera</i>			
Westphalian	Asturian		Canonbie Bridge Sandstone				
			Eskbank Wood	<i>tenuis</i>		High Coal	
	Bolsovian	Pennine Upper Coal Measures	<i>phillipsii</i>	Cambriense Aegiranum	=Riddings =Skelton		
	Duckmantian	Pennine Middle Coal Measures	U. <i>similis-pulchra</i> Lower <i>similis-pulchra</i>	Vanderbeeki	Archerbeck Coal Nine Foot Coal =Queenslie/Solway		
Langsettian	Pennine Lower Coal Measures		<i>modiolaris</i> <i>communis</i> <i>lenisulcata</i>				
		Namurian	Pendleian to Yeadonian	Yoredale	Stainmore		Subcrenatum
			Alston		<i>C. leion</i>	Catsbit (=Great) Lst	

Figure 5. Lithostratigraphical classification of Pennsylvanian strata in the Canonbie area (based on Trueman & Weir, 1946; Ramsbottom *et al.* 1978 and Waters *et al.* 2007). The base of the Westphalian at Canonbie is marked by the presence of a non-sequence locally in parts of the basin centre and an angular unconformity on basin margins. C. – *Cravenoceras*; Lst – Limestone; U. – Upper.




Figure 6. Summary of the main features of the Warwickshire Group from the Canonbie Coalfield (Becklees Borehole). The side panel shows distribution plots of gamma-ray values for the Canonbie Bridge Sandstone and Becklees Sandstone formations and clearly shows that the Canonbie Bridge Sandstone is dominated by higher gamma-ray values, linked to the presence of abundant rock fragments. API – American Petroleum Institute; BHCS – Borehole Compensated Sonic log; GR – Gamma-Ray log; MB – Marine Band; PMCM – Pennine Middle Coal Measures Formation; PUCM – Pennine Upper Coal Measures Formation.




Figure 7. Correlation of the Warwickshire Group and upper part of the Pennine Coal Measures Group from boreholes in the Canonbie area. See Figure 2 and Table 1 for borehole locations. BHCS – Borehole Compensated Sonic log; cong – conglomerate; CPDL – Compensated Photo-density log; Fm. – Formation; Gp. – Group; GR – Gamma-Ray; mst – mudstone; SDU – BPB standard density unit; sst – sandstone.



Figure 8. Oxidized gley palaeosol overlain by remnants of oxidized coal. Pennine Upper Coal Measures Formation [3902 7714]. LM – lacustrine mudstone; OC – oxidized coal; OG – oxidized gley.

effects of reddening, the degree of biostratigraphical control is reduced compared with the Pennine Coal Measures Group (Besly & Cleal, 1997; Powell *et al.* 2000).

The basis for the recognition of the group is that primary red-beds must be present. The group typically comprises red mudstones and sandstones, with varying amounts of pebbly sandstone, conglomerate, grey mudstone, thin coals, lacustrine limestone (*Spirorbis* limestone) and calcrete palaeosols. The lithologies of the Warwickshire Group formed predominantly within oxygenated surface conditions within a drier climate.

The Warwickshire Group has not previously been recognized from the Canonbie area, although earlier work had recorded the presence of red strata in the ‘Upper Coal Measures’ of the area (Peach & Horne, 1903; Simpson & Richey, 1936; Barrett & Richey, 1945; Trotter, 1953; Lumsden *et al.* 1967; Day, 1970; Picken, 1988). Picken (1988) described the succession as largely comprising interbedded sandstones and mudstones, generally calcareous. Marker beds include *Spirorbis* limestones and a persistent coal that was named the High Coal (Fig. 6).

The detailed stratigraphy in the Canonbie area has been determined by a study of the outcrops along the River Esk between [3915 7706] and [3981 7576] which form the type section (Fig. 9), together with the relevant borehole information. As a result of this work, three subdivisions of the group have been identified, described below from oldest to youngest. These have been given formation status and were named and defined by Jones & Holliday (2007). Due to the extensive Devonian and Holocene cover in the area it is difficult to map these formations in the field, hence they should be regarded as informal names. However, they are readily identifiable at outcrop along the River Esk and from detailed descriptions of borehole core.

4.a. Eskbank Wood Formation

This formation is 145–175 m thick, although the full thickness of the formation is not always preserved on

the northwestern flank of the coalfield due to erosion at the sub-Permian unconformity surface. The formation dominantly comprises mudstone (60–70%) with the remainder being largely sandstone. Red mudstones are interbedded with fine- to medium-grained sandstones, calcrete palaeosols, thin beds of *Spirorbis* limestone and *Estheria*-bearing mudstones. Sparse thin coals and grey mudstones are present in the lower part of the formation; some of these coals have been oxidized and altered to limestone. One thick coal (the High Coal), locally up to about 1.2 m thick, is known only from boreholes (Figs 6, 7). The outcrop section [NY 392 770] proves up to 75 m of the formation, including mudstones that contain at several localities a non-marine fauna, particularly bivalves, identified as *Anthraconauta phillipsii*, *A. aff. phillipsii*, *A. cf. tenuis*, *A. cf. wrightii*, possibly *Anthracomya pruvosti* and also ostracods (A. E. Trueman in Barrett & Richey, 1945, p. 39; Lumsden *et al.* 1967, p. 178), and indicative of an Asturian age (Fig. 5).

The conformable, slightly diachronous base of the formation is taken at the first major primary red-bed strata overlying the grey Pennine Upper Coal Measures Formation (Fig. 6) and is difficult to locate in uncored boreholes. Its exact position differs between boreholes but it generally occurs close to the High Coal. The gradational nature of the boundary indicates there was local variation and alternation between oxidizing and reducing conditions. As the High Coal can be identified in open-hole from suitable well-logs (e.g. sonic and density logs), it is suggested that, in the absence of core data, the top of the coal is taken to mark the boundary between the Pennine Upper Coal Measures and the Eskbank Wood formations (Figs 6, 7).

4.b. Canonbie Bridge Sandstone Formation

This formation is more or less continuously exposed along the River Esk in the immediate area north and south of Canonbie Bridge (Fig. 2) and its base is taken at the lowermost sandstone bed in a thick (20–30 m) multi-storey sandstone complex at [3919 7695] (Fig. 9). The formation ranges in thickness from 131 m in Broadmeadows Borehole up to a maximum of 168 m in the Becklees Borehole (Figs 6, 7). On the northwestern flank of the Canonbie Coalfield the full thickness of the formation is not always preserved due to erosion at the sub-Permian unconformity surface.

The formation comprises interbedded reddish brown to greenish grey, moderately to poorly sorted, fine- to coarse-grained lithic arenite sandstones (50–70%) (Fig. 10), reddish brown mudstones and reddish brown mudstone palaeosols including calcretes. Sandstones are generally cross-bedded, forming individual sharp to erosively based channellized units that range in thickness from 2 to 10 m. In some places these amalgamate into multi-storey sandbodies up to 40–50 m thick (Fig. 11). Intraformational mudstone conglomerates occur scattered throughout the sandstones but are more common lining channel bases. In hand specimen the




Figure 9. Composite sedimentological log of the Warwickshire Group succession exposed along the River Esk, Canonbie between [3917 7705] and [3930 6726].




Figure 10. Ternary diagram to show the classification of the Warwickshire Group sandstones. Classification scheme after Pettijohn, Potter & Siever (1987).

sandstones can be micaceous and contain a noticeable component of greenish grey grains that, in thin-section, can be identified as lithic clasts, mainly comprising fine foliated or schistose low-grade pelitic and psammitic metasedimentary clasts (Table 4; Fig. 12). The high lithic component gives the sandstones higher gamma-ray log values compared to the overlying Becklees Sandstone (Fig. 6).

4.c. Becklees Sandstone Formation

The Becklees Sandstone Formation is the uppermost unit recognized and its top is truncated at the sub-Permian unconformity. Thus its full thickness is not known, but boreholes show that at least 200 m of probable late Carboniferous strata conformably overlie those of the river section; seismic reflection surveys indicate that this thickness could be up to 700 m along the axis of the Solway Syncline (Chadwick *et al.* 1995) (Fig. 4). Although these higher strata cannot be directly examined, their well-log and seismic reflection

character suggest that they are similar to the youngest Carboniferous red-beds seen at outcrop.

The formation comprises interbedded subfeldspathic arenitic sandstones (70–90 %) (Fig. 10), reddish brown mudstones and reddish brown mudstone palaeosols including calcretes. Limestones and thin coals are also recorded in cuttings from the Becklees Borehole, although these are uncommon and were not seen at outcrop.

Sandstones are fine- to medium-grained, orange brown to bright reddish brown and pinkish brown, moderately to well sorted and typically lack mica. They are generally cross-bedded and occur in thick, sharp-based successions (10–30 m), some showing fining upwards. The sandstones contain a noticeably lower proportion of lithic clasts and more rounded grains than the underlying Canonbie Bridge Sandstone Formation (Table 4; Fig. 13). At outcrop this lithological change is abrupt; in well-logs it can be seen as a lower response on the gamma-ray log (Fig. 6). Sand-filled polygonal cracks penetrating the Canonbie Bridge Sandstone Formation have been recorded at the base of the formation [3936 7631]. These may be joints which, if this is the case, would point to a significant time gap between the deposition of the two formations, marked by lithification of the Canonbie Bridge Sandstone.

5. Warwickshire Group sedimentology and depositional environments

Four facies associations are recognized in the Warwickshire Group succession at Canonbie (Table 5). Detailed description of the sedimentology of the succession can be found in the online Appendix at <http://journals.cambridge.org/geo>.

Initially sedimentation in the lower part of the Warwickshire Group (Eskbank Wood Formation) occurred on a poorly drained alluvial plain in which reducing conditions dominated, marked by the presence of peat mires and floodplain lakes. Grey lacustrine mudstones (Fig. 14) were also deposited, and these can contain




Figure 11. (a) Example of cross-bedding within the channel facies; (b) Line drawing of (a). Cross-bedding foresets dip to the right and set bounding surfaces, marked by thicker lines, dip to the left. There is a structural dip of about 18° down to the south at this locality. Canonbie Bridge Sandstone Formation, west bank of the River Esk [3918 7690]. View looking SW. Height of visible part of person is 1.7 m.

Table 4. Petrographic and heavy mineral characteristics of the two main sandstone types from the Warwickshire Group of the Canonbie area

Stratigraphy (formation)	Classification, grain size and textural characteristics	Detrital components	Authigenic components	Heavy mineral components
BSF	Subfeldspathic-arenite. Fine- to medium-grained; poorly to moderately sorted; bimodal texture present in places; grains typically subangular to subrounded, some larger grains are rounded or well rounded.	Dominantly quartz (78–90%) – generally monocrystalline, less common polycrystalline quartz and chert. Inclusion ‘bubbles’ seen in some quartz grains. Potassium feldspar forms a significant constituent (8–16%). Rock fragments less common (2–5%) but comprise low grade metasedimentary clasts, intraformational mudstone clasts and some igneous rock fragments. Minor sodium feldspar and mica grains.	Main component is blocky, pore filling dolomite. Quartz overgrowths and authigenic illite also present. Clay/iron oxide ‘dust’ rims common & iron oxide occurs as pore filling cement locally. Generally poorly cemented with a porous, open framework & common oversized pores, linked to extensive grain and cement dissolution.	RuZi variable (17–34) and ATi index 42–45. All other indices (GZi, MZi and CZi) are low (0–7) with the CBSF characterised by slightly higher MZi, but similar CZi and GZi, in comparison to the BSF.
CBSF	Lithic-arenite. Medium- to medium- to coarse-grained, poorly to moderately sorted with subangular to subrounded grains.	Dominantly quartz (45–52%) – generally monocrystalline, although polycrystalline quartz is common. Rock fragments a major constituent (36–42%) – mainly foliated or schistose low grade metasedimentary clasts with aligned micas and stretched quartz. Intraformational mudstone clasts and some igneous rock fragments also occur. K-feldspar grains common and, in lesser quantities, elongate mica grains and Na-feldspar. Tourmaline present as a trace.	Main component is quartz, forming overgrowths on detrital quartz and feldspars. Authigenic illite present as grain replacive and pore filling cement, some pore-filling kaolinite ‘books’. Clay/iron oxide ‘dust’ rims common and iron oxide forms pore filling cement in places. Compactional textures are common, particularly indented micas and rock fragments. Porosity usually quite high, linked to grain and cement dissolution.	RuZi index characterised by low values (11–19); higher ATi (62–78) than the BSF. All other indices (GZi, MZi and CZi) are low (0–7). Garnets from 2 samples have high almandine +spessartine compositions (~0.8–1.0; spessartine averages ~0.3) with low grossular (0–0.2) and minor pyrope (<0.1). One garnet has considerably higher pyrope (~0.35).

BSF – Becklees Sandstone Formation; CBSF – Canonbie Bridge Sandstone Formation. ATi – apatite–tourmaline; CZi – Chrome-spinel–zircon; GZi – garnet–zircon; MZi – monazite–zircon; RuZi – rutile–zircon.




Figure 12. Thin-section photomicrographs of the Canonbie Bridge Sandstone Formation (Sample NJN101) in plane polarized light to show the common occurrence of lithic rock fragments (RF). Also present are quartz grains (Q). Scale bar is 0.5 mm across.




Figure 13. Thin-section photomicrograph of the Becklees Sandstone Formation (Sample NJN99) in plane polarized light. Quartz (Q) forms the dominant detrital component, with minor feldspar (F) and rock fragments (RF). Rhombs of dolomite cement (D) can also be seen in places. Scale bar is 0.5 mm across.

a varied fauna including *Estheria* and non-marine bivalves (e.g. *Anthraconauta*, *Modiolaris* and *Naiadites*). The deposition of thin fresh-water limestones (Fig. 15) suggests the periodic development of shallow, well-oxygenated fresh-water lakes, possibly characterized by high rates of surface-water evaporation. Into these lakes small lacustrine deltas formed, fed from river systems. The presence of thick channel sandstones with blocky gamma-ray profiles and thin-bedded crevasse splays indicates that, at times, the environment became more akin to an alluvial plain rather than a wetland. Poorly drained soils were typically formed at this time, although periods of better-drained conditions were characterized by the development of semi-gley, pseudogley and brunified alluvial types, with common red and green mottling of the soil profiles (Table 5; Figs 16, 17). Incipient calcimorphic soils, represented as isolated carbonate nodules within brunified alluvial

soils, are also recorded in the lower part of the Warwickshire Group.

The upper part of the Eskbank Wood Formation shows an increasing development of better-drained floodplain conditions marked by shallow lakes and floodplains that often dried out, leading to the formation of desiccation cracks and better-drained soils. Brunified alluvial and calcimorphic palaeosols become dominant, some of which can be quite well developed (Fig. 18). Lacustrine facies are also present, but are less common than lower in the formation. These are generally red but lack features indicative of well-drained conditions (e.g. desiccation cracks), suggesting periods of subaqueous deposition, possibly in reducing conditions. These were later oxidized and reddened, probably early in the diagenetic process. Channels are also known, but they are typically minor and do not form a significant percentage of the formation.

The Canonbie Bridge Sandstone Formation shows a marked and apparently abrupt change in sedimentation patterns, with the establishment of an alluvial plain in which large fluvial systems dominated (Fig. 11). Typically these channels carried a coarse sediment load marked by presence of common lithic detritus, recycled from an earlier metasedimentary terrain. Such fluvial systems record the delicate interplay between subsidence, base-level change and sediment supply, with a high sediment load combined with increasing rates of accommodation space generation suggested. Palaeocurrent analysis indicates that these channel systems flowed from south to north, with low sinuosity channel systems suggested by the low spread of the palaeocurrent data (Fig. 19). Locally, overbank and well-drained palaeosol facies are known, with brunified alluvial and calcimorphic palaeosols dominant (Figs 20, 21). This indicates that the floodplain was generally well-drained, although a thin coal is reported from cuttings from high in the Canonbie Bridge Sandstone Formation in the Becklees Borehole, which would record a temporary reversal to poorly drained conditions.

The Becklees Sandstone Formation also comprises a high proportion of sandstone, and deposition in alluvial channel systems is again indicated. Too few palaeocurrent measurements were measured at outcrop to be confident of channel trends, although dipmeter data suggest flow towards both the south and the NW (Fig. 19). Detailed analysis was not possible for this formation, although a provenance change is suggested by the change in palaeocurrent data and the reduction in both grain size and lithic clast content.

During the late Carboniferous it is known that there was a change in climate from humid or sub-humid conditions in Bolsovian times to semi-arid conditions by Asturian times. This was brought about by the rain-shadow effect of the continuously rising Variscan mountain chain to the south during the formation of the Pangaeian supercontinent (Parrish, 1982; Rowley *et al.* 1985; Besly, 1987). This led to the development of a monsoonal circulation pattern with a marked seasonality to the climate (Parrish, 1982; Rowley *et al.*

Table 5. Characteristics of the facies associations and facies recognized from the Warwickshire Group of the Canonbie area

Facies association	Facies	Lithology and colour	Sedimentary structures	Thickness and geometry	Other characteristics	Interpretation
Channel-Belt						
	Channel-fill	Reddish-brown sandstone (typically 80%); very fine to coarse-grained, commonly micaceous; subordinate beds of conglomerate, silty sandstone, siltstone and claystone also occur, generally <0.2 m thick.	Unidirectional sedimentary structures, typically trough and planar tabular cross-bedding, sets generally 0.2–0.3 m in thickness. Also present are current ripple cross-lamination, climbing ripple cross-lamination, plane beds, flat lamination and soft-sediment deformation.	Erosively-based channelized sandbodies 2–10 m thick.; thicker sandbodies with multiple, internal (laterally extensive) erosion surfaces are <50 m thick; bases generally conglomeratic with quartzite and red intraformational mudstone clasts.	Carbonaceous debris and some plant fragments (e.g. <i>Cordaites</i>) are present locally; rock fragments common to abundant in Canonbie Bridge Sandstone Formation; low spread of palaeocurrent data, generally towards the North.	Deposited within braided fluvial channels. Multiple erosion surfaces indicate repeated episodes of channel scouring forming channel belts likely to be many kilometres in width. Unidirectional sedimentary structures indicate downstream migrating current generated sandy bedforms, formed from turbulent flows. The low spread of palaeocurrents suggests a low channel sinuosity. Heavy mineral, garnet geochemistry and zircon age data indicate a Variscan source area that lay far to the south.
Subaqueous lacustrine and alluvial floodplain						
	Lacustrine limestone	Grey, pinkish grey and greenish grey calcareous mudstones to muddy limestones.	Generally massive or weakly laminated.	Typically 0.05–0.5 m thick; laterally continuous sheet-like beds with sharp bases and tops.	Non-marine fauna including <i>Spirorbis</i> , ostracods, fish scales and bivalves, e.g. <i>Anthraconauta</i> .	Non-marine fauna indicates deposition in shallow, well-oxygenated fresh-water lakes. Carbonate precipitation can be a common product of fresh-water lakes in humid environments. This facies has been described from contemporaneous deposits elsewhere in the UK where they are termed <i>Spirorbis</i> or Entomostracan limestones.
	Lacustrine mudstone	Reddish brown, pinkish grey and greenish grey laminated claystones, silty claystones and siltstones, rare sandy lenses.	Laminated, with rare cross-laminated and less common wave rippled sandy lenses; local desiccation cracks.	1–6 m in thickness; laterally continuous; generally gradational tops and sharp or gradational bases.	Bivalves including <i>Anthraconauta</i> and <i>Anthracomya</i> occur, as well as ostracods, rare burrows and plant fragments; small hematite nodules (?after siderite).	Lacustrine deposition indicated by fine-grain size, horizontal lamination and non-marine bivalves. Deposition from suspension, with lamination formed by subtle fluctuations in sediment supply, into perennial bodies of water, generally a few metres deep. Desiccation cracks indicate that the lakes sometimes dried out. Cross-laminated sandy lenses formed from sediment transported by unidirectional tractional currents. In some lakes wave-fetch was large enough to rework bottom sediment. Reducing conditions indicated by the presence of siderite, although later oxidation led to its transformation into hematite.

Table 5. Cont.

Facies association	Facies	Lithology and colour	Sedimentary structures	Thickness and geometry	Other characteristics	Interpretation
	Lacustrine delta	Grey to purple grey laminated micaceous siltstones and sandy siltstones in their lower parts, passing upwards into silty sandstones and very fine- and fine-grained sandstones in their upper parts	Lower, siltier parts generally contain parallel lamination and current and wave ripple cross-laminated sandy lenses; upper part comprises current ripple cross-lamination, with minor wave rippling and low-angle cross-bedding	< 2.8 m thick; laterally continuous in sheet-like beds; can be either sharply based or coarsen upwards	Upwards-coarsening is characteristic; not common at outcrop but, where seen is underlain by lacustrine mudstone facies and overlain by red brunified alluvial and calcimorphic palaeosols	Coarsening-upwards character suggests a small, fluviially-dominated delta that prograded into a lake. Deposition from bedload-transporting tractional currents that flowed down the delta front, generating ripples and small dunes. At times waves reworked the lower parts of the delta. Shallow water is suggested by the limited thickness (< 3 m) deposit, with lake infilling and emergence indicated by the development of a palaeosol on its top.
	Overbank	Pinkish, purplish and reddish brown silty claystones, siltstones and rarely silty sandstones	Usually poorly bedded or massive	Typically a few metres in thickness, up to a maximum of 23 m; unknown geometry, but likely to be laterally extensive or tapering away from nearby channel; gradational tops and bases	Common plant material is diagnostic, including <i>Calamites</i> , <i>Sigillaria</i> and <i>Neuropteri</i> ; common reduction spots	Formed adjacent to the bank of a river channel, following overtopping of the bank during flood stage. Suspended sediment was dumped proximal to the channel as a result of a decrease in turbulence during flow expansion. Rapid sedimentation rates inhibited dynamic sorting resulting in a structureless deposit.
	Crevasse splay	Pinkish to reddish brown beds of sandstone or silty sandstone, generally very fine- to fine-grained, rarely medium-grained or conglomeratic	Dominantly current ripple cross-lamination and climbing ripples, although small scale sets of cross-bedding can occur	Characteristically form individual, upwards fining tabular beds that are generally <0.6 m in thickness; beds are usually sharp or erosively based, often with a thin basal intraformational conglomerate; individual beds can stack vertically to form thicker sandbodies < ~ 2 m in thickness	This facies typically occurs interbedded either with lacustrine or palaeosol facies	Each sandstone bed is interpreted as the deposit of a sand-laden crevasse splay that originated by breaching or crevasse of a channel bank. This allowed floodwater from the channel to flow into an adjacent interchannel area as an unconfined, unidirectional turbulent flow capable of forming ripples and small dunes. Interbedded association with lacustrine facies and palaeosols suggests they were episodically deposited on the floodplain.
Palaeosol	Brunified alluvial palaeosol	Reddish brown to purplish red beds of claystone, silty claystone and siltstone, local greenish grey mottling	Typically structureless to weakly laminated with no obvious horizonation; desiccation cracks and small-scale curved and slickensided 'listric' surfaces are common and a pseudobrecciated fabric sometimes seen on bedding surfaces	Bed thickness 0.5–3 m; generally gradational tops and bases	Roots are present but are not common; they form thin red subvertical traces up to a few centimetres in length typically associated with green reduction halos; rare plant debris and small pedogenic carbonate (calcrete) nodules	Presence of roots indicates a preserved soil, but their scarcity suggests repeated sediment aggradation, probably on a muddy floodplain. Desiccation cracks and the associated pedobrecciated fabric formed during drying out of the sediment. Red colour results from the oxidation of ferrous iron to hematite under free draining conditions. Greenish grey mottling reflects temporary change to poorly-drained conditions, allowing localized reduction to occur. This type of palaeosol is similar to modern brunified alluvial soils.

Vertic brunified palaeosol	Brick red to reddish brown claystone, silty claystone and siltstone, local greenish grey mottling	Destratified, commonly with a rubbly, fragmented texture; listric surfaces and desiccation cracks	Beds 0.5–1 m thick; generally gradational tops and bases	Can be characterized by concave upward planes ('slickensides'), <1.1 m across and 0.15 m in height; calcrete nodules in minor amounts; nodules either scattered randomly or concentrated in the lower part of the bed	This palaeosol resulted from repeated wetting and drying, causing the sediment to break up. This formed the rubbly texture and also the concave upward planes, which are interpreted as pseudo-anticlines (= the subsurface expression of surface mounds known as gilgai). Successive wetting and drying also led to the deposition of CaCO ₃ lower in the soil profile. Locally reduced zones represent 'drab haloes', formed by the removal of iron by dissolution. This is favoured along flow paths such as desiccation cracks, ped boundaries and in the presence of organic matter. Has many features similar to those described from modern soils known as vertisols.
Calcimorphic palaeosol	Reddish brown claystone, silty claystone and siltstone, scattered to coalesced white to cream calcium carbonate nodules; local greenish grey mottling	Structureless, commonly with a rubbly texture; pedogenic listric surfaces and desiccation cracks common	Beds <~0.5 m in thickness; generally gradational tops and bases	Characterized by the ubiquitous presence of displacive nodules (glaebules) ranging from weakly developed, isolated glaebules to coalesced glaebules that form <80 % of the bed.; range from sub-spherical to ellipsoidal in form and from 0.01–0.13 m in size; ellipsoidal nodules tend to be vertically inclined; rare preserved (reddened) roots	The glaebules formed by displacive growth of calcium carbonate in vadose zone of the soil, linked to evapo-transpiration of carbonate-saturated pore waters. Desiccation cracks indicate well-drained conditions were prevalent and this type of palaeosol is generally taken as an indicator of formation in a semi-arid climate. Palaeosols described here range from Stage 1 (more isolated calcrete glaebules forming an aggradational calcimorphic palaeosol) to Stage 3 (coalesced nodules).
Semi-gley palaeosol	Grey to brownish grey claystone, silty claystone and siltstone, locally with reddish brown mottling	Structureless	Beds <1 m in thickness; generally gradational tops and bases	Characteristic features are the colour and the common occurrence of sphaerosiderite; coaly laminae, roots and reduction spots can also be common; typically more common in the lower part of the red-bed succession and in the Pennine Upper Coal Measures Group	Mainly shows features of poor drainage, and formation under reducing conditions. However, sphaerosiderite indicates mobilisation of iron by capillary action during periods of better drainage linked to temporary lowering of the water table to allow some oxygenation of the soil. Reddish brown and brown colours also indicate slightly better drained conditions. Brown colour is related to a hydrated iron oxide, stable in conditions where there is not enough organic matter to cause reduction and not enough time for complete oxidation to hematite.

Table 5. Cont.

Facies association	Facies	Lithology and colour	Sedimentary structures	Thickness and geometry	Other characteristics	Interpretation
	Pseudogley palaeosol	Bright brick red to reddish brown silty claystone to claystone; some grey, green and yellow-white mottling	Totally destratified, and contains abundant listric surfaces and desiccation cracks	Beds <0.5 m in thickness; generally gradational tops and bases	Not common; roots including <i>Stigmara</i> typical, preserved either as black, carbonaceous forms or as oxidized traces; colour mottling (reduction) generally concentrated along the fringes of the roots	Shows similarities to modern pseudogleys. The soils are not saturated with water during most of the year and much of the pore system is filled with air. Hence, dissolved ferrous iron oxidizes to Fe ^(III) and converts to the insoluble Fe ^(III) -(oxy)-hydrates. Red colour is a primary pedogenic feature, formed under oxygenated conditions. Any water that penetrates the soil moves through larger pores, cracks and along root channels, resulting in iron reduction (pseudogleying) being concentrated along these zones, giving rise to the colour mottling.
	Oxidized gley palaeosol	Pinkish grey to reddish grey brown claystone, silty claystone and siltstone; some colour mottling; reddening is non-pervasive	Totally destratified and contains abundant listric surfaces	Beds of <2 m in thickness; generally gradational tops and bases	Other features include partially or completely oxidized siderite concretions; rooting, including <i>Stigmara</i> , is a common feature, although many have clearly undergone some oxidation and are present only as traces; red, yellow and green colour mottling typical, both as haloes along former roots and as reduction spots	This is interpreted as an alluvial hydromorphic soil (gleysol) that originally accumulated under poorly drained, reducing conditions and was later oxidized during early burial. This is indicated by the patchy reddening which indicates incomplete oxidation of the sediment. The presence of <i>Stigmara</i> and relict siderite nodules indicates formation originally under waterlogged conditions. The siderite is later partially replaced by hematite as a result of oxidation during early burial.
	Histosol (coal)	Bituminous coal, inferior coal and coaly shale; can occur with interbedded black mudstone beds	Lacks sedimentary structures, but can be banded	Beds from a few centimetres <1.2 m; sheet like form and can be extensive over many square kilometres	A volumetrically insignificant facies within the Warwickshire Group; best known is the High Coal; not seen at outcrop but described from core and can be interpreted from well-logs	Result from the authochthonous accumulation of organic material in a peat-forming mire. Thick, siliciclastic-free coals probably formed as ombrotrophic bog deposits, with interbedded mudstones representing the development of rheotrophic swamp conditions in which the surface of the mire is flooded, allowing sediment input and the development of a siliciclastic palaeosol.




Figure 14. Example of the lacustrine mudstone facies. Note the presence of parallel lamination marked by preferential splitting of laminae. Lens cap for scale is 5 cm across. Eskbank Wood Formation, River Esk [3915 7705].



Figure 15. Lacustrine limestone facies forming the resistant, lighter coloured bed. Lens cap for scale is 5 cm across. Eskbank Wood Formation, River Esk [3915 7705]. LL – lacustrine limestone; LM – lacustrine mudstone.



Figure 16. Example of a pseudogley palaeosol with common pedogenic listric surfaces and lighter coloured goethitic mottling (arrowed) along former roots. Eskbank Wood Formation, River Esk [3922 7693]. Lens cap for scale is 5 cm across. For a colour version of this figure see online Appendix at <http://journals.cambridge.org/geo>.

1985). The presence of more evolved, red palaeosols and reddening in general in the Warwickshire Group succession of Canonbie is indicative of a gradual change to a more arid climate. The Warwickshire Group




Figure 17. Example of a vertic brunified palaeosol (B horizon). Note the prominent curved 'slickensided' surfaces (arrowed) overlying nodular accumulations of pedogenic carbonate (calcrete) in the underlying C horizon. A brunified alluvial palaeosol (BA) is present above. Eskbank Wood Formation, River Esk [339174 557705]. Lens cap for scale is 5 cm across. For a colour version of this figure see online Appendix at <http://journals.cambridge.org/geo>.



Figure 18. Example of a calcimorphic palaeosol (calcrete), with coalesced carbonate glaebules. Eskbank Wood Formation, River Esk [3915 7705]. Lens cap for scale is 5 cm across. For a colour version of this figure see online Appendix at <http://journals.cambridge.org/geo>.

also shows good evidence for significant variability in sediment flux, with large volumes of coarse sediment being supplied into the basin at times, with channels suggested to be flowing to the north. It is thought likely that the abrupt increase in grain size is linked to these Variscan tectonic effects in the south, leading to major rejuvenation of the source area, although the cannibalism of earlier metasedimentary terrains probably also occurred.

6. Comparisons with other UK areas

In addition to the extensive developments in the English Midlands and the southern North Sea, several other areas in northern England and Scotland also preserve strata broadly equivalent to those present at Canonbie.




Figure 19. Palaeocurrent rose diagrams for channel facies from (a) Canonbie Bridge Sandstone Formation and (b) Becklees Sandstone Formation. These data are restored with respect to structural dip. Data for the Becklees and Glenzierfoot boreholes are derived from analysis of dipmeter data. No. – number of readings; PCL – primary current lineation; VM – vector mean.

6.a. Southern Scotland and Northern England

6.a.1. Southern Scotland

In southern Scotland, thick successions of late Carboniferous strata, that is, above the Aegiranum Marine Band, are locally known from the Sanquhar, SW Ayrshire, Lanarkshire (along the Central Coalfield Syncline), Thornhill, Douglas and East Fife areas (Mykura, 1967; Davies, 1970; McMillan & Brand, 1995; Forsyth, Hall & McMillan, 1996) (Fig. 1). Most authors document the presence of red-beds in these areas but typically their descriptions also include coals,



Figure 20. Example of a vertic brunified palaeosol, with complex brecciated, desiccated fabric (arrowed). This view is subparallel to bedding. Canonbie Bridge Sandstone Formation, River Esk [3944 7636]. Lens cap for scale is 5 cm across. For a colour version of this figure see online Appendix at <http://journals.cambridge.org/geo>.



Figure 21. Example of a calcimorphic palaeosol (calcrete), showing scattered, subvertical carbonate glaebules (white). Becklees Sandstone Formation, River Esk [3930 7626]. Hammer head is 0.2 m in length. For a colour version of this figure see online Appendix at <http://journals.cambridge.org/geo>.

siderite nodules, numerous rooted horizons, as well as coal replaced by limestone, indicating that later secondary reddening was an important process in their formation. Drilled approximately 6 km SSE of Glasgow, the BGS Hallside Borehole [NS 6694 5975] contains a few hundred metres of Scottish Upper Coal Measures Formation strata, comprising interbedded red sandstone, mottled mudstone and micaceous siltstone, with some purple and red palaeosols and rare coals replaced by limestone. Interestingly, desiccation cracks have been locally recorded in the core, indicating the start of better-drained conditions.

The most compelling evidence for the presence of Warwickshire Group strata comes from SW Ayrshire, where plant fossils have been suggested to be either Asturian or even Stephanian in age (Wagner, 1966). Mykura (1967) recorded up to 450 m of strata above the Aegiranum Marine Band in boreholes in this area. Of these, the uppermost ~ 120 m of strata are predominantly red-beds, and include *Spirorbis* limestones,

desiccation-cracked mudstones, mud-clast breccias and beds with calcareous concretions (Mykura, 1967), which are likely to be calcrete palaeosols. These higher strata invite comparison with the Eskbank Wood Formation at Canonbie (Fig. 22). It would appear that there is no evidence for strata equivalent to the Canonbie Bridge and Becklees Sandstone formations.

6.a.2. West Cumbria

The West Cumbrian Coalfield lies to the SSW of Canonbie and the two areas are connected in the subsurface via the Solway Syncline beneath Permo-Triassic cover (Chadwick *et al.* 1995) (Figs 1, 3). Akhurst *et al.* (1997) assigned the youngest Carboniferous strata of west Cumbria, of *phillipsii* to *tenuis* Chronozone (late Bolsovian to Asturian) age, to the Whitehaven Sandstone Formation, divisible into the Whitehaven Sandstone and the overlying Millyeat Beds members (Fig. 22).

The Whitehaven Sandstone Member is an approximately 100 m thick succession, largely comprising a reddened, locally coarse-grained to pebbly, quartz-dominant sandstone, deposited by large braided fluvial river systems that flowed from NE to SW (Jones, 1993). No calcrete palaeosols are known, although evidence for incipient primary reddening and desiccation cracks occur, suggesting that the member could be included in the Warwickshire Group. This agrees with the recent work of Waters *et al.* (2007). The Whitehaven Sandstone Member lacks the large quantities of lithic grains seen in the Canonbie Bridge Sandstone Formation and hence there would appear to be no direct lithological correlatives of this member at Canonbie (Fig. 22).

The overlying Millyeat Beds Member contains a greater proportion of mudstones than in the underlying member, comprising an alternating grey and red-bed succession of mudstone, less common sandstone, thin coals, locally present gley palaeosols that show some incipient reddening, and *Spirorbis*-bearing limestones (Brockbank, 1891; Kendall, 1896). These characteristics are similar to the facies described from the Eskbank Wood Formation of Canonbie. If this is the case, this suggests that Warwickshire Group sedimentation, represented by the Whitehaven Sandstone Member, began earlier in West Cumbria than Canonbie (Fig. 22). It is notable that there appear to be no preserved equivalents of the higher Canonbie Bridge Sandstone and Becklees Sandstone formations preserved in west Cumbria.

Other nearby coalfields, such as at Midgeholme and Stainmore, can be excluded from this discussion because only strata up to the lowermost Duckmantian are preserved (Ramsbottom *et al.* 1978; Stone *et al.* 2010). The Northumberland–Durham Coalfield preserves a sequence of Bolsovian rocks, up to 300 m thick, locally with Pennine Upper Coal Measures Formation strata, for example, in the Boldon Syncline

(Smith, 1994), but no primary red-beds are known there.

6.a.3. Ingleton

The Ingleton Coalfield forms an isolated area of Upper Carboniferous rocks preserved in a syncline at the northern margin of the Craven Basin (Fig. 1). Upper Carboniferous strata are generally not well exposed due locally to unconformable Permian cover and, more generally, an extensive blanket of Quaternary deposits. Ford (1954) provided the only modern account of the succession and recognized a lower grey, coal-bearing unit (Pennine Lower and Middle Coal Measures formations) and an upper dominantly red-bed succession, about 580 m thick, believed to be separated by a pronounced unconformity (Ford, 1954; Ramsbottom *et al.* 1978) (Fig. 22).

The ‘Red Measures’ above this unconformity have been divided into lower and upper units, with the ‘Lower Red Measures’ comprising up to 275 m of strata in the Holden House Borehole [SD 6579 7397]. These strata typically consist of red conglomerate and sandstone, mudstone, locally variegated and mottled, calcareous mudstone, ‘seatearth’ palaeosols and thin coals, and contain a bivalve fauna indicating the basal part of the *phillipsii* Chronozone (upper Bolsovian age).

The overlying ‘Upper Red Measures’, about 240 m thick, were separated from the ‘Lower’ red-beds by a series of faults (Ford, 1954). The succession was described as generally redder, dominantly comprising sandstone, but also including several beds of breccia, rooted horizons, ‘seatearth’ palaeosols and three thin limestone beds (Ford, 1954). The non-marine bivalve fauna indicates a late Bolsovian, or even possibly early Asturian age.

The descriptions of the ‘Lower Red Measures’ suggest that they largely comprise secondarily reddened Upper Carboniferous strata. The coarseness of the succession invites comparison with the similarly aged Whitehaven Sandstone Formation of West Cumbria and thus it may belong to the lowermost part of the Warwickshire Group, although this is extremely speculative. The ‘Upper Red Measures’ are tentatively, though more certainly, assigned to the Warwickshire Group and may be equivalent to the Eskbank Wood Formation of Canonbie (Fig. 22).

6.b. English Midlands, southern England and South Wales

The English Midlands saw the diachronous spread of red-beds northwards during the late Duckmantian onwards (Besly, 1998; Powell *et al.* 2000). The earliest unit within the Warwickshire Group, the Etruria Formation, comprises a mottled red mudstone-dominated succession with subordinate lenticular coarse sandstones (‘espleys’), with volcanic and lithic clasts in places (B. M. Besly, unpub. Ph.D. thesis, Keele Univ. 1983; Glover, Powell & Waters, 1993;




Figure 22. Proposed chronostratigraphical correlation of the Warwickshire Group from the Canonbie area with other areas mentioned in the text. The stratigraphy presented here is taken from a compilation of data presented in Powell *et al.* (2000) and Waters *et al.* (2007). Grey area shows the proposed correlation between the Boulton, Pennant Sandstone and Halesowen formations with the Canonbie succession. Fm. – Formation; Gp. – Group; MB – Marine Band; MBF – Millyeat Beds Member; Mbr. – Member; PMCM – Pennine Middle Coal Measures Formation; PUCM – Pennine Upper Coal Measures Formation; SST – sandstone; SMCM – Scottish Middle Coal Measure Formation; SUCM – Scottish Upper Coal Measures Formation; SWUCM – South Wales Upper Coal Measures Formation; WAWK – Warwickshire Group; WSF – Whitehaven Sandstone Formation; WSM – Whitehaven Sandstone Member.

Powell *et al.* 2000). The formation is up to 300 m thick and is known from central and northern England, North Wales and Lancashire (Powell *et al.* 2000). It typically lacks calcrete palaeosols, although well-drained palaeosols such as brunified alluvial and ferruginous palaeosols have been described, as well as post-depositionally oxidized and hydromorphic gleys (B. M. Besly, unpub. Ph.D. thesis, Keele Univ. 1983; Besly & Fielding, 1989; Besly & Cleal, 1997).

The Halesowen Formation overlies the Etruria Formation and typically comprises grey-green or red micaceous sandstones (litharenite) and mudstone with thin coals, beds of *Spirorbis* limestone and calcrete (Powell *et al.* 2000). Glover & Powell (1996) and Besly & Cleal (1997) described sandstones from the Halesowen Formation as metasediment-dominated litharenites (Association B of Besly & Cleal, 1997). These sandstones contain up to 60% low-grade metasedimentary grains, mainly foliated psammites and pelites rich in muscovite, chlorite and biotite; this gives fresh sandstones a distinctive green colour (Glover & Powell, 1996). The Halesowen Formation sandstones also contain abundant garnet, and a source rich in chrome spinel and chloritoid is suggested (Hallsworth, 1992). This metasedimentary detritus is thought to be largely recycled, derived from the Rheno-Hercynian Zone of the Variscan Orogenic belt (Besly & Cleal, 1997; Besly, 1998).

Warwickshire Group strata are also known to extend between the South Wales, Forest of Dean, Bristol, Oxfordshire, Berkshire and Kent coalfields (Kelling, 1974; Besly, 1988; Shephard-Thorn, 1988; Foster *et al.* 1989; Sherlock, Jones & Jones, 2000; Waters *et al.* 2007) (Fig. 1). Here they predominantly comprise grey, coal-bearing successions with thick lithic ('Pennant' type) sandstones. The sandstones are typically coarse-grained, locally conglomeratic and comprise feldspathic and micaceous lithic arenites with some recycled Upper Palaeozoic sedimentary and volcanic clasts (Kelling, 1974). Locally, red-beds are also known; these become more common upwards through the group (Waters *et al.* 2007). The petrological similarities between the Halesowen Formation of the Midlands and the Pennant Sandstone Formation of South Wales suggests a similar derivation, and braidplain systems sourced from the Hercynian mountains to the south have been proposed (Besly, 1988; Kelling & Collinson, 1992; Jones & Hartley, 1993; Sherlock, Jones & Jones, 2000).

Spanning the interval between the late Asturian and early Stephanian is the Salop Formation (Powell *et al.* 2000). This comprises an interbedded succession of red and red-brown mudstones and sandstones, some pebbly, with thin *Spirorbis* limestone and calcrete palaeosols in the lower part (Powell *et al.* 2000). Sandstones are mostly sublitharenite (Association C of Besly & Cleal, 1997), largely comprising monocrystalline and polycrystalline quartz and minor potassium feldspar. These sandstones also typically contain detrital carbonate grains, formed

either through reworking of calcrete or representing detritally reworked Carboniferous Limestone grains (Glover & Powell, 1996; Besly & Cleal, 1997). According to Besly & Cleal (1997), sandstones of the Salop Formation (Association C) are identifiable in the field by their distinctive orange-brown colour. The Salop Formation consists of recycled Carboniferous and perhaps Devonian lithic material and also contains calcrete palaeosols.

It is difficult to be confident of any correlation of Upper Carboniferous rocks between Canonbie and the English Midlands, due to the limits of the faunal and floral control, the large distances involved and the likelihood of lateral facies changes. Although both dominantly comprise mudstone-rich successions, the Eskbank Wood Formation is younger than the Etruria Formation. If they are equivalent this would indicate a strong element of diachroneity; hence it would appear that there is no equivalent to the Eskbank Wood Formation in the Midlands. The Halesowen Formation is of much the same age as both the Eskbank Wood and the Canonbie Bridge Sandstone formations and has a similar litharenitic composition to the latter. Heavy mineral data from the rocks at Canonbie are more restricted in their compositional variations and, in general, are characterized by lower Monazite–Zircon, Chrome-spinel–Zircon and Rutile–Zircon compositions than the rocks from the Pennine Basin (Table 3; Figs 23, 24). When the data are compared to samples from the Mexborough suite (late Duckmantian to Bolsovian) and the Halesowen Formation (Asturian), the Canonbie rocks show more similarity to the Halesowen Formation; the Mexborough suite is more heterogeneous and is characterized by a much larger range of Monazite–Zircon, Chrome-spinel–Zircon (Fig. 24) and Rutile–Zircon values (Hallsworth & Chisholm, 2000). Overall, the heavy minerals suggest that the rocks from Canonbie may contain material derived from the Variscan orogenic belt, but the data do not exclude contributions from other sources. The garnet compositions from the Canonbie Bridge Sandstone Formation also show some similarities to those measured from the Halesowen Formation (Hallsworth *et al.* 2000), but are more restricted in their compositional range (Fig. 25). However, recent zircon age dating from a sample (NHN100) from the Canonbie Bridge Sandstone Formation shows unequivocally that a significant proportion of the sediment was derived from the Variscides of western or central Europe, supporting the proposed correlation with the Halesowen Formation and the Pennant Sandstone Formation (Morton, Fanning & Jones, 2010). It also means that the idea of a local, contemporaneous source for the Canonbie red-beds, proposed by Chadwick *et al.* (1995), can be discounted.

The overlying Salop Formation also appears to have some features in common with the Becklees Sandstone Formation, particularly its distinctive orange-brown colour and lower proportion of lithic material. However,




Figure 23. Heavy mineral cross-plots for (a) Apatite–Tourmaline with Rutile–Tourmaline (ATi–RuTi), (b) Chrome-spinel–Zircon with Monazite–Zircon (CZi–MZi) and (c) Garnet–Zircon with Monazite–Zircon (GZi–MZi) for the Warwickshire Group from the Canonbie area. Data were obtained from samples NJN98 and 99 (Becklees Sandstone) and NJN100, 101 and 102 (Canonbie Bridge Sandstone). There is considerable variation in the ATi values (11–78), reflecting variable amount of apatite dissolution during sediment transport and storage. The rocks of the Canonbie Bridge Sandstone have higher ATi (62–78) than the Becklees Sandstone (42–45). All other indices GZi, MZi and CZi are low (0–7) with the Canonbie Bridge Sandstone characterized by slightly higher MZi, but similar CZi and GZi, in comparison to the Becklees Sandstone.

the Salop Formation clearly has some locally reworked material (Carboniferous Limestone grains), which is not seen in the restricted exposures of the Becklees Sandstone Formation. These correlations are extremely tentative and more work would be needed on the petrography and provenance indicators of the Canonbie succession in order to confirm any relationship.

The Upper Stephanian to Autunian successions of the Midlands are not likely to be correlatives of the Canonbie succession because they are typically quite restricted in their geographical distribution and have local source areas. For example, the Clent Formation contains a distinctive suite of clasts derived largely from Uriconian (Precambrian) rocks of the Welsh Borderlands, and the Kenilworth Sandstone Formation is a distinctive unit formed from Precambrian and Lower Palaeozoic rocks of the Lickey Ridge (Glover & Powell, 1996; Powell *et al.* 2000).

6.c. Southern North Sea

Successions chronostratigraphically equivalent to Canonbie are present offshore in the southern North Sea. The Bolsovian-aged Cleaver Formation is a grey-bed succession, suggested to have been derived from a combination of detritus sourced from the




Figure 24. Cross-plot of Chrome-spinel–Zircon with Monazite–Zircon (CZi–MZi) to compare the data from Canonbie with other examples of Westphalian sandstones from the onshore and offshore UK (data derived from Hallsworth, 1992; Hallsworth, pers. comm. 2001; Hallsworth & Chisholm, 2000; Hallsworth *et al.* 2000; Morton, Hallsworth & Moscariello, 2005). The Canonbie data are distinct from the southern North Sea Ketch and Cleaver formations (which generally have high CZi, low MZi values) and the onshore Mexborough suite (which is characterized by a much larger range of MZi and CZi), suggesting that the Canonbie rocks are not derived from the same source. There would appear to be some similarity between the Canonbie rocks and the Halesowen Formation (Daleswood Borehole). Bh – Borehole; CZi – Chrome-spinel–Zircon; Fm. – Formation; MZi – Monazite–Zircon; sst – Sandstone; SNS – southern North Sea.

Rinkøbing Fyn High to the northeast and from the Variscan orogenic belt (Saxo-Thuringian Zone) to the south (Morton, Hallsworth & Moscariello, 2005). The overlying red-bed facies of the Boulton Formation range from Bolsovian to Asturian in age and form the topmost Westphalian unit present in the UK sector of the southern North Sea (Moscariello, 2003; Besly, 2005).

The Ketch Formation largely comprises fluvial single and multi-storey braided channel deposits, with rooted overbank deposits and ferruginous palaeosols (Besly, Burley & Turner, 1993). Although the base of the Ketch Formation is defined as the lowest occurrence of primary red-bed facies (Cameron, 1993, p. 22), an angular unconformity is present at the base of the formation in condensed sections near sub-basin margins, for example, in blocks 44/13 and 44/14 (Besly, 1998; Moscariello, 2003; Jones, Allen & Morrison, 2005). Further south, in blocks 44/26 to 44/28, the Cambriense Marine Band is present and a thick coal-bearing section occurs above it (B. Besly, pers. comm. 2009).

Initially it was proposed that the ‘Barren Red Measures’ in the southern North Sea represented the diachronous spread of well-drained alluvium locally




Figure 25. Garnet compositions expressed in terms of the relative abundances of almandine plus spessartine (AS), pyrope (P) and grossular (G) end-members for (a) the Canonbie Bridge Sandstone Formation, (b) Mexborough Rock (data from Hallsworth & Chisholm 2000), (c) Halesowen Formation (data from Hallsworth *et al.* 2000) and (d) combined plot where only samples with greater than 5% spessartine are shown to distinguish garnet populations with relatively high spessartine from those which are almandine-rich. The Canonbie Bridge Sandstone Formation is predominantly characterized by high almandine+spessartine compositions with low grossular and minor pyrope. There is one anomalous garnet with considerably higher pyrope, suggesting that there is some heterogeneity in the garnet population. The garnet compositions compare favourably with those published for the Mexborough Rock and Halesowen Formation. (a) Canonbie: 22 samples from NJN101 & CAN1; (b) Mexborough Rock: 50 samples Eh14773 (Dalton Rock); (c) Halesowen Formation: 50 samples from the Daleswood Farm Borehole, at 216 m downhole depth, and (d) Canonbie (> 5% spessartine = 21 out of 22), Halesowen (> 5% spessartine = 23 out of 50), Mexborough (> 5% spessartine = 16 out of 50). CAN1, Eh14773 and NJN101 are samples registered at the British Geological Survey.

sourced from adjacent developing growth anticlines (e.g. Murdoch and Ravenspurh anticlines), which formed as a result of late Carboniferous regional compressional tectonics (Leeder & Hardman, 1990). Besly, Burley & Turner (1993) suggested that the Ketch Formation formed contemporaneously with the evolving Variscan foreland basin, with sediment derived as a result of Variscan uplift. More recent work on the heavy mineral assemblages suggests that the Ketch Formation comprises a single source area draining two distinct lithologies, one from an ultramafic source and the other from Al-poor metasedimentary rocks; the Rinkøbing Fyn High is favoured as the likely source area (Morton, Hallsworth & Moscariello, 2005).

The Asturian Boulton Formation comprises finer-grained channel facies, as well as non-marine limestones, calccrete palaeosols and some grey, coal-bearing facies (Besly, 1998, 2005). The formation is less feldspathic and richer in metamorphic lithic material than the Ketch Formation and a Variscan source is suggested for this unit (Besly, 1998; Morton, Hallsworth & Moscariello, 2005). Jones, Allen & Morrison (2005, p. 186) also described the presence of low-grade metasedimentary rock fragments in sandstones of the Boulton Formation, which they believed are indicative of a southerly derivation.

In terms of relating the onshore Canonbie succession with that of the southern North Sea, Besly (1998, p. 123) has suggested that the Ketch Formation is equivalent to that of the Canonbie succession. The Eskbank Wood Formation is lithologically similar to the Ketch Formation, but is younger (Fig. 22). Besly (1998) has also suggested that the lithic-rich Boulton Formation is equivalent to the Halesowen Formation. Chronostratigraphically it would appear to be equivalent to both the Eskbank Wood and Canonbie Bridge Sandstone formations, and lithologically it has much in common with both the onshore Halesowen and Canonbie Bridge Sandstone formations. Unfortunately, the heavy mineral character of the Boulton Formation is currently unknown. If it is comparable with the onshore Halesowen and Canonbie Bridge Sandstone formations then there would appear to be no equivalent to the overlying Salop and Becklees Sandstone formations in the currently available well penetrations in the UK sector of the southern North Sea due to sub-Permian erosion. The probable equivalents of these sections may be present in the youngest Carboniferous penetrated in the Dutch sector of the southern North Sea (the Step Graben Formation, e.g. in wells K07–08 (van Adrichem Boogaert & Kouwe, 1995) and F10–02).

7. Conclusions

Our understanding of the youngest Carboniferous rocks (Bolsovian–Asturian) in the UK is hampered by their limited preservation and the common reddening of the strata. One such Upper Carboniferous succession is present at outcrop and in boreholes around the Canonbie area of SW Scotland and adjacent parts of northern England, preserved within the NNE-trending Solway Syncline. This is one of the few areas in the UK outside of the English Midlands where such strata can be directly examined at outcrop and hence it is an important scientific site. Early workers placed these reddened Upper Carboniferous strata at Canonbie into the Upper Coal Measures and were uncertain as to their origin. From the work presented here, it is clear that these strata can be assigned to the Warwickshire Group on the basis that red-bed facies occur which show evidence for early oxidation (primary reddening) of the strata. At outcrop, about 290 m of the Warwickshire Group are almost continuously exposed along the banks of the River Esk. In the subsurface, up to about 530 m of the group is known from the Becklees Borehole, and seismic reflection data indicate it could be up to about 700 m thick in the centre of the Solway Syncline.

Three informal formations have been recognized within the Warwickshire Group: from base upwards, the Eskbank Wood, Canonbie Bridge Sandstone and Becklees Sandstone formations. These three formations have distinctive well-log signatures and hence are readily correlatable in the subsurface around Canonbie. The base of the group is diachronous, marked by the repeated alternations of grey and primary red-bed strata. Initially (Eskbank Wood Formation), sedimentation was dominated by muddy facies, with lacustrine mudstones dominant, but with common development of palaeosols, including calcretes, and thin beds of *Spirorbis* limestone. The overlying Canonbie Bridge Sandstone Formation marks the incoming of thick units of medium- and coarse-grained cross-bedded channel sandstones. These represent large, braided-river systems flowing northwards across a well-drained alluvial plain. A noticeable feature of these sandstones is their greenish grey colouration, which can be related to the presence of abundant lithic grains. Overbank and floodplain fines were commonly deposited lateral to channels and soils were able to form during intervals of low sediment aggradation.

The Becklees Sandstone Formation is the youngest unit recognized from the Warwickshire Group of Canonbie, overlain unconformably by Permian strata. Its full thickness is not known, but up to 200 m is proved in the Becklees Borehole. This sandstone is finer-grained, has a distinct orange-brown colour and contains a significantly lower proportion of lithic components.

Comparisons with other areas suggest that these are the lateral equivalents of some of the formations recognized elsewhere in Scotland, the English Midlands and in the southern North Sea. The published

descriptions of the red-beds in Ayrshire are similar to those of the Eskbank Wood Formation. The Eskbank Wood Formation has some lithological similarities to the Etruria and Cleaver formations, although it is much younger and hence it would appear that there is no equivalent to the Eskbank Wood Formation in the Midlands and southern North Sea.

Metasediment-dominated litharenites from the Halesowen Formation of the English Midlands and the Pennant Sandstone Formation of South Wales appear to be similar to those of the Canonbie Bridge Sandstone. This is largely thought to have been derived from the Variscan orogenic belt. Heavy mineral data, garnet geochemistry and zircon age dating supports this conclusion. Offshore, the Boulton Formation also contains lithic-rich sandstones that are believed to be Variscan sourced, and hence this formation could also be equivalent to the Canonbie Bridge Sandstone.

Finally, the Becklees Sandstone has much in common with the Salop Formation of the English Midlands, particularly in terms of its distinctive orange-brown colour and lower proportion of lithic material. It would appear that there is no equivalent in the UK sector of the southern North Sea, although candidates exist in the Dutch sector.

Further work is needed to confirm these conclusions and there is still much to learn about the provenance of these units. However, it is clear that southerly-derived Variscan detritus was able to travel considerable distances across the southern North Sea, UK and adjacent areas, to a position a few hundred kilometres north of what has previously been recognized. Thus, despite their somewhat limited present extent, the Canonbie Warwickshire Group strata are probably representative of the sedimentary facies laid down over all or much of northern England and its environs towards the end of the Carboniferous and are indicative of the palaeogeographical conditions prevalent there at that time. Their absence elsewhere in this area is a measure of the extent and depth of erosion beneath the sub-Permian unconformity and suggests that this is of greater magnitude than commonly assumed. This conclusion has major implications for studies attempting to model burial history and the timing of generation of hydrocarbons and mineralizing fluids from Carboniferous strata in and around northern England.

Acknowledgements. Dave Millward and Andrew McMillan from the BGS Edinburgh Office are thanked for supporting this work and they, together with Colin Waters, provided helpful comments on an early draft. Sam Holloway kindly supplied the interpretation of seismic line ED-86-02, the use of which was originally granted by Enterprise Oil. Much of the heavy mineral data were derived from reports that Claire Hallsworth compiled for BGS, the use of which is gratefully acknowledged. We also would like to express our gratitude to Andy Morton for obtaining zircon age dates to test our Variscan provenance hypothesis. Bernard Besly and an anonymous reviewer are thanked for their helpful and

constructive reviews of the manuscript. The authors publish with the permission of the Executive Director, BGS (NERC).

References

- AKHURST, M. C., CHADWICK, R. A., HOLLIDAY, D. W., McCORMAC, M., McMILLAN, A. A., MILLWARD, D. & YOUNG, B. 1997. *Geology of the west Cumbria district*. Memoir of the British Geological Survey, England and Wales, Sheets 28, 37 and 47, 138 pp.
- BAILEY, E. B. 1926. Subterranean penetration by a desert climate. *Geological Magazine* **63**, 276–80.
- BARRETT, B. H. & RICHEY, J. E. 1945. *Economic geology of Canonbie Coalfield (Dumfriesshire & Cumberland)*. Geological Survey of Great Britain Wartime Pamphlet no. 42, 1–51.
- BESLY, B. M. 1987. Sedimentological evidence for Carboniferous and Early Permian palaeoclimates of Europe. *Extrait des Annales de la Société Géologique du Nord* **151**, 131–43.
- BESLY, B. M. 1988. Palaeogeographic implications of late Westphalian to early Permian red-beds, Central England. In *Sedimentation in a synorogenic basin complex: the Upper Carboniferous of northwest Europe* (eds B. M. Besly & G. Kelling), pp. 200–21. London: Blackie.
- BESLY, B. M. 1998. Carboniferous. In *Petroleum Geology of the North Sea: basic concepts and recent advances, Fourth Edition* (ed. K. W. Glennie), pp. 104–36. Oxford: Blackwell.
- BESLY, B. M. 2005. Late Carboniferous redbeds of the UK southern North Sea, viewed in a regional context. In *Carboniferous hydrocarbon geology: the southern North Sea and surrounding onshore areas* (eds J. D. Collinson, D. J. Evans, D. W. Holliday & N. S. Jones), pp. 225–6. Yorkshire Geological Society Occasional Publication no. 7.
- BESLY, B. M. & CLEAL, C. J. 1997. Upper Carboniferous stratigraphy of the West Midlands (UK) revised in the light of borehole geophysical logs and detrital compositional suites. *Geological Journal* **32**, 85–118.
- BESLY, B. M. & FIELDING, C. R. 1989. Palaeosols in Westphalian coal-bearing and red-bed sequences, Central and Northern England. *Palaeogeography, Palaeoclimatology, Palaeoecology* **70**, 303–30.
- BESLY, B. M. & TURNER, P. 1983. Origin of red-beds in a moist tropical climate (Etruria Formation, Upper Carboniferous, UK). In *Residual Deposits* (ed. R. C. L. Wilson), pp. 131–47. Geological Society of London, Special Publication no. 11.
- BESLY, B. M., BURLEY, S. D. & TURNER, P. 1993. The late Carboniferous ‘Barren Red-bed’ play of the Silver Pit area, Southern North Sea. In *Petroleum Geology of Northwest Europe: Proceedings of the 4th Conference* (ed. J. R. Parker), pp. 727–40. London: Geological Society.
- BRITISH GEOLOGICAL SURVEY. 1999. *Coal Resources Map of Britain*. G. R. Chapman, Natural Environment Research Council and the Coal Authority.
- BROCKBANK, W. 1891. On the occurrence of the Permians, Spirorbis Limestones, and Upper Coal Measures at Frizington Hall in the Whitehaven District. *Memoirs of the Proceedings of the Manchester Literary and Philosophical Society* **4**, 418–26.
- CAMERON, T. D. J. 1993. 5. Carboniferous and Devonian of the Southern North Sea. In *Lithostratigraphic nomenclature of the UK North Sea* (eds R. W. O’B. Knox & W. G. Cordey), 93 pp. Nottingham: British Geological Survey.
- CHADWICK, R. A., HOLLIDAY, D. W., HOLLOWAY, S. & HULBERT, A. G. 1995. *The structure and evolution of the Northumberland–Solway Basin and adjacent areas*. Subsurface Memoir of the British Geological Survey, 90 pp.
- CORFIELD, S. M., GAWTHORPE, R. L., GAGE, M., FRASER, A. J. & BESLY, B. M. 1996. Inversion tectonics of the Variscan foreland of the British Isles. *Journal of the Geological Society, London* **153**, 17–32.
- DAVIES, A. 1970. Carboniferous rocks of the Sanquhar outlier. *Bulletin of the Geological Survey of Great Britain* **31**, 37–88.
- DAY, J. B. W. 1970. *Geology of the country around Bewcastle*. Memoir of the Geological Survey of Great Britain England and Wales, 357 pp.
- DE PAOLA, N., HOLDSWORTH, R. E., MCCAFFREY, K. J. W. & BARCHI, M. R. 2005. Partitioned transtension: an alternative to basin inversion models. *Journal of Structural Geology* **27**, 607–25.
- FORD, T. D. 1954. The Upper Carboniferous rocks of the Ingleton Coalfield. *Quarterly Journal of the Geological Society of London* **110**, 231–65.
- FORSYTH, I. H., HALL, I. H. S. & McMILLAN, A. A. 1996. *Geology of the Airdrie district*. Memoir of the British Geological Survey, Sheet 31W (Scotland), 94 pp.
- FOSTER, D., HOLLIDAY, D. W., JONES, C. M., OWENS, B. & WELSH, A. 1989. The concealed Upper Palaeozoic rocks of Berkshire and South Oxfordshire. *Proceedings of the Geological Association* **100**, 395–407.
- GALEHOUSE, J. S. 1971. Point-counting. In *Procedures in Sedimentary Petrology* (ed. R. E. Carver), pp. 385–407. New York: Wiley-Interscience.
- GLOVER, B. W., POWELL, J. H. & WATERS, C. N. 1993. Etruria Formation (Westphalian C) palaeoenvironments and volcanicity on the southern margins of the Pennine Basin, South Staffordshire, England. *Journal of the Geological Society, London* **150**, 737–50.
- GLOVER, B. W. & POWELL, J. H. 1996. Interaction of climate and tectonics upon alluvial architecture: Late Carboniferous–Early Permian Sequences at the southern margin of the Pennine Basin. *Palaeogeography, Palaeoclimatology and Palaeoecology* **121**, 13–34.
- HALLSWORTH, C. R. 1992. *Stratigraphic variations in the heavy minerals and clay mineralogy of the Westphalian to ?Early Permian succession from the Daleswood Farm borehole, and the implications for sand provenance*. British Geological Survey Technical Report WH/92/185R.
- HALLSWORTH, C. R. & CHISHOLM, J. I. 2000. Stratigraphic evolution of provenance characteristics in Westphalian sandstones of the Yorkshire Coalfield. *Proceedings of the Yorkshire Geological Society* **53**, 43–72.
- HALLSWORTH, C. R., MORTON, A. C., CLAUDE-LONG, J. & FANNING, C. M. 2000. Carboniferous sand provenance in the Pennine Basin, UK: constraints from heavy mineral and detrital zircon age data. *Sedimentary Geology* **137**, 147–85.
- JONES, C. M., ALLEN, P. J. & MORRISON, N. H. 2005. Geological factors influencing gas production in the Tyne field (Block 44/18a), southern North Sea, and their impact on future infill well planning. In *Carboniferous hydrocarbon geology: the southern North Sea and surrounding onshore areas* (eds J. D. Collinson, D. J. Evans, D. W. Holliday & N. S. Jones), pp. 183–94. Yorkshire Geological Society Occasional Publication no. 7.

- JONES, J. A. & HARTLEY, A. J. 1993. Reservoir characteristics of a braid-plain depositional system: the Upper Carboniferous Pennant Sandstone of South Wales. In *Characterization of Fluvial and Aeolian Reservoirs* (eds C. P. North & D. J. Prosser), pp. 143–56. Geological Society of London, Special Publication no. 73.
- JONES, N. S. 1993. *Stratigraphical and sedimentological characteristics of the 'Whitehaven Sandstone Series', Westphalian C, West Cumbrian Coalfield*. British Geological Survey Technical Report WH/93/102R.
- JONES, N. S. & HOLLIDAY, D. W. 2007. *The stratigraphy and sedimentology of Upper Carboniferous Warwickshire Group red-bed facies in the Canonbie area of S.W. Scotland*. British Geological Survey Report IR/06/043.
- KELLING, G. 1974. Upper Carboniferous sedimentation in South Wales. In *The Upper Palaeozoic and post-Palaeozoic rocks of Wales* (ed. T. R. Owen), pp. 185–224. University of Wales Press.
- KELLING, G. & COLLINSON, J. D. 1992. Silesian. In *Geology of England and Wales* (eds P. M. D. Duff & A. J. Smith), pp. 239–73. London: Geological Society.
- KENDALL, J. D. 1896. The Whitehaven Sandstone Series. *Transactions of the Institution of Mining Engineers* **10**, 204–24.
- LEEDER, M. R. & HARDMAN, M. 1990. Carboniferous geology of the Southern North Sea Basin and controls on hydrocarbon prospectivity. In *Tectonic Events Responsible for Britain's Oil and Gas Reserves* (eds R. F. P. Hardman & J. Brooks), pp. 87–105. Geological Society of London, Special Publication no. 55.
- LUMSDEN, G. I., TULLOCH, W., HOWELLS, M. F. & DAVIES, A. 1967. *The Geology of the Neighbourhood of Langholm*. Memoirs of the Geological Survey, Scotland, 255 pp.
- MCMILLAN, A. A. & BRAND, P. J. 1995. Depositional setting of Permian and Upper Carboniferous strata of the Thornhill Basin, Dumfriesshire. *Scottish Journal of Geology* **31**, 43–52.
- MORTON, A. C. & HALLSWORTH, C. R. 1994. Identifying provenance-specific features of detrital heavy mineral assemblages in sandstones. *Sedimentary Geology* **90**, 241–56.
- MORTON, A. C. & HALLSWORTH, C. R. 1999. Processes controlling the composition of heavy mineral assemblages in sandstones. *Sedimentary Geology* **124**, 3–29.
- MORTON, A. C., CLAOUÉ-LONG, J. C. & HALLSWORTH, C. R. 2001. Zircon age and heavy mineral constraints on provenance of North Sea Carboniferous sandstones. *Marine and Petroleum Geology* **18**, 319–37.
- MORTON, A. C., HALLSWORTH, C. R. & MOSCARIELLO, A. 2005. Interplay between northern and southern sediment sources during Westphalian deposition in the Silverpit Basin, southern North Sea. In *Carboniferous hydrocarbon geology: the southern North Sea and surrounding onshore areas* (eds J. D. Collinson, D. J. Evans, D. W. Holliday & N. S. Jones), pp. 135–46. Yorkshire Geological Society Occasional Publication no. 7.
- MORTON, A. C., FANNING, M. & JONES, N. S. 2010. Variscan sourcing of Westphalian (Pennsylvanian) sandstones in the Canonbie Coalfield, UK. *Geological Magazine*, doi: 10.1017/S0016756810000014, published online 12 Feb 2010.
- MOSCARIELLO, A. 2003. The Schooner Field, Blocks 44/26a, 43/30a, UK North Sea. In *United Kingdom Oil and Gas Fields, Commemorative Millennium Volume* (eds J. G. Gluyas & H. M. Hichins), pp. 811–24. Geological Society of London, Memoir no. 20.
- MYKURA, W. 1960. The replacement of coal by limestone and the reddening of Coal Measures in the Ayrshire Coalfield. *Bulletin of the Geological Survey of Great Britain* **16**, 69–109.
- MYKURA, W. 1967. The Upper Carboniferous rocks of south-west Ayrshire. *Bulletin of the Geological Survey of Great Britain* **26**, 23–98.
- PARRISH, J. T. 1982. Upwelling and petroleum source beds, with reference to the Paleozoic. *American Association of Petroleum Geologists Bulletin* **66**, 750–74.
- PEACH, B. N. & HORNE, J. 1903. The Canonbie Coalfield: its Geological Structure and Relations to the Carboniferous Rocks of Northern England and Central Scotland. *Transactions of the Royal Society of Edinburgh* **40**, 835–77.
- PETTIJOHN, F. J., POTTER, P. E. & SIEVER, R. 1987. *Sand and Sandstone, 2nd edition*. New York: Springer-Verlag, 553 pp.
- PICKEN, G. S. 1988. The concealed coalfield at Canonbie: an interpretation based on boreholes and seismic surveys. *Scottish Journal of Geology* **24**, 61–71.
- POWELL, J. H., CHISHOLM, J. I., BRIDGE, D. MCC., REES, J. G., GLOVER, B. W. & BESLY, B. M. 2000. *Stratigraphical framework for Westphalian to Early Permian red-bed successions of the Pennine Basin*. British Geological Survey Research Report RR/00/01.
- RAMSBOTTOM, W. H. C., CALVER, M. A., EAGER, R. M. C., HODSON, F., HOLLIDAY, D. W., STUBBLEFIELD, C. J. & WILSON, R. B. 1978. *A correlation of Silesian rocks in the British Isles*. Geological Society of London, Special Report no. 10, 81 pp.
- ROWLEY, D. B., RAYMOND, A., PARRISH, J. T., LOTTES, A. L., SCOTESE, C. R. & ZIEGLER, A. M. 1985. Carboniferous palaeogeographic, phytogeographic and palaeoclimatic reconstruction. *International Journal of Coal Geology* **5**, 7–42.
- SHEPHARD-THORN, E. R. 1988. *Geology of the country around Ramsgate and Dover*. Memoir of the British Geological Survey, Sheets 274 and 279 (England and Wales), 49 pp.
- SHERLOCK, S. C., JONES, K. A. & JONES, J. A. 2000. A central European Variscide source for Upper Carboniferous sediments in SW England: ⁴⁰Ar/³⁹Ar detrital white mica ages from the Forest of Dean. *Journal of the Geological Society, London* **157**, 905–8.
- SIMPSON, J. B. & RICHEY, J. E. 1936. *The geology of the Sanquhar Coalfield*. Memoirs of the Geological Survey, Scotland, 97 pp.
- SMITH, D. B. 1994. *Geology of the country around Sunderland*. Memoir of the British Geological Survey, sheet 21 (England and Wales), 161 pp.
- STONE, P., MILLWARD, D., YOUNG, B., MERRITT, J. W., CLARKE, S. M., MCCORMAC, M. & LAWRENCE, D. J. D. 2010. *British Regional Geology Northern England (5th edition)*. Keyworth, Nottingham: British Geological Survey.
- TROTTER, F. M. 1953. Reddened beds of Carboniferous age in North-West England and their origin. *Proceedings of the Yorkshire Geological Society* **29**, 1–20.
- TROTTER, F. M. 1954. Reddened beds in the Coal Measures of South Lancashire. *Bulletin of the Geological Survey of Great Britain* **5**, 61–80.
- TRUEMAN, A. E. & WEIR, J. 1946–58. *A monograph on British Carboniferous non-marine Lamellibranchia*. Palaeontographical Society Monograph, 449 pp.

- TURNER, P. 1980. *Continental Red Beds*. Developments in Sedimentology, vol. 29. Amsterdam: Elsevier, 562 pp.
- VAN ADRICHEM BOOGAERT, H. A. & KOUWE, W. F. P. 1995. Stratigraphic Nomenclature of the Netherlands, revision and update by RGD and NOGEPa, Section C: Silesian (Limburg Group). *Mededelingen Rijks Geologische Dienst* **50**, 40 pp.
- WAGNER, R. H. 1966. On the presence of probably Upper Stephanian beds in Ayrshire. *Scottish Journal of Geology* **2**, 122–3.
- WATERS, C. N., BROWNE, M. A. E., DEAN, M. T. & POWELL, J. H. 2007. *Lithostratigraphical framework for Carboniferous successions of Great Britain (Onshore)*. British Geological Survey Research Report RR/07/01.