

Psychological characteristics and motivation of women seeking labiaplasty

D. Veale^{1*}, E. Eshkevari¹, N. Ellison¹, A. Costa¹, D. Robinson², A. Kavouni³ and L. Cardozo²

¹NIHR Specialist Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and the Institute of Psychiatry, King's College London, London, UK

²Department of Urogynaecology, King's College London, London, UK

³Cosmetic Solutions, London, UK

Background. Labiaplasty is an increasingly popular surgical intervention but little is known about the characteristics and motivation of women who seek the procedure or the psychosexual outcome.

Method. A total of 55 women seeking labiaplasty were compared with 70 women who did not desire labiaplasty. Various general measures of psychopathology as well as specific measures (Genital Appearance Satisfaction; Cosmetic Procedure Screening for labiaplasty) were used. Labia measurements of the women seeking labiaplasty were also obtained.

Results. Women seeking labiaplasty did not differ from controls on measures of depression or anxiety. They did, however, express increased dissatisfaction towards the appearance of their genitalia, with lower overall sexual satisfaction and a poorer quality of life in terms of body image. Women seeking labiaplasty reported a significantly greater frequency of avoidance behaviours on all the domains assessed, and greater frequency of safety-seeking behaviours for most of the domains. Key motivations reported for labiaplasty were categorized as cosmetic, functional or sexual. Of the 55 women seeking labiaplasty, 10 met diagnostic criteria for body dysmorphic disorder.

Conclusions. This is the first controlled study to describe some of the characteristics and motivations of women seeking labiaplasty. We identified a wide range of avoidance and safety-seeking behaviours, which occurred more frequently in the labiaplasty group than the control group. These could be used clinically as part of a psychological intervention for women seeking labiaplasty.

Received 18 December 2012; Revised 1 April 2013; Accepted 11 April 2013; First published online 10 May 2013

Key words: Body dysmorphic disorder, female genital cosmetic surgery, labia, labiaplasty.

Introduction

Labiaplasty is a surgical procedure in women that usually reduces the degree of protrusion of the labia minora. The desire for labiaplasty is becoming increasingly common, with the incidence in the National Health Service (NHS) of 1726 in the year 2010–2011 (Health and Social Care Information Centre, 2012). The number of labiaplasties conducted in the private sector is unknown, but the procedure is often discussed in the media and marketed on the Internet. Liao *et al.* (2010) identified 18 publications covering 937 case reports or series of labiaplasty worldwide up to March 2009. We conducted a similar search of reports published after March 2009 and found a further six publications and 64 additional cases.

Social and cultural factors might be driving demand for general cosmetic surgery. Some women may be

seeking flat vulvas with no protrusion beyond the labia majora, from seeing porn actresses on the Internet or desiring a prepubescent aesthetic ideal that is seen in advertisements (Liao & Creighton, 2007) or women's magazines (Bramwell, 2002). However there is still a lack of knowledge regarding the motivation and psychological characteristics of women who seek labiaplasty. Bramwell *et al.* (2007) have conducted a qualitative study, involving a thematic analysis of a structured interview, on six women who had previously undergone labiaplasty. They reported that all the women described feeling their genital appearance as 'weird' or abnormal prior to the surgery. Women reported the appearance of their genitals as having a negative impact on their sex lives, including inhibition of sexual relationships. Braun (2009) has also reviewed the reasons for undergoing labiaplasty in previous case reports provided by surgeons and found that they fell broadly into either aesthetic or functional categories.

Some women seeking labiaplasty may have body dysmorphic disorder (BDD). This is characterized by a preoccupation with a perceived defect that is not observable or appears slight to others while the

* Address for correspondence: D. Veale, Centre for Anxiety Disorders and Trauma, The Maudsley Hospital, 99 Denmark Hill, London SE5 8AZ, UK.
(Email: David.Veale@kcl.ac.uk)

person's concern is markedly excessive. Crouch *et al.* (2011) have described the size of the labia of women seeking labiaplasty to be within normal published limits. To fulfill the diagnostic criteria for BDD, however, the perceived defect must be either significantly distressing or cause impairment in social, occupational or other important areas of functioning. The most common preoccupations in BDD are the facial skin, nose, eyes, eyelids, mouth and chin – or just being ugly in general (Neziroglu & Yaryura-Tobias, 1993; Phillips *et al.* 1993; Veale *et al.* 1996 *a,b*; Phillips & Diaz, 1997). However, a preoccupation with the size or shape of the labia being abnormal or too large is an uncommon presentation of BDD in a mental health setting.

The current study was therefore exploratory and designed to discover more about the characteristics and motivation of women seeking labiaplasty by comparing such a population with a control group of women from the community who were not seeking labiaplasty. We hypothesized that women seeking labiaplasty, in comparison with control women, would report significantly worse symptoms of anxiety and depression, reduced sexual satisfaction, greater negative effect of body image on quality of life, lower genital appearance satisfaction, and greater frequency of BDD. Another aim of the study was to explore the motivation for seeking labiaplasty and to examine the extent to which women's concerns with their labia interfered in their life. We also wanted to identify the specific avoidance and safety-seeking behaviours that may be relevant for developing a psychological intervention. A safety-seeking behaviour is any behaviour performed in order to prevent or minimize a feared catastrophe (Salkovskis, 1991). For women seeking labiaplasty the catastrophe may include being humiliated or rejected by others. We hypothesized that the form of the behaviours may be similar to those seen in BDD (Lambrou *et al.* 2012). Thus the safety strategies may be designed to verify whether the labia are abnormal or to camouflage the genitalia during sexual encounters. Avoidance behaviours might include avoidance of intimacy or public changing rooms.

Method

Participants

We recruited 125 women who were categorized into two groups:

- (1) Women seeking labiaplasty. We recruited 55 women seeking labiaplasty from the following sources: (a) 31 (56.4%) from a private cosmetic clinic. These were recruited from a total of 73 women who had labiaplasty in the recruitment period who were given written information about the study; (b) 19 (34.5%)

from an NHS gynaecology clinic. These were drawn from a total of 35 women who had a labiaplasty and were given information about the study (c); five (9.1%) by an email from a research volunteer database of individuals in the community (Mind Search) at the Institute of Psychiatry, King's College London. The Mind Search database contains details of over 3500 individuals who have volunteered to participate in psychological or psychiatric research. The five women who volunteered were not seen in a clinic but were either seeking labiaplasty or indicated that they would seek labiaplasty if they could afford it in the future.

- (2) Controls. We recruited 70 women for the control group from the following sources: (a) 31 (44.3%) from a gynaecology clinic in the state sector, where the individuals recruited by the surgeon were having a non-cosmetic gynaecological surgical procedure on the NHS; and (b) 39 (55.7%) by an email to the research volunteer database (Mind Search, described above). The women in the control group were asked to participate in a study that aimed to explore women's attitudes towards their external genitalia. They were characterized by not requesting or wanting labiaplasty.

In order to take part, all participants were required to be between 18 and 60 years of age, and proficient in English (in order to provide consent and complete the questionnaires). There were no significant differences between the two groups in age, sexual orientation, marital status, education, ethnicity or parity (Table 1). There was a trend towards Black or Black British women in the control group but this was not statistically significant.

Procedure

Participants in the gynaecology or cosmetic surgery setting were recruited after they had been assessed by a surgeon and invited to participate in the study. Participants were also recruited from Mind Search whereby their contact details (emails) were provided to us for 225 people. Of these volunteers invited to participate by email, 51 responded and 41 participated to completion. Participants from both groups completed the questionnaires listed below, either online or in a pen-and-paper format. Participants who scored more than the cut-off score on the Cosmetic Procedure Screening questionnaire (Veale *et al.* 2011) were interviewed to determine whether they might fulfil the diagnostic criteria of BDD. A trained research worker experienced in BDD conducted the interview.

Only participants in the labiaplasty group were asked about the duration of their problem and their motivation for seeking labiaplasty.

Table 1. Participant demographics and comparisons between the labiaplasty ($n=55$) and control ($n=70$) groups

	Labiaplasty	Control	Comparison
Median age, years (IQR)	30.00 (20.5–39.5)	28.50 (18.5–38.5)	$U=1902.50$ $Z=-0.11$ $p=0.911$
Sexual orientation, n (%)			
Only to the opposite sex	39 (73.6)	52 (74.3)	$\chi^2=1.129$ $df=2$ $p=0.609$
Mainly to the opposite sex	11 (20.8)	11 (15.7)	
Equally to both sexes	3 (5.7)	7 (10.0)	
Mainly to the same sex	0	0	
Only to the same sex	0	0	
Marital status, n (%)			
Single	35 (63.6)	38 (54.3)	$\chi^2=3.870$ $df=3$ $p=0.286$
Separated/divorced	7 (12.7)	5 (7.1)	
Married/cohabiting	12 (21.8)	26 (37.1)	
Widowed	1 (1.8)	1 (1.4)	
Children, n (%)			
Yes	15 (27.3)	22 (31.4)	$\chi^2=0.255$ $df=1$ $p=0.613$
No	40 (72.7)	48 (68.6)	
Ethnicity, n (%)			
White	48 (87.3)	53 (75.7)	$\chi^2=7.17$ $df=3$ $p=0.067$
Black/Black British	0	8 (11.4)	
Mixed	2 (3.6)	4 (5.7)	
Other	3 (5.5)	3 (4.3)	
Higher education, n (%)			
Secondary	19 (34.5)	21 (30)	$\chi^2=0.253$ $df=1$ $p=0.615$
Tertiary	35 (63.6)	47 (67)	

IQR, Interquartile range; df, degrees of freedom.

Materials

Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983)

The HADS is a 14-item self-report instrument used to examine the severity of anxiety and depressive symptoms in two separate subscales. Each subscale comprises seven items, and the possible range of scores is from 0 to 21 on each subscale. A score between 8 and 10 identifies borderline cases, 11 and 15 moderate cases, and 16 or above severe cases.

The Prolapse–Urinary Incontinence Sexual Function Questionnaire (PISQ; Rogers et al. 2001)

The PISQ has 31 items. Each item has a range of 0–4 (except for item 5, which has a range of 0–5) and scores range from 0 to 125. Despite the name, the PISQ covers a broad measure of sexual satisfaction in women. Higher scores represent increasing sexual satisfaction.

Body Image Quality of Life Inventory (BIQLI; Cash & Fleming, 2002; Hrabosky et al. 2009)

The BIQLI is a 19-item self-report assessment scale that measures the impact of body image concerns on a broad range of life domains (e.g. sense of self, social functioning, sexuality, emotional well-being, eating, exercise, grooming). Each item is rated by the participant on a seven-point Likert scale, ranging from –3 (very negative effect) to +3 (very positive effect). The BIQLI is scored as an average numeric score of the 19 items, where a more negative score reflects a more negative body image affecting the quality of life.

Genital Appearance Satisfaction (GAS) scale (Bramwell & Morland, 2009)

The GAS scale contains 11 statements about attitudes towards genital appearance to be rated by the participant. Each item is scored between 0 and 3 on a Likert frequency scale. Total scores range from 0 to 33.

Sample items include 'I feel discomfort around my genitalia when I wear tight clothes'; 'I feel that my genital area looks asymmetric or lop-sided'; and 'I feel my labia are too large'. Higher scores represent greater dissatisfaction with the genitalia. Cronbach's α for the overall scale is 0.82.

Cosmetic Procedure Screening Scale for BDD (COPS; Veale et al. 2011) and Yale–Brown Obsessive Compulsive Scale for BDD (BDD-YBOCS; Phillips et al. 1997)

The COPS is a nine-item questionnaire that is validated for screening for BDD in people with general appearance concerns. Participants are asked to list the features of their body in the order for which they had concerns. The items follow the diagnostic features of BDD. The total scores range from 0 to 72. The questionnaire has acceptable internal consistency (Cronbach's α is 0.91). Participants who scored more than the cut-off score of 40 were interviewed using a module within the Structured Clinical Interview for DSM-IV disorders (SCID; First et al. 1995). We did not use the SCID to determine other psychiatric disorders unless body image symptoms were better explained by another diagnosis (e.g. anorexia nervosa). Those found to have a diagnosis of BDD were then interviewed with the BDD-YBOCS (Phillips et al. 1997). This is a 12-item observer-rated scale. Items are rated on a scale from 0 to 4, and the range for total score is from 0 to 48. Higher scores indicate greater BDD symptomatology. A clinical case is likely to score 24 or more on the BDD-YBOCS.

Cosmetic Procedure Screening for labiaplasty (COPS-L; Veale et al. 2013)

The COPS has been modified to focus on concerns about the appearance of the labia rather than general appearance. The domains follow the diagnostic criteria for BDD and include items such as perceived abnormality, effect on sexual relationship, interference with leisure activities, noticeability in public). The COPS-L is thus a general measure of genital appearance distress and overlaps to a certain extent with the GAS scale. The measure consists of nine items, with total scores ranging from 0 to 72 and higher scores indicating greater impairment. Cronbach's α is 0.91.

Avoidance and safety-seeking behaviours

A checklist of avoidance and safety-seeking behaviours related to the genitalia was drawn from those commonly occurring in BDD (Lambrou et al. 2012). We modified items to relate to the genitalia and generated additional items after interviewing six women seeking labiaplasty, before commencing the study. The final

avoidance checklist consisted of eight items and the final safety-seeking behaviour checklist consisted of 13 items that participants rated on a Likert scale of frequency ranging from 0 (never) through to 4 (always). Respondents had the opportunity to add any avoidance and safety-seeking behaviours that they did which were not listed.

Past cosmetic procedures

Participants were asked an open question: 'Have you had any cosmetic procedure(s) in the past (including Botox injections, dermabrasion, cellulite treatment, chemical peels, collagen injections, laser hair removal or laser skin resurfacing, as well as surgery like breast implants, nose reshaping or liposuction)?' They were then requested to report up to five previous procedures.

Interference in life

We asked specific questions concerning the extent to which the labia interfere with life. The scale was based on the Work and Social Adjustment Scale (Mundt et al. 2002) in which participants are asked to rate the degree of interference in: (a) relationships in general or dating; (b) sexual life; (c) social life; (d) leisure; (e) discomfort; (f) hygiene; and (g) distress in general. Interference and distress were rated on an eight-point Likert scale from 0 ('not at all') to 8 ('extremely').

Labiaplasty group only

A surgeon measured the degree of protrusion of the labia minora and width of each labium with a disposable tape measure in the labiaplasty group. All measurements were made in the lithotomy position with minimal stretching of the labia. The width was measured anterior-posteriorly from the clitoral hood and the lower aspect of the labia minora. We took the average of left and right measurements.

The labiaplasty group completed two additional measures:

- (1) Reported motivation for labiaplasty. We explored the motivation for seeking labiaplasty by asking an open-ended question: 'Please describe your main reason for seeking labiaplasty'. The responses were analysed independently by two raters (D.V. and E.E.) and coded into three categories, which emerged from the participants' accounts: aesthetic, functional and sexual. The researchers discussed the process by which the themes were derived to ensure the analysis was logical and valid. There was also a 'combination' category where two or three reasons were provided. The category 'aesthetic' was defined as any response that included

the evaluation of the appearance of the genitalia, such as being ugly or unfeminine, sapping confidence or causing self-consciousness. The category 'functional' was defined as any response that referred to physical symptoms such as irritation, discomfort or pain or interference in physical activities (other than sexual ones). The category 'sexual' was defined as any reason grounded in a sexual context, including avoidance and interference in intimacy and sexual relationships. Two of the authors rated participants' responses independently ($n=48$).

- (2) Duration of the problem. Participants were asked to report the duration of the problem with their labia with the following questions: (a) 'For how long have you felt there was a problem with the appearance or function of your labia?'; (b) 'How long is it since you wished to have a procedure to reduce the size of your labia?'

Statistical analysis

Data were analysed using SPSS v20 (IBM, USA). Given the non-normal distribution of most of these variables, as demonstrated from Kolmogorov–Smirnov tests, non-parametric parameters (e.g. median and interquartile range; IQR) and comparison tests (χ^2 , Mann–Whitney U tests) are reported. All tests were two-tailed and α was set at 0.05%.

Ethics

The Joint South London and Maudsley Trust and the Institute of Psychiatry NHS Research Ethics Committee granted ethics permission (09/H0807/33). Consent to contact was obtained by the surgeon. Informed consent was obtained over the telephone.

Results

Standardized questionnaires

Table 2 reports the differences between the two groups on the standardized questionnaires. There were no significant differences in the severity of symptoms of anxiety or depression on the HADS.

The control group had significantly higher body image quality of life than the labiaplasty group as evident on the BIQLI, with a small effect size. As expected, the labiaplasty group had significantly higher dissatisfaction towards the appearance of their genital area compared with the control group as evident on the GAS and the COPS-L total scores, with a large effect size. The labiaplasty group had significantly lower overall sexual satisfaction than the control group as evident on the PISQ, with a small effect size.

Avoidance and safety-seeking behaviours

Women seeking labiaplasty reported a significantly greater frequency of avoidance behaviours compared with the control group on all the domains assessed (Table 3). Large effect sizes were found for avoidance of the sight of their own genitalia; going to public changing rooms; sexual intercourse or intimacy; and wearing certain types of clothes (e.g. underwear or tight-fitting clothes or a swimming costume).

The frequency of safety-seeking behaviours was also significantly greater for all the domains, except for those that involve reassurance seeking or checking by photographing of the labia (Table 4).

Interference in life

There were significant differences between the labiaplasty group and the control group for all the domains assessed (Table 5). The highest effect size was found for interference in sexual life, discomfort and distress in general. Examples of qualitative responses for the questions on interference in life are shown below:

- (1) Relationships in general: 'My husband has no empathy regarding my labia-concerns, and I can't understand why he doesn't think I'm a freak and therefore why he isn't forcing me to get a labiaplasty'; 'It doesn't affect the dating so much, just the build up to sleeping with the person isn't exciting – the relationship is stressful rather than fun'.
- (2) Sexual relationships: 'The labia can get trapped in my vagina during intercourse. It's uncomfortable and then I have to release them'; 'I won't date and I have been celibate for 5 years'; 'I feel very uncomfortable about my partner giving oral sex'.
- (3) Social life: 'It restricts what I can wear so certain things I cannot go to as I cannot get comfortable which then ruins the outing'; '...when I walk or wear tight clothes and it isn't always convenient to re-adjust myself'.
- (4) Leisure: 'Any kind of sports – running, bike riding – are all uncomfortable. The labia rub more and are more uncomfortable'; 'I used to swim weekly, however since private showers are no longer available I will not go'.
- (5) Discomfort: 'It gets trapped between my underwear and my leg and is very painful, I am constantly trying to readjust myself to get comfortable all the time'; 'Labia are long and get in the way, I sit on them sometimes'.
- (6) Hygiene: 'More skin generates sweat therefore feels more dirty'; 'I feel I have to be more particular when washing and drying myself – it's sometimes difficult to get my vagina dry properly after a

Table 2. Comparisons of the labiaplasty ($n=55$) and control ($n=70$) groups on standardized questionnaires

	Labiaplasty: median (interquartile range)	Control: median (interquartile range)	Comparison	Effect size: Cohen's d
HADS – anxiety	9.0 (5.0 to 13.0)	8.0 (4.0 to 12.0)	$U=1479.00$ $Z=-1.124$ $p=0.263$	0.20
HADS – depression	3.0 (0.5 to 5.5)	4.0 (1.5 to 6.5)	$U=1541.50$ $Z=-0.783$ $p=0.436$	0.14
BIQLI	-0.21 (-1.21 to 0.79)	0.42 (-0.68 to 1.63)	$U=1284.00$ $Z=-2.041$ $p=0.041$	0.37
GAS scale	23.5 (19.88 to 27.63)	6.00 (3.14 to 8.88)	$U=66.50$ $Z=-9.026$ $p<0.001$	2.87
COPS	24.0 (16.75 to 31.25)	14.00 (7.0 to 21.0)	$U=908.50$ $Z=-3.393$ $p=0.001$	0.68
COPS-L	42.0 (29.5 to 54.5)	4.0 (4.0 to 12.0)	$U=99.00$ $Z=-9.093$ $p<0.001$	3.09
PISQ	90.0 (78.5 to 101.5)	99.0 (89.5 to 108.5)	$U=1195.00$ $Z=-2.461$ $p=0.014$	0.47

HADS, Hospital Anxiety and Depression Scale; BIQLI, Body Image Quality of Life Inventory; GAS, Genital Appearance Satisfaction; COPS, Cosmetic Procedure Screening Scale for body dysmorphic disorder; COPS-L, Cosmetic Procedure Screening for labiaplasty; PISQ, Prolapse–Urinary Incontinence Sexual Function Questionnaire.

shower'; 'Sometimes after going to the toilet, it drips as there is so much of it'.

- (7) General distress: 'I feel like a "freak". I know it doesn't bother other women but to me they were very ugly'; 'It is disgusting, not natural and certainly not something a 22 year old should have. I get very upset over it'; 'I feel they are unsightly and look like an elephant's vagina! Big and grey – ough!'

Labia measurements

Comparisons of the average width of the labia minora of the private patients (mean=28.09 mm, s.d.=6.04 mm, $n=23$, range 17–41.5 mm) and the NHS patients (mean=40.27 mm, s.d.=6.99 mm, $n=11$, range 30–52.5 mm) in a non-parametric independent-samples comparison test demonstrated that the NHS patients appeared to have significantly greater labia minora width than the private patients ($U=20.50$, $Z=-3.91$, $p<0.001$). It was not possible to check on the inter-rater reliability of the two surgeons. All the women were, however, in

the normal range found for women in the community. For example, Lloyd *et al.* (2005) found that women had a mean width of 21.8 mm (s.d.=9.4 mm, range 7–50 mm, $n=50$).

The association between the degree of protrusion of the labia minora and the degree of dissatisfaction with the labia (as reflected on the GAS scores) was explored using Spearman's rho. A non-significant correlation was found between measurements of the protrusion of the labia minora and the GAS scores in the private patients ($r=-0.405$, $p=0.068$, $n=21$) and in the NHS patients ($r=0.455$, $p=0.160$, $n=11$).

Motivation for seeking labiaplasty

The mean duration of the problem with the labia reported by the women seeking labiaplasty was 13.4 (s.d.=10.0) years, with responses ranging from 18 months to 37 years. The mean time taken to decide to make a labiaplasty request was 2.9 (s.d.=2.96) years. A total of 48 responses were provided for motivations for seeking labiaplasty and were coded. Table 6 summarizes the motivations and includes some

Table 3. Avoidance behaviours reported by women seeking labiaplasty ($n=55$) and controls ($n=70$)^a

Item	Labiaplasty: median (interquartile range)	Control: median (interquartile range)	Comparison	Effect size: Cohen's <i>d</i>
1. I avoid having a medical examination or treatment because of my labia	1.0 (0.5–1.5)	0 (0)	$U=1105.00$ $Z=-4.906$ $p<0.001$	0.98
2. I avoid going to public changing rooms because of my labia	1.0 (1.0–3.0)	0 (0)	$U=843.50$ $Z=-6.451$ $p<0.001$	1.42
3. I would avoid exercising in a gym or playing a sport because of my labia	0 (0.5–1.5)	0 (0)	$U=1305.00$ $Z=-4.478$ $p<0.001$	0.87
4. I avoid wearing a swimming costume because of my labia	0.5 (0–1.0)	0 (0)	$U=1032.00$ $Z=-5.375$ $p<0.001$	1.12
5. I avoid making love or being intimate because of my labia	2.0 (1.0–3.0)	0 (0)	$U=527.00$ $Z=-7.668$ $p<0.001$	1.91
6. I avoid certain types of clothes because of my labia, for example certain types of underwear or tight-fitting clothes	2.0 (1.0–3.0)	0 (0)	$U=556.00$ $Z=-7.306$ $p<0.001$	1.76
7. I avoid looking at pictures of naked women in magazines or on the Internet	1.0 (0–3.0)	0 (0)	$U=1114.00$ $Z=-4.705$ $p<0.001$	0.93
8. I avoid looking at my own labia	1.0 (1.0–3.0)	0 (0)	$U=915.50$ $Z=-5.544$ $p<0.001$	1.15

^a Additional avoidance behaviour reported by the labiaplasty group in an open-ended question included 'being naked in front of my own children'; 'having a bikini wax'; 'partner giving oral sex'; 'having any light on during sex'; 'being naked when I am on my own'; 'anything that might irritate the genital area (e.g. scented soaps or panty liners, certain types of condom)'; 'using tampons'; 'wearing any underwear'.

examples from the open question. For two discordant responses, consensus was reached following discussion. However, one response was not coded, as it did not fit coherently with any of the other reasons. ('I wanted to use some of my money to do something that might make a positive impact on my life.')

Previous cosmetic procedures

In the labiaplasty group, 37 (71.15%) women reported no cosmetic procedures in the past; 15 (28.9%) women reported having had at least one cosmetic procedure. Of these, five reported having two cosmetic procedures and one of these had had three procedures. Of these 15 women with a history of cosmetic procedures, nine (60%) were private patients, four (26.7%) were NHS patients, and two (13.3%) were community controls from Mind Search. The procedures included breast augmentation ($n=6$), dermafiller ($n=2$), Botox ($n=3$), rhinoplasty ($n=2$), laser resurfacing ($n=2$),

pinnaplasty/otoplasty ($n=1$), laser hair removal ($n=2$), scar treatment ($n=1$), mole removal ($n=1$) and cellulite treatment ($n=1$). None had had previous genital cosmetic surgery.

A total of three controls (5.88%) reported having had cosmetic procedures in the past, and 64 (94.12%) reported no cosmetic procedures in the past. Of these three participants, two also reported a second and third procedure. The procedures included breast augmentation ($n=2$), laser hair removal ($n=2$), dermabrasion ($n=1$) and liposuction ($n=1$). A Pearson's χ^2 test demonstrated a significant difference between the groups, with more cosmetic procedures in the labiaplasty group ($\chi^2=11.66$, degrees of freedom=1, $p=0.001$).

BDD

A total of 10 women in the labiaplasty group and four control women scored above the clinical cut-off on the

Table 4. Safety-seeking behaviours in women seeking labiaplasty ($n=55$) and controls ($n=70$)

Item	Labiaplasty: median (interquartile range)	Control: median (interquartile range)	Comparison	Effect size: Cohen's d
1. I check my labia in mirrors	1.0 (0.5–1.5)	1.0 (0.5–1.5)	$U=916.00$ $Z=-5.282$ $p<0.001$	1.09
2. I check my labia directly by looking at them without a mirror	2.0 (1.0–3.0)	0 (0–1.0)	$U=611.00$ $Z=-6.589$ $p<0.001$	1.50
3. I check my labia by taking photographs of myself	0 (0)	0 (0)	$U=1650.00$ $Z=-1.494$ $p=0.144$	0.28
4. I check my labia by feeling them with my finger(s)	2.0 (1.5–2.5)	1.0 (0.5–1.5)	$U=771.50$ $Z=-5.810$ $p<0.001$	1.25
5. I compare my labia to other women in magazines or on film or Internet	1.0 (0.5–1.5)	0 (0)	$U=542.00$ $Z=-7.317$ $p<0.001$	1.75
6. I compare my labia to medical illustrations and photographs	1.0 (0–2.0)	0 (0)	$U=627.00$ $Z=-6.970$ $p<0.001$	1.62
7. I compare my labia to other women directly (e.g. other women in changing rooms or seeing another women naked for some reason)	0.5 (0–1.0)	0 (0)	$U=981.00$ $Z=-5.848$ $p<0.001$	1.25
8. I change my posture to avoid my labia being seen at a certain angle	1.0 (0–2.0)	0 (0)	$U=830.50$ $Z=-6.235$ $p<0.001$	1.39
9. I hide my labia with something when I am with others (e.g. my hand)	1.0 (0–3.0)	0 (0)	$U=671.50$ $Z=-6.655$ $p<0.001$	1.54
10. I try to convince others about how unattractive my labia are	0 (0)	0 (0)	$U=1638.00$ $Z=-2.054$ $p=0.027$	0.39
11. I ask others to confirm the existence of my defective labia	0 (0)	0 (0)	$U=1650.00$ $Z=-2.116$ $p=0.030$	0.39
12. I seek reassurance about whether the size of my labia has got worse	0 (0)	0 (0)	$U=1669.00$ $Z=-1.626$ $p=0.169$	0.30
13. I seek reassurance about whether my labia are hidden	0 (0)	0 (0)	$U=1687.00$ $Z=-1.790$ $p=0.119$	0.32

COPS screening scale and were thus identified as having possible BDD. Clinical interviews were performed on these participants, except for two control participants. A total of 10 individuals (18.2%, 95% confidence interval $\pm 7.9\%$) in the labiaplasty group and no controls were subsequently diagnosed with BDD (seven women reported labia-only preoccupation and three had a preoccupation about their general appearance

that included the labia as their main feature of preoccupation). All of these participants had labia minora within the normal range according to the surgeon's measures.

Of the labiaplasty participants with BDD, six (60%) had specific concerns with their genitalia only, while the remaining four (40%) participants with BDD had concern with both their genitalia and other features

Table 5. Interference in life in women seeking labiaplasty ($n=55$) and controls ($n=70$)

Domain	Labiaplasty		Control		Comparison	Effect size: Cohen's d
	Median (IQR)	n	Median (IQR)	n		
Relationship in general	4 (1.5–6.5)	47	0 (0)	67	$U=414.00$ $Z=-4.536$ $p<0.001$	0.93
Sexual relationship	4 (2–6)	49	0 (0)	67	$U=245.0$ $Z=-8.255$ $p<0.001$	2.41
Social life	0 (0–2)	55	0 (0)	69	$U=1072.50$ $Z=-5.672$ $p<0.001$	1.17
Leisure	2 (0.5–3.5)	55	0 (0)	69	$U=586.50$ $Z=-6.986$ $p<0.001$	1.19
Discomfort	4 (2–6)	55	0 (0)	68	$U=353.50$ $Z=-8.152$ $p<0.001$	2.2
Hygiene	1 (0–5)	55	0 (0)	70	$U=1006.50$ $Z=-5.399$ $p<0.001$	1.09
Distress	5 (4–6)	55	0 (0)	69	$U=283.50$ $Z=-8.521$ $p<0.001$	2.41

IQR, Interquartile range.

of their body. Also, eight of the participants with BDD listed their labia as their first feature of concern, while the remaining two listed another body part as their first feature of concern and their labia as their second.

The individuals with BDD in the labiaplasty group were compared with the remainder of the labiaplasty group. On the COPS-L total score, those with BDD (median COPS-L score=56.5, IQR=47.62–65.38) scored higher than the remainder of the labiaplasty group (median COPS-L score=32.5, IQR=21.62–43.38) and this was statistically significant ($U=54.50$, $Z=-3.69$, $p<0.001$, effect size $r=-0.50$). It was possible to complete the BDD-YBOCS in six of the 10 participants with BDD, and scores obtained ranged from 21 to 40 (the individual scores were 21, 23, 26, 27, 30 and 40; median=26.5, IQR=21.5–31.5). Therefore five out of six were in the mild to moderate range on a standardized observer-rated measure of BDD symptomatology.

Discussion

We have conducted the first descriptive study of women seeking labiaplasty compared with a control group who had no significant difference in age, marital

status, education level, ethnicity and parity. There were no significant differences between women seeking labiaplasty and controls in terms of symptoms of anxiety or depression. The hypothesized differences between the labiaplasty group and the control group occurred in the specific measures of the GAS scale and on preoccupation, distress and interference with life. The labiaplasty group experienced reduced sexual satisfaction and quality of life related to body image compared with the control group. The labiaplasty group was also more likely to have previously undergone cosmetic procedures, suggesting that, for at least for some of the women, it was a life-style choice and they placed greater value on enhancing their appearance. Another reason is the increased likelihood of a diagnosis of BDD for the labiaplasty group than for the control group.

Participants' specific beliefs about their genitalia can be identified on the GAS scale and the distress item on the interference with life scale. These included feelings that the labia were asymmetrical, abnormal and 'too large'; or that they were causing discomfort, self-consciousness and/or an enormous sense of shame. It is possible that some of the distress occurs from a

Table 6. Reported motivation for seeking labiaplasty^a

Reason	n (%)	n (%) as sole reason	Examples
Aesthetic	31 (70.8)	10 (20.8)	'I do not like the asymmetrical appearance or the way the inner lips protrude from the outer lips' 'I am very comfortable with my body but the only thing that spoils it, is the fact that my labia are visible when standing naked' 'For a more feminine appearance. I think I currently look (and feel) untidy'
Functional (physical)	30 (62.5)	9 (18.75)	'My labia cause me a lot of irritation in my clothes and can become very sore. It can then lead to a great amount of discharge, which can be embarrassing' 'Constant irritation; external thrush; the labia are large enough to hold urine, so for a long time I thought I had urinary incontinence; discomfort when sitting' 'As the pain when I exercise and just walking can be awful'
Sexual	18 (37.5)	4 (8.33)	'Sex is uncomfortable and sometimes painful' 'It has caused several issues but perhaps the main reason is that I could never relax around any partner. I have always covered myself, and it has been my uttermost fear' 'I am fed up of being embarrassed about my body. It prevents me having a sex life, or means I jump into bed with men to not have to anticipate their reaction to my labia for weeks on end'
Combination	24 (50.0)	24 (50.0)	'To improve appearance and to increase confidence in body image and in sexual situations. To physically be more comfortable when wearing certain clothes and doing certain sports' (cosmetic, functional, sexual) 'Reduce discomfort when walking to feel "normal" and improve sex drive with husband as felt embarrassed with look and wouldn't let him go near the area' (cosmetic, functional, sexual) 'I don't feel confident in changing rooms and I believe it plays a part in restricting myself from getting close to boys. I also sometimes find it uncomfortable' (cosmetic, functional, sexual)

^aNote that percentages do not add up to 100%, as participants were able to provide reasons that fulfilled multiple categories, and these are included in the coding.

lack of awareness regarding the normal variation in labia size. This suggests a role for healthcare professionals, especially gynaecologists, in providing reassurance and psycho-education to some patients (Lloyd *et al.* 2005) but it is not known how effective this is in reducing genital dissatisfaction. Women may accept or know that their labia minora are normal in size but still feel they are abnormal or ugly. Reassurance may also be ineffective at reducing symptoms such as discomfort or pain and is ineffective for cosmetic interventions in people with BDD.

Women on the NHS had all been referred by their general practitioner who acted as a 'gate keeper' for more functional reasons for labiaplasty. Those who requested labiaplasty in the private sector were more likely to be self-referrals who reported more cosmetic reasons than in the NHS. Functional reasons might be related to social expectations to wear tight-fitting spandex clothing and underclothing that may

cause increased discomfort. If there is loose clothing and no obvious cause then the physical symptoms may fall into the category of being medically unexplained.

The checklist of avoidance and safety-seeking behaviours enabled the authors to identify a wide range of behaviours that occurred more frequently in the labiaplasty group. This information could be used clinically for a psychological intervention for some women seeking labiaplasty. Frequent comparisons with other women, pictures in the media, or checking with or without a mirror are all likely to increase preoccupation and distress (Veale *et al.* 1996b; Neziroglu *et al.* 2008). It may be possible to develop a psychological intervention that involves behavioural experiments and graded exposure to public situations (e.g. changing rooms). However it will be difficult to test out worries in sexual situations if the woman does not have a sexual partner.

It was a surprising finding that the duration of concern in women seeking labiaplasty had a mean of 10 years. It suggests that the onset of the problem including possible links to specific aversive memories (e.g. teasing) may be worth exploring.

The limitations of this study include the sample size and whether it is representative, the type of labiaplasty performed, and the measurements of the labia. A future larger sample size would allow the exploration of subgroup analyses (e.g. those with BDD *versus* those without BDD; those presenting privately *versus* those presenting in the NHS; those presenting for different motivations). However, the present study was designed to be exploratory. It is not known how representative the women in the present study are with respect to women seeking labiaplasty, as we do not have information on those who declined to participate. The sample was predominantly heterosexual, with no women in either group sexually oriented only or mainly to the same sex. Our question on cosmetic procedures was open and we did not systematically ask about all types of cosmetic procedures. Lastly, limiting the number of cosmetic procedures that could be coded to five may have influenced the result. Another limitation is that the sample was one of convenience, and the results may not be generalizable to other groups of women (e.g. in other practice settings or countries).

There were possible variations in the measurement of the size of the labia by site and it is not known if this reflects a true difference between the NHS and private patients or by surgeon. It would therefore be important to obtain inter-rater reliability scores before a study commences in future research.

It is also not possible to comment on the prevalence of BDD, as this was an opportunistic and small sample. However, the estimate is slightly higher than other studies on the prevalence of BDD in cosmetic settings (Crerand *et al.* 2010). However, the focus of the pre-occupation was specific to their genitalia. In other areas, the diagnosis of BDD may be associated with poor outcome (Phillips *et al.* 2001; Veale *et al.* 2003; Tignol *et al.* 2007; Crerand *et al.* 2010). Further research is required to repeat the measures after labiaplasty to determine the satisfaction and psychosexual outcomes of this procedure, particularly in those with BDD and to report on any complications.

Acknowledgements

We acknowledge support from the National Institute for Health Research (NIHR) Biomedical Research Centre for Mental Health at South London and Maudsley NHS Foundation Trust and the Institute of Psychiatry, King's College London. This paper

presents independent research funded by the NIHR. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Declaration of Interest

None.

References

- Bramwell R** (2002). Invisible labia: the representation of female external genitals in women's magazines. *Sexual and Relationship Therapy* 17, 187–190.
- Bramwell R, Morland C** (2009). Genital appearance satisfaction in women: the development of a questionnaire and exploration of correlates. *Journal of Reproductive and Infant Psychology* 27, 15–27.
- Bramwell R, Morland C, Garden AS** (2007). Expectations and experience of labial reduction: a qualitative study. *British Journal of Obstetrics and Gynaecology* 114, 1493–1499.
- Braun V** (2009). Female genital cosmetic surgery: a critical review of current knowledge and contemporary debates. *Journal of Women's Health* 19, 1393–1407.
- Cash TF, Fleming EC** (2002). The impact of body-image experiences: development of the Body Image Quality of Life Inventory. *International Journal of Eating Disorders* 31, 455–460.
- Crerand CE, Menard W, Phillips KA** (2010). Surgical and minimally invasive cosmetic procedures among persons with body dysmorphic disorder. *Annals of Plastic Surgery* 65, 11–16.
- Crouch NS, Deans R, Michala L, Liao LM, Creighton SM** (2011). Clinical characteristics of well women seeking labial reduction surgery: a prospective study. *British Journal of Obstetrics and Gynaecology* 118, 1507–1510.
- First MB, Spitzer RL, Gibbon M, Williams JBW** (1995). *Structured Clinical Interview for DSM-IV Axis I Disorders*. Biometric Research Department: New York.
- Health and Social Care Information Centre** (2012). Hospital Episode Statistics (www.hesonline.nhs.uk). Accessed 4 April 2012.
- Hrabosky JI, Cash TF, Veale D, Neziroglu F, Soll EA, Garner DM, Phillips KA** (2009). Multidimensional body image comparisons among patients with eating disorders, body dysmorphic disorder, and clinical controls: a multisite study. *Body Image* 6, 155–163.
- Lambrou C, Veale D, Wilson G** (2012). Appearance concerns comparisons among persons with body dysmorphic disorder and nonclinical controls with and without aesthetic training. *Body Image* 9, 86–92.
- Liao LM, Creighton SM** (2007). Requests for cosmetic genitoplasty: how should healthcare providers respond? *British Medical Journal* 334, 1090–1092.
- Liao LM, Michala L, Creighton SM** (2010). Labial surgery for well women: a review of the literature. *British Journal of Obstetrics and Gynaecology* 117, 20–25.

- Lloyd J, Crouch NS, Minto CL, Liao L-M, Creighton SM** (2005). Female genital appearance: 'normality' unfolds. *British Journal of Obstetrics and Gynaecology* **112**, 643–646.
- Mundt JC, Marks IM, Shear MK, Greist JH** (2002). The Work and Social Adjustment Scale: a simple measure of impairment in functioning. *British Journal of Psychiatry* **180**, 461–464.
- Neziroglu F, Khemlani-Patel S, Veale D** (2008). Social learning theory and cognitive behavioral models of body dysmorphic disorder. *Body Image* **5**, 28–38.
- Neziroglu F, Yaryura-Tobias JA** (1993). Body dysmorphic disorder: phenomenology and case descriptions. *Behavioural Psychotherapy* **21**, 27–36.
- Phillips KA, Diaz SF** (1997). Gender differences in body dysmorphic disorder. *Journal of Nervous and Mental Disease* **185**, 570–577.
- Phillips KA, Grant JD, Siniscalchi J, Albertini RS** (2001). Surgical and nonpsychiatric treatment of patients with body dysmorphic disorder. *Psychosomatics* **42**, 504–510.
- Phillips KA, Hollander E, Rasmussen SA, Aronowitz BR, Decaria C, Goodman WK** (1997). A severity rating scale for body dysmorphic disorder: development, reliability, and validity of a modified version of the Yale–Brown Obsessive Compulsive Scale. *Psychopharmacology Bulletin* **33**, 17–22.
- Phillips KA, McElroy SL, Keck PE Jr, Pope HG Jr, Hudson JI** (1993). Body dysmorphic disorder: 30 cases of imagined ugliness. *American Journal of Psychiatry* **150**, 302–308.
- Rogers RG, Kammerer-Doak D, Villarreal A, Coates K, Qualls C** (2001). A new instrument to measure sexual function in women with urinary incontinence or pelvic organ prolapse. *American Journal of Obstetrics and Gynecology* **184**, 552–558.
- Salkovskis PM** (1991). The importance of behaviour in the maintenance of anxiety and panic. *Behavioural Psychotherapy* **19**, 6–19.
- Tignol J, Biraben-Gotzamanis L, Martin-Guehl C, Grabot D, Aouizerate B** (2007). Body dysmorphic disorder and cosmetic surgery: evolution of 24 subjects with a minimal defect in appearance 5 years after their request for cosmetic surgery. *European Psychiatry* **22**, 520–524.
- Veale D, Boocock A, Gournay K, Dryden W, Shah F, Willson R, Walburn J** (1996a). Body dysmorphic disorder. A survey of fifty cases. *British Journal of Psychiatry* **169**, 196–201.
- Veale D, De Haro L, Lambrou C** (2003). Cosmetic rhinoplasty in body dysmorphic disorder. *British Journal of Plastic Surgery* **56**, 546–551.
- Veale D, Ellison N, Werner TG, Dodhia R, Serafty M, Clarke A** (2011). Development of a cosmetic procedure screening questionnaire (COPS) for body dysmorphic disorder. *Journal of Plastic, Reconstructive and Aesthetic Surgery* **65**, 530–532.
- Veale D, Eshkevari E, Ellison N, Cardozo L, Robinson D, Kavouni A** (2013). Validation of genital appearance satisfaction scale and the cosmetic procedure screening scale for women seeking labiaplasty. *Journal of Psychosomatic Obstetrics and Gynecology* **34**, 46–52.
- Veale D, Gournay K, Dryden W, Boocock A, Shah F, Willson R, Walburn J** (1996b). Body dysmorphic disorder: a cognitive behavioural model and pilot randomised controlled trial. *Behaviour Research and Therapy* **34**, 717–729.
- Zigmond A, Snaith RP** (1983). The Hospital Depression and Anxiety Scale. *Acta Psychiatrica Scandinavica* **67**, 361–370.