

Maternal responsiveness and sensitivity reconsidered: Some is more

MARC H. BORNSTEIN AND NANMATHI MANIAN

Eunice Kennedy Shriver National Institute of Child Health and Human Development

Abstract

Is it always or necessarily the case that common and important parenting practices are better, insofar as they occur more often, or worse, because they occur less often? Perhaps, less is more, or some is more. To address this question, we studied mothers' microcoded contingent responsiveness to their infants ($M = 5.4$ months, $SD = 0.2$) in relation to independent global judgments of the same mothers' parenting sensitivity. In a community sample of 335 European American dyads, videorecorded infant and maternal behaviors were timed microanalytically throughout an extended home observation; separately and independently, global maternal sensitivity was rated macroanalytically. Sequential analysis and spline regression showed that, as maternal contingent responsiveness increased, judged maternal sensitivity increased to significance on the contingency continuum, after which mothers who were even more contingent were judged less sensitive. Just significant levels of maternal responsiveness are deemed optimally sensitive. Implications of these findings for typical and atypical parenting, child development, and intervention science are discussed.

The perfect is the enemy of the good. —Voltaire

Parenting practices are consensually believed to contribute to the course and outcome of child development. For example, parental practices such as involvement, devotion, investment, and responsiveness are commonly assumed to be good things for parents and children. Here we asked how the performance of a commonly understood important and potent parenting practice is judged with respect to its frequency of occurrence. Perhaps *more is better*, or alternatively, *less is more* or *some is more*. Just as a parent might not be sufficiently involved, devoted, invested, or sensitive, a parent might be too involved, too devoted, too invested, or too responsive.

We address this basic theoretical and practical question related to individual variation in parenting infants. Specifically, we asked, *How are global judgments of maternal parenting sensitivity assigned to levels of specific maternal contingent responsiveness to specific infant behaviors?* In a large community sample, we counted and timed infant and maternal behaviors microanalytically throughout an extended video-recorded observation, and we separately and independently rated the same mothers macroanalytically for their global parenting sensitivity. For infants, exploration, facial affect, and vocalization (the behaviors we studied) serve as principal ex-

pressions of state of arousal and affect, and cognitive, communicative, emotional, and social functioning. They are frequent and prominent behaviors in the first half-year of life (the age we studied), and they are behaviors that mothers monitor closely and to which they respond. Maternal responsiveness (the reactions to young children their mothers display in the context of everyday dyadic interactions) has often been singled out as especially significant in child development. For mothers, therefore, we microanalytically counted and timed their most frequent and prominent responses, engaging their infants one-on-one socially, encouraging their infants to attend to the environment, and speaking to their infants. To evaluate the global quality of maternal parenting, we separately and independently rated maternal sensitivity using a standardized, validated, and reliable macroanalytic scale. Associating maternal responsiveness with maternal sensitivity yielded novel insights into parenting practices, perceptions that may have telling consequences in children's normal and atypical development because they expose basic processes underlying adaptation and maladaptation (Cicchetti & Toth, 2009). Identifying and exploring prominent processes that motivate developmental trajectories constitute basic goals of developmental science (Cicchetti & Pogge-Hesse, 1982).

Responsive Parenting

Maternal responsiveness has attracted the attention of developmental scientists for many reasons. Following Drillien (1957), Bowlby (1969) asserted that one of the principal antecedents of secure attachment in children was the attachment

We thank A. Bradley and A. Dovidio. This research was supported by the Intramural Research Program of the National Institutes of Health, Eunice Kennedy Shriver National Institute of Child Health and Human Development.

Address correspondence and reprint requests to: Marc H. Bornstein, Child and Family Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, Suite 8030, 6705 Rockledge Drive, Bethesda, MD 20892-7971; E-mail: Marc_H_Bornstein@nih.gov.

figure's responsiveness to child distress. Ainsworth, Blehar, Waters, and Wall (1978, p. 152) later concluded that "the most important aspect of maternal behavior commonly associated with the security–anxiety dimension of infant attachment is manifested in different specific ways in different situations, but in each it emerges as sensitive responsiveness to infant signals and communications." Responsiveness reflects a key parent element of recurring and meaningful sequences in everyday exchanges between child and parent that involve child action and parent reaction that generalizes across caregiving contexts (e.g., laboratory and home; Lohaus, Keller, Ball, Elben, & Völker, 2001). Responsive parenting fosters a broad array of highly valued developmental outcomes in children, including awareness of the caregiver's availability and reliability, thus promoting a sense of security and trust, behavioral independence, social facility, symbolic competence, verbal ability, and intellectual achievement (Ainsworth, Bell, & Stayton, 1974; Bornstein, Tamis-LeMonda, Hahn, & Haynes, 2008; Bus & van IJzendoorn, 1992; De Wolff & van IJzendoorn, 1997; MacDonald, 1992).

Despite its consensual familiarity, centrality, robustness, and predictive validity, certain assumptions about maternal responsiveness (and parenting more generally) have gone untested. One such assumption concerns perceived caregiving sensitivity of different levels of responsiveness.

Undercontingency and Overcontingency

Parents vary in responding to their young children. In Western cultures, mothers reportedly respond to only approximately 30%–50% of their infants' babbling (e.g., Gros-Louis, West, Goldstein, & King, 2006) and 50%–75% of their infants' expressions of distress (van IJzendoorn & Hubbard, 2000). What meaning should we ascribe to amount of parental responsiveness? If meaningfulness is judged in predictive validity terms (e.g., for children's development), undercontingency as well as overcontingency appear to be less than optimal parenting strategies. A number of macro- and microanalytic studies converge on an "optimum midrange model" of parenting contingency, where both higher and lower degrees of contingency are deemed problematic.

Undercontingency

From an early age, infants are sensitive to a lack of contingency in normal social interactions with their parents, as evidenced by their vivid (and unhappy) response when mothers become still-faced (Goldstein, Shwade, & Bornstein, 2009; Tronick, Als, Adamson, Wise, & Brazelton, 1978) or when synchronous interactions with mothers are replaced via video with noncontingent actions (Bigelow, MacLean, & MacDonald, 1996). In neither case is the mother attempting to elicit negative affect, but rather merely not responding as she normally would. That even young infants react negatively to noncontingent maternal behavior attests to the importance of contingency rules by which infants and mothers normally interact (Moore et al., 2009). Maternal failures to respond

to infant signals result in poorly timed, mutually unsatisfying interactions and ultimately undesirable child outcomes: maternal unresponsiveness at 3 and 9 months predicts insecure attachment at 12 months (Isabella & Belsky, 1991), aggressive and disruptive behavior at 3 years (Shaw, Keenan, & Vondra, 1994), and externalizing problems at 10 years (Wakschlag & Hans, 1999).

Overcontingency

In the general context of the foregoing literature, it would seem that more consistent responsiveness would be primed to foster the development of more positive outcomes in children (Van Egeren, Barratt, & Roach, 2001). Some have theorized or posited that more contingent parents create more optimal developmental environments for children (Chapple, 1970; Cohn & Elmore, 1988; Dunham & Dunham, 1994; Tarabulsky, Tessier, & Kappas, 1996; Van Egeren et al., 2001); for example, on learning principles, consistent contingency increases the predictability of interaction. Hsu and Lavelli (2005) reported that more responsive Italian and Canadian mothers had more verbal children and, although specific associations emerged in each culture, each association turned on presumed linear relations between maternal responsiveness and child language outcomes. However, high levels of contingency may be intrusive or inappropriate and are associated with communicative stress and hypervigilance (Beebe et al., 2008). Malatesta, Culver, Tesman, and Shepard (1989, p. 29) concluded "that even if maternal responses were in some sense 'appropriate' (concern to sad faces, smiling to happy faces), the sheer amount of stimulating activity might exceed the infants' arousal tolerance." Similarly, van den Boom (1994) contended that infant acquisition of self-reliance might develop under a mother who is responsive but who is not continuously attentive to every infant signal; a parent being totally responsive leaves little room for the development of child self-sufficiency. Overcontingent parenting is not focused on being appropriately responsive to a child, but rather on indiscriminately responding to all of the child's needs and cries for attention.

"Goldilocks"

Thus, the literature suggests that undercontingency and overcontingency to young infants may both be detrimental to development. Therefore, it is possible to find in the literature behavioral indicators that point to a curvilinear relation between maternal responsiveness (or, for that matter, stimulation or involvement) and positive child outcomes. Belsky, Rovine, and Taylor (1984) reported that an intermediate amount of reciprocal interaction at 9 months was associated with secure attachment in infants at 12 to 13 months, whereas insecure-resistant infants had engaged in less reciprocal interaction and insecure-avoidant infants had engaged in more reciprocal interactions. Malatesta et al. (1989); Lewis and Feiring (1989); Isabella, Belsky, and von Eye (1989); Leyendecker,

Lamb, Fracasso, Scholmerich, and Larson (1997); and Beebe and colleagues (Beebe et al., 2000, 2010; Jaffe, Beebe, Feldstein, Crown, & Jasnow, 2001) subsequently reported similar findings. However, these studies all focused on relations between levels of maternal responsiveness and later child attachment classification.

This Study

The focus of the extant literature on the differential predictive validity of levels of maternal responsiveness (or stimulation) in relation to child (usually attachment) outcome is valuable and instructive. That said, this attachment research emphasizes the interpersonal over the intrapersonal, and it also oversteps several key intermediary questions that otherwise address process and mechanism, and so refine and explain the observed predictions. Thus, the present study addresses the following questions: How does maternal contingency vary by different responses and to different infant behaviors? How can levels of maternal contingency be quantified in ways that go beyond the usual qualitative attributions of, for example, “low” versus “midrange” versus “high”? What quantitative levels of maternal contingency are judged to be optimally sensitive? With the preceding review and these questions in mind, we wished to discern how different maternal responses to different infant behaviors distribute themselves, to bring a quantitative perspective to bear on the distribution of maternal responsiveness and to determine how levels of mothers’ contingency to their infants’ behaviors in situations of natural interactions might be evaluated in terms of mothers’ perceived global parenting sensitivity. Guided by the existing literature, we hypothesized that maternal over- and undercontingency would be rated as generally less sensitive, whereas levels surrounding significant maternal contingency would be rated as optimally sensitive. In this study, we parsed two key modalities of interaction (gaze and vocalization) in infant–mother dyads. The literature on maternal responsiveness has included responsiveness to both positive and negative infant behaviors; however, the present study focuses on responsiveness to positive infant behaviors.

Maternal Sensitivity

To test the study’s main hypothesis, we evaluated the sensitivity of maternal responsiveness. To do so, we separately and independently assessed global dimensions of the same mothers’ emotional relationships with their infants using a validated and reliable instrument, the Emotional Availability Scales (EA Scales; Biringen, Robinson, & Emde, 1998). The EA Sensitivity Scale assesses overall acceptance, appropriateness, flexibility, affect regulation, and variety and creativity of behavior displayed toward the infant. To appraise the generalizability of our findings, we also coded a second maternal EA Scale, Structuring. Maternal structuring captures the extent to which the mother successfully organizes or scaffolds in a way that is received by the child and that still offers

opportunities for the child to exercise his or her autonomy. The results for sensitivity and structuring ($r = .84$) were the same. In the balance of this paper, when we refer to sensitivity, our results apply equally to structuring (more information is available from the authors).

The traditional approach to analyzing infant–mother interaction has applied correlation (e.g., Bornstein & Tamis-LeMonda, 1990). This approach to characterizing interaction tells us about the relative standing of dyads in a group on a pair of variables. Although informative, correlation is relatively mute about the nature and mechanics of interaction of individual dyads, and it gains no purchase on causality. It is sometimes assumed that behaviors that are correlated are also contingent, but in actuality, correlation is not necessarily related to interactional contingency. Exclusive reliance on tests of simple association therefore sells short an understanding of temporal or causal relations that are believed to structure infant–mother interactions and predict child outcomes. Our design and analyses were also geared to redress this shortcoming. We supplemented correlations with microanalytic measurement that is requisite to deconstruct contingencies in interaction over time and then applied sequential analysis to investigate which behaviors tend to lead and lag temporally when infants and mothers interact. Sequential analysis lays bare the underlying dynamic contingent structure of interactions (Bornstein, Toda, Azuma, Tamis-LeMonda, & Ogino, 1990; Damast, Tamis-LeMonda, & Bornstein, 1996). One supplementary hypothesis of this study was that specific infant and mother behaviors would covary, and a second supplementary hypothesis was that mother behaviors would be contingent on infant behaviors.

Method

Participants

The sample consisted of 335 European American mother–infant dyads (154 mothers and daughters, and 181 mothers and sons). Mothers were recruited from hospital birth notifications, mass mailings, and newspaper advertisements and were primiparous with healthy infants. Infants ranged from 4.7 to 6.5 months ($M = 5.4$, $SD = 0.2$) at birth, and 99.10% were term (M birth weight = 3510.19 g, $SD = 489.04$). Mothers’ ages ranged from 13.91 to 42.48 years ($M = 28.27$, $SD = 6.13$). Of participating mothers, 11.6% had not completed high school, 12.8% had completed high school, 21.2% had partial college, 29.9% had completed college, and 24.5% had completed university graduate programs. Family socioeconomic status (SES; Hollingshead, 1975; see also Bornstein, Hahn, Suwalsky, & Haynes, 2003) varied across nearly the full range of social class, 14–66, with a mean of 48.37 ($SD = 13.80$). Thus, ours is a socioeconomically heterogeneous (not homogenous middle-class) community sample in terms of maternal education and family SES, but we recruited an ethnically homogenous European American sample because they are currently the majority cultural group in the United States (~75% of the population of the United States self-identifies as European

American in descent; Humes, Jones, & Ramirez, 2011; Tilton-Weaver & Kakiyama, 2008; US Census Bureau, 2008) and because parenting processes and child behavior alike are known to vary with ethnicity (e.g., Graham, 1992; Parke, 2000; Tomlinson & Swartz, 2003). An ethnically homogeneous community sample constitutes a first step in understanding the matrix of associations surrounding infant behavior, maternal contingent interactions, and maternal sensitivity that logically antecedes embarking on more complex studies and analyses with ethnically diverse samples. This study therefore intentionally avoids ethnicity confounds that have plagued the parenting literature and would cloud our findings.

Procedure

Each mother–infant dyad was visited at home by a single filmer to make an hour-long audiovisual record of the dyad's naturalistic behavior. After a conventional period of acclimation (McCune-Nicolich & Fenson, 1984), filming commenced. The filmer resisted talking to the mother or making eye contact or otherwise interacting with the infant or mother during filming. Dyads were recorded at a time when the infant was awake and alert ($M = 99.66\%$ of the session), and mothers were in view of their infant ($M = 94.06\%$ of the session) and solely responsible for the baby. The filmer explained that she was interested in the typical activities of young infants, and the mother was asked to go about her normal routine and to disregard the filmer insofar as possible. In general, we attempted to remain faithful to a principle of ecological validity by focusing on naturalistic interactions between mothers and infants in their own homes. By allowing infants and mothers to be observed in the surroundings most familiar to them and in which they are most comfortable, home study yields behavioral data that are as natural, uninhibited, and valid as possible. Observations encompassed a variety of routine mother–infant daily activities, such as feeding, diapering, bathing, and playing. The filmer focused on the infant, but included the infant and mother in the same frame

when they were proximal, panned the environment with any changes in physical setting, indicated if mother left the room, and briefly captured any object or person on which the infant fixated (Vibbert & Bornstein, 1989).

Data coding and scoring

All records were coded independently by two separate coding teams, one team of coders for microlevel coding of individual behaviors and another team of coders for macrolevel coding of sensitivity. The two teams of coders were unaware of each other's coding, and coding was fully separate and independent. All coders also were blind to the hypotheses of the study (Bornstein, Azuma, Tamis-LeMonda, & Oginio, 1990).

Microcoding of infant and maternal behaviors. Mutually exclusive and exhaustive independent coding schemes were brought to bear for different infant and maternal behaviors. Coders made separate passes through the records for infant behaviors (look at mother, look at object, nondistress vocalization, and smile) and maternal behaviors (encourage attention to mother, encourage attention to object, and speech to the infant). The first 50 min of the record were used to accommodate occasional momentary interruptions and early endings of recordings. For all cases, the total length of the record that could be coded was more than 48 min. (If codable time totaled less than 50 min, data for the case were prorated. There were two cases with codable time less than 50 min, and for both cases the total duration was ≥ 2947 s.) Coders were trained to reliability on consensus codings, and between 18% and 26% of the sample was coded independently by alternate coders (or coders other than those who were trained on reliability coding). None of these coders participated in EA Scale coding, and they were not aware of the EA Scale scores. Table 1 and Table 2 provide operational definitions and reliability indices for all behavior codes.

Onsets and offsets of maternal and infant behaviors were coded to the nearest 0.1 s, resulting in timed-sequential data

Table 1. Operational definitions and reliability indices of infant behaviors

| Behavior: Operational Definition | κ |
|---|-------------------|
| <i>Look at mother:</i> Infant looks at the mother's face or head. Focused fixation must be evident. An active behavior component often accompanies clear and focused fixation (e.g., brightening of the face, widening of the eyes, stilling, increased motor excitement, positive vocalizations, or reaching). A change in fixation is coded after the infant has looked away from the target for 1 s. | 0.67 |
| <i>Look at object:</i> Infant looks at any discrete object or maternal body part other than the face. Focused fixation must be evident. An active behavior component often accompanies clear and focused fixation (e.g., brightening of the face, widening of the eyes, stilling, increased motor excitement, positive vocalizations, or reaching). A change in fixation is coded after the infant has looked away from the target for 1 s. | 0.71 |
| <i>Nondistress vocalization:</i> Infant emits any positively or neutrally toned vocalization that is clearly audible. Included are babbling, cooing, laughing, vocal play, shrieking, and sighs or grunts not indicative of distress. Vocalizations of all durations are coded. | 0.70 |
| <i>Smile:</i> The infant emits a clear, unambiguous smile. The corners of the infant's mouth are extended outward and upward, the eyes "brighten" and are focused, and the eyebrows are relaxed or raised. | 0.48 ^a |

^aInfant smiling was an infrequent (and very brief) event, accounting for <5% of the events coded in the mode for infant facial expression.

Table 2. Operational definitions and reliability indices of maternal behaviors

| Behavior: Operational Definition | κ |
|--|----------|
| <i>Encourage attention to herself:</i> Mother attempts to draw the infant into face-to-face social interaction with herself. Physical attempts include intentionally moving her face toward the infant or moving the infant toward her face. Verbal attempts include making very specific comments about herself that are clearly designed to capture the infant's interest. Pauses of 2 s or longer are coded as terminations of an ongoing behavior. | 0.71 |
| <i>Encourage attention to object:</i> Mother physically moves the infant or an object so that the infant can see or touch it, or the mother verbally refers to an object or an object-related event or activity. Pauses of 2 s or longer are coded as terminations of an ongoing behavior. | 0.73 |
| <i>Speech to the infant:</i> Mother directs words and speechlike sounds to the infant. Included are syllable sounds, parts of words, single words, conversations, and singing. Changes and pauses in vocalization lasting less than 1 s are not recorded. | 0.69 |

(Bakeman, 2004; Bakeman, Deckner, & Quera, 2005). The data were then formatted for sequential analysis using the Sequential Data Interchange Standard. For each dyad, codes for infant behaviors were followed by codes for maternal behaviors, along with the onset time for each. We used the Generalized Sequential Quierer program (version 4.1.2; Bakeman & Quera, 2004; <http://www.gsu.edu/~psyab/sg.htm>) to conduct simple and sequential statistical analyses. We calculated an odds ratio, which is a descriptive measure of effect size (Bakeman et al., 2005), of lead-lag interactions for each dyad for 12 behavioral sequences:

1. infant look at mother—mother encourage attention to herself;
2. infant look at mother—mother encourage attention to object;
3. infant look at mother—mother speech to the infant;
4. infant look at object—mother encourage attention to herself;
5. infant look at object—mother encourage attention to object;
6. infant look at object—mother speech to the infant;
7. infant nondistress vocalization—mother encourage attention to herself;
8. infant nondistress vocalization—mother encourage attention to object;
9. infant nondistress vocalization—mother speech to the infant;
10. infant smile—mother encourage attention to herself;
11. infant smile—mother encourage attention to object; and
12. infant smile—mother speech to the infant.

Coding of maternal and infant behaviors was mutually exclusive, except that some behaviors were coded both in mother speech to the infant and mother encouraging attention to object (e.g., “Baby, look at this toy”), and between mother speech to the infant and mother encouraging attention to herself (e.g., “Baby, look at mommy”). However, different aspects of behavior were coded in mother encouraging attention (to object or to herself) and mother speech. Maternal encouragement describes a goal-directed form of maternal behavior that can include many different component behaviors (e.g.,

talking, looking, facial expression, or offering objects), whereas coding maternal speech exclusively comprised sounds mothers emitted to the infant. Empirically, mother speech to the infant and mother encouraging attention to object, and mother speech to the infant and mother encouraging attention to herself, shared little common variance, 3.6% and 7.8%, respectively.

In addition, the contingency analyses we planned focused on a particular maternal response to a particular infant behavior. For example, the mother may respond in different ways to an infant vocalization. We analyzed the contingency of each maternal response to each infant behavior.

Contingency was operationally defined in terms of lags of time units (Bakeman & Gnisci, 2005), that is, the length of time after a given onset or offset that a target response could occur and constitute a contingent behavior rather than a spontaneous or isolated behavior. Based on an extensive review of existing parametric research (see Beebe et al., 2010; Cote, Bornstein, Haynes, & Bakeman, 2008; Gratier, 2003; Van Egeren et al., 2001) and features of our own data set (durations, gaps between onsets, and visual plots of infant–mother behaviors), we determined that a 3-s time period best captured contingencies for these mother behaviors in response to these infant behaviors in naturalistic settings. That is, the onset of a target maternal behavior had to occur within 3 s of the onset of a given infant behavior for the target to be considered contingent on the given. Because we were interested in contingent turn taking in infant–mother speech, the only exception was that maternal speech to the infant had to occur within 3 s of the offset of infant nondistress vocalization to be contingent.

Time units were tallied separately for each dyad in 2×2 tables for each behavioral sequence (1–12 above), and an odds ratio was computed for each table (see Bakeman et al., 2005, p. 415). Odds ratios (*ORs*) of >1 indicate that bouts of the target behavior were more likely to begin within 3 s of the onset of the given behavior than at other times, whereas *ORs* of 0–1 indicate less likelihood. If fewer than five occurrences of the given behavior were counted during the entire observation, we regarded the value of the odds ratios as missing for that dyad because there was not a sufficient sample of the behavior from that dyad to draw conclusions about behav-

ioral contingency. With the exception of infant smile–mother speech to the infant, data for fewer than 3% of dyads were insufficient for any given analysis.

Macrocoding of maternal sensitivity. Sensitivity for each mother was separately and independently evaluated by a different team of coders from the same recorded observations using the Sensitivity Scale of the Emotional Availability Scales: Infancy to Early Childhood Version (EA Scales, 3rd ed.; Biringen et al., 1998). The EA Scales were developed as a global assessment of parenting quality based on observations and ratings of parent–child interaction (Bornstein, Suwalsky, & Breakstone, 2012). The EA Sensitivity Scale has been used with a wide variety of populations and applied successfully across a range of settings (Biringen & Easterbrooks, 2012; Robinson, Little, & Biringen, 1993). The general construct of sensitivity, initially developed by Ainsworth (1967) as normative and presumably universal, has proved to be a robust predictor of myriad positive child characteristics, including attachment, social responsibility, self-confidence, self-esteem, and behavioral adjustment in socioemotional development as well as language and play and school readiness in cognitive development (Isley, O’Neil, & Parke, 1996; Pettit & Bates, 1989; van IJzendoorn, Dijkstra, & Bus, 1995). Maternal sensitivity was coded on a 1 (*highly insensitive*) to 9 (*highly sensitive*) scale in half points. All coders were trained on the EA Scales to obtain satisfactory reliability with one of the authors of the EA Scales and with one another. Inter-coder reliability was assessed using average absolute agreement intraclass correlation coefficients (ICCs; as recommended by Shrout & Fleiss, 1979) in a two-way random-effects model. For reliability, 23% of randomly selected records were double coded (ICC for sensitivity = 0.92).

Results

Preliminary analyses

Prior to formal analysis, univariate distributions of infant and maternal behaviors and maternal sensitivity were evaluated for normality and outliers. Bivariate plots were screened between the infant/maternal behaviors and sensitivity ratings; residuals were examined for influential cases using scatter plots, standardized residuals, studentized deleted residuals, and Cook *D* statistics. The frequencies of maternal and infant behaviors were normally distributed, with the exception of infant smile, which required square-root transformation. Because the odds ratio has a lower limit of 0 and no upper limit, it is known to have a skewed distribution. Hence, we used a square root transformation and excluded outliers (maximum of four per analysis) to normalize the distribution for each pair of infant–mother behaviors. Sensitivity was raised to the second power to approximate normality and reduce outliers. Transformed variables were used in analyses; for clarity, untransformed data are presented as descriptive statistics.

No gender differences emerged for any infant or maternal behaviors, infant–mother contingencies, or EA Scales; therefore, the data are reported for girls and boys combined. We examined correlations between the study variables and maternal age and family SES. Maternal age was correlated with mother speech, $r(333) = .42, p < .01$, and sensitivity, $r(333) = .35, p < .01$, and with infant looks at objects, $r(333) = -.12, p < .05$. SES was correlated with mother encouraging attention to herself, $r(333) = .14, p < .05$, speech, $r(333) = .35, p < .01$, and sensitivity, $r(333) = .25, p < .01$, and with infant looks at objects, $r(333) = -.16, p < .05$. Because sensitivity was strongly correlated with maternal age, and because maternal age and SES were highly correlated, $r(333) = .70, p < .001$, sensitivity was regressed on maternal age and residualized scores were used in analyses.

Descriptive statistics

Frequencies of infant and mother behaviors are reported in Table 3. Mother speech had the highest frequency; infant smile had the lowest. For sensitivity, mothers were rated high on average but showed nearly the full range (2.0 to 8.5), suggesting that in a prolonged observation it was possible to discern wide individual differences among mothers of infants as young as 5 months. These findings compare favorably with ratings of other studies (Kogan & Carter, 1996).

Intercorrelations between infant and mother behaviors and sensitivity

Intercorrelations between infant and mother behaviors were calculated to examine whether mothers or infants who engaged in a given behavior more frequently had partners who engaged in a complementary behavior more frequently (Table 4). Mothers who encouraged attention to themselves and to ob-

Table 3. Descriptive statistics for infant and mother behaviors and Emotional Availability (EA) Scale ($N = 335$)

| Infant Behaviors | Frequency |
|--------------------------------|-----------------|
| Look at mother | 51.73 (27.45) |
| Look at object | 134.82 (44.19) |
| Nondistress vocalization | 146.41 (76.14) |
| Smile | 11.78 (12.41) |
| Mother Behaviors | |
| Encourage attention to herself | 22.28 (15.81) |
| Encourage attention to object | 32.21 (21.37) |
| Speech to the infant | 309.40 (167.31) |
| EA Scale | Rating |
| Sensitivity | 6.18 (1.38) |

Note: All values are means (standard deviations).

Table 4. Correlations between infant and mother behaviors ($N = 335$)

| | Mother Encourages Attention to Herself | Mother Encourages Attention to Object | Mother Speech to Infant |
|---------------------------------|--|---------------------------------------|-------------------------|
| Infant looks at mother | .40*** | .11* | .27*** |
| Infant looks at object | -.01 | .23*** | -.01 |
| Infant nondistress vocalization | .08 | .05 | .08 |
| Infant smiles | .24*** | .17** | .03 |
| EA Scale sensitivity | .14** | .00 | .27** |

* $p < .05$. ** $p < .01$. *** $p < .001$.

jects more frequently had infants who looked at them and smiled more frequently. Mothers who spoke more frequently had infants who looked at them more frequently. Mothers who encouraged infant attention to objects more frequently had infants who looked at objects more frequently. In addition, mothers who encouraged attention to themselves and spoke more to their infants were rated as more sensitive (Table 4).

Contingency of mother behaviors

Student t tests were calculated to determine whether pairs of behaviors were contingent (i.e., whether odds ratios differed significantly from 1; Wickens, 1993). Eleven of 12 pairs of mothers' contingent behaviors to their infants were significant (Table 5). For example, it was significantly likely that mothers responded within 3 s of the onset of their infants looking at them by encouraging their infants to attend to the mothers themselves. Only one odds ratio, infant vocalize—mother encourage attention to object, was not significant; it

was as likely that mothers responded by encouraging attention as by any other behavior within 3 s of the offset of infant vocalizing. Only the odds ratio for infant look at object—mother encourage attention to herself was < 1 and significant; it was significantly unlikely that mothers responded within 3 s of the onset of their infants looking at an object by encouraging attention to themselves.

Contingency in infant–mother behaviors and maternal sensitivity

Only two correlations between (transformed) odds ratios for the 12 pairs of infant–mother behaviors and (transformed and residualized) ratings of maternal sensitivity were significant (range $r_s = -.22$ to $.15$). Mothers who were more likely to contingently speak to infants in response to their infants' looking at them were rated as less sensitive, $r(330) = -.22$, $p < .001$. Mothers who were more likely to encourage their infants' attention to objects contingent on infants' looking at them were rated as more sensitive, $r(330) = .15$, $p < .01$. Only one other study has compared a microanalytic approach to contingency with a macroanalytic measure of maternal sensitivity. Keller, Lohaus, Völker, Cappenberg, and Chasiotis (1999) analyzed the relation between contingency (assessed in a microanalytical, exclusively face-to-face procedure) and sensitivity (rated on the Ainsworth scale) in German middle-class mothers interacting with their 3-month-olds in face-to-face exchange, play with objects, and situations with an extradyadic orientation. As here, they found no zero-order associations between maternal sensitivity and contingency indexes (see also Lohaus et al., 2001).

However, zero-order correlations mask potential nonlinear associations between maternal responsiveness to infants and maternal sensitivity. Nonlinear relations cannot be detected, represented, or studied within the confines of zero-order correlations or linear regressions. A nonsignificant correlation may belie curvilinear or other nonlinear relations (such as

Table 5. Odds ratios (OR) for infant–mother behaviors

| Given Infant Behavior | Time Window for Given | Target Mother Response | N | Mean OR (95% CI) | t (df) |
|-----------------------------|-----------------------|---------------------------------|------------------|------------------|---------------------|
| 1. Look at mother | Onset + 3 s | Encourages attention to herself | 326 | 8.52 (7.48–9.56) | $t(325) = 19.89***$ |
| 2. Look at mother | Onset + 3 s | Encourages attention to object | 332 | 1.32 (1.13–1.52) | $t(331) = 3.20**$ |
| 3. Look at mother | Onset + 3 s | Speech to the infant | 332 | 2.10 (1.97–2.23) | $t(331) = 19.61***$ |
| 4. Look at object | Onset + 3 s | Encourages attention to herself | 331 | 0.79 (0.69–0.99) | $t(330) = 3.99**$ |
| 5. Look at object | Onset + 3 s | Encourages attention to object | 333 | 3.08 (2.75–3.41) | $t(332) = 15.52***$ |
| 6. Look at object | Onset + 3 s | Speech to the infant | 331 | 1.07 (1.03–1.11) | $t(330) = 2.71**$ |
| 7. Nondistress vocalization | Offset + 3 s | Encourages attention to herself | 331 | 1.82 (1.58–2.04) | $t(330) = 7.03***$ |
| 8. Nondistress vocalization | Offset + 3 s | Encourages attention to object | 335 | 1.03 (0.99–1.04) | $t(334) = 0.55, ns$ |
| 9. Nondistress vocalization | Offset + 3 s | Speech to the infant | 333 | 1.35 (1.30–1.40) | $t(332) = 15.41***$ |
| 10. Smile | Onset + 3 s | Encourages attention to herself | 230 ^a | 4.50 (3.56–5.44) | $t(229) = 7.31***$ |
| 11. Smile | Onset + 3 s | Encourages attention to object | 232 ^a | 1.49 (1.05–1.93) | $t(231) = 2.19*$ |
| 12. Smile | Onset + 3 s | Speech to the infant | 230 ^a | 1.68 (1.54–1.82) | $t(229) = 6.87***$ |

^aBecause infant smile was infrequent, 103 dyads had missing odds ratios.

* $p < .05$. ** $p < .01$. *** $p < .001$.

Table 6. Results of spline regressions for infant–mother behaviors and sensitivity

| Given Infant Behavior | Target Mother Response | N | Estimation of Knot | | Estimation of First Slope | | Estimation of Second Slope | |
|-----------------------------|---------------------------------|-----|--------------------|---------------------------|---------------------------|---------------------------|----------------------------|---------------------------|
| | | | Knot | F | B ₁ (95% CI) | F | B ₂ (95% CI) | F |
| 1. Look at mother | Encourages attention to herself | 326 | 4.14 | $F(2, 323) = 3.03^*$ | 1.14 (−0.38, 2.65) | $F(1, 323) = 2.19$ | −5.02 (−9.04, −0.99) | $F(1, 323) = 6.01^*$ |
| 2. Look at mother | Encourages attention to object | 332 | 1.56 | $F(2, 329) = 6.30^{**}$ | 5.00 (2.23, 7.77) | $F(1, 329) = 12.60^{***}$ | −9.18 (−16.92, −1.45) | $F(1, 329) = 5.45^*$ |
| 3. Look at mother | Speech to the infant | 332 | 1.10 | $F(2, 329) = 12.51^{***}$ | 14.39 (−1.90, 30.68) | $F(1, 329) = 3.02$ | −26.15 (−44.02, −8.29) | $F(1, 329) = 8.30^{**}$ |
| 4. Look at object | Encourages attention to herself | 331 | 1.32 | $F(2, 328) = 1.15$ | 2.17 (−1.36, 5.70) | $F(1, 328) = 1.46$ | −9.31 (−22.28, 3.65) | $F(1, 328) = 1.99$ |
| 5. Look at object | Encourages attention to object | 333 | 1.88 | $F(2, 330) = 0.93$ | −2.61 (−6.57, 1.35) | $F(1, 330) = 1.68$ | 2.78 (−3.84, 9.41) | $F(1, 330) = 0.68$ |
| 6. Look at object | Speech to the infant | 331 | 1.20 | $F(2, 328) = 3.54^*$ | 13.36 (−0.30, 27.02) | $F(1, 328) = 3.70$ | −50.40 (−87.89, −12.91) | $F(1, 328) = 6.99^{**}$ |
| 7. Nondistress vocalization | Encourages attention to herself | 331 | 0.95 | $F(2, 328) = 2.02$ | 5.19 (−0.08, 10.47) | $F(1, 328) = 3.75$ | −7.48 (−15.02, −0.47) | $F(1, 323) = 4.39^*$ |
| 8. Nondistress vocalization | Encourages attention to object | 335 | 1.02 | $F(2, 332) = 2.53$ | 5.56 (0.68, 10.43) | $F(1, 332) = 5.03^*$ | −8.59 (−18.14, 0.97) | $F(1, 332) = 3.13$ |
| 9. Nondistress vocalization | Speech to the infant | 333 | 1.17 | $F(2, 330) = 6.53^{**}$ | 25.88 (5.53, 46.22) | $F(1, 330) = 6.26^*$ | −51.82 (−81.81, −21.82) | $F(1, 330) = 11.55^{***}$ |
| 10. Smile | Encourages attention to herself | 230 | 1.59 | $F(2, 227) = 6.54^{**}$ | 5.37 (2.33, 8.40) | $F(1, 227) = 12.14^{**}$ | −8.70 (−13.49, −3.91) | $F(1, 27) = 12.81^{**}$ |
| 11. Smile | Encourages attention to object | 232 | 0.59 | $F(2, 228) = 1.17$ | 7.06 (−2.51, 16.62) | $F(1, 228) = 2.11$ | −9.53 (−21.89, 2.82) | $F(1, 228) = 2.31$ |
| 12. Smile | Speech to the infant | 230 | 1.02 | $F(2, 227) = 4.01^*$ | 8.94 (.56, 17.31) | $F(1, 227) = 4.42^*$ | −17.21 (−29.48, −4.95) | $F(1, 227) = 7.65^{**}$ |

* $p < .05$. ** $p < .01$. *** $p < .001$.




Figure 1. Estimated knot and slopes from spline regression of contingency of infant nondistress vocalization for mother speech to the infant (transformed odds ratios) and maternal sensitivity (transformed and residualized).

lines with different slopes) that better describe the association between the variables. To investigate nonlinear associations between maternal contingency and sensitivity, we examined the bivariate scatter plots with locally weighted fitting and smoothing scatterplots (LOWESS) smoothers between contingency odds ratios and sensitivity. LOWESS fits curves and surfaces to noisy data by a multivariate smoothing procedure (Cleveland, 1979). Examination of several scatterplots with LOWESS smoothers at 0.50 tension revealed no relations, linear relations, or linear relations with different slopes before and after a given point (“knot”) on the contingency continuum. We therefore performed a series of spline regressions (Marsh & Cormier, 2002) to assess whether changes in slopes at knots were significant and, if significant, to obtain linear rates of change leading to and following the knot. Spline regression does not assume a curvilinear (e.g., \cap -shaped) pattern but instead allows for different linear trend lines in segments of the sample. For each pair of contingency odds ratio–EA Scale relations, the location of a single knot was estimated using nonlinear least squares regression. The start value needed for the nonlinear least squares regression was provided by visual inspection of the scatterplots with the LOWESS line.

Estimations of the knots, sample sizes, and results of spline regressions are presented separately for each of the 12 contingency odds ratio–EA Scale pairs in Table 6. For example, for the contingency of infant nondistress vocalization—mother speech to the infant and maternal sensitivity (Pair 9), spline regression estimated the knot at 1.17 to be significant. Figure 1 shows the negative quadratic regression lines changed slopes at a specific point on the contingency continuum (at a transformed $OR = 1.17$). The linear relation between this dyadic contingency and maternal sensitivity is positive until the knot, after which the linear relation is

negative. In English, as maternal responsive speech to infants contingent on infants’ nondistress vocalization increased, separate and independent ratings of maternal sensitivity also increased, until a level of contingency where the odds of mothers responding were just significantly greater than 1; at greater levels of contingency, mothers whose speech was more and more contingent were rated as less and less sensitive. Eleven of 12 contingency odds ratio–EA Scale relations followed this general pattern. Seven of 12 contingency–sensitivity relations showed significant knots; for each, the linear relation between contingency and sensitivity before the knot was positive (significant or nonsignificant), and the linear relation after the knot was negative and significant. Five of 12 contingency–sensitivity relations showed no significant knot; however, 4 of those 5 followed the same negative quadratic trend (positive slope followed by negative slope). Only 1 relation in 12 pairs failed to follow the general pattern (Pair 5); examination of the graph showed no relation between sensitivity and contingency of mother encouraging attention to object in response to infant looking at object. Finally, Tables 5 and 6 show that neither odds ratios nor knots were overdetermined (ORs range = 0.79–8.52, knots range = 0.59–4.14).

Discussion

This study focused on relations between maternal responsiveness, based on microanalytic measures of contingency, and maternal sensitivity, based on separate and independent macroanalytic global ratings. We found, in a nutshell, that, as contingency in maternal responses to infant behaviors increased, judged maternal sensitivity also increased, but only until a just-significant degree of contingency, after which mothers who were more and more contingent were rated as less and less sensitive. Mothers who adhered to levels of responsive-

ness just greater than $OR = 1$ across a range of infant-behavior—mother-response combinations were judged to be optimally sensitive. Eleven of 12 different contingency–sensitivity relations followed this general pattern.

Correlation and contingency

Six of 12 (50%) correlations between raw frequencies of mother and infant behaviors were significant; however, 11 of 12 (92%) possible contingency pairs were significant. If our analyses of infant–mother interactions had focused on correlation alone, we would incorrectly conclude that about one half of infant–mother interaction domains are coordinated. Moreover, even if dyads correspond in how relatively frequently each engages in a behavior pair, correlation provides no information on whether their behaviors are synchronized and occur contingently on one another in real time. Contingency analyses add temporal and (quasi-)causal information about moment-to-moment processes in interactions that exceed correlational analysis. (A contingency is a regular temporal relation between the occurrences of two events. A contingency exists between events A and B when the probability of occurrence of event B, given the previous occurrence of A, is greater than that of B without A. The relation between B and A may or may not be causal because the existence of a contingency is a necessary, but not a sufficient, condition for inferring causality.) Moreover, two behaviors can be correlated but not contingent, just as they can be uncorrelated but contingent (as our data demonstrate). Our approach to interaction via sequential analysis permits fine-grained assessments of infant–mother transactional processes as well as stronger inferences about causal relations between them (Bornstein, Tamis-LeMonda, & Haynes, 1999).

Sensitivity–contingency relations

Zero-order bivariate correlations between contingency odds ratios and EA Scale sensitivity showed no linear relations, positive linear relations, or negative linear relations. However, further analyses revealed consistently significant nonlinear relations of contingency with EA Scale sensitivity, wherein at just-significant levels of contingency patterns of relations changed from positive (or no relation) to negative. Specifically, when the likelihood of mothers' responses to infants' behaviors was determined to fall at a just-significant level on the contingency continuum, mothers were independently judged to be optimally sensitive. Mothers could fall at the lower end of the contingency continuum because they are nonresponsive to their infants' behaviors, are noncontingent, or respond outside the 3-s time window. Low-contingency mothers were independently rated to be less sensitive. As the likelihood of mothers' contingent responsiveness increased, independent ratings of their sensitivity increased, with the highest ratings awarded to mothers just above a significant level on the contingency continuum. There are also large differences in the levels of the knots

across the 12 pairs of contingencies. All of the knots (except for Pair 8) fall outside the 95% confidence interval for the mean odds ratios. The grand mean of knots, or points on the contingency continuum with highest sensitivity ratings for all pairs of infant–mother behaviors, was 1.46, showing that optimal maternal sensitivity occurs when the likelihood of mothers' responding to infants just exceeds the likelihood of not responding within a time window. However, the sensitivity of mothers who responded more and more to their infants' behaviors was independently judged to diminish. In briefer words, both under- and overcontingency in mothers are deemed less generally sensitive than just-significant levels of maternal contingency. Most of the work on maternal sensitivity has used rating scales (like the EA Scale); rating scales capture qualitative aspects of interactional sequences, but they fail to assess specific behavioral linkages in responsiveness, as does contingency.

It is noteworthy that the relation between contingency of maternal behavior and sensitivity was independent of the relation between amount of maternal behavior and sensitivity. For instance, how frequently mothers encourage infant attention to objects and their sensitivity was unrelated at a group level, even if mothers' encouraging attention to themselves as well as their speech to infants and sensitivity were positively related (Table 4). However, at the dyadic level, the contingency of these behaviors showed a more telling pattern of nonlinear relations with sensitivity, where moderate levels of contingency were judged highest in sensitivity (Table 6).

Implications for normative development

In interaction, contingencies generate expectancies of predictable partner reactions in relation to one's own behaviors, and vice versa. Mastering contingencies permits children to predict events and organize their behavior coherently, either to elicit desirable outcomes or to avoid aversive consequences. The ability to detect contingencies is one of the quintessential features of adaptation (Canfield & Haith, 1991), and learning contingencies between one's own behavior and environmental events is a key adaptation in childhood (Millar & Weir, 1992).

Human infants normally develop in a responsive, social environment and quickly come to perceive contingent regularities in others' behavior (Bornstein et al., 1990; Feldman, 2003). Contingency detection may be present from birth (Gewirtz & Palàez-Nogueras, 1992) and is thought to be especially salient during interactions (Rovee-Collier, 1987; Stern, 1985; Tarabulsky et al., 1996). Gergely and Watson (Gergely, 2003; Gergely & Watson, 1996, 1999) proposed an innate "contingency detection module," which analyzes the temporal conditional probabilities affording infants the capacity to detect contingent relations in the environment. Bloom (1988) and Bigelow (1998; Bigelow & Rochat, 2006), respectively, demonstrated that infants as young as 2 to 3 months discriminate between adult behaviors that occur contingently and behaviors that take place randomly and that infants identify and adapt to levels of contingency they experience frequently. Furthermore, experi-

ences with contingent responsiveness appear to instill in infants an awareness of caregiver availability and reliability and so promote senses of security, trust, and effectance (MacDonald, 1992; Watson & Ramey, 1972); anticipatory control, which is inherent in contingency, is also reputedly enjoyable, stimulating, and rewarding to infants (Gergely & Watson, 1999).

Our study showed that European American mothers who are moderately but significantly contingent to their young infants are judged to be the most sensitive, and mothers who are either noncontingent to less contingent or more contingent to consistently contingent are judged to be less sensitive. Social responsiveness between people (mothers and infants included) is rarely, if ever, perfect. Rather, variability in social (and maternal) responsiveness is common. Mothers' quantitatively as well as qualitatively imperfect contingency may be adaptive for normal infant development in several ways. Experiences with imperfect contingency have implications for understanding the nature and structure of infant learning and interactions. For example, 3-month-olds display an attentional preference for moderate contingency over perfect contingency, and the fastest learning of novel actions occurs in situations that offer moderate levels of contingency (Watson, 1979). Watson (1985) further observed that behavioral arousal in infants was reduced when no contingency existed or when experienced contingency approached perfection; rather, infant arousal was maximized at moderate levels of contingency. This observation and our findings articulate with the classic Yerkes–Dodson law, the empirical relation between arousal and performance that shows that performance increases with physiological or mental arousal, but only up to a point; when levels of arousal become too high, performance decreases (Yerkes & Dodson, 1908). For mothers, a certain level of arousal may be optimal; when levels are too high or too low, responsiveness will be disrupted or deficient (Henry & Wang, 1998; Wang, 1997). Thus, focusing attention on imperfect contingencies that are present in the environment or in others' behavior, rather than on consistent contingency, is adaptive. Infants' early ability to detect differences between perfect and imperfect contingencies has been hypothesized as an initial basis for distinguishing self from other. Even younger 2-month-olds detect response contingencies in interactions with mothers and prefer intermittently contingent "normal" interactions over noncontingent responses (Murray & Trevarthen, 1985; Nadel, Carchon, Kervella, Marcelli, & Réserbat-Plantey, 1999; Reddy, Chisholm, Forrester, Conforti, & Maniatopoulou, 2007). Markova and Legerstee (2006) tested 5- and 13-week-old infants in face-to-face interactions with their mothers in three types of contingent situations: one involving intermittent and imperfect contingencies (a natural situation), one involving near contingencies (an imitative situation), and one involving noncontingent interactions (a yoked situation). Infants of affectively attuned mothers preferred the intermittent contingencies of natural interactions. By the end of the first month of life, infants may already prefer "normal" intermittent contingencies over noncontingent or perfectly contingent interactions with

mothers. Imperfect contingency, such as we found naturalistically, fits infants' normal interactions and expectations.

Clinical implications

Parental responsiveness is best understood in relation to children's needs and proclivities, and these relations vary at different ages and in different contexts. However, hypo- and hyperlevels of parental contingency can translate into "withdrawn" and "vigilant" parenting, respectively, engendering nonoptimal child development. As Ainsworth et al. (1974) observed, high maternal contingency often is judged as overstimulating, insensitive, hypervigilant, or hovering, and constant, rapid parental responses may be nonproductive. Overcontingent parenting risks failing to instill a child with the confidence and ability to self-regulate and explore, and prevent or inhibit the proper development of coping skills generally deemed as interfering with children's autonomy and individuation. By the other side of the same token, unresponsive parenting risks poor development, as is the case with children of depressed mothers (Cohn & Tronick, 1989; Cooper, Tomlinson, Swartz, Woolgar, & Murray, 1999; Feldman, 2003; Murray & Cooper, 1997). Interactive contingency that is insufficient or excessive predicts or engenders insecure-type attachments. Where "tightly coupled" mother–infant dyads experience disruptions as psychologically "shattering," relatively "loosely coupled" (midrange) dyads are more resilient (Sander, 1995).

By contrast, moderate levels of maternal contingency have been associated with secure attachment relationships (Belsky et al., 1984; Hane, Feldstein, & Dernetz, 2003; Isabella & Belsky, 1991; Jaffe, Beebe, Feldstein, Crown, & Jasnow, 2001; Leyendecker et al., 1997; Malatesta et al., 1989; Tobias, 1995; Völker, Keller, Lohaus, Cappenberg, & Chasiotis, 1999; Warner, 1992). Jaffe (Jaffe et al., 2001) and Beebe (Beebe et al., 2011), for example, found that midrange levels of coordination between maternal and infant vocal exchanges were associated with security of attachment, whereas either very low or very high levels of coordination were associated with insecure attachment. Sensitive mothers may choose not to respond to each and every infant behavior, thereby allowing their infants the opportunity to act in a more autonomous fashion, a hypothesis that we plan to test. Because our program of research is longitudinal, our own next steps also include examining the relative predictive validity of maternal odds ratios for children's long-term developmental outcomes.

Our pattern of findings is consistent with the notion that undercontingency and overcontingency in mother–infant interactions are suggestive of maternal psychopathology. Clinically depressed mothers, although not a homogenous group, tend to be unresponsive to infant regulatory signals and fail to provide infants with appropriate and contingent stimulation (Cohn & Tronick, 1989; Feldman, 2003; Jameson, Gelfand, Kulcsar, & Teti, 1997; Reck et al., 2011), thereby making for longer response latencies, larger and longer mismatches, and few reparations. As a consequence, the infants of chroni-

cally depressed mothers experience negative affect and, over time, develop a depressed negative mood (Reck et al., 2011). By contrast, overcontingency overrides appropriateness, a key ingredient of responsiveness (Ainsworth et al., 1974), and seems to index communicative stress and hypervigilance (Beebe et al., 2008; Kaitz & Mayal, 2005) that may inhibit the development of security and autonomy (Belsky et al., 1984). Thus, it seems that nonoptimal interactions indicative of distress are characterized by both heightened interactive contingency (in some modalities) and lowered (in others; Belsky et al., 1984; Hane et al., 2003; Leyendecker et al., 1997; Lewis & Feiring, 1989; Malatesta et al., 1989; Roe, Roe, Drivas, & Bronstein, 2006).

In the mutual regulation model (Tronick, 2003, 2007; Tronick & Cohn, 1989), mother–infant interplay is typically a moderately coordinated affair, in which affective interactive “matches” (coordinated behavioral and affective states) and “mismatches” (uncoordinated behavioral and affective states) occur, with mismatches frequently and quickly being repaired (“interactive repair”) back to interactional matching states (Reck et al., 2011; Tronick & Reck, 2009). Reparation or its failure has important developmental consequences. In normal interactions, reparations over time can instill a sense of mastery and control (Tronick & Reck, 2009). Unrepaired interactive mismatches can accumulate negative affect in the infant and eventuate in the development of psychopathology (Tronick & Reck, 2009).

Limitations and future directions

Although the characteristics of our sample may limit broad generalizability of the findings, our participants were socioeconomically heterogeneous, recruited over wide age, education, and SES ranges, and represented a demographically significant portion of the US population. Nonetheless, it may be revealing to explore the generalizability of contingency–sensitivity relations in more diverse and even clinical populations. In addition, the potential long-term consequences of individual variation in parental responsiveness are not well understood. Avoiding simple “dose–response” or “more is better” approaches to responsiveness (and perhaps to other forms of parenting) will be essential in future theorizing. Our measure of contingency focused on promptness between microcoded behaviors; however, sensitivity is a different, and global, construct that includes appropriateness.

The sensitivity construct developed in the context of theory and research in maternal caregiving responses to infant distress (Ainsworth et al., 1978; Bowlby, 1969). Here we studied sensitivity and contingency to nondistressed infant behaviors. Future work could look at contingency responsiveness in mothers to distress (infant cry) as well as incorporate other nondistressed interactive behaviors (touch). In addition, the complex curvilinear contingency–sensitivity associations we identified in this study are by no means unique in developmental science (see Hane et al., 2003; Leerkes & Crockenberg, 2002). This pattern of findings suggests that innumer-

able weak to moderate correlations between parenting and child development could be revisited to discern nonlinear (η) versus linear (r) associations.

According to Cicchetti and Toth (2009, p. 16), “developing and evaluating methods for preventing and ameliorating maladaptive and psychopathological outcomes” is a key goal of developmental psychopathology. Sensitive responsiveness is widely identified with positive parenting and is often the target of intervention because variations from optimal responsiveness may distort children’s experiences with subsequent negative alterations in cognition or social interaction (Cicchetti & Tucker, 1994). One illustrative trial was conducted in the Netherlands by van den Boom (1994). Her study targeted 100 6-month-old infants who were selected shortly after birth for being irritable and therefore at risk for developing insecure attachment. Individually tailored interventions consisted of three 2-hr sessions over 3 months that were meant to prevent attachment disorders by improving maternal responsiveness. Mothers of 50 treatment infants were encouraged to imitate infant behaviors, repeat infant verbal expressions, and responsively soothe infant cries. Van den Boom assessed mother–infant interaction and infant exploration both before and after the intervention as well as attachment security 3 months after the intervention concluded. Immediately postintervention, mothers were more responsive and infants were more sociable, self-soothing, and engaged in more sophisticated exploration. In the follow-up 3 months later, intervention infants were more securely attached than controls: 62% of treatment infants were classified as secure compared to 28% of control infants. Significant effects on maternal responsiveness and child cooperation persisted until at least the children’s third year.

On the assumption that more (responsiveness) is better, a usual goal of clinical family interventions is to increase parental practices like responsiveness (e.g., Anisfeld, Casper, Nozyce, & Cunningham, 1990; De Wolff & van IJzendoorn, 1997; Dickie & Gerber, 1980; Gardner, Walker, Powell, Grantham-McGregor, 2003; Kendrick et al., 2000; Mahoney & Powell, 1988; Riksen-Walraven, 1978; van IJzendoorn, Juffer, & Duyvesteyn, 1995). The consensus from a large number of early reviews held that (a) interventions can enhance parental responsiveness and (b) enhancing responsiveness might brighten child development (e.g., Geeraert, Van den Noortgate, Grietens, & Onghena, 2004; Kendrick et al., 2000; Peltó, Dickin, & Engle, 1999; van IJzendoorn, Juffer, et al., 1995). For example, Keys to Caregiving (Speitz, Johnson-Crowley, Sumner, & Barnard, 1990) is an intervention program designed to help parents understand and practice appropriate patterns of responding to infant behaviors that occur commonly in daily life by focusing on what mothers do with their infants. Practice sessions intend to render maternal responsiveness habitual. However, consistent with our findings, two recent meta-analyses on the effectiveness of parenting programs challenge the commonly held assumption that more is better (Kaminski, Vallew, Filene, & Boyle, 2008; Lundahl, Risser, & Lovejoy, 2006).

Besides augmenting our understanding of the nature and structure of responsiveness, this study may therefore have practical implications for clinical interventions. In a nutshell, if interventions are operationalized behaviorally to move parents away from nonresponding or low levels of responding, they need to do so toward goals of just contingent (not extreme) levels of responsiveness.

Conclusions

Parenting children is a two-way street. Children are active partners in their socialization, and parents regularly respond to their children's cues: parents feed and diaper their crying baby and opt to play when the same infant is awake and alert. A transactional perspective on development underscores that parents and children influence one another through time

(Bornstein, 2009); hence, a fuller picture of parent–child mutuality can be developed only when domains of child activity and dimensions of parental responsiveness are jointly considered. Responsiveness has been thought about (and assessed) more broadly within a monotonic framework of parenting. Psychoanalysts, personality theorists, ethologists, and attachment theorists historically have conceptualized parenting as traitlike and (uni)dimensional (Holden, 2010; Martin, 1989). In an operational sense, the “warm,” “adequate,” and “good” parent is *more* involved, devoted, invested, and sensitive. However, an optimum “midrange model” of interactive contingency constitutes an important antidote to the widespread simplistic assumption in the parenting literature that more (contingency) is better.

In good responsive parenting (and perhaps in many domains in parenting), *some is more*.

References

- Ainsworth, M. D. S. (1967). *Infancy in Uganda: Infant care and the growth of love*. Baltimore, MD: Johns Hopkins University Press.
- Ainsworth, M. D. S., Bell, S. M., & Stayton, D. J. (1974). Infant–mother attachment and social development. In M. P. Richards (Ed.), *The introduction of the child into a social world* (pp. 99–135). London: Cambridge University Press.
- Ainsworth, M. D. S., Blehar, M. C., Waters, E., & Wall, S. (1978). *Patterns of attachment: A psychological study of the strange situation*. Hillsdale, NJ: Erlbaum.
- Anisfeld, E., Casper, V., Nozyce, M., & Cunningham, N. (1990). Does infant carrying promote attachment? An experimental study of the effects of increased physical contact on the development of attachment. *Child Development*, *61*, 1617–1627.
- Bakeman, R. (2004). Sequential analysis. In M. Lewis-Beck, A. E. Bryman, & T. F. Liao (Eds.), *The SAGE encyclopedia of social science research methods* (Vol. 3, pp. 1024–1026). Thousand Oaks, CA: Sage.
- Bakeman, R., Deckner, D. F., & Quera, V. (2005). Analysis of behavioral streams. In D. M. Teti (Ed.), *Handbook of research methods in developmental psychology* (pp. 394–420). Oxford: Blackwell.
- Bakeman, R., & Gnisci, A. (2005). Sequential observational methods. In M. Eid & E. Diener (Eds.), *Handbook of multimethod measurement in psychology* (pp. 127–140). Washington, DC: American Psychological Association.
- Bakeman, R., & Quera, V. (2004). General sequential querier: GSEQ for Windows, version 4.1.2 [Computer software]. Atlanta, GA: Georgia State University and Universidad de Barcelona.
- Beebe, B., Jaffe, J., Buck, K., Chen, H., Cohen, P., Feldstein, S., et al. (2008). Six-week postpartum maternal depressive symptoms and 4-month mother–infant self- and interactive contingency. *Infant Mental Health Journal*, *29*, 442–471.
- Beebe, B., Jaffe, J., Lachmann, F., Feldstein, S., Crown, C., & Jasnow, J. (2000). Systems models in development and psychoanalysis: The case of vocal rhythm coordination and attachment. *Infant Mental Health Journal*, *21*, 99–122.
- Beebe, B., Jaffe, J., Markese, S., Buck, K., Chen, H., Cohen, P., et al. (2010). The origins of 12-month attachment: A microanalysis of 4-month mother–infant interaction. *Attachment & Human Development*, *12*, 3–141.
- Belsky, J., Rovine, M., & Taylor, D. (1984). The Pennsylvania infant and family development project: III. The origins of individual differences in infant–mother attachment: Maternal and infant contributions. *Child Development*, *55*, 718–728.
- Bigelow, A. E. (1998). Infants' sensitivity to familiar imperfect contingencies in social interaction. *Infant Behavior and Development*, *21*, 149–162.
- Bigelow, A. E., MacLean, B. K., & MacDonald, D. (1996). Infants' response to live and replay interactions with self and mother. *Merrill–Palmer Quarterly*, *42*, 596–611.
- Bigelow, A. E., & Rochat, P. (2006). Two-month-old infants' sensitivity to social contingency in mother–infant and stranger–infant interaction. *Infancy*, *9*, 313–325.
- Biringen, Z., & Easterbrooks, M. A. (2012). Emotional availability: Concept, research, and window on development and psychopathology. *Development and Psychopathology*, *24*, 1–8.
- Biringen, Z., Robinson, J. L., & Emde, R. N. (1998). *The Emotional Availability Scales* (3rd ed.). Fort Collins, CO: Colorado State University, Department of Human Development and Family Studies.
- Bloom, K. (1988). Quality of adult vocalizations affects the quality of infant vocalizations. *Journal of Child Language*, *15*, 469–480.
- Bornstein, M. H. (2009). Toward a model of culture↔parent↔child transactions. In A. Sameroff (Ed.), *The transactional model of development: How children and contexts shape each other* (pp. 139–161). Washington, DC: American Psychological Association.
- Bornstein, M. H., Azuma, H., Tamis-LeMonda, C. S., & Ogino, M. (1990). Mother and infant activity and interaction in Japan and in the United States: I. A comparative macroanalysis of naturalistic exchanges. *International Journal of Behavioral Development*, *13*, 267–287.
- Bornstein, M. H., Hahn, C.-S., Suwalsky, J. T. D., & Haynes, O. M. (2003). Socioeconomic status, parenting, and child development: The Hollingshead Four-Factor Index of Social Status and the Socioeconomic Index of Occupations. In M. H. Bornstein & R. H. Bradley (Eds.), *Monographs in parenting series: Socioeconomic status, parenting, and child development* (pp. 29–82). Mahwah, NJ: Erlbaum.
- Bornstein, M. H., Suwalsky, J. T. D., & Breakstone, D. A. (2012). Emotional relationships between mother and infants: Knowns, unknowns, and unknown unknowns. *Development and Psychopathology*, *24*, 137–142.
- Bornstein, M. H., & Tamis-LeMonda, C. S. (1990). Activities and interactions of mothers and their firstborn infants in the first six months of life: Covariation, stability, continuity, correspondence, and prediction. *Child Development*, *61*, 1206–1217.
- Bornstein, M. H., Tamis-LeMonda, C. S., Hahn, C.-S., & Haynes, O. M. (2008). Maternal responsiveness to very young children at three ages: Longitudinal analysis of a multidimensional modular and specific parenting construct. *Developmental Psychology*, *44*, 867–874.
- Bornstein, M. H., Tamis-LeMonda, C. S., & Haynes, O. M. (1999). First words in the second year: Continuity, stability, and models of concurrent and predictive correspondence in vocabulary and verbal responsiveness across age and context. *Infant Behavior and Development*, *22*, 65–85.
- Bornstein, M. H., Toda, S., Azuma, H., Tamis-LeMonda, C. S., & Ogino, M. (1990). Mother and infant activity and interaction in Japan and in the United States: II. A comparative microanalysis of naturalistic exchanges focused on the organization of infant attention. *International Journal of Behavioral Development*, *13*, 289–308.
- Bowlby, J. (1969). *Attachment and loss: Vol. 1. Attachment*. New York: Basic Books.
- Bus, A. G., & van IJzendoorn, M. H. (1992). Patterns of attachment in frequently and infrequently reading mother–child dyads. *Journal of Genetic Psychology*, *153*, 395–403.

- Canfield, R. L., & Haith, M. M. (1991). Young infants' visual expectations for symmetric and asymmetric stimulus sequences. *Developmental Psychology*, 27, 198–208.
- Chapple, E. (1970). *Culture and biological man*. New York: Holt, Rinehart and Winston.
- Cicchetti, D., & Pogge-Hesse, P. (1982). Possible contributions of the study of organically retarded persons to developmental theory. In E. Zigler & D. Balla (Eds.), *Mental retardation: The developmental-difference controversy* (pp. 277–318). Hillsdale, NJ: Erlbaum.
- Cicchetti, D., & Toth, S. L. (2009). The past achievements and future promises of developmental psychopathology: The coming of age of a discipline. *Journal of Child Psychology and Psychiatry*, 50, 16–25.
- Cicchetti, D., & Tucker, D. (1994). Development and self-regulatory structures of the mind. *Development and Psychopathology*, 6, 533–549.
- Cleveland, W. S. (1979). Robust locally weighted regression and smoothing scatterplots. *Journal of the American Statistical Association*, 74, 829–836.
- Cohn, J., & Elmore, M. (1988). Effect of contingent changes in mothers' affective expression on the organization of behavior in 3-month-old infants. *Infant Behavior and Development*, 11, 493–505.
- Cohn, J. F., & Tronick, E. (1989). Specificity of infants' response to mothers' affective behavior. *Journal of the American Academy of Child & Adolescent Psychiatry*, 28, 242–248.
- Cooper, P., Tomlinson, M., Swartz, L., Woolgar, M., & Murray, L. (1999). Post-partum depression and the mother–infant relationship in a South African peri-urban settlement. *British Journal of Psychiatry*, 175, 554–558.
- Cote, L. R., Bornstein, M. H., Haynes, O. M., & Bakeman, R. (2008). Mother–infant person- and object-directed interactions in Latino immigrant families: A comparative approach. *Infancy*, 13, 338–365.
- Damast, A. M., Tamis-LeMonda, C. S., & Bornstein, M. H. (1996). Mother–child play: Sequential interactions and the relation between maternal beliefs and behaviors. *Child Development*, 67, 1752–1766.
- De Wolff, M. S., & van IJzendoorn, M. H. (1997). Sensitivity and attachment: A meta-analysis on parental antecedents of infant attachment. *Child Development*, 68, 571–591.
- Dickie, J. R., & Gerber, S. C. (1980). Training in social competence: The effect on mothers, fathers, and infants. *Child Development*, 51, 1248–1251.
- Drillien, C. M. (1957). The social and economic factors affecting the incidence of premature birth: I. Premature births without complications of pregnancy. *Journal of Obstetrics and Gynecology of the British Empire*, 64, 161–184.
- Dunham, P., & Dunham, F. (1994). Optimal social structures and adaptive infant development. In C. Moore & P. Dunham (Eds.), *Joint attention: Its origins and role in development* (pp. 159–188). Hillsdale, NJ: Erlbaum.
- Feldman, R. (2003). Infant–mother and infant–father synchrony: The coregulation of positive arousal. *Infant Mental Health Journal*, 24, 1–23.
- Gardner, J. M., Walker, S. P., Powell, C. A., & Grantham-McGregor, S. (2003). A randomized controlled trial of a home-visiting intervention on cognition and behaviour in low birth weight infants. *Journal of Pediatrics*, 143, 634–639.
- Geeraert, L., Van den Noortgate, W., Grietens, H., & Onghena, P. (2004). The effects of early prevention programs for families with young children at risk for physical child abuse and neglect: A meta-analysis. *Child Maltreatment*, 9, 277–291.
- Gergely, G. (2003). The development of understanding self and agency. In U. Goswami (Ed.), *Blackwell's handbook of child cognitive development* (pp. 26–46). Oxford: Blackwell.
- Gergely, G., & Watson, J. (1996). The social biofeedback model of parental affect-mirroring: The development of emotional self-awareness and self-control in infancy. *International Journal of Psychoanalysis*, 77, 1181–1212.
- Gergely, G., & Watson, J. (1999). Early socio-emotional development: Contingency perception and the social–biofeedback model. In P. Rochat (Ed.), *Early social cognition* (pp. 101–136). Mahwah, NJ: Erlbaum.
- Gewirtz, J. L., & Palàez-Nogueras, M. (1992). B. F. Skinner's legacy to human infant behavior and development. *American Psychologist*, 47, 1411–1422.
- Goldstein, M. H., Schwade, J. A., & Bornstein, M. H. (2009). The value of vocalizing: Five-month-old infants associate their own noncry vocalizations with responses from caregivers. *Child Development*, 80, 636–644.
- Graham, S. (1992). Most of the subjects were white and middle class: Trends in published research on African Americans in selected APA Journals, 1970–1989. *American Psychologist*, 47, 629–639.
- Gratier, M. (2003). Expressive timing and interactional synchrony between mothers and infants: Cultural similarities, cultural differences, and the immigration experience. *Cognitive Development*, 18, 533–554.
- Gros-Louis, J., West, M. J., Goldstein, M. H., & King, A. P. (2006). Mothers provide differential feedback to infants' prelinguistic sounds. *International Journal of Behavioral Development*, 30, 509–516.
- Hane, A., Feldstein, S., & Dernetz, V. (2003). The relation between coordinated interpersonal timing and maternal sensitivity in four-month-old infants. *Journal of Psycholinguistic Research*, 32, 525–539.
- Henry, J. P., & Wang, S. (1998). Effects of early stress on adult affiliative behavior. *Psychoneuroendocrinology*, 23, 863–875.
- Holden, G. W. (2010). *Parenting: A dynamic perspective*. Thousand Oaks, CA: Sage.
- Hollingshead, A. B. (1975). *The Four Factor Index of Social Status*. Unpublished manuscript, Yale University, Department of Sociology. Retrieved from http://www.yale-university.com/sociology/faculty/docs/hollingshead_socStat4factor.pdf
- Hsu, H.-C., & Lavelli, M. (2005). Perceived and observed parenting behavior in American and Italian first-time mothers across the first 3 months. *Infant Behavior and Development*, 28, 503–518.
- Humes, K. R., Jones, N. A., & Ramirez, R. R. (2011). Overview of race and Hispanic origin: 2010. In US Census Bureau, *2010 Census Briefs* (pp. 1–24). Retrieved from <http://www.census.gov/prod/cen2010/briefs/c2010br-02.pdf>
- Isabella, R. A., & Belsky, J. (1991). Interactional synchrony and the origins of infant–mother attachment: A replication study. *Child Development*, 62, 373–384.
- Isabella, R. A., Belsky, J., & von Eye, A. (1989). Origins of infant–mother attachment: An examination of interactional synchrony during the infant's first year. *Developmental Psychology*, 25, 12–21.
- Isley, S., O'Neil, R., & Parke, R. D. (1996). The relation of parental affect and control behaviors to children's classroom acceptance: A concurrent and predictive analysis. *Early Education and Development*, 7, 7–23.
- Jaffe, J., Beebe, B., Feldstein, S., Crown, C., & Jasnou, M. (2001). Rhythms of dialogue in infancy: Co-ordinated timing in development. *Monographs of the Society for Research in Child Development*, 66, vi–131.
- Jameson, P. B., Gelfand, D. M., Kulcsar, E., & Teti, D. M. (1997). Mother–toddler interaction patterns associated with maternal depression. *Development and Psychology*, 9, 537–550.
- Kaitz, M., & Maytal, H. (2005). Interactions between anxious mothers and their infants: An integration of theory and research findings. *Infant Mental Health*, 26, 570–597.
- Kaminski, J. W., Vallew, L. A., Filene, J. H., & Boyle, C. L. (2008). A meta-analytic review of components associated with Parent Training program effectiveness. *Journal of Abnormal Child Psychology*, 36, 567–589.
- Keller, H., Lohaus, A., Völker, S., Cappenberg, M., & Chasiotis, A. (1999). Temporal contingency as an independent component of parenting behavior. *Child Development*, 70, 474–485.
- Kendrick, D., Elkan, R., Hewitt, M., Dewey, M., Blair, M., Robinson, J., et al. (2000). Does home visiting improve parenting and the quality of the home environment? A systematic review and meta-analysis. *Archives of Disease in Childhood*, 82, 443–451.
- Kogan, N., & Carter, A. S. (1996). Mother–infant reengagement following the still-face: The role of maternal emotional availability in infant affect regulation. *Infant Behavior and Development*, 19, 359–370.
- Leerkes, E. M., & Crockenberg, S. C. (2002). The development of maternal self-efficacy and its impact on maternal behavior. *Infancy*, 3, 227–247.
- Lewis, M., & Feiring, C. (1989). Infant, mother and mother–infant interaction behavior and subsequent attachment. *Child Development*, 60, 831–837.
- Leyendecker, B., Lamb, M., Fracasso, M., Scholmerich, A., & Larson, D. (1997). Playful interaction and the antecedents of attachment: A longitudinal study of Central American and Euro-American mothers and infants. *Merrill-Palmer Quarterly*, 43, 24–47.
- Lohaus, A., Keller, H., Ball, J., Elben, C., & Völker, S. (2001). Maternal sensitivity: Components and relations to warmth and contingency. *Parenting*, 1, 267–284.
- Lundahl, B., Risser, H. J., & Lovejoy, M. C. (2006). A meta-analysis of parent training: Moderators and follow-up effects. *Clinical Psychology Review*, 26, 86–104.
- MacDonald, K. (1992). Warmth as a developmental construct: An evolutionary analysis. *Child Development*, 63, 753–773.
- Mahoney, G., & Powell, A. (1988). Modifying parent–child interaction: Enhancing the development of handicapped children. *Journal of Special Education*, 22, 82–96.

- Malatesta, C. Z., Culver, C., Tesman, J. R., & Shepard, B. (1989). The development of emotion expression in the first two years of life. *Monographs of the Society for Research in Child Development*, 54(1–2), 1–104.
- Markova, G., & Legerstee, M. (2006). Contingency, imitation, and affect sharing: Foundations of infants' social awareness. *Developmental Psychology*, 42, 132–141.
- Marsh, L. C., & Cormier, D. R. (2002). *Spline regression models*. Thousand Oaks, CA: Sage.
- Martin, J. A. (1989). Personal and interpersonal components of responsiveness. *New Directions for Child and Adolescent Development*, 43, 5–14.
- McCune-Nicolich, L., & Fenson, L. (1984). Methodological issues in studying early pretend play. In T. D. Yawkey & A. D. Pelligrini (Eds.), *Child's play: Developmental and applied* (pp. 81–124). Hillsdale, NJ: Erlbaum.
- Millar, W. S., & Weir, C. G. (1992). Relations between habituation and contingency learning in 5- to 12-month-old infants. *Cahiers de Psychologie Cognitive/Current Psychology of Cognition*, 12, 209–222.
- Moore, G. A., Hill-Soderlund, A. L., Propper, C. B., Calkins, S. D., Mills-Koonce, W. R., & Cox, M. J. (2009). Mother–infant vagal regulation in the face-to-face still-face paradigm is moderated by maternal sensitivity. *Child Development*, 80, 209–223.
- Murray, L., & Cooper, P. J. (Eds.). (1997). *Postpartum depression and child development*. New York: Guilford Press.
- Murray, L., & Trevarthen, C. (1985). Emotional regulation of interaction between two-month-olds and their mothers. In T. Field & N. Fox (Eds.), *Social perception in infants* (pp. 101–125). Norwood, NJ: Ablex.
- Nadel, J., Carchon, I., Kervella, C., Marcelli, D., & Réserbat-Plantey, D. (1999). Expectancies for social contingency in 2-month-olds. *Developmental Science*, 2, 164–173.
- Parke, R. D. (2000). Beyond white and middle class: Cultural variations in families—Assessments, processes, and policies. *Journal of Family Psychology*, 14, 331–333.
- Pelto, G. H., Dickin, K., & Engle, P. L. (1999). A critical link: Interventions for physical growth and psychological development. *Nutritional Anthropology*, 2, 21–27.
- Pettit, G. S., & Bates, J. E. (1989). Family interaction patterns and children's behavior problems from infancy to 4 years. *Developmental Psychology*, 25, 413–420.
- Reck, C., Noe, D., Stefenelli, U., Fuchs, T., Cenciotti, F., Stehle, E., et al. (2011). Interactive coordination of currently depressed inpatient mothers and their infants during the postpartum period. *Infant Mental Health Journal*, 32, 542–562.
- Reddy, V., Chisholm, V., Forrester, D., Conforti, M., & Maniatiopoulou, D. (2007). Facing the perfect contingency: Interactions with the self at 2 and 3 months. *Infant Behavior and Development*, 30, 195–212.
- Ricksen-Walraven, J. M. (1979). Effects of caregiver behavior on habituation rate and self-efficacy in infants. *International Journal of Behavioral Development*, 1, 105–130.
- Robinson, J., Little, C., & Biringen, Z. (1993). Emotional communication in mother–toddler relationships: Evidence for early gender differentiation. *Merrill–Palmer Quarterly*, 39, 496–517.
- Roe, K. V., Roe, A., Drivas, A., & Bronstein, R. (2006). A curvilinear relationship between maternal vocal stimulation and three-month-olds' cognitive processing: A cross-cultural phenomenon. *Infant Mental Health Journal*, 11, 175–189.
- Rovee-Collier, C. (1987). Learning and memory in infancy. In J. D. Osofsky (Ed.), *Handbook of infant development* (pp. 98–148). New York: Wiley.
- Sander, L. W. (1995). Identity and the experience of specificity in a process of recognition: Commentary on Seligman and Shanok. *Psychoanalytic Dialogues*, 5, 579–593.
- Shaw, D. S., Keenan, K., & Vondra, J. I. (1994). Developmental precursors of externalizing behavior: Ages 1 to 3. *Developmental Psychology*, 30, 355–364.
- Shrout, P. E., & Fleiss, J. L. (1979). Intraclass correlations: Uses in assessing raters reliability. *Psychological Bulletin*, 86, 420–428.
- Speitz, A., Johnson-Crowley, N., Sumner, G., & Barnard, K. (1990). *Keys to caregiving manual*. Seattle, WA: NCAST Publications.
- Stern, D. N. (1985). *The interpersonal world of the infant: A view from psychoanalysis and development psychology*. New York: Basic Books.
- Tarabulsky, G., Tessier, R., & Kappas, A. (1996). Contingency detection and the contingent organization of behavior interactions: Implications for socioemotional development in infancy. *Psychological Bulletin*, 120, 25–41.
- Tilton-Weaver, L. C., & Kakihara, F. (2008). United States of America. In J. J. Arnett (Ed.), *International encyclopedia of adolescence* (Vol. 1, pp. 1061–1076). New York: Routledge.
- Tobias, K. (1995). *The relation between maternal attachment and patterns of mother–infant interaction at four months*. Unpublished doctoral dissertation, City University of New York.
- Tomlinson, M., & Swartz, L. (2003). Imbalances in the knowledge about infancy: The divide between rich and poor countries. *Infant Mental Health Journal*, 24, 547–556.
- Tronick, E., Als, H., Adamson, L., Wise, S., & Brazelton, T. B. (1978). The infant's response to entrapment between contradictory messages in face-to-face interaction. *Journal of the American Academy of Child Psychiatry*, 17, 1–13.
- Tronick, E., & Reck, C. (2009). Infants of depressed mothers. *Harvard Review of Psychiatry*, 17, 147–156.
- Tronick, E. Z. (2003). “Of course all relationships are unique”: How co-creative processes generate unique mother–infant and patient–therapist relationships and change other relationships. *Psychological Inquiry*, 23, 473–491.
- Tronick, E. Z. (2007). *The neurobehavioral and social–emotional development of infants and children*. New York: Norton.
- Tronick, E. Z., & Cohn, J. F. (1989). Infant–mother face-to-face interaction: Age and gender differences in coordination and the occurrence of miscoordination. *Child Development*, 60, 85–92.
- US Census Bureau. (2008). *2008 American community survey*. Retrieved from www.factfinder.census.gov
- van den Boom, D. C. (1994). The influence of temperament and mothering on attachment and exploration: An experimental manipulation of sensitive responsiveness among lower class mothers with irritable infants. *Child Development*, 65, 1457–1477.
- Van Egeren, L. A., Barratt, M. S., & Roach, M. A. (2001). Mother–infant responsiveness: Timing, mutual regulation, and interactional context. *Developmental Psychology*, 37, 684–697.
- van IJzendoorn, M. H., Dijkstra, J., & Bus, A. G. (1995). Attachment, intelligence, and language: A meta-analysis. *Social Development*, 4, 115–128.
- van IJzendoorn, M. H., & Hubbard, F. O. A. (2000). Are infant crying and maternal responsiveness during the first year related to infant–mother attachment at 15 months? *Attachment and Human Development*, 2, 371–391.
- van IJzendoorn, M. H., Juffer, F., & Duyvesteyn, M. G. C. (1995). Breaking the intergenerational cycle of insecure attachment: A review of the effects of attachment-based interventions on maternal sensitivity and infant security. *Journal of Child Psychology and Psychiatry*, 36, 225–248.
- Vibbert, M., & Bornstein, M. H. (1989). Specific associations between domains of mother–child interaction and toddler referential language and pretense play. *Infant Behavior and Development*, 12, 163–184.
- Völker, S., Keller, H., Lohaus, A., Cappenberg, M., & Chasiotis, A. (1999). Maternal interactive behavior in early infancy and later attachment. *International Journal of Behavioral Development*, 23, 921–936.
- Wakschlag, L. S., & Hans, S. L. (1999). Relation of maternal responsiveness during infancy to the development of behavior problems in high-risk youths. *Developmental Psychology*, 35, 569–579.
- Wang, S. (1997). Traumatic stress and attachment. *Acta Physiologica Scandinavica*, 161, 164–169.
- Warner, R. (1992). Cyclicity of vocal activity increases during conversation: Support for a nonlinear systems model of dyadic social interaction. *Behavioral Science*, 37, 128–138.
- Watson, J. S. (1979). Perception of contingency as a determinant of social responsiveness. In E. Thoman (Ed.), *The origins of social responsiveness*. Hillsdale, NJ: Erlbaum.
- Watson, J. S. (1985). Contingency perception in early social development. In T. M. Field & N. A. Fox (Eds.), *Social perception in infants* (pp. 157–176). Norwood, NJ: Ablex.
- Watson, J. S., & Ramey, C. T. (1972). Reactions to response-contingent stimulation in early infancy. *Merrill–Palmer Quarterly*, 18, 219–227.
- Wickens, T. D. (1993). Analysis of contingency tables with between-subjects variability. *Psychological Bulletin*, 113, 191–204.
- Yerkes, R. M., & Dodson, J. D. (1908). The relation of strength of stimulus to rapidity of habit-formation. *Journal of Comparative Neurology and Psychology*, 18, 459–482.