

## Season of birth modifies puberty in female and male goats raised under subtropical conditions

J. A. Delgado<sup>1†</sup>, M. A. De Santiago-Miramontes<sup>1</sup> and E. Carrillo<sup>2</sup>

<sup>1</sup>Centro de Investigación en Reproducción Caprina, Universidad Autónoma Agraria Antonio Narro, Periférico Raúl López Sánchez y Carretera a Santa Fe, CP 27054, Torreón, Coahuila, México; <sup>2</sup>Instituto Tecnológico de Torreón, Carretera Torreón-San Pedro km 7.5, Torreón, Coahuila, México

(Received 24 October 2006; Accepted 29 March 2007)

*In seasonal goats and sheep breeds, onset of puberty is modified by the season of birth. As adult does and bucks from subtropical Mexico display seasonal variation in their reproductive behaviour, this study was carried out to determine the effect of season of birth on puberty. Three groups of each sex born in January, May and October were used. During the seasons, does and bucks were weaned at an age of 30 days and offered ad libitum alfalfa hay and 100 g of commercial concentrate. In the female kids, the onset of ovulatory activity was determined by progesterone plasma concentrations once in a week from 3 months of age until the onset of puberty. In the male kids, the onset of puberty was individually recorded by observing the ability to mount and intromit an induced oestrous female goat aged 3 months and the presence of spermatozoa in the ejaculate obtained in an artificial vagina 1 week after the first mount. In female kids, there was an effect of the season on the date of first ovulation ( $P < 0.001$ ). In the May group, ovulatory activity commenced at an earlier age ( $201 \pm 3$  days) compared with January ( $264 \pm 5$  days) and October ( $344 \pm 5$  days) groups ( $P < 0.001$ ). In the January group also, the ovulatory activity commenced earlier than the October group ( $P < 0.001$ ). In males, an effect of the season of birth on the first mounting was observed ( $P < 0.001$ ). The male kids that were born in May ( $111 \pm 3$ ) and October ( $112 \pm 5$  days) attained puberty earlier than those born in January ( $131 \pm 4$  days;  $P < 0.001$ ). The time of onset of puberty did not differ between groups of May and October. All males showed the presence of spermatozoa in the first ejaculate obtained 1 week after the first mount. The spermatozoa in all ejaculates were immobile. It was concluded that the season of birth modified the onset of puberty in both genders, but these modifications were more pronounced in the female than in the male kid goats.*

**Keywords:** goats, puberty, seasonal variation, subtropics

### Introduction

Puberty can be defined as the first ovulation and/or first oestrous behaviour in females and the first mount and/or ejaculation with the release of sperm in males. Puberty is influenced by several environmental factors (Foster, 1994). There is an interbreed difference in the onset of sexual activity in does and bucks that are well fed. In does of the Shiba breed and in local female goats from the Caribbean Guadeloupe Island, puberty starts from 5.6 to 6.7 months of age, whereas the Saanen does attain puberty at 7.8 months of age (Amoah and Bryant, 1984; Chemineau, 1993; Sakurai *et al.*, 2004). Tokara and Damascus bucks attain puberty at approximately 4 and 17 months of age, respectively (Elwisy and Elsayaf, 1971; Nishimura *et al.*, 2000). In goats and sheep breeds displaying seasonal

reproduction, the onset of puberty in does occur only during the breeding season (Ricoardeau *et al.*, 1984; Papachristoforou *et al.*, 2000). The onset of puberty is influenced by the month of birth and also by photoperiodic changes (Greyling and Van Niekerk, 1990; Foster, 1994). Spring-born Suffolk ewes and Alpine does attain puberty at approximately 30 weeks of age, but autumn-born females attain puberty at approximately 1 year of age (Ricoardeau *et al.*, 1984; Foster, 1994). In contrast, in lambs of seasonal breeds, the season of birth does not appear significantly to affect the onset of puberty (Wood *et al.*, 1991; Herbosa *et al.*, 1995). Testicular growth, an index of spermatogenesis, was not modified in Suffolk rams subjected to an increasing or decreasing photoperiod from birth (Delgado *et al.*, 1995; Herbosa *et al.*, 1995). In spring-born gonadectomised oestradiol-treated rams raised in natural-simulated photoperiod or reversed-simulated photoperiod, the rise in the level of pubertal LH was similar in both groups (Herbosa *et al.*,

<sup>†</sup> E-mail: joaldesa@yahoo.com

1995) or delayed only for approximately 3 weeks in reverse natural-simulated photoperiod (Wood *et al.*, 1991). These results strongly suggest that female sheep and goats of seasonal breeds appear to be more affected by the season of birth compared with males. The local female and male goats from subtropical Mexico display marked seasonal variations in their sexual activity. The breeding season of females lasts from September to March, whereas the sexual activity of males lasts from May to December (Delgadillo *et al.*, 1999; Delgadillo-Sánchez *et al.*, 2003). Therefore, this study was carried out to determine the effect of the season on puberty in female and male goats from subtropical Mexico.

**Material and methods**

*Study site and management of experimental groups*

This study was performed in Torreón, State of Coahuila, Mexico (latitude 26°23'N; longitude 104°47'W). Photoperiod in this region varies from 13 h 41 min of light at summer solstice to 11 h 19 min of light at winter solstice. Three groups of local female and male goats born in different months of the year were used. In the first groups, mean ( $\pm$ s.e.) birth dates of females ( $n=9$ ) and males ( $n=12$ ) were January  $3 \pm 0.5$  days and  $5 \pm 0.4$  days, respectively. In the second groups, females ( $n=18$ ) were born in May  $25 \pm 1.9$  days and males ( $n=20$ ) in May  $29 \pm 1.7$  days. In the third groups, females ( $n=10$ ) and males ( $n=10$ ) were born in October  $30 \pm 0.8$  days and  $28 \pm 1.3$  days, respectively. In each season, animals were weaned at the age of 30 days and fed *ad libitum* with alfalfa hay and 100 g of a commercial concentrate containing 14% of crude protein, 10% of fibre, 2.5 Mj/kg and 62% of total digestible nutrients. Animals had free access to mineral blocks and water. Female and males were separated and allocated in shaded open pens from weaning to puberty.

*Measurements of reproductive activity and body and testicular weights*

**Does.** In all the three experimental groups, live weight was determined every 2 weeks from birth until the onset of puberty. The determination of the onset of ovarian activity commenced from 3 months of age. This determination was carried out once in a week by measurement of the progesterone plasma concentration (Terqui and Thimonier, 1974). Progesterone levels greater than 1 ng/ml in two consecutive samples were considered as indicative of luteal activity, which in turn indicates the onset of puberty.

**Bucks.** In the three experimental groups, live weight was determined every 2 weeks from birth until the onset of puberty. The testicular weight, an index of spermatogenic activity (Delgadillo *et al.*, 1995), was determined every 2 weeks from birth to the end of the study by comparative palpation with an orchidometer (Oldham *et al.*, 1978). Puberty can be defined as the ability of males to mount and intromit an oestrus doe and the presence of spermatozoa in the ejaculate. From 3 months of age,

individual sexual behaviour was assessed once in a week. Males had 5 min following presentation to mount an intact doe (same body size as that of males) artificially induced into oestrus by weekly intramuscular injections of 100  $\mu$ g oestradiol benzoate. One week after the occurrence of mounting, males were allowed to ejaculate into an artificial vagina to determine the presence of spermatozoa.


*Statistical analyses*

Changes in body weight of females and males and testicular weight in males were analysed by a two-way ANOVA with repeated measurements (month of birth and time of experiment). Body weight at birth, average daily gain, body and testicular weights, and age at puberty were analysed by a one-way ANOVA (month of birth). When differences between groups were found, these variables were compared by a Fisher's protected LSD test.

**Results**

*Females*

**Body weight.** The evolution of body weight of the three groups from birth to the onset of puberty is shown in Figure 1. The data for body weight and age at the time



**Figure 1** Increase (mean  $\pm$  s.e.) of body weight of female goats from subtropical Mexico, which were born in January (●), May (■) and October (○) (\* $P < 0.05$ ).

**Table 1** Body weights and age at puberty (mean ± s.e.) of female goats from subtropical Mexico, which were born in January, May and October

	Body weight at birth (kg)	Average daily gain (g)	Body weight at puberty (kg)	Age at puberty (days)
January	3.1 <sup>a</sup> ± 0.17	94 <sup>a</sup> ± 4	27 <sup>a</sup> ± 1.0	264 <sup>a</sup> ± 5
May	3.1 <sup>a</sup> ± 0.10	122 <sup>b</sup> ± 3	28 <sup>a</sup> ± 0.8	201 <sup>b</sup> ± 3
October	2.7 <sup>b</sup> ± 0.5	83 <sup>a</sup> ± 4	32 <sup>b</sup> ± 1.3	344 <sup>c</sup> ± 5


<sup>a,b,c</sup> Means with different superscripts in the same column are significantly different ( $P < 0.05$ ).

of puberty are shown in Table 1. Season of birth had a strong effect on live weight at birth ( $P < 0.05$ ). Females born in October weighed less than those born in January and May ( $P < 0.05$ ). No difference was found between the females born in January and May ( $P > 0.05$ ). Effect of time of experiment ( $P < 0.001$ ) and an interaction between time and group ( $P < 0.001$ ) on live weight were observed, which indicated that groups showed varying responses with time. Pair-wise comparisons showed that the evolution of body weight of females born in May was different from those born in January and October ( $P < 0.001$ , both comparisons). In contrast, no difference was found between the January and October groups. Differences in body weight recorded on a monthly basis are indicated in Figure 1. Live weight at puberty varied according to season ( $P < 0.01$ ). The weights of females of the October group were higher at puberty than those of January and May groups ( $P < 0.01$ , both comparisons). The weight of females of the January group did not differ ( $P > 0.05$ ) from that of the May group. The average daily gain from birth to puberty was also different between groups ( $P < 0.001$ ). This average daily gain was greater in animals born in May than in those born in October and January ( $P < 0.001$ ). The average daily gain in animals born in October did not differ from those born in January ( $P > 0.05$ ).


**Onset of ovarian activity.** The percentages of females attaining puberty are shown in Figure 2. The data concerning the age at puberty are shown in Table 1. There was a highly significant effect of the season on age at first ovulation ( $P < 0.001$ ). In females of the May group, ovulatory activity commenced earlier compared with females of the January and October groups ( $P < 0.001$ ). In females born in January, ovulatory activity commenced earlier compared with females of the October group ( $P < 0.001$ ). The onset of puberty was observed in September, December and October for January, May and October groups, respectively.

**Bucks**

**Body weight.** The evolution of body weight of males of the three groups from birth to puberty is shown in



**Figure 2** Cumulative percentage of female goats from subtropical Mexico attaining puberty, which were born in January (●), May (■) and October (○).



**Figure 3** Increase (mean ± s.e.) of body weight of male goats from subtropical Mexico, which were born in January (●), May (■) and October (○) ( $P < 0.05$ ).

Figure 3. The data for body weight are shown in Table 2. Season had a strong effect on live weight at birth ( $P < 0.001$ ). The body weights of males born in October were less than those born in January ( $P < 0.01$ ) and in May ( $P < 0.001$ ). No difference was found between males of these two groups ( $P > 0.05$ ). An effect of time of

**Table 2** Body and testicular weights, and age at puberty (mean  $\pm$  s.e.) of the local male goats from subtropical Mexico, which were born in January, May and October


	Body weight at birth (kg)	Average daily gain (g)	Body weight at puberty (kg)	Testicular weight at puberty (g)	Age at puberty (days)
January	3.4 <sup>a</sup> $\pm$ 0.2	132 <sup>a</sup> $\pm$ 3	20 <sup>a</sup> $\pm$ 0.7	40 <sup>a</sup> $\pm$ 1.9	131 <sup>a</sup> $\pm$ 4
May	3.6 <sup>a</sup> $\pm$ 0.1	141 <sup>a</sup> $\pm$ 7	19 <sup>a</sup> $\pm$ 0.6	32 <sup>ab</sup> $\pm$ 3.7	111 <sup>b</sup> $\pm$ 3
October	2.6 <sup>b</sup> $\pm$ 0.2	111 <sup>b</sup> $\pm$ 9	15 <sup>b</sup> $\pm$ 0.8	22 <sup>b</sup> $\pm$ 3.9	112 <sup>b</sup> $\pm$ 5

<sup>a,b</sup> Means with different superscripts in the same column are significantly different ( $P < 0.05$ ).

experiment ( $P < 0.001$ ) and an interaction between time and group ( $P < 0.05$ ) on live weight were observed, which indicated that groups showed varying responses with time. The evolution of body weight of the males when compared two by two groups was different throughout the experiment ( $P < 0.05$ ). Pair-wise comparisons showed that body weight of males of the May group differed from those of the January and October groups ( $P < 0.05$ ). There were no differences between the January and October groups ( $P > 0.05$ ). The differences that were observed on a monthly basis are indicated in Figure 3. Live weight at puberty varied according to season ( $P < 0.001$ ). The live weights of males of the October group at puberty were less than those of the January ( $P < 0.01$ ) and May groups ( $P < 0.001$ ). The live weights of males of January did not differ ( $P > 0.05$ ) from those of May group. The average daily gain from birth to puberty was also different between groups ( $P < 0.05$ ). This average was greater in animals born in May than in animals born in October. The average daily gain did not differ ( $P > 0.05$ ) between males born in January and May.

**Testicular weight.** The evolution of testicular weight of males of the three groups from birth to puberty is shown in Figure 4. The data for testicular weight are shown in Table 2. An effect of time of experiment ( $P < 0.001$ ) and an interaction between time and group ( $P < 0.01$ ) on testicular weight were observed, indicating that groups showed varying responses with time. Pair-wise comparisons showed that the evolution of testicular weight differed only between the January and May groups ( $P < 0.01$ ). The differences recorded on a monthly basis are indicated in Figure 4. Testicular weight at puberty varied according to season ( $P < 0.05$ ). Males of the October group that attained puberty had smaller testicular weights compared with those born in January ( $P < 0.01$ ). No difference was registered between males born in October *v.* May, or between males born in May *v.* January.

**Attainment of puberty.** Age at first mounting with intromission varied significantly with the season of birth



**Figure 4** Increase (mean  $\pm$  s.e.) of unpaired testicular weight of male goats from subtropical Mexico, which were born in January (●), May (■) and October (○) (\*\* $P < 0.01$ ).

( $P < 0.001$ ; Table 2). The onset of puberty occurred earlier in May- and October-born males than in January-born males ( $P < 0.001$ ). Puberty did not differ ( $P > 0.05$ ) between May- and October-born males. All males showed presence of spermatozoa in the first ejaculate obtained 1 week after the first mount. The spermatozoa did not show mobility in any of the ejaculate.

**Discussion**

The results of this study show that in male and female goats from the local areas of subtropical Mexico, the season of birth considerably modified the onset of puberty. The effect of season of birth on puberty was more pronounced in females than in males. Regardless of the season of birth, male goats attained puberty at a younger age than did females.

Females in the January and May groups attained puberty in September and December (the same year of birth), respectively. The body weight at puberty was similar in both groups (about 27 kg) but females in the January group were about 9 weeks older than those in the May group (approx. 38 *v.* 29 weeks old). Onset of puberty in both groups

occurred probably because they had perceived enough increasing days, allowing the onset of puberty during the decreasing days of the autumn, after having reached a sufficient body weight. The difference of age in attaining puberty in the January and May groups was probably because of the difference in body weight at the beginning of the natural breeding season, which lasts from September to March (Delgadillo *et al.*, 2004a). Females in the January group with higher mature body weight attained puberty in September, whereas those in the May group should have attained a body size so that puberty commenced in December, similar to those reported in Suffolk ewes and Alpine goats (Ricoardeau *et al.*, 1984; Foster, 1994). October-born females also reached puberty in the autumn (October) of the subsequent year. In fact, they were approximately 20 and 10 weeks older than females in the January and May groups, respectively, with highest (32 kg) body weight. Delayed puberty in this group was probably attributed to the fact that they required to perceive the increasing days from spring and summer to start puberty during the short days of the autumn (Foster, 1994). In females, the effect of season on puberty observed in our study agrees with data reported in reproductive seasonal breeds of goats and sheep, in which puberty starts only during the breeding season of the adult females (Ricoardeau *et al.*, 1984; Forcada *et al.*, 1991; Greyling, 2000). In these seasonal breeds, the long days of spring and summer followed by short days of autumn provide the stimulatory signal to initiate puberty, when the animals have achieved the appropriate body growth to initiate ovulation (Foster, 1994; Adam *et al.*, 1998).

In our study, despite the fact that all animals were fed *ad libitum*, there was an effect of season on average body daily gain. This was already reported in *ad libitum*-fed young or adult animals when subjected to increasing or long days, compared with those raised in decreasing or short days (Greyling and Van Niekerk, 1990; Wood *et al.*, 1991; Delgadillo *et al.*, 2004b). The differences in growth rates cannot explain the difference in puberty between the January and October groups because they had a similar growth rate, but puberty appeared later in the October group. In spring-born Soay ewe lambs receiving a diet *ad libitum* or at maintenance, puberty was not delayed and both groups started ovulatory cycles in autumn in spite of a different body weights (Adam *et al.*, 1998). In our study, the difference in age and body weight at puberty were probably because of the season of birth, then the different photoperiodic conditions in which they were raised: January- and May-born groups received enough long days during winter and spring and attained puberty during the short days of autumn in the same year. In contrast, the October group must first perceive increasing days to reach puberty during the decreasing days of autumn of the following year, as reported in other seasonal breeds (Foster, 1994). In Alpine does, for example, the onset of puberty occurs at about 45 and 36 weeks for females born in October and January, respectively (Ricoardeau *et al.*, 1984). The age at the onset of

puberty of female goats from subtropical Mexico born in winter, spring or autumn is similar to that reported in different breeds of small ruminants from temperate and subtropical latitudes (Foster, 1994; Lassoued and Reikik, 2001; Freitas *et al.*, 2004).

In males, puberty was slightly modified by the season of birth. May- and October-born males attained puberty at the same age with a different body weight and similar testicular weight despite being born during increasing or decreasing day length, respectively. These two groups attained puberty earlier than January-born males, which were born during increasing days. This weight of January group at puberty was higher than that of May and October groups. The lower body weight at birth and the lower average body daily gain of October-born males does not appear to have influenced the onset of puberty because the males of this group attained puberty at the same age as the May group, which showed a higher growth rate. This conclusion is reinforced by their similar average daily gain: the January group was older at puberty than the May group having similar body and testicular weights. These data suggest that in subtropical Mexican male goats, puberty was not strongly dependent on photoperiodic changes, although adult males display a reproductive seasonality controlled mainly by photoperiodic changes (Delgadillo *et al.*, 1999 and 2004b). In contrast to females, puberty of Mexican male goats does not appear to be controlled by photoperiodic changes, as reported previously in other breeds. In gonad-intact Ile-de-France and Suffolk rams, changes in testicular weight, an index of spermatogenesis, or the rise in LH secretion are not modified when subjected after birth to increasing or decreasing photoperiod, or to constant long days (Colas *et al.*, 1987; Herbosa and Foster, 1996). In spring-born gonadectomised oestradiol-treated lambs raised in natural or artificially reversed photoperiod, rise of pubertal LH is observed to be similar or slightly delayed for approximately 3 weeks in reverse natural simulated-photoperiod (Wood *et al.*, 1991; Herbosa *et al.*, 1995). Considered together, these results suggest that at least in the above-mentioned breeds and in young males from subtropical Mexico, puberty started indifferently during increasing or decreasing days, which indicates that the influence of photoperiod on the onset of puberty was less. More generally, the age at puberty in local male goats from subtropical Mexico born in winter, spring or autumn is lower than that reported in various breeds of small ruminants from temperate and subtropical latitudes (Elwishy and Elswaf, 1971; Chakraborty *et al.*, 1989; Özsar *et al.*, 1990; Ahmad and Noakes, 1996).

The difference in puberty between males and females may be because of a different sensibility of the hypothalamo-pituitary axis to steroid negative feedback. In both the sexes, the onset of puberty originates in a decrease in the sensitivity to the negative feedback of oestradiol in females and testosterone in males (Claypool and Foster, 1990; Foster, 1994). It was shown that this decreased sensitivity to steroids occurred later in females than in males,

provoking an earlier onset of puberty in males (Claypool and Foster, 1990; Foster, 1994; Herbosa *et al.*, 1995). This difference is probably because of a different response of males and females to photoperiodic information. In females, decreasing day length is necessary to attain puberty, whereas in males, it appears that decreasing day length is not necessary to attain the puberty (Wood *et al.*, 1991).

Results of this study show that in these subtropical conditions, the season of birth affects the onset of puberty. This fact must be taken into account because it is usually hypothesised that in subtropical conditions, level of nutrition is the main environmental factor controlling the time of the breeding season rather than that of puberty (Walkden-Brown and Bocquier, 2000). Our data challenged this conclusion by showing that the onset of puberty is dependent on season of birth and that this effect of season is not obtained through seasonal changes in availability of food (birth is reached at different weights depending on the season). Our data, therefore, clearly indicate that, in addition to the availability of food, other environmental factors such as photoperiod must be considered.

In conclusion, these data show that in local goats from subtropical Mexico, season of birth influences the onset of puberty. In does, puberty depends on photoperiodic changes, and the onset of puberty occurs only during the short days of autumn during the natural breeding season. In contrast, puberty in male goats does not appear to depend on photoperiodic changes. Bucks attain puberty either during the long or during the short days.

### Acknowledgments

This work was supported by the International Foundation of Science (Grant B/2071-3F). We express our thanks to the Assay Laboratory of the Station de Physiologie de la Reproduction et des Comportements of Nouzilly, France, for carrying out the radio-immunoassays, to B. Malpoux and P. Chemineau for their comments to improve the manuscript, and to D. López for her secretarial help. M.A. De Santiago was supported by a CONACyT scholarship for her Postgraduate studies.

### References

Adam CL, Findlay PA, Kyle CE and Young P 1998. Effect of restricted nutrition on timing of puberty in female Soay sheep. *Journal of Reproduction and Fertility* 112, 31–37.

Ahmad N and Noakes DE 1996. Sexual maturity in British breeds of goat kids. *British Veterinary Journal* 152, 93–103.

Amoah EA and Bryant MJ 1984. A note on the effect of contact with male goats on occurrence of puberty in female goat kids. *Animal Production* 38, 141–144.

Chakraborty PK, Stuart LD and Brown JL 1989. Puberty in the male Nubian goat: serum concentrations of LH, FSH and testosterone from birth through puberty and semen characteristics at sexual maturity. *Animal Reproduction Science* 20, 91–101.

Chemineau P 1993. Reproducción de las cabras originarias de las zonas tropicales. *Revista Latinoamericana de Pequeños Ruminantes* 1, 2–14.

Claypool LE and Foster DL 1990. Sexual differentiation of the mechanism controlling pulsatile secretion of luteinizing hormone contributes

to sexual differences in the timing of puberty in sheep. *Endocrinology* 126, 1206–1215.

Colas G, Guerin Y, Brios M and Ortavant R 1987. Photoperiodic control of testicular growth in the ram lamb. *Animal Reproduction Science* 13, 255–262.

Delgadillo JA, Hochereau-de Reviers MT, Daveau A and Chemineau P 1995. Effect of short photoperiodic cycles on male genital tract and testicular parameters in male goats (*Capra hircus*). *Reproduction Nutrition Development* 35, 549–558.

Delgadillo JA, Canedo GA, Chemineau P, Guillaume D and Malpoux B 1999. Evidence for an annual reproductive rhythm independent of food availability in male creole goats in subtropical Northern Mexico. *Theriogenology* 52, 727–737.

Delgadillo-Sánchez JA, Flores-Cabrera JA, Véliz-Deras FG, Duarte-Moreno G, Vielma-Sifuentes J, Poindron-Massot P and Malpoux B 2003. Control de la reproducción de los caprinos del subtropico mexicano utilizando tratamientos fotoperiódicos y efecto macho. *Veterinaria México* 34, 69–79.

Delgadillo JA, Fitz-Rodríguez G, Duarte G, Véliz FG, Carrillo E, Flores JA, Vielma J, Hernández H and Malpoux B 2004a. Management of photoperiod to control caprine reproduction in the subtropics. *Reproduction Fertility and Development* 16, 471–478.

Delgadillo JA, Cortez ME, Duarte G, Chemineau P and Malpoux B 2004b. Evidence that photoperiod controls the annual changes in testosterone secretion, testicular and body weight in subtropical male goats. *Reproduction Nutrition and Development* 44, 183–193.

Elwishi AB and Elsawaf SA 1971. Development of sexual activity in male Damascus goats. *Indian Journal of Animal Science* 41, 350–356.

Forcada F, Abecia JA and Zarazaga L 1991. A note on attainment of puberty of September-born early-maturing ewe lambs in relation to the level of nutrition. *Animal Production* 53, 407–409.

Foster DL 1994. Puberty in sheep. In *The physiology of reproduction* (eds E Knobil and JD Neill), pp. 411–451. Raven Press Ltd, New York, USA.

Freitas VJF, Lopes-Junior ES, Rondina D, Salmato-Vanderley CSB, Salles HO, Simplicio AA, Baril G and Saumande J 2004. Puberty in Anglo-Nubian and Saanen female kids raised in the semi-arid of north-eastern Brazil. *Small Ruminant Research* 53, 167–172.

Greyling JPC 2000. Reproduction traits in the Boer goat doe. *Small Ruminant Research* 36, 171–177.

Greyling PC and Van Niekerk CH 1990. Puberty and induction of puberty in female Boer goat kids. *South African Journal of Animal Science* 20, 193–200.

Herbosa CG and Foster DL 1996. Defeminization of the reproductive response to photoperiod occurs early in prenatal development in the sheep. *Biology of Reproduction* 54, 420–428.

Herbosa CG, Wood RI and Foster DL 1995. Prenatal androgens modify the reproductive response to photoperiod in the developing sheep. *Biology of Reproduction* 52, 163–169.

Lassoued N and Rezik M 2001. Differences in reproductive efficiency between female sheep of the Queue Fine de l'Ouest purebred and their first cross with the D'Man. *Animal Research* 50, 373–381.

Nishimura S, Okano K, Yasukouchi K, Gotoh T, Tabata S and Iwamoto H 2000. Testis developments and puberty in the male Tokara (Japanese native) goat. *Animal Reproduction Science* 64, 127–131.

Oldham CM, Adams NR, Gherardi PB, Lindsay DR and McKintosh JB 1978. The influence of level of feed intake on sperm-producing capacity of testicular tissue in the ram. *Australian Journal of Agricultural Research* 29, 173–179.

Özsar S, Güeven B, Celebi M, Kalkandelen G and Van de Wiel DFM 1990. Testosterone and LH concentrations in the male Angora goat during puberty. *Animal Reproduction Science* 23, 319–326.

Papachristoforou C, Koumas A and Photiou C 2000. Seasonal effect on puberty and reproductive characteristics of female Chios sheep and Damascus goats born in autumn or in February. *Small Ruminant Research* 38, 9–15.

Ricordeau G, Bouillon J, Gaillard A, Lajous A and Lajous D 1984. Modalités et caractéristiques de reproduction chez les caprins aspects génétiques. *Bulletin Technique d'Insemination* 391, 367–382.

Sakurai K, Ohkura S, Matsuyama S, Katoh K, Obara Y and Okamura H 2004. Body growth and plasma concentrations of metabolites and metabolic

## Delgadillo, De Santiago-Miramontes and Carrillo

hormones during the pubertal period in the female Shiba goats. *Journal of Reproduction and Development* 50, 197–205.

Terqui M and Thimonier J 1974. Nouvelle méthode radio-immunologique rapide pour l'estimation du niveau de progestérone plasmatique. Application pour le diagnostic précoce de la gestation chez la brebis et la chèvre. *CR Academie de Sciences, Paris, D* 279, 1109–1112.

Walkden-Brown SW and Bocquier F 2000. Nutritional regulation of reproduction in goats. *Proceedings of the Seventh International Conference on Goats, Tours, France*, pp. 389–395.

Wood RI, Ebling FJP, l'Anson H and Foster DL 1991. The timing of neuroendocrine sexual maturity in the male lamb by photoperiod. *Biology of Reproduction* 45, 82–88.