

Coccidial oocyst release: once a day or all day long? Tropical bird hosts shed new light on the adaptive significance of diurnal periodicity in parasite output

Research Article

Cite this article: Biard C, Monceau K, Teixeira M, Motreuil S, Bettencourt-Amarante S, Develay L, Moreau J (2022). Coccidial oocyst release: once a day or all day long? Tropical bird hosts shed new light on the adaptive significance of diurnal periodicity in parasite output. *Parasitology* **149**, 469–481. <https://doi.org/10.1017/S003118202100202X>

Received: 5 September 2021

Revised: 9 November 2021

Accepted: 9 November 2021

First published online: 24 November 2021




Keywords:

Birds; circadian rhythms; coccidia; host–parasite interactions; temperate forest; tropical forest

Author for correspondence:

Jérôme Moreau,

E-mail: jerome.moreau@u-bourgogne.fr

Clotilde Biard¹ , Karine Monceau² , Maria Teixeira³, Sébastien Motreuil³, Soline Bettencourt-Amarante³, Lucie Develay³ and Jérôme Moreau^{2,3} 

¹Sorbonne Université, UPEC, Paris 7, CNRS, INRA, IRD, Institut d'Écologie et des Sciences de l'Environnement de Paris, F-75005 Paris, France; ²Centre d'Études Biologiques de Chizé, UMR 7372, CNRS and La Rochelle Université, 79360 Villiers-en-Bois, France and ³UMR CNRS 6282 Biogéosciences, Université Bourgogne Franche-Comté, 6 Boulevard Gabriel, 21000 Dijon, France

Abstract

Many parasites spend part of their life cycle as infectious forms released from an infected host in the external environment, where they may encounter and infect new hosts. The emergence of infectious life stages often occurs once a day to minimize mortality in adverse environments. In bird hosts, intestinal parasites such as coccidia are generally released with feces in the late afternoon. This dynamic is adaptive since it allows avoiding desiccation and ultraviolet (UV) radiation, thus reducing mortality of oocysts in the environment until transmission to the next host. If this circadian rhythm is the result of natural selection to increase oocyst survival, we may hypothesize that oocysts will appear in feces at different times depending on the environment where hosts live. Particularly, in an environment where UV radiation and desiccation are very low, we may expect oocyst circadian release to disappear since the main selective pressure would be relaxed. We sampled different species of birds in tropical and temperate forests in spring and investigated coccidian oocyst output. A strong circadian variation in the prevalence of hosts shedding coccidian oocyst was detected for species caught in the temperate forest with an increase in prevalence in the late afternoon, whereas prevalence of birds shedding oocysts was constant over the course of the day for most species sampled in the tropical rain forest. These results are consistent with the hypothesis that oocysts' circadian output is maintained by natural selection to increase oocyst survival. We discuss the adaptive significance of diurnal periodicity in parasite output.

Introduction

From an evolutionary perspective, natural selection should favour any trait increasing the probability of parasite transmission from one host to another (Bush *et al.*, 2001; Combes, 2001). For example, many parasite species with complex life cycles (i.e. parasites with successive host species), have evolved strategies of releasing infective forms at times of the day when the definitive host is most likely to be present in the environment (Théron, 1984; Combes *et al.*, 1994; Combes, 2001 and references therein). Such a phenomenon of diurnal periodicity is also observed in parasites with simple life cycles (i.e. parasites with a single host species). Coccidia from the genus *Isospora* (Protozoa, Apicomplexa) are intracellular intestinal parasites of birds, and typically peak oocyst release in the late afternoon (Boughton, 1933; Kruszewicz, 1995; Brawner and Hill, 1999; Dolnik, 1999a, 1999b; Hudman *et al.*, 2000; Brown *et al.*, 2001; Misof, 2004). These protozoans exhibit both sexual and asexual phases. In most species, one or more asexual multiplication cycles are completed in the intestinal epithelium of the host, before sexual development occurs (Olsen, 1974). Zygotes produced during the sexual phase transform into oocysts that are then released into the host's intestinal tract and excreted within feces. To complete their development, oocysts must sporulate outside the host to become infective before being ingested by their new host (see e.g. Abd-Al-Aal *et al.*, 2000).

In birds, coccidian oocyst shedding shows a strong circadian pattern that appears to be ubiquitous since in all species tested to date (16 species), oocyst shedding has been found to peak in the afternoon (Knight *et al.*, 2018), whatever the latitude considered, from the tropics zone to arctic area (Dolnik *et al.*, 2011). Even in an extreme environment with permanent daylight like the arctic, this periodicity is conserved despite the absence of diurnal periodicity in hosts' activity (Dolnik *et al.*, 2011). The peak in oocyst shedding late in the afternoon is considered an adaptive trait providing resistance to desiccation and ultraviolet (UV) radiation for oocysts in the external environment (Martinaud *et al.*, 2009). Indeed, a single hour of exposure of the host's feces to natural sunlight reduces the viability of oocysts by ca. 50%, which further decreases with continued exposure (Martinaud *et al.*, 2009).



Fig. 1. Location of the open area inside the Tropical Forest 1 – French Guyana, Nouragues French research, Saut-Pararé station (copyright: Florian Jeanne).

Therefore, releasing oocysts throughout the day would result in a large proportion quickly dying due to sunlight exposure, and a decrease in transmission probability to the next host. Interestingly, all bird species showing temporal variation in the shedding of coccidia oocysts with a peak late in the afternoon roughly have in common one characteristic of their environment: a ground exposed to strong solar radiation including the arctic area (Knight *et al.*, 2018). However, an exciting question that remains open is whether oocysts shedding in an environment where the sun reaching the ground is dramatically reduced throughout the year still shows diurnal periodicity. Indeed, in such environments, the selective pressure of desiccation and UV exposure of oocysts during the daytime would be relieved. One would, therefore, predict periodicity in oocyst shedding either not to evolve or be lost as it would reduce transmission probability without the benefits of increasing oocysts viability and should thus be selected against.

Although tropical rainforests receive 12 h of sunlight daily, it is generally assumed that <2% of radiation ever reach the ground. Tropical rainforests have dense vegetation, often forming three different layers including a dense canopy that blocks out ca. 95% of sunlight (Brinkmann, 1971; Bourgeron, 1983; Chazdon and Fetcher, 1984; Vazquez-Yanes *et al.*, 1990). Therefore, the rainforest ground rarely receives direct sunlight. Also, tropical rainforests are characterized by high air humidity, ranging on average from 80 to 90% relative humidity (RH; Freiberg, 1997; Bruno *et al.*, 2006) and soil moisture. By combining low direct sunlight and high RH, this environment provides feces shed by birds with very favourable conditions to allow oocyst sporulation and survival. If temporal variation in shedding of coccidial

oocysts is under natural selection to avoid desiccation and UV radiation as hypothesized, we may predict that no circadian rhythm should be observed in tropical forests contrary to temperate forests. Indeed, temperate forests are exposed to a stronger irradiance and lower soil moisture compared to tropical forests since the vegetation canopy is moderately dense and allows light to penetrate (Szwagrzyk *et al.*, 2001). However, to our knowledge, temporal variation in the shedding of coccidia oocysts has never been investigated in a tropical rain forest environment.

To investigate whether periodically synchronized release of coccidia oocysts from the genus *Isospora* may be absent in tropical forests, consistent with the hypothesis of relaxed natural selection linked to UV radiation and desiccation, we compared patterns of oocyst shedding from birds living in the primary rain forest in French Guyana (low luminosity and high humidity and moisture) and that from birds living in open temperate forests in mainland France (high ground luminosity and low humidity and moisture). The generality of this expected difference in the patterns of oocyst shedding across the diversity in host ecology and life history was assessed by sampling a variety of bird species belonging to different families within both habitat types. We expected marked temporal variation in the shedding of coccidia oocysts for host species living in open forest and no or weak temporal variation for hosts living inside the dark rain forest. Besides, we took advantage of the spatial heterogeneity in vegetation cover existing around the field station located in the tropical rain forest to study variation in oocyst-shedding patterns at a smaller scale. Indeed, a large area has been maintained open at the edge of the field station for more than 30 years (Fig. 1) and has selected bird species specialists of open habitats. We, therefore, expected temporal

variation in the shedding of coccidia oocysts from host species inhabiting this open area, contrary to those inhabiting the close by rain forest.

Materials and methods

Trapping birds and sampling feces

Adult males and females of different bird species (1-year-old and older, identified by plumage characteristics) were caught using mist-nets from sunrise to sunset in both tropical and temperate areas. Species and age were determined according to handbooks (Hilty, 2002; Restall *et al.*, 2006a, 2006b). The full list of species and the number of individuals caught are given in Table 1. In the temperate forest habitat, we sampled birds in a young deciduous forest located in Sologne (thereafter called Temperate Forest, France, 47°26'N, 2°13'E) during spring 2019 (10 sampling days). At that time of the season, sunrise occurred at 06:00 am and sunset around 09:00 pm. In the tropical forest, we sampled birds in the Amazonian primary rain forest during July and August 2018 and October 2019 (15 and 11 sampling days in total) in the two field sites of the Nouragues French research station: Saut-Pararé (thereafter Tropical Forest 1, French Guyana, 4°02'N, 52°41'W, 8 sampling days in 2018 and 11 sampling days in 2019) and Inselberg (thereafter Tropical Forest 2, French Guyana, 4°05'N, 52°41'W, 7 sampling days in 2018) where sunrise occurred at 06:00 am and sunset at 06:00 pm. In the first site, we sampled birds both within the forest undergrowth and at its edge in the open area close to the field camp (Fig. 1). On one typical sunny day, we measured abiotic parameters at the ground level in these two areas (open and undergrowth): temperature (°C), UV index and luminosity intensity (in lux). These measures were performed five times over the course of the day (08:00 am, 10:00 am, 12:00 am, 15:00 pm and 17:30 pm), using a CROSCAAL phone – Trekker – X4 equipped with software X4 sensors in ten different spots randomly chosen to take spatial heterogeneity into account.

For birds that can be kept in captivity throughout their lives such as poultry, the preferred method to track temporal patterns of oocysts release as accurately as possible is to collect feces every 2 h from sunrise to sunset for each bird maintained in an individual cage (Filipiak *et al.*, 2009). However, under field conditions, enclosing wild birds in a cage for an entire day is not acceptable from an ethical point of view, due to the stress of capture and prolonged restraint, which may alter both their subsequent survival and ability to reproduce, as birds might have been breeding at the time of capture. To limit the stress imposed on birds, we followed the protocol adopted in snow bunting (Dolnik *et al.*, 2011). Birds were placed in individual cages, the bottom of which was lined with aluminium foil, at any time of capture during the day, for a maximum of 1 h before being released at their site of capture, had the birds shed feces or not. All fresh feces present on the aluminium foil were collected and homogenized with a small spoon and stored at 4°C in 15 mL Falcon tubes labelled with the date, time of the day and bird species. To avoid repeated sampling of the same individuals, birds having shed feces were marked with a drop of non-toxic nail varnish on the tarsus and in the case of recapture, were not subjected again to feces collection and were thus immediately released.

Prevalence and intensity of parasite infection by *Isospora* spp.

At the end of each field session in the two habitats (temperate/tropical), feces samples were weighted to the nearest 0.01 g using an electronic balance and a solution of Sheater (45%

sugar in water) was added to a concentration of 1 mg of feces for 4 mL of sheater, after which samples were gently homogenized. We sampled 600 µL of this solution on a McMaster chamber and allowed oocysts to float to the top for 10 min. All the oocysts of *Isospora* spp. under the grid of each chamber in the McMaster were counted using 10× magnification. Two counts were obtained for each individual fecal sample, and the average concentration, expressed as the number of oocysts per gram of fecal sample, was used as an estimate of oocyst production rate.

Statistical analyses

The time of feces collection was first calculated for each bird as the average hour between entrance and release from the cage. This time was standardized in the proportion of daily sunshine duration to take differences in daylength between tropical and temperate sites into account as follows:

$$h_{\text{standardized}} = \frac{h_{\text{average}} - h_{\text{sunrise}}}{h_{\text{sunset}} - h_{\text{sunrise}}}$$

with h_{average} the average hour of collection and h_{sunrise} and h_{sunset} the sunrise and sunset time. Additionally, to compare morning and afternoon prevalence and infection intensity, daylength was divided into two parts: morning: standardized hour <0.5 and afternoon: standardized hour >0.5. All statistical analyses were performed with R software (v.4.0.4, R Core Team, 2021) implemented with packages *MASS*, *car* for model comparison and for unbalanced design and *emmeans* for *post hoc* pairwise comparison tests including the Benjamini and Hochberg' correction for multiple comparisons (Benjamini and Hochberg, 1995).

Differences in prevalence and parasite load between temperate (Temperate Forest) and tropical environments (Tropical Forest 1 + Tropical Forest 2) in birds sampled in the forest (birds caught in the open area in Tropical Forest 1 were excluded from this analysis) were analysed using a generalized linear mixed-effects model with binomial distribution (B-GLMM, using *glmer* function) and negative binomial distribution (NB-GLMM, using *glmmTMB* function), respectively, including either the standardized hour or part of the day, habitat (temperate vs tropical) and their interaction as fixed effects. Species identity was included as a random effect. The slopes of fitted lines/curves were provided assorted with their 95% confidence interval (CI) from the models. Similar models were run to compare the two sites in the tropical habitat (i.e. Tropical Forest 1 vs Tropical Forest 2).

The same procedures were used to investigate differences in prevalence and parasite load in birds from open area vs surrounding undergrowth in Tropical Forest 1.

Differences in abiotic factors during the day (temperature, UV radiation and luminosity) between these two habitats were analysed including the standardized hour of the day as a quadratic effect to test for a potential maximal effect at midday and the location and their interaction. The temperature was analysed using a Gaussian GLM (*glm* function) whereas UV radiation and luminosity were analysed using NB-GLM (*glm.nb* function).

For all GLMs and GLMMs, model comparisons were made using a deviance analysis for unbalanced design based on the likelihood ratio χ^2 -statistics or *F*-ratio for Gaussian GLM to test the contribution of the variable. Interactions were discarded when non-significant.

Table 1. Complete list of bird species included in this study, showing prevalence of infection by *Isoospora* spp. (total number of birds caught and number of birds infected in brackets) separately for birds caught in the morning (hours standardized <0.5) and in the afternoon (hours standardized >0.5)

Species	Standardized hour		
	(0:0.5)	(0.5:1)	
Temperate forest – Souesmes	99 (14)	112 (95)	
Acrocephalidae			
<i>Hippolais polyglotta</i>	Melodious warbler	1 (1)	4 (4)
Aegithalidae			
<i>Aegithalos caudatus</i>	Long-tailed tit	1 (0)	4 (4)
Certhiidae			
<i>Certhia brachydactyla</i>	Short-toed tree creeper	4 (0)	2 (1)
Emberizidae			
<i>Emberiza citrinella</i>	Yellowhammer	0	1 (1)
Fringillidae			
<i>Fringilla coelebs</i>	Common chaffinch	5 (1)	7 (7)
<i>Carduelis cannabina</i>	Common linnet	1 (0)	0
<i>Pyrrhula pyrrhula</i>	Eurasian bullfinch	2 (0)	5 (3)
<i>Carduelis chloris</i>	European greenfinch	3 (0)	0
Locustellidae			
<i>Locustella naevia</i>	Common grasshopper-warbler	2 (0)	0
Motacillidae			
<i>Anthus trivialis</i>	Tree pipit	0	2 (2)
Muscicapidae			
<i>Phoenicurus ochruros</i>	Black redstart	1 (0)	0
<i>Luscinia megarhynchos</i>	Common nightingale	2 (0)	2 (2)
<i>Phoenicurus phoenicurus</i>	Common redstart	0	1 (1)
<i>Erithacus rubecula</i>	European robin	11 (1)	4 (4)
<i>Saxicola torquatus</i>	European stonechat	1 (0)	1 (1)
Paridae			
<i>Parus caeruleus</i>	Eurasian blue tit	9 (1)	14 (11)
<i>Parus major</i>	Great tit	6 (3)	8 (4)
<i>Parus palustris</i>	Marsh tit	2 (1)	3 (2)
Passeridae			
<i>Passer domesticus</i>	House sparrow	3 (0)	11 (11)
Phylloscopidae			
<i>Phylloscopus collybita</i>	Common chiffchaff	10 (0)	8 (6)
<i>Phylloscopus trochilus</i>	Willow warbler	0	1 (0)
Picidae			
<i>Jynx torquilla</i>	Eurasian wryneck	1 (0)	2 (2)
Prunellidae			
<i>Prunella modularis</i>	Dunnoek	3 (1)	3 (1)
Sittidae			
<i>Sitta europaea</i>	Eurasian nuthatch	4 (1)	1 (1)
<i>Sturnus vulgaris</i>	Common starling	2 (0)	0
Sylviidae			
<i>Sylvia communis</i>	Common whitethroat	4 (0)	9 (9)
<i>Sylvia atricapilla</i>	Eurasian blackcap	17 (4)	13 (12)
<i>Sylvia borin</i>	Garden warbler	3 (0)	4 (4)

(Continued)

Table 1. (Continued.)

Species		Standardized hour	
		(0:0.5)	(0.5:1)
Troglodytidae			
<i>Troglodytes troglodytes</i>	Eurasian wren	0	1 (1)
Turdidae			
<i>Turdus philomelos</i>	Song thrush	1 (0)	1 (1)
Tropical Forest 1 – Pararé		79 (17)	74 (17)
Cardinalidae			
<i>Cyanocopsa cyanooides</i>	Blue-black grosbeak	1 (0)	0
Columbidae			
<i>Leptotila rufaxilla</i>	Grey-fronted dove	0	1 (0)
Cotingidae			
<i>Lipaugus vociferans</i>	Screaming piha	1 (0)	1 (0)
Formicariidae			
<i>Formicarius analis</i>	Black-faced antthrush	1 (0)	0
Furnariidae		14 (3)	14 (5)
<i>Xiphorhynchus pardalotus pardalotus</i>	Chestnut-rumped woodcreeper	1 (0)	1 (1)
<i>Automolus infuscatus</i>	Olive-backed foliage-gleaner	4 (1)	0
<i>Xenops minutus</i>	White-throated xenops	0	1 (0)
<i>Philydor erythrocerum</i>	Rufous-rumped foliage-gleaner	1 (0)	0
<i>Sclerurus rufigularis</i>	Short-billed leaf-tosser	2 (1)	0
<i>Glyphorhynchus spirurus spirurus</i>	Wedge-billed woodcreeper	6 (1)	12 (4)
Grallariidae			
<i>Hylopezus macularius</i>	Spotted antpitta	0	1 (0)
Pipridae		9 (1)	14 (0)
<i>Pipra erythrocephala</i>	Golden-headed manakin	1 (0)	0
<i>Manacus manacus manacus</i>	White-bearded manakin	1 (1)	2 (0)
<i>Dixiphia pipra pipra</i>	White-crowned manakin	2 (0)	7 (0)
<i>Lepidothrix serena</i>	White-fronted manakin	2 (0)	3 (0)
<i>Corapipo gutturalis</i>	White-throated manakin	3 (0)	2 (0)
Poliophtilidae			
<i>Ramphocaenus melanurus</i>	Long-billed gnatwren	1 (0)	0
Thamnophilidae		29 (8)	26 (8)
<i>Hypocnemoides melanopogon</i>	Black-chinned antbird	0	1 (0)
<i>Percnostola rufifrons rufifrons</i>	Black-headed antbird	4 (0)	0
<i>Frederickena viridis</i>	Black-throated antshrike	1 (0)	0
<i>Myrmotherula (epinecrophylla) gutturalis</i>	Brown bellied antwren	4 (2)	1 (1)
<i>Thamnomanes caesius</i>	Cinereous antshrike	1 (0)	3 (1)
<i>Willisornis poecilinotus</i>	Common scale-backed antbird	2 (1)	1 (0)
<i>Thamnomanes ardesiacus</i>	Dusky-throated antshrike	0	2 (2)
<i>Cercomacra cinerascens</i>	Grey antbird	0	1 (0)
<i>Myrmotherula menestriesii</i>	Grey antwren	1 (0)	4 (2)
<i>Myrmotherula longipennis</i>	Long-winged antwren	0	5 (0)
<i>Thamnophilus murinus</i>	Mouse-coloured antshrike	2 (0)	0
<i>Isleria guttata</i>	Rufous-bellied antwren	2 (1)	1 (0)
<i>Hylophylax naevius naevius</i>	Spot-backed antbird	2 (1)	0

(Continued)

Table 1. (Continued.)

Species		Standardized hour	
		(0:0.5)	(0.5:1)
<i>Hypocnemis cantator</i>	Warbling-antbird	1 (1)	0
<i>Myrmotherula axillaris</i>	White-flanked antwren	2 (0)	2 (0)
<i>Pithys albifrons albifrons</i>	White-plumed antbird	7 (2)	5 (2)
Thraupidae		4 (0)	0
<i>Tachyphonus surinamus surinamus</i>	Fulvous-crested tanager	2 (0)	0
<i>Lanio fulvus</i>	Fulvous shrike-tanager	2 (0)	0
Tityridae		5 (0)	3 (1)
<i>Onychorhynchus coronatus</i>	Royal Flycatcher	0	1 (0)
<i>Terentotriccus erythrurus</i>	Ruddy-tailed flycatcher	1 (0)	0
<i>Myiobius barbatus</i>	Sulphur-rumped flycatcher	4 (0)	2 (1)
Troglodytidae			
<i>Pheugopedius coraya</i>	Coraya wren	3 (1)	0
Turdidae			
<i>Turdus albicollis</i>	White-necked thrush	4 (1)	0
Tyrannidae		6 (2)	13 (3)
<i>Platyrhynchus saturatus</i>	Cinamon-crested spadebill	0	1 (0)
<i>Platyrhynchus coronatus</i>	Golden-crowned spadebill	1 (0)	1 (0)
<i>Mionectes macconnelli</i>	McConnell's flycatcher	5 (2)	6 (2)
<i>Rhynchocyclus olivaceus</i>	Olivaceous flatbill	0	3 (1)
<i>Corythopsis torquatus</i>	Ringed antpipit	0	2 (0)
Vireonidae			
<i>Hylophilus ochraceiceps</i>	Tawny-crowned greenlet	1 (1)	1 (0)
Tropical Forest 1 open area - Pararé		9 (0)	16 (10)
Fringillidae		0	4 (3)
<i>Euphonia cayennensis</i>	Golden-sided euphonia	0	1 (1)
<i>Euphonia violacea</i>	Violaceous euphonia	0	3 (2)
Thraupidae		6 (0)	9 (7)
<i>Coereba flaveola</i>	Bananaquit	0	1 (1)
<i>Thraupis episcopus</i>	Blue-grey tanager	0	1 (0)
<i>Saltator maximus</i>	Buff-throated saltator	1 (0)	0
<i>Sporophila lineola</i>	Lined seedeater	0	1 (0)
<i>Ramphocelus carbo carbo</i>	Silver-beaked tanager	5 (0)	6 (6)
Tyrannidae		2 (0)	2 (0)
<i>Pitangus sulphuratus</i>	Great kiskadee	1 (0)	0
<i>Tyrannus melancholicus</i>	Tropical kingbird	1 (0)	2 (0)
Vireonidae			
<i>Cyclarhis gujanensis</i>	Rufous-browed peppershrike	1 (0)	1 (0)
Tropical Forest 2 - Inselberg		49 (5)	40 (9)
Caprimulgidae			
<i>Caprimulgus nigrescens</i>	Blackish nightjar	0	1 (1)
Cardinalidae			
<i>Cyanocompsa cyanooides</i>	Blue-black grosbeak	1 (0)	0
Furnariidae		8 (3)	4 (4)
<i>Xiphorhynchus pardalotus pardalotus</i>	Chestnut-rumped woodcreeper	1 (0)	0

(Continued)

Table 1. (Continued.)

Species		Standardized hour	
		(0:0.5)	(0.5:1)
<i>Deconychura longicauda</i>	Northern long-tailed woodcreeper	1 (1)	1 (1)
<i>Certhiasomus stictolaemus</i>	Spot-throated woodcreeper	0	1 (1)
<i>Glyphorhynchus spirurus spirurus</i>	Wedge-billed woodcreeper	6 (2)	2 (2)
Galbulidae			
<i>Galbula albirostris</i>	Yellow-billed jacamar	1 (0)	0
Pipridae			
		17 (0)	10 (0)
<i>Manacus manacus manacus</i>	White-bearded manakin	1 (0)	0
<i>Dixiphia pipra pipra</i>	White-crowned manakin	7 (0)	4 (0)
<i>Lepidothrix serena</i>	White-fronted manakin	7 (0)	2 (0)
<i>Corapipo gutturalis</i>	White-throated manakin	2 (0)	4 (0)
Poliophtilidae			
<i>Microbates collaris</i>	Collared gnatwren	1 (0)	0
Thamnophilidae			
		12 (2)	15 (3)
<i>Pernostola rufifrons rufifrons</i>	Black-headed antbird	1 (0)	0
<i>Willisornis poecilinotus</i>	Common scale-backed antbird	1 (0)	4 (0)
<i>Thamnomanes ardesiacus</i>	Dusky-throated antshrike	2 (1)	1 (1)
<i>Myrmotherula menetriesii</i>	Grey antwren	1 (0)	1 (0)
<i>Myrmotherula longipennis</i>	Long-winged antwren	1 (0)	4 (2)
<i>Thamnophilus murinus</i>	Mouse-coloured antshrike	1 (0)	0
<i>Hylophylax naevius naevius</i>	Spot-backed antbird	1 (0)	0
<i>Myrmotherula axillaris</i>	White-flanked antwren	0	1 (0)
<i>Pithys albifrons albifrons</i>	White-plumed antbird	4 (1)	4 (0)
Thraupidae			
		0	3 (0)
<i>Tachyphonus surinamus surinamus</i>	Fulvous-crested tanager	0	2 (0)
<i>Saltator grossus</i>	Slate-coloured grosbeak	0	1 (0)
Tityridae			
		2 (0)	1 (0)
<i>Schoffornis turdina</i>	Brown-winged schiffornis	1 (0)	0
<i>Myiobius barbatus</i>	Sulphur-rumped flycatcher	1 (0)	1 (0)
Turdidae			
<i>Turdus albicollis</i>	White-necked thrush	4 (0)	1 (1)
Tyrannidae			
<i>Mionectes macconnelli</i>	McConnell's flycatcher	3 (0)	4 (0)
Vireonidae			
<i>Hylophilus ochraceiceps</i>	Tawny-crowned greenlet	0	1 (0)

Birds were caught in three different environments: Temperate Forest (Souesmes), Tropical Forest 1 (Nouragues, Pararé) undergrowth and open area and Tropical Forest 2 (Nouragues, Inselberg). Differences in parasite prevalence in host feces between morning and afternoon differed between tropical and temperate species (B-GLMM: effect of morning vs afternoon: $\chi^2_1 = 76.21$, $P < 0.0001$, effect of the site: $\chi^2_1 = 55.41$, $P < 0.0001$, effect of their interaction: $\chi^2_1 = 38.03$, $P < 0.0001$). In the Temperate Forest, the prevalence of *Isospora* was higher in the afternoon than in the morning, whereas prevalence did not differ between morning and afternoon in the Tropical Forest (Table 2).

Results

Differences in temporal patterns of parasite prevalence in species from temperate and tropical forests

The prevalence of *Isospora* spp. oocysts in bird feces increased dramatically as the day progressed for species inhabiting temperate forest to reach a maximum of 90% of infected individuals [slope_{Temperate-Forest} (95% CI) = 6.89 (5.27; 8.50)] whereas in the same time the prevalence for species living in tropical forest remained unchanged around 25% of infected individuals [slope_{Tropical-Forest}

(95% CI) = 0.86 (−0.40; 2.12)] (B-GLMM: effect of time: $\chi^2_1 = 69.76$, $P < 0.0001$, effect of location: $\chi^2_1 = 6.65$, $P < 0.01$, effect of their interaction: $\chi^2_1 = 33.72$, $P < 0.0001$) (Fig. 2). The prevalence of *Isospora* spp. oocysts in bird feces did not differ between the two tropical forest sites [B-GLMM: effect of time: $\chi^2_1 = 2.47$, $P = 0.12$, effect of sites: $\chi^2_1 = 1.70$, $P = 0.19$, effect of their interaction: $\chi^2_1 = 0.98$, $P = 0.32$, slope_{Tropical-Forest 1} (95% CI) = 0.43 (−1.18; 2.05) and slope_{Tropical-Forest 2} (95% CI) = 1.87 (−0.42; 4.20)]. Complementary analyses of parasite prevalence contrasting morning and afternoon data are presented in Table 1 and yielded the same results.

Table 2. Post hoc tests based on the B-GLMM for the comparisons of morning (am) and afternoon (pm) prevalence in temperate and tropical areas (corrected for multiple comparisons, see text for details)

Contrasts	Estimate	Standard error	Z ratio	P
Temperate pm – am	3.70	0.42	8.73	<0.0001
Tropical pm – am	0.37	0.34	1.10	0.32
Temperate am – tropical am	–0.27	0.42	–0.63	0.53
Temperate pm – tropical pm	3.06	0.41	7.45	<0.0001
Temperate pm – tropical am	3.43	0.42	8.23	<0.0001
Tropical pm – temperate am	0.64	0.42	1.51	0.20

Significant differences are in bold.




Fig. 2. Prevalence of *Isospora* spp. (i.e. probability of infection, 1: infected, 0: not infected) in fecal samples from different bird species as a function of time elapsed through the day (standardized time of the day from 0 corresponding to sunrise to 1 corresponding to sunset) and of sampling site (Temperate Forest in black and Tropical Forest 1+2 in grey). Data points show actual prevalence, and predicted values fitted with the GLM model are in plain lines assorted with 95% CI in light grey.

Differences in temporal patterns of infection intensity between species of temperate and tropical forests

Pre-analyses showed differences in infection intensity, measured by parasite load in the host feces, between morning and afternoon in Temperate Forest (Wilcoxon test: $W = 983.5$, $P = 0.004$), and in Tropical Forest 1 (Wilcoxon test: $W = 206.5$, $P = 0.01$) but not in Tropical Forest 2 (Wilcoxon test: $W = 29.5$, $P = 0.38$) (mean \pm s.e. of number of *Isospora* spp. shed in feces in the morning vs in the afternoon respectively: Temperate Forest: 35.36 ± 15.95 vs 222.81 ± 46.01 , Tropical Forest 1: 29.71 ± 24.88 vs 91.06 ± 44.13 and Tropical Forest 2: 27.20 ± 23.25 vs 62.33 ± 37.92). Therefore, analyses of differences in parasite load were restricted to data from infected birds sampled in the afternoon (excluding birds from the open area in Tropical Forest 1). Parasite load remained constant all the afternoon in species living in the temperate habitat [slope_{Temperate-Forest} (95% CI) = -0.002 (-3.01 ; 3.01); mean \pm s.e. parasite load: 222.81 ± 46.01 oocysts per bird]. In contrast, parasite load increased over time from 0 to around 200 oocysts per bird at sunset [slope_{Tropical-Forest} (95% CI) = 5.98 (2.27 ; 9.69), mean \pm s.e. parasite load: 80.72 ± 30.97 oocysts per bird] (NB-GLMM: effect of time: $\chi^2_1 = 0.00$, $P = 1.00$, effect of location: $\chi^2_1 = 7.86$, $P < 0.01$, effect of their interaction: $\chi^2_1 = 5.84$, $P < 0.05$) (Fig. 3). In the Tropical Forest, the number of oocysts released in bird feces

increased through the afternoon and did not differ between the two sites and [NB-GLMM: effect of time: $\chi^2_1 = 6.94$, $P < 0.01$, effect of sites: $\chi^2_1 = 0.14$, $P = 0.71$, effect of their interaction: $\chi^2_1 = 0.02$, $P = 0.89$, slope_{Tropical-Forest 1} (95% CI) = 6.18 (1.73 ; 10.6) and slope_{Tropical-Forest 2} (95% CI) = 6.64 (1.36 ; 11.9)].

Differences in temporal patterns of prevalence and intensity between open field and undergrowth in the Tropical Forest 1

Prevalence of *Isospora* infection increased in the open area but remained stable inside the Tropical Forest throughout the day [B-GLMM: effect of time: $\chi^2_1 = 4.90$, $P < 0.05$; effect of area: $\chi^2_1 = 2.66$, $P = 0.10$, effect of their interaction: $\chi^2_1 = 4.30$, $P < 0.05$; slope_{open area} (95% CI) = 8.87 (1.02 ; 16.72) vs slope_{undergrowth} (95% CI) = 0.47 (-1.09 ; 2.02)] (Fig. 4). For example, in the most abundant species captured in this open area (silver-beaked tanager), all individuals caught in the afternoon shed oocyst in their feces whereas no individual caught in the morning did (Table 1). Considering only infected birds in the afternoon (standardized hour > 0.5), parasite load was higher in the open area than that in the undergrowth (NB-GLMM: effect of time: $\chi^2_1 = 1.08$, $P = 0.30$; effect of area: $\chi^2_1 = 5.35$, $P < 0.05$, mean \pm s.e. parasite load open area: 1028.70 ± 441.20 oocysts/bird and forest undergrowth: 91.06 ± 44.13 oocysts/bird) (Fig. 5).




Fig. 3. Parasite load of *Isospora* spp. (number of oocysts per gram of feces) in different bird species as a function of time elapsed through the afternoon (standardized time of the day ranging from 0.5 corresponding to midday to 1 corresponding to sunset) and of sampling site (Temperate Forest in black and Tropical Forest 1 + 2 in grey). Data points show actual parasite loads, and predicted values fitted with the GLM model are in plain lines assorted with 95% CI in light grey. Data presented here is restricted to infected birds sampled after a standardized hour of 0.5 (see text for more details).




Fig. 4. Prevalence of *Isospora* spp. (probability of infection, 1: infected, 0: not infected) in fecal samples for different bird species as a function of time elapsed through the day (standardized time of the day from 0 corresponding to sunrise to 1 corresponding to sunset) and of sampling location within the Tropical Forest 1 site (open area in black, surrounding undergrowth in grey). Data points show actual prevalence (1: infected, 0: not infected), and predicted values fitted with the GLM model are in plain lines assorted with 95% CI in lighter grey.

Comparison of abiotic factors between the forest undergrowth and open area

Variation in temperature showed a curvilinear relationship with time elapsed since sunrise, with a peak around 35°C in open area and only around 30°C in the forest undergrowth between 12:00 and 15:00 (Gaussian GLM: time \times area: $F_{1,94} = 4.88$, $P = 0.03$, time² \times area: $F_{1,94} = 5.20$, $P = 0.02$, Fig. 6a). In open area, UV radiation and luminosity showed a similar relationship with time from the morning to midday to reach a high intensity of UV radiation and luminosity intensity before a strong decrease until the sunset. In contrast, in the forest undergrowth UV radiation and luminosity intensity remains at very low level throughout the day (NB-GLM for UV radiation: time: $\chi^2_1 = 56.39$, $P < 0.0001$; time²: $\chi^2_1 = 59.99$, $P < 0.0001$; location: $\chi^2_1 = 169.08$, $P < 0.0001$, Fig. 6b and NB-GLM for luminosity: time: $\chi^2_1 = 52.01$, $P < 0.0001$; time²: $\chi^2_1 = 57.30$, $P < 0.0001$; location: $\chi^2_1 = 188.73$, $P < 0.0001$, Fig. 6c).

Discussion

In this study, we expected circadian variation in coccidial oocysts shedding to be found in temperate, but not in tropical forest habitat, consistent with the hypothesis that this trait in parasites is maintained by natural selection to avoid desiccation and UV radiation, to which oocysts are sensitive before sporulation. We predicted that in an environment where solar radiation reaching the ground is dramatically reduced, like in the dense undergrowth of a tropical forest, the circadian output of oocysts typically found in bird coccidia would disappear. Our results clearly confirmed this prediction since circadian variation in coccidial oocysts was high in the temperate forest (i.e. 90% of birds were infected in the late afternoon and almost none in the morning) and absent in the tropical forest (approximately 20% of birds shed oocyst in their faces whatever the hour of trapping). Furthermore, at a smaller spatial scale in the Tropical Forest 1




Fig. 5. Parasite load of *Isospora* spp. (number of oocysts per gram of feces) in different bird species as a function of time elapsed through the afternoon (standardized time of the day from 0.5 corresponding to midday to 1 corresponding to sunset) and of sampling location within the Tropical Forest 1 site (open area in black and surrounding undergrowth in grey). Data points show actual parasite loads, and predicted values fitted with the GLM model are in plain lines assorted with 95% CI in light grey. Data presented here is restricted to infected birds sampled after a standardized hour of 0.5 (see text for more details).

site, a similar result was found, as birds captured in the open area showed a strong circadian variation contrary to bird species from the surrounding undergrowth of the rain forest.

The main goal of this study was to go one step further regarding the adaptive significance of circadian variation in coccidial oocysts shedding in birds, demonstrated in all tested species including birds living in the arctic area (see Knight *et al.*, 2018 for a review). Ten years earlier, Martinaud *et al.* (2009) experimentally demonstrated that the release of oocysts in the late afternoon was an adaptive trait to avoid desiccation and UV radiation, thus reducing mortality of oocysts in the external environment. Before our study, such a circadian pattern appeared to be a ubiquitous phenomenon in intestinal bird parasites, since all studies showed that oocyst shedding peaks in the late afternoon (reviewed in Knight *et al.*, 2018). To our knowledge, this study is the first to document the absence of circadian variation in coccidial oocysts shedding in a large set of species inhabiting the undergrowth of a rain forest. This environment is characterized by low luminosity and low UV radiation levels reaching the ground, combined with high soil and air moisture (Chazdon and Fetcher, 1984; Vazquez-Yanes *et al.*, 1990). All these abiotic characteristics are favourable for the survival of oocysts in host feces (Allen and Fetterer, 2002). If temporal variation in the shedding of coccidial oocysts is a trait that has evolved under a natural selection to avoid desiccation and UV radiation as commonly thought (Martinaud *et al.*, 2009), such a pattern should not be observed in an environment where these abiotic selective pressures are relaxed. Contrary to the continuous multi-layered vegetation canopy of rain forests, the canopy in temperate forests is moderately dense and allows greater and seasonally varying light penetration (Anderson, 1964; Koizumi and Oshima, 1985; Szwagrzyk *et al.*, 2001). Moreover, most temperate forests are managed and affected by human activities, and light intensities reaching the ground may differ greatly according to the stage in the forest management (Canham and Marks, 1985; Buckley, 1992; McElrone *et al.*, 2013). For example, in most recent cuts (1-year-old), light intensity was 3.5–5 times higher than that in uncut plots, and rapidly decreased over time following selection cutting, especially at the ground level (Beaudet *et al.*, 2004). Therefore,

temperate forests are exposed to a stronger and more variable irradiance compared to tropical rain forest. We observed marked temporal variation in the shedding of coccidial oocysts in birds from the temperate forest, consistent with previous studies (Knight *et al.*, 2018). On the contrary, bird species living in the tropical forest did not show this diurnal variation. These results suggest that natural selection (evolutionary costs and benefits) may indeed shape release rhythms in *Isospora* spp. of birds like in many other parasite species (Reece *et al.*, 2017).

In addition, comparing circadian variation in coccidial oocysts shedding within the tropical habitat between species exclusively living in the open area and those living in the surrounding undergrowth further supports this hypothesis. This open area was created at the construction of the field camp, more than 30 years ago, and has since selected bird species specialized in open habitats. Indeed, species captured in the open field were never caught in the undergrowth, like silver-beaked tanager (*Ramphocelus carbo*), golden-sided euphonia (*Euphonia cayennensis*), violaceous euphonia (*Euphonia violacea*), tropical kingbird (*Tyrannus melancholicus*), great kiskadee (*Pitangus sulphuratus*), buff-throated saltator (*Saltator maximus*) and rufous-browed peppershrike (*Cyclarhis gujanensis*). A review of the literature (Handbook of the Birds of the World Series) confirms that these bird species are specialists of open habitats. Unlike the surrounding undergrowth, this open area is characterized by strong level of luminosity and UV radiation reaching the ground. Indeed, during a typical day in the tropical environment, light intensity and UV radiation reaching the ground measured in the open area were on average 100 times higher compared to the ground in the undergrowth, at any time but in particular at midday. Furthermore, the field camp is located near the Equator, a place where the annual UV doses reach their maximum (Marks and Plewig, 1991), increasing the selective pressure towards a marked circadian variation in coccidial oocysts shedding. Therefore, *Isospora* sp. infecting birds living in this open area have likely been selected to evolve strong temporal variation in oocysts shedding allowing oocyst survival in such a harsh external environment.

Regarding variation in the intensity of *Isospora* spp. infection, we found that parasite load remained stable in temperate species




Fig. 6. (a) Temperature ($^{\circ}\text{C}$), (b) UV radiation index and (c) luminosity intensity (lux) as a function of time elapsed through the day and of sampling location within Tropical Forest 1 site (open area in black and surrounding undergrowth in grey). At the time of sample collection, sunrise occurred at 06:00 am and sunset around 06:00 pm. Data points show actual values, and predicted values fitted with the GLM model are in plain lines assorted with 95% CI in light grey.

through the day while it increased in tropical areas, to reach by the end of the afternoon approximately the same parasite load as in temperate species. This increase in oocyst release during the day confirms previous results on two species (*Serinus serinus* and *Sylvia borin*) in which coccidial parasite load in the feces peaked in the late afternoon (Lopez *et al.*, 2007). The mechanisms underlying this process, however, are still unclear. After different phases of multiplication inside the intestinal tractus of its host, oocysts are released from its host cell and shed into the environment with the feces. The proximal factors triggering coccidial oocyst discharge are not identified yet but it is known that they are linked to host physiology (Boughton, 1933; Dolnik, 1999a,

1999b) and potentially mediated by melatonin, which regulates daily activity patterns in vertebrates. Indeed, outside polar regions with constant daylight, avian *Isopora* species use cyclic changes in melatonin concentration in their host blood to synchronize oocyst output (Dolnik *et al.*, 2011 and references therein). Therefore, we may hypothesize that the threshold of physiological signals inducing oocyst discharge is very low for tropical species inhabiting the dense undergrowth, allowing oocysts discharge early in the morning. The accumulation of these signals as the day progressed may induce the slight but continuous increase in parasite load observed in our data over the afternoon. Another but not exclusive hypothesis would be that the amount of feces

produced could be higher in the late afternoon allowing a higher discharge in oocyst in the same time.

Finally, our results meet an interesting methodology issue for researchers interested in working on intestinal parasites like *Isoospora* spp. in wild birds' species (Biard et al., 2015, 2010). Due to circadian rhythm in oocyst shedding in temperate habitat, there is a large difference in the number of *Isoospora* sp. oocysts shed between the morning and afternoon for a given individual (Filipiak et al., 2009). Hence, to accurately assess, and make meaningful comparisons in, the prevalence of *Isoospora* infection, sampling of feces must be standardized at the same time of the day, as already demonstrated in previous studies (Filipiak et al., 2009; Dolnik et al., 2010). In the tropical rain forest where this circadian rhythm in oocyst shedding is absent, prevalence of infection may be assessed from feces collected at any time of the day. This makes it easier to work on the effects of intestinal parasites on bird host life-history traits such as sexually selected traits for instance (Brawner et al., 2000; Hórák et al., 2004). However, as parasite load increases as the day progresses, we emphasize the need to precisely record the time of day at which samples are collected. If *Isoospora* spp. parasite loads are to be compared, then sampling time should be standardized at the same time of day.

Concluding remarks

We believe that studying tropical birds improved our knowledge of the ecology of *Isoospora* spp. in natural populations of hosts and the adaptive significance of daily rhythm in oocyst shedding. The available information on intestinal parasites in tropical birds to date is scarce as research has often focused on blood parasites (Ricklefs, 1992; Durrant et al., 2006; Ricklefs and Sheldon, 2007; Merino et al., 2008; Svensson-Coelho et al., 2014). Our study clearly showed that the knowledge on the interaction between parasites and hosts in the temperate area cannot be transposed for birds in tropical area. Therefore, future investigations are needed to reveal more information on host–parasite interactions concerning the probability to be infected in the two different ecosystems. A comparative analysis with a large set of species would be very useful to investigate the ecological factors and host life-history traits underlying parasite diversity across different ecological contexts (tropical vs temperate).

Data. Once the paper has been accepted, data will be deposited in the Dryad.

Acknowledgements. This study has been conducted within the French natural reserve of Nouragues. We acknowledge the Nouragues CNRS research field station (USR mixte LEEISA, CNRS; Cayenne) and the staff dedicated. We thank all the people who contributed to the fieldwork.

Author contributions. CB, KM and JM conceived and designed the study. CB, MT, SM, SBA, LD and JM conducted data gathering. KM and JM performed statistical analyses. CB, KM and JM wrote the article.

Financial support. We acknowledge the Nouragues CNRS research field station, which benefits from 'Investissement d'Avenir' grants managed by Agence Nationale de la Recherche (AnaEE France ANR-11-INBS-0001; Labex CEBA ANR-10-LABX-25-01).

Conflict of interest. The authors declare there are no conflicts of interest.

Ethical standards. This work conforms to French legal requirements, and accepted international ethical standards, including those relating to conservation and welfare, and to the journal's policy on these matters. We thank the Direction Régionale de l'Environnement, de l'Aménagement et du Logement (DREAL) de la Guyane, de la Nouvelle-Aquitaine et du Loir et Cher for delivering legal authorizations of catching birds (authorization nos.: R03-2018-07-17-002, R03-2019-03-14-005, 128/2017, 2019D/2323 and 41-2018-04-03-002).

References

- Abd-Al-Aal Z, Ramadan NF and Al-Hoot A (2000) Life-cycle of *Isoospora mehlhornii* sp. nov. (Apicomplexa: Eimeriidae), parasite of the Egyptian swallow *Hirundo rubicola savignii*. *Parasitology Research* **86**, 270–278.
- Allen PC and Fetterer RH (2002) Recent advances in biology and immunobiology of *Eimeria* species and in diagnosis and control of infection with these coccidian parasites of poultry. *Clinical and Microbiology Review* **15**, 58–65.
- Anderson MC (1964) Studies of the woodland light climate: II. Seasonal variation in the light climate. *Journal of Ecology* **52**, 643–663.
- Beaudet M, Messier C and Leduc A (2004) Understorey light profiles in temperate deciduous forests: recovery process following selection cutting. *Journal of Ecology* **92**, 328–338.
- Benjamini Y and Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B* **57**, 289–300.
- Biard C, Saulnier N, Gaillard M and Moreau J (2010) Carotenoid-based bill colour is an integrative signal of multiple parasite infection in blackbird. *Naturwissenschaften* **97**, 987–995.
- Biard C, Monceau K, Motreuil S and Moreau J (2015) Interpreting immunological indices: the importance of taking parasite community into account. An example in blackbirds *Turdus merula*. *Methods in Ecology and Evolution* **6**, 960–972.
- Boughton DC (1933) Diurnal gametic periodicity in avian *Isoospora*. *American Journal of Epidemiology* **18**, 161–184.
- Bourgeron PS (1983) Spatial aspects of vegetation structure. In Golley FB (eds), *Tropical Rain Forest Ecosystems: Structure and Function. Ecosystems of the World*. Amsterdam and New York: Elsevier Scientific Pub. Co., pp. 29–47.
- Brawner WR and Hill GE (1999) Temporal variation in shedding of coccidian oocysts: implications for sexual-selection studies. *Canadian Journal of Zoology* **77**, 347–350.
- Brawner WR, Hill GE and Sundermann CA (2000) Effects of coccidian and mycoplasmal infections on carotenoid-based plumage pigmentation in male House finches. *The Auk* **117**, 952–963.
- Brinkmann WLF (1971) Light environment in a tropical rain forest of central Amazonia. *Acta Amazonica* **1**, 37–49.
- Brown M, Ball S and Holman D (2001) The periodicity of isosporan oocyst discharge in the greenfinch (*Carduelis chloris*). *Journal of Natural History* **35**, 945–948.
- Bruno RD, Rocha HR, Freitas HC, Goulden ML and Miller SD (2006) Soil moisture dynamics in an eastern Amazonian tropical forest. *Hydrology Processes* **20**, 2477–2489.
- Buckley GP (1992) *Ecology and Management of Coppice Woodlands*. London, UK: Chapman and Hall.
- Bush AO, Fernández JC, Esch GW and Seed JR (2001) *Parasitism: The Diversity and Ecology of Animal Parasites*. Cambridge, UK: Cambridge University Press.
- Canham CD and Marks PL (1985) The response of woody plants to disturbance: patterns of establishment and growth. In Pickett STA and White PS (eds), *The Ecology of Natural Disturbance and Patch Dynamics*. New York, USA: Academic Press, pp. 197–216.
- Chazdon RL and Fetcher N (1984) Photosynthetic light environments in a lowland tropical rain forest in Costa Rica. *Journal of Ecology* **72**, 553–564.
- Combes C (2001) *Parasitism: The Ecology and Evolution of Intimate Interactions*. London, UK: University of Chicago Press and London.
- Combes C, Fournier A, Moné H and Théron A (1994) Behaviours in trematode cercariae that enhance parasite transmission: patterns and processes. *Parasitology* **109**, S3–S13.
- Dolnik OV (1999a) Diurnal periodicity of oocysts release of *Isoospora dilatata* (Sporozoa: Eimeriidae) from the common Starling (*Sturnus vulgaris*) in nature. *Parasitologia* **33**, 74–80.
- Dolnik OV (1999b) Diurnal periodicity in appearance of *Isoospora* (Protozoa: Coccidia) oocysts from some passerine birds. *Proceedings of the Zoological Institute RAS* **281**, 113–118.
- Dolnik OV, Dolnik VR and Bairlein F (2010) The effect of host foraging ecology on the prevalence and intensity of coccidian infection in wild passerine birds. *Ardea* **98**, 97–103.
- Dolnik OV, Metzger BJ and Loonen MJJE (2011) Keeping the clock set under the midnight sun: diurnal periodicity and synchrony of avian *Isoospora* parasites cycle in the High Arctic. *Parasitology* **138**, 1077–1081.

- Durrant KL, Beadell JS, Ishtiaq F, Graves GR, Olson SL, Gering E, Peirce MA, Milensky CM, Schmidt BK, Gebhard C and Fleischer RC (2006) Avian hematozoa in South America: a comparison of temperate and tropical zones. *Ornithological Monographs* **60**, 98–111.
- Filipiak L, Mathieu F and Moreau J (2009) Caution on the assessment of intestinal parasitic load in studying parasite-mediated sexual selection: the case of blackbirds coccidiosis. *International Journal for Parasitology* **39**, 741–746.
- Freiberg M (1997) Spatial and temporal pattern of temperature and humidity of a tropical premontane rain forest tree in Costa Rica. *Selbyana* **18**, 77–84.
- Hilty S (2002) *Birds of Venezuela*, 2nd Edn, Princeton, USA: Princeton University Press.
- Hörak P, Saks L, Karu U, Ots I, Surai PF and McGraw KJ (2004) How coccidian parasites affect health and appearance of greenfinches. *Journal of Animal Ecology* **73**, 935–947.
- Hudman SP, Ketterson ED and Nolan V (2000) Effects of time of sampling on oocyst detection and effects of age and experimentally elevated testosterone on prevalence of coccidia in male dark-eyed juncos. *The Auk* **117**, 1048–1051.
- Knight A, Ewen JG, Brekke P and Santure AW (2018) Chapter two – The evolutionary biology, ecology and epidemiology of coccidia of passerine birds. In Rollinson D and Stothard JR (eds), *Advances in Parasitology*. London, UK: Academic Press, pp. 35–60.
- Koizumi H and Oshima Y (1985) Seasonal changes in photosynthesis of four understory herbs in deciduous forests. *Botanical Magazine* **98**, 1–13.
- Kruszewicz AG (1995) The occurrence of *Isospora lacazei* (Coccidia: Eimeriidae) and its influence on nestling growth in house sparrows (*Passer domesticus*) and tree sparrows (*Passer montanus*). In Pinowski J, Kavanagh BP and Pinowska B (eds), *Nestling Mortality of Granivorous Birds Due to Microorganisms and Toxic Substances: Synthesis*. Warszawa, Poland: Polish Scientific Publishers, pp. 291–305.
- Lopez G, Figuerola J and Soriguer R (2007) Time of day, age and feeding habits influence coccidian oocyst shedding in wild passerines. *International Journal for Parasitology* **37**, 559–564.
- Marks R and Plewig G (1991) *The Environmental Threat to the Skin*. London, UK: Taylor and Francis Ltd.
- Martinaud G, Billaudelle M and Moreau J (2009) Circadian variation in shedding of the oocysts of *Isospora turdi* (Apicomplexa) in blackbirds (*Turdus merula*): an adaptative trait against desiccation and ultraviolet radiation. *International Journal for Parasitology* **39**, 735–739.
- McElrone AJ, Choat B, Gambetta GA and Brodersen CR (2013) Water uptake and transport in vascular plants. *Nature Education Knowledge* **4**, 6.
- Merino S, Moreno J, Vásquez RA, Martínez J, Sánchez-Monsálvez I, Estades CF, Ippi S, Sabat P, Rozzi R and McGehee S (2008) Haematozoa in forest birds from southern Chile: latitudinal gradients in prevalence and parasite lineage richness. *Austral Ecology* **33**, 329–340.
- Misof K (2004) Diurnal cycle of *Isospora* spp. oocyst shedding in Eurasian blackbirds (*Turdus merula*). *Canadian Journal of Zoology* **82**, 764–768.
- Olsen OW (1974) *Animal Parasites: Their Life Cycles and Ecology*. Baltimore, USA: University Park Press.
- R Core Team (2021) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at <https://www.R-project.org/>.
- Reece SE, Prior KF and Mideo N (2017) The life and times of parasites: rhythms in strategies for within-host survival and between-host transmission. *Journal of Biological Rhythms* **32**, 516–533.
- Restall R, Rodner C and Lentino M (2006a) *Birds of Northern South America: An Identification Guide. Species Accounts*, vol. **1**, 1st Edn. London, UK: Helm Field Guides.
- Restall R, Rodner C and Lentino M (2006b) *Birds of Northern South America: An Identification Guide. Plates and Maps*, vol. **2**, 1st Edn. London, UK: Helm Field Guides.
- Ricklefs RE (1992) Embryonic development period and the prevalence of avian blood parasites. *Proceedings of the National Academy of Sciences* **89**, 4722–4725.
- Ricklefs RE and Sheldon KS (2007) Malaria prevalence and white-blood-cell response to infection in a tropical and in a temperate thrush. *The Auk* **124**, 1254–1266.
- Svensson-Coelho M, Ellis VA, Loiseau BA, Blake JG and Ricklefs RE (2014) Reciprocal specialization in multihost malaria parasite communities of birds: a temperate-tropical comparison. *American Naturalist* **184**, 624–635.
- Szwagrzyk J, Szweczyk J and Bodzarczyk J (2001) Dynamics of seedling banks in beech forest: results of a 10-year study on germination, growth and survival. *Forest Ecology and Management* **141**, 237–250.
- Théron A (1984) Early and late shedding patterns of *Schistosoma mansoni* Cercariae: ecological significance in transmission to human and murine hosts. *Journal of Parasitology* **70**, 652–655.
- Vazquez-Yanes C, Orozco-Segovia A, Rincon E, Sanchez-Coronado ME, Huante P, Toledo JR and Barradas VL (1990) Light beneath the litter in a tropical forest: effect on seed germination. *Ecology* **71**, 1952–1958.