

# Immune cell arrival kinetics to peritoneum and role during murine-experimental trichomoniasis

F. J. Rangel-Mata<sup>1</sup>, E. E. Ávila-Muro<sup>1</sup> , J. E. Reyes-Martínez<sup>1</sup> ,  
L. M. Olmos-Ortiz<sup>1</sup> , M. E. Brunck<sup>2</sup>, L. A. Arriaga-Pizano<sup>3</sup> and P. Cuéllar-Mata<sup>1</sup> 

## Research Article

**Cite this article:** Rangel-Mata FJ, Ávila-Muro EE, Reyes-Martínez JE, Olmos-Ortiz LM, Brunck ME, Arriaga-Pizano LA, Cuéllar-Mata P (2021). Immune cell arrival kinetics to peritoneum and role during murine-experimental trichomoniasis. *Parasitology* **148**, 1624–1635. <https://doi.org/10.1017/S0031182021001311>

Received: 17 March 2021

Revised: 9 June 2021

Accepted: 15 July 2021

First published online: 9 August 2021

### Key words:

Host–parasite interaction; NETs; trichomoniasis

### Author for correspondence:

P. Cuéllar-Mata, E-mail: [mata@ugto.mx](mailto:mata@ugto.mx)

<sup>1</sup>Departamento de Biología, Universidad de Guanajuato, Guanajuato, Mexico; <sup>2</sup>Escuela de Ingeniería y Ciencias, Tecnológico de Monterrey, Monterrey, Mexico and <sup>3</sup>Unidad de Investigación Médica en Inmunoquímica, IMMS, Ciudad de México, Mexico

### Abstract

*Trichomonas vaginalis* causes trichomoniasis, an inflammatory process related to an increased rate of HIV transmission. In order to study *T. vaginalis* infection response in a microorganism-free environment, an infection model was established providing a host–parasite interaction system useful to study the interplay between immune cells and the parasite. Infected mice peritoneal cells were immunophenotyped at different times after infection using flow cytometry. Neutrophils and macrophages showed the most relevant increase from third to 12th day post-infection. A high number of B lymphocytes were present on 15th day post-infection, and an increase in memory T cells was observed on sixth day post-infection. The levels of NO increased at day 10 post-infection; no significant influence was observed on *T. vaginalis* clearance. Increased viability of *T. vaginalis* was observed when the NETs inhibitors, metformin and Cl<sup>−</sup> amidine, were administered, highlighting the importance of this mechanism to control parasite infection (43 and 86%, respectively). This report presents a comprehensive cell count of the immune cells participating against trichomoniasis in an *in vivo* interaction system. These data highlight the relevance of innate mechanisms such as specific population changes of innate immune cells and their impact on the *T. vaginalis* viability.

## Introduction

*Trichomonas vaginalis* is a protozoan parasite with a cosmopolitan presence which may invade the human urogenital tract and cause trichomoniasis. The incidence of trichomoniasis is increasing worldwide, with 50% of asymptomatic cases reported in North America (Twu *et al.*, 2013; Kissinger, 2015). Trichomoniasis induces vaginitis, cervicitis, urethritis, pelvic inflammatory disease and diverse adverse effects during pregnancy resulting in social, medical and economic implications (Swygard *et al.*, 2004; Twu *et al.*, 2013). Women infected during pregnancy are predisposed to early rupture of placental membranes and premature birth (Petrin *et al.*, 1998). In addition, trichomoniasis is related to increased incidence of cervical-uterine and prostate cancer, infertility, and higher susceptibility to HIV infection (Kharsany *et al.*, 1993; McClelland *et al.*, 2007; Kissinger *et al.*, 2009; Twu *et al.*, 2013; Smith and Garber, 2015).

Innate immunity is the first line of defence against microorganisms (Fichorova, 2009) including *T. vaginalis*. Epithelial cells, vaginal mucus, antimicrobial peptides, chemokines and immune cells such as neutrophils and macrophages provide the innate response in the female genital tract (Menezes and Tasca, 2016). On the other hand, adaptive immunity is a pathogen-specific mechanism that requires the activation of B and T lymphocytes, mediated by antigen-presenting cells (Nemati *et al.*, 2018).

An endometrial inflammatory response is characterized by the presence of different cell populations, mainly neutrophils, macrophages and lymphocytes (Alderete and Garza, 1985; Song *et al.*, 2008; Reighard *et al.*, 2011; Nam *et al.*, 2012). Furthermore, an increase in CD4<sup>+</sup> T cells has been described in *T. vaginalis*-infected women and murine models (Yano *et al.*, 1983; Levine *et al.*, 1998; Reighard *et al.*, 2011; Smith and Garber, 2015), and parasite-specific IgG and IgM have been observed (Menezes and Tasca, 2016). The persistence of the parasite in hosts that have developed humoral and cell-mediated responses suggests an incomplete response. As neutrophils and macrophages are relevant in the endometrial response, this prompts further investigations of the innate arm of the immune response. Usually, neutrophils battle and kill pathogens through phagocytosis, degranulation, reactive oxygen species (ROS) production and neutrophil extracellular traps (NETs) formation. Nevertheless, the mechanism in *T. vaginalis* infection has not been fully elucidated and the participation of other cell types has not been considered. The use of *in vivo* interaction systems overcomes several disadvantages of cell culture, allowing the study of the integration of a complex cellular diversity and its role during the pathogenesis, permitting the observation of cell migration to determine cell function in the infection process. As in the gut tract, the female urogenital tract is populated by microorganisms, either commensal or pathogenic bacteria which could be masking the specific defence mechanisms established by local immune cells. This report aimed to describe the recruitment kinetics of immune cells to the infection site in order to define

their role in an *in vivo* infection system. Besides, we show that NETs contributed to reduce the parasite viability, whereas reactive oxygen or nitrogen species have a limited impact.

## Materials and methods

### Parasite culture

*Trichomonas vaginalis* strain GT-21 was donated by Dr Anaya-Velazquez and Dr Padilla-Vaca (University of Guanajuato, Mexico); the strain was isolated from a symptomatic female patient and its virulence was demonstrated (Olmos-Ortiz *et al.*, 2017). The parasite was routinely cultured at 37°C in TYI-S-33 medium supplemented with 6% adult bovine serum and 1.5% Diamond vitamins. Potential mycoplasma contamination was investigated using a Venor GeM mycoplasma kit (Sigma-Aldrich, Cat. # MP0025), according to the manufacturer's protocol. The GT-21 strain was further investigated for the presence of any of the four known *T. vaginalis* virus (TVV 1–4) according to the previously described protocols (Figs S1 and S2) (Goodman *et al.*, 2011; Fraga *et al.*, 2012).

### Mice maintenance conditions

BALB/c mice between 6 and 8 weeks of age (weight from 25 to 30 g) were used. Animals were bred in polycarbonate cages administering water and food *ad libitum*, in a temperature-controlled environment at 25 ± 3°C and handled according to the Mexican norm NOM-062-ZOO-1999.

### *Trichomonas vaginalis* infection

Six animals per group were inoculated with 0, 1, 3, 5 or 8 million trophozoites diluted in 500 µL of sterile PBS. The peritoneal injection was performed using a 1 mL syringe with a 0.5 × 16 mm needle. The mice were sacrificed 15 days after inoculation to confirm the establishment of the infection and the peritoneal lavages were performed using 2 mL of cold PBS, recovering 500 µL of peritoneal wash from each mouse. Aliquots of the peritoneal washes (100 µL) were inoculated in LB liquid medium and Muller Hinton solid medium to identify any bacterial contamination during the process. After 3 days of inoculation with 8 million trophozoites, this group showed high mortality (40%). The evaluation of the infection was performed in the rest of the animals, according to the following criteria.

- Peritoneal washes cell count. The cells contained in peritoneal washes were stained with propidium iodide (PI) and the total count of cells including trophozoites and peritoneal cells was performed using a Neubauer chamber. Results were expressed as cells mL<sup>-1</sup> for each experimental condition.
- Live parasites recovery. Also, the viability of the parasite recovered from the peritoneal cavity was verified as reported (Marcel *et al.*, 2001). That is, the presence of live trophozoites was evaluated by the addition of 20 µL of peritoneal washes into 5 mL of TYI-S-33 medium added with penicillin/streptomycin (100 U mL<sup>-1</sup> and 100 µg mL<sup>-1</sup>, respectively), cultured at 37°C during 24 h and the absorbance to 450 nm was measured.
- Necropsy. The animals were sacrificed by cervical dislocation 15 days after parasite inoculation. Liver, stomach, kidneys, bladder, lungs, heart and encephalon were examined for anatomic damage. In addition, spleens were resected and weighed for splenomegaly assessment.

### Flow cytometry

Six groups of animals were inoculated with 5 million trophozoites. Mice from each group were sacrificed at different times, 0, 3, 6, 9, 12 and 15 days after inoculation. Peritoneal washes were performed, and cells were resuspended at 1 million into 200 µL sterile PBS supplemented with 5 mM EDTA, 3% FBS (Biowest, Nuaillé, France) and 2% BSA (Gold Biotechnology, Sant Louis, MO, USA). The cells were incubated with the following rat antibodies: anti-CD8-AF647 (Santa Cruz Biotechnology, Dallas, TX, USA), anti-CD19-APC/Cy7, anti-F4/80-PerCP and anti-Ly-6c-PE/Cy7 (Biolegend, San Diego, CA, USA). Besides, Syrian hamster anti-CD3-PE and rat anti-CD4-FITC antibodies were used. After 60 min of incubation on ice, the cells were washed twice, resuspended in 500 µL of sterile PBS and incubated with PI for 15 min (Biolegend). After washing under the same conditions, the cells were fixed with 4% paraformaldehyde for 30 min, washed three times with PBS containing glycine, resuspended in 100 µL of sterile PBS and analysed by flow cytometry. Cells were acquired on FACScanto (BD Biosciences) in the Hospital of Specialities 'Centro Medico Nacional Siglo XXI' and data were analysed with FlowJo software (FlowJo V10, Ashland, Oregon, USA).

### Reactive nitrogen species quantification

Two iNOS inhibitors, N(w)-nitro-L-arginine methyl ester (L-NAME) or aminoguanidine (AG) (Sigma Aldrich, San Luis, MO, USA), were independently administered to mice at 1% in the drinking water to investigate the participation of nitric oxide in the immune response against *T. vaginalis*. Nine days after parasite inoculation, when the peak of infection occurred, the mice were sacrificed (Barajas-Mendiola *et al.*, 2019), and peritoneal washes were obtained. Reactive nitrogen species (RNS) were determined by nitrite quantification according to the modified Griess method (Arias-Negrete *et al.*, 2004).

### NETs inhibition


To investigate the role of NETs to control *T. vaginalis* growth, the NADPH-oxidase inhibitor metformin and the peptidyl arginine deiminase-4 (PAD4) inhibitor Cl-amidine were used; these compounds interfere in the signalling cascade for NETosis. Mice were inoculated with 5 million trophozoites by intraperitoneal injection, and 2% metformin in drinking water along the infection period, or Cl-amidine (0.3 mg per 200 µL sterile PBS/day/mice, intraperitoneally) during 3 days after infection. The dose of inhibitors was adapted according to previous reports (Knight *et al.*, 2013; Wang *et al.*, 2015). Mice were sacrificed, and the peritoneal washes were used for extracellular DNA (NETs) quantification assays.

### Extracellular DNA quantification

Extracellular DNA present in peritoneal washes was digested with micrococcal nuclease (500 mU mL<sup>-1</sup>, New England Biolabs) for 15 min, and the nuclease activity was stopped with 5 mM EDTA. Supernatants were collected and centrifuged at 2500 g for 5 min. DNA was calculated using Quant-it DNA PicoGreen (Invitrogen) under a standard curve using λ-DNA, as indicated by the provider.

### Statistical analysis

Statistical analysis of three independent experiments was performed with the GraphPad Prism 5 software, using analysis of variance of Kruskal–Wallis and Dunn's post-hoc test. Each experiment included six mice per condition, and the significance



**Fig. 1.** Signs of intraperitoneal infection. Mice were intraperitoneally inoculated with 1, 3 or 5 million trophozoites, and group control was inoculated with sterile PBS. Mice were sacrificed 15 days after inoculation for analysis of infection signs such as (A) cell density (B) trophozoites count, (C) splenomegaly analyses, spleen weight (mg)/body weight (g) ratio in each group; and (D) leucocytes count in blood samples. The experiments were performed by triplicate and statistical analysis carried out by Kruskal–Wallis and Dunn's post-test using  $n=6$ ,  $P<0.001$ .

$P < 0.01$  was indicated with \*,  $P < 0.001$  was indicated with \*\*, and  $P < 0.0001$  was indicated with \*\*\*.

## Results


### *Trichomonas vaginalis* intra-peritoneal infection

To establish an *in vivo* interaction system that allowed the study of the immune response, both parasite viability and host survival were evaluated, after trophozoite introduction at different quantities into the mice peritoneum. Parasite cultures were free of mycoplasma contamination and TVV (see Figs S1 and S2). Mice were inoculated with 1, 3, 5 or 8 million GT-21 *T. vaginalis* trophozoites and then sacrificed 15 days after inoculation to perform the necropsy and analyses of infection signs. Total cell count, immune cells and trophozoites, was determined by Neubauer chamber count and it revealed a considerable increase of peritoneal cells  $\text{mL}^{-1}$  with respect to uninfected mice, up to  $17.4 \times 10^6$  or  $57.4 \times 10^6$  cells in mice inoculated with 3 or 5 million trophozoites, respectively. The inoculation of 1 million trophozoites did not induce significant changes in the cell

population ( $5.7 \times 10^6$  cells  $\text{mL}^{-1}$ ). As expected, uninfected mice showed an invariable number of cells per millilitre ( $5.2 \times 10^6$  cells  $\text{mL}^{-1}$ ) (Fig. 1A). As more than 40% mortality on mice was observed in the group inoculated with 8 million trophozoites, the data obtained from this were excluded from the study.

To determine the parasite viability, 20  $\mu\text{L}$  of the peritoneal washes were inoculated in TYI-S-33 medium, and 24 h after inoculation, the parasite viability was evaluated as described (Marcel *et al.*, 2001; Olmos-Ortiz *et al.*, 2017; Barajas-Mendiola *et al.*, 2019). The results confirm the infection establishment with an increasing number of parasites isolated,  $1.6 \times 10^5 \pm 0.4 \times 10^5$  or  $6.9 \times 10^5 \pm 0.9 \times 10^5$  trophozoites  $\text{mL}^{-1}$  when 3 or 5 million trophozoites were inoculated, respectively (Fig. 1B). Peritoneal washes obtained from mice inoculated with 1 million trophozoites did not show parasite growth.

Besides, we observed that spleens from infected mice increased in size compared to those from uninfected animals (2.3 or 1.9 cm, respectively). We decided to evaluate splenomegaly determining the spleen/body weight ratio ( $\text{mg g}^{-1}$ ). This ratio was 4.78, 7.12 and 6.74 in mice inoculated with 1, 3 or 5 million trophozoites, respectively, while the uninfected mice ratio was 3.97 (Fig. 1C).




**Fig. 2.** Gates selection for flow cytometry analysis. (A) Single cells. (B) Living cells. (C) Ly6C-F4/80 markers and Ly6C<sup>medium</sup>F4/80<sup>-</sup> for granulocytes; Ly6C<sup>high</sup>F4/80<sup>-</sup> for monocytes; Ly6C<sup>-</sup>F4/80<sup>+</sup> for resident macrophages; and Ly6C<sup>+</sup>F4/80<sup>+</sup> for inflammatory macrophages. (D) Double-negative cells were gated for T cells (CD3<sup>+</sup>) and B cells (CD19<sup>+</sup>). (E) CD3<sup>+</sup> cells were gated by Th cells (CD4<sup>+</sup>) and Tc cells (CD8<sup>+</sup>).

On day 15 post-infection, the differential leukocyte count in blood showed lymphocytosis and neutropenia similar to a well-established infection process (Fig. 1D) (Malla *et al.*, 2004; Lazenby *et al.*, 2013; Smith and Garber, 2015). Moreover, the pathologic findings revealed the presence of a whitish nodule in the peritoneal tissue of infected mice, more abundant in those inoculated with 5 million trophozoites (Fig. S3).

Taken together the data, the increase of peritoneal cell total count, long-term parasite survival, splenomegaly, differential leukocyte count and pathologic analysis indicate an intense immune response to a well-established infection process. The inoculum of 5 million trophozoites was used for the next experiments.

### *Trichomonas vaginalis* induces intense cellular changes in response to infection

A flow cytometry analysis was performed in peritoneal washes obtained from infected mice, at 3, 6, 9, 12 and 15 days after parasite inoculation, and the cell population changes were determined. PI was used for gating out dead cells and gates were designed as Ly6C<sup>-</sup> and F4/80<sup>+</sup> for resident macrophages, Ly6C<sup>+</sup> F4/80<sup>+</sup> for inflammatory macrophages, Ly6C<sup>medium</sup>F4/80<sup>-</sup> for polymorphonuclears (PMN) and Ly6C<sup>high</sup>F4/80<sup>-</sup> for monocytes (Fig. 2). The results show a significant increase in the average of different cell lineages, such as PMN, which increase from 7.35 to 42.9%




**Fig. 3.** Cell population changes during *T. vaginalis* infection. Mice were inoculated with 5 million trophozoites and the peritoneal washes were obtained at 0, 3, 6, 9, 12 and 15 days after parasite inoculation. Dot plots indicate Ly6C vs F4/80 profiles of live cells obtained from peritoneal washes. 1 = Resident macrophages, 2 = inflammatory macrophages, 3 = polymorphonuclear cells and 4 = monocytes. Data are representative of three independent experiments.

from day 0 to 3, and monocytes from 0.24% to 26.9% on the same post-infection period (Fig. 3). In addition, a decrease in the resident macrophage population (Ly6C<sup>-</sup>, F4/80<sup>+</sup>) was observed,

which was replaced by an important increase in the inflammatory macrophage population (Ly6C<sup>+</sup>, F4/80<sup>+</sup>), 9–12 days after inoculation (52%). Interestingly, at 15 days post-infection, the results

**Table 1.** Total count of cell populations in peritoneal washes from mice infected with *T. vaginalis*

Cell type	Phenotype	Days post-infection					
		0	3	6	9	12	15
Resident macrophages ( $10^3$ )	Ly6C <sup>-</sup> F4/80 <sup>+</sup>	2500	3	8	30	200	23 400
Inflammatory macrophages ( $10^3$ )	Ly6C <sup>+</sup> F4/80 <sup>+</sup>	10	80	600	9500	39 500	1500
Polymorphonuclear cells ( $10^3$ )	Ly6C <sup>medium</sup> F4/80 <sup>-</sup>	400	9500	17 500	12 500	11 000	4500
Monocytes ( $10^3$ )	Ly6C <sup>high</sup> F4/80 <sup>-</sup>	55	7500	7000	4000	1500	150

**Fig. 4.** Lymphocytes number determined by flow cytometry analysis. Changes in B cell (white bars) and T cell (grey bars) population during the infection period. Data are representative of three independent experiments expressed as the mean  $\pm$  s.d. Significance was determined by Kruskal-Wallis analysis and Dunn's test ( $n=6$ , \*\*\* $P < 0.001$ ). \*\*\*Data were compared against time 0

showed a similar trend to uninfected mice (control group) and to those shown at day 0 (Fig. 3).


An increase in absolute numbers for PMN (Ly6C<sup>medium</sup> F4/80<sup>-</sup>) was observed, which gradually increases along the infection. A similar trend was observed in inflammatory macrophages lineage (Ly6C<sup>+</sup> F4/80<sup>+</sup>). Nine days after inoculation, the changes were evident (Table 1). These results suggest that PMN (Ly6C<sup>medium</sup> F4/80<sup>-</sup>) and activated macrophages (Ly6C<sup>+</sup> F4/80<sup>+</sup>) can be participating in the immune response against the parasite.

Next, the presence of CD3 and CD19 corresponding to the T and B lymphocytes (T and B cells) was analysed in the double-negative population (Ly6C<sup>-</sup> F4/80<sup>-</sup>). The results reveal that B cells decreased 3 days after parasite inoculation, from 2.4 to 1.4 million, but subsequently showed steady increases having a maximum peak 15 days after inoculation (21.4 million cells). On the other hand, T cells showed a rapid increase 3 days after infection, they were from 0.3 to 2.2 million cells, and a gradual decrease along the infection until reached the baseline values (Fig. 4).

According to flow cytometry analysis, neutrophils and inflammatory macrophages showed an increase in cell percentage since day 3 and day 9 post-infection, respectively, reaching the maximum at day 12 for macrophages (Fig. 5). Then, we studied the defence mechanisms proposed for those cell lineages in the physiopathology of the infection.

### Trichomonas vaginalis is resistant to nitric oxide production

Phagocytic cells have an arsenal of antimicrobial activities, including the production of RNS and NETs formation. To know the

**Fig. 5.** Timeline for cellular analyses during *T. vaginalis* infection. Mice were intraperitoneally inoculated with the parasite, image shows the sampling times for: flow cytometry analyses, NETs quantification and NO determination. It includes the graph of the main cell population changes observed during early infection.


impact of the RNS produced by these cells on the *T. vaginalis* viability, we used two inhibitors: AG and a nitro-L-arginine (L-NAME). As expected, mice treated with AG or L-NAME produced lower NO quantities than uninfected mice (Fig. 6A). Then, we measured the parasite viability in the presence of these inhibitors. The results show that the inhibitor AG allowed the growth of  $746\,355 \pm 229\,699$  trophozoites per mL whereas in its absence we determined  $707\,844 \pm 178\,775$  trophozoites per mL, which does not represent a significant difference in the parasite viability. With respect to the effect of L-NAME inhibitor on the parasite viability on infected mice,  $350\,881 \pm 234\,913$  trophozoites were quantified but the statistical analysis showed no significant difference in the parasite growth obtained from the peritoneal washes (Fig. 6B).

### NETs participate in the parasite control in peritoneal infection

To investigate the participation of extracellular traps induced by neutrophils on the parasite control, mice were treated with the NADPH inhibitors, metformin (Wang *et al.*, 2015; Carestia *et al.*, 2016) or Cl-amidine (Rein *et al.*, 1980; Candeias *et al.*, 1993; Luo *et al.*, 2006; Song *et al.*, 2010; Kusunoki *et al.*, 2016), blocking the deimination of arginine residues to citrulline in the histones during NETs formation (Wang *et al.*, 2009; Leshner *et al.*, 2012). Thus, the mice were inoculated with the parasite and free DNA was quantified in the peritoneal washes using the Quant-it DNA Pico green kit (Fig. 6C). Also, we determined viable trophozoites in the peritoneal washes extracted from infected mice treated either with metformin or Cl-amidine. Notably, parasite viability increased to 43% and 86% when they were treated with metformin and Cl-amidine, respectively, compared to the control group (Fig. 6D). These results strongly suggest that the formation of extracellular traps contributes significantly to parasite elimination.

### Peritoneal CD4:CD8 lymphocyte ratio increases during the peritoneal infection

In murine intravaginal infection and human vaginal trichomoniasis, CD4 T lymphocytes are usually associated with an



**Fig. 6.** Impact of NO production and NETs formation in the parasite viability. Mice were inoculated with 5 million trophozoites; (A) nitrite was quantified by modified Griess assay (Material and methods) in the presence or absence of NOS2 enzyme inhibitors, aminoguanidine (AG) or L-nitro arginine methyl ester (L-NAME); (B) parasite viability in peritoneal washes from AG or L-NAME-treated or untreated infected mice; (C) NETs were quantified by Quant it DNA kit in peritoneal washes from metformin-treated or Cl-amidine-treated infected mice; (D) parasite viability in peritoneal washes from metformin or Cl-amidine-treated mice. Data are representative of three independent experiments expressed as the mean  $\pm$  s.d. Significance was determined by Kruskal–Wallis analysis and Dunn’s post-test ( $n=6$ ,  $***P<0.001$ ). ns = non-significant.


inflammatory process (Critchlow *et al.*, 1995; Levine *et al.*, 1998; Smith and Garber, 2015). CD4 cells are important for B-cell germinal centre formation (MacLennan *et al.*, 1997) and adaptive immunity coordination that help CD8 T lymphocyte and B lymphocyte responses (Gasper *et al.*, 2014). To investigate the CD4:CD8 lymphocyte ratio changes during the infection, we analysed the proportion of these populations in the positive CD3 gate of peritoneal cells obtained from *T. vaginalis*-infected group. The flow cytometry analysis revealed that infected mice presented an average increase of CD3<sup>+</sup>CD4<sup>+</sup> cell population reaching 87.1% at day 12, thus representing a significant increase compared with mice cells at 0 time of 55.7%. Later, CD3<sup>+</sup>CD8<sup>+</sup> cells did not exhibit significant changes along the entire infection period as it is appreciated because we determined a 7.7% ratio of this population in uninfected mice and a 7.15% ratio in infected mice (Fig. 7A).

#### Intense proliferation of memory T cells is induced during trichomoniasis

We have previously reported a proinflammatory profile prompted in the host by *T. vaginalis* infections, which include the

production of nitric oxide and various cytokines such as TNF- $\alpha$ , IL-6, IL-10 that may suggest a Th1-type response (Olmos-Ortiz *et al.*, 2017). To clarify the participation of T cells, we analysed cell-specific surface markers, to reveal *T. vaginalis*-induced activation and differentiation of T effector cells. CD4 T cells differentiate into Th1 cells and the effector Th1 cells express Ly6c marker in specific response to viral infection (Matsuda *et al.*, 2006; Marshall *et al.*, 2011). We determined for the first time if effector CD4<sup>+</sup>Ly6C<sup>+</sup> T cells arise during trichomoniasis. Our results show an increased CD4<sup>+</sup>Ly6C<sup>+</sup> cell population during early infection in peritoneal washes, 3 and 6 days post-infection but not 9 days post-infection (Fig. 7B, up).

In addition, it is known that CD8<sup>+</sup>T cells primarily differentiate into cytotoxic T lymphocytes (CTLs) or memory T cells (Tcm). F4/80 marker expression has been observed on activated CD8<sup>+</sup>T cells from mice in acute viral infections (Lin *et al.*, 2003), which indicates the activation of these cells. Because its presence has not been investigated in trichomoniasis, the presence of these cells was explored, and the expression of F4/80 on CD8<sup>+</sup>T cells was analysed. The peritoneal cells obtained from uninfected mice exhibited a high percentage of F4/80<sup>+</sup>CD8<sup>+</sup>T cells (8%) whereas peritoneal cells from infected mice expressed only 2.6



**Fig. 7.** Analysis of lymphocytes populations and subpopulations. Mice were inoculated with 5 million trophozoites and peritoneal washes were obtained at 0, 3, 6, 9, 12 and 15 days after infection. (A) Dot plot of CD4:CD8 T cells, (B) Th1 cells were gated as a CD4<sup>+</sup>Ly6C<sup>+</sup> population (up); activated T cells were gated like a CD8<sup>+</sup>F4/80<sup>+</sup> population (middle), and Tcm cells were gated for the presence of Ly6C marker (CD8<sup>+</sup>Ly6C<sup>+</sup>) (down). Images are representative of three independent experiments using  $n = 6$ .

or 1.5% of that cell type, 3 or 6 days after infection, respectively. The F4/80 expression in T cells emerged as 10%, 9 days after infection but reduced until 5% at 15 days after infection (Fig. 7B, down), just after the peak of the CTL response.

On the other hand, Ly6C has been referenced as a characteristic cell-surface marker of Tcm cells and is involved in the migration properties of lymphocytes (Hänninen *et al.*, 2011). To know if CD8<sup>+</sup>T cells matured towards Tcm, we analysed the presence of the Ly6c marker in CD8<sup>+</sup>T cells. The results showed a notable increase (49.6%) of this marker 3 days after infection, while uninfected cells did not (12.4%); the maximum peak of this expression is observed 6 days after infection (80.3%) and show a decreasing trend 9, 12 and 15 days after infection (48.7, 25.7 and 13.29%, respectively) (Fig. 7B, middle). Those results are according to

the previous reports that describe the function of Ly6C marker on the subsets of memory T cells (Hänninen *et al.*, 2011). The above means that even though the lymphocytes lineage does not lose its proliferative capacity, most T cells differentiate into Tcm and fewer develop a cytotoxic activated cell (T cytotoxic).

## Discussion

We established an intraperitoneal murine model of trichomoniasis for the study of the immune cell populations during their interaction with *T. vaginalis* allowing the survival for both host and parasite. Our interest was to determine the earliest events of the immune response, rather than the infection mechanism. For that, we assured the infection according to Gomez-Barrio *et al.*



(2002) observing pathologic damages such as visceral necrosis spots and ascites, but we also assured the host survival to allow the manifestation of the intense immune response induced (Nogal Ruiz *et al.*, 1997). Also, we evaluated other infection signs such as weight gain of the animal model, physical condition and changes of lymphocyte number in blood samples (Fig. 1D), as well as the presence and viability of the parasite along all the infection period for our experiments. The spleen–body weight ratio was greater in our evaluation than the difference of previously reported values for the subcutaneous inoculation (Mason and Gwanzura, 1990). Our peritoneal model showed a spleen–body weight ratio of 7.1, whereas this ratio was 5.44 for the subcutaneous inoculation. Previously reported spleen–body weight ratios were obtained during a subcutaneous mouse challenge, with various pieces of information missing, such as the dose and strain of patient-collected *T. vaginalis* used. Then, we can conclude that our intraperitoneal model induces a strong immune response. It should be noted that in healthy mice the spleen–body weight ratio did not show significant differences in either intraperitoneal or subcutaneous models (3.96 and 4.36, respectively). Furthermore, immunophenotyping performed in peritoneal washes obtained from mice infected with *T. vaginalis* (Fig. 2) allowed us to establish a dynamic scheme of the immune cell recruitment in response to this parasitic invasion. Our model opens the possibility to perform additional studies about the *in vivo* host–parasite interaction free from the influence of hormones and vaginal microbiota.

PMN cell and monocyte populations were kept in constant increase during early infection, and they were replaced gradually later by a high presence of inflammatory macrophages 9 days after infection. As IL-8 is a known mediator of the early immune response to *T. vaginalis*, these could be responsible for the initial neutrophil accumulation observed (Shaio *et al.*, 1992, 1994, 1995; Ryu *et al.*, 2004; Nam *et al.*, 2012).

The inverse correlation observed between the gradual decrease in the numbers of Ly6C<sup>+</sup> monocytes and the increase in the numbers of inflammatory macrophages is expected and the result of inflamed-site infiltration by stimulated monocytes which differentiate into inflammatory macrophages and dendritic cells (Nahrendorf *et al.*, 2007; Misharin *et al.*, 2014; Morias *et al.*, 2015). Although these two populations express similar surface markers, an adequate panel of antibodies for flow cytometry let us perform an accurate identification of individual cellular subsets (Misharin *et al.*, 2013) but it is still necessary to perform a precise and complete identification of the relative and absolute composition of inflammatory cell populations in response to trichomoniasis.

Also, an increase in the CD4:CD8 T lymphocytes ratio has been observed in cervical biopsies from infected women using FACS analyses (Levine *et al.*, 1998; Reighard *et al.*, 2011). Even if this is a different model, it is possible in human cells to activate the defence in a similar way. Therefore, we propose this host–parasite interaction system to study *T. vaginalis* early infection and quantify several cell types participating in this process, also to define mechanisms that affect the parasite survival.

The relation between *T. vaginalis* infection and HIV susceptibility is not clear yet. Several hypotheses have been proposed to explain it, for example, the increase in susceptible leucocytes in the infection site including CD4 lymphocytes, monocytes and Langerhans cells (Edwards and Morris, 1985; Levine *et al.*, 1998; Smith and Garber, 2015). Although it has been reported that there is an increase in CD4<sup>+</sup> over the development of the infection, the role of those cells in parasite clearance is currently unclear. CD4<sup>+</sup> T cells give rise to distinct functional cell types, such as Th1, Th2, Th17 and Treg cells depending on the infection type and cytokines produced. In our work group, the presence of IL-13 and IL-17 has been determined in vaginal washes of

infected mice (Olmos-Ortiz *et al.*, 2017). Our observed increase in the CD3<sup>+</sup>CD4<sup>+</sup> T cell compartment up to 3 days post-infection is consistent with low IL-3 and IL-17 at earlier time points.

To the best of our knowledge, there is currently scarce information regarding the role of memory T cells during trichomoniasis infections. Therefore, our results showing the development of memory T cells during the investigated timeframe are highly novel. These memory T cells could provide protection or a faster response upon a reinfection event. Memory cells arise from the differentiation of naïve cells incorporating surface molecules such as Ly6C in T lymphocytes in different systems (Hänninen *et al.*, 2011; Marshall *et al.*, 2011; Terrazas *et al.*, 2017). Ly6C is a haematopoietic mouse differentiation marker and sends signals for T-cell activation and cytokine production. Similarly, a role of Ly6C marker has been attributed in the T-cell immunity as a T-cell development, and in the helper-T-cell functional division. Ly6C is expressed in both mature CD8 and CD4 T cells and is involved in the homing of memory T cells (Tcm) (Walunas *et al.*, 1995; Henderson *et al.*, 2002; Hänninen *et al.*, 2011).

Our results showed an early increase in CD4<sup>+</sup>Ly6C<sup>+</sup> population, 3 and 6 days post-inoculation, followed by an increase in CD8<sup>+</sup>Ly6C<sup>+</sup> cells that remain increased during the time of our research. These results are in accordance with a previous report using a model of acute lymphocytic choriomeningitis virus (Marshall *et al.*, 2011), in the same way that we observed an increase in B lymphocytes at 15 days post-infection, and consistent with reports that refer to the presence of specific immunoglobulins against *T. vaginalis* (Smith and Garber, 2015). Nevertheless, trichomoniasis results in a constant reinfection (Abraham *et al.*, 1996; Petrin *et al.*, 1998; Smith and Garber, 2015). It will be important to investigate the type of CD4<sup>+</sup> or CD8<sup>+</sup> T cells generated and differentiated into memory T cells during infection. In addition, it will be important to elucidate the signalling route followed to differentiate into memory T cells and if these memory T cells remain for a long time.

Monocyte-macrophage lineage cells showed an increased presence in infected mice, as was revealed by flow cytometry data. These cells possess diverse mechanisms to eliminate the pathogens, such as the RNS production. Our experiments showed a high production of NO in infected mice as other researchers have reported for *in vitro* assays (Park *et al.*, 1997; Han *et al.*, 2009; Bogdan, 2015). Nevertheless, the cytotoxic activity was not efficient *in vivo* to reduce parasite viability, because we still observed that NO increase up to eight times in infected mice in comparison to the control group. The above suggests that NO produced by iNOS is not a definitive tool of defence against *T. vaginalis* although it has been described that NO have cytotoxic activity against different parasites such as *Schistosomas mansoni*, *Leishmania major* and *Toxoplasma gondii* (James and Glaven, 1989; Green *et al.*, 1990; James and Hibbs, 1990). *In vitro* experiments using murine peritoneal macrophages have shown guidance to decrease the *T. vaginalis* viability by NO cytotoxic activity (Park *et al.*, 1997). However, our research showed null activity of NO against the parasite probably due to the different concentrations of NO released by macrophages (18 and 8 μM, respectively). It is known that NO cause toxicity through the inhibition of hydrogenosomal enzyme activities (Ryu and Lloyd, 1995). Nevertheless, it has been widely reported that the microorganisms have developed various mechanisms to resist cytotoxic activity by NO, such as the consumption of arginine to deprive host cells of nitric oxide synthase 2 (NOS2) (Stadelmann *et al.*, 2013; Cusumano *et al.*, 2014). In our model, NO can act as a second messenger for *T. vaginalis* that trigger the cyclic guanosine monophosphate (cGMP) production to stabilize the hydrogenosomal membrane, especially in iron depletion conditions (Cheng *et al.*, 2015), promoting the intraperitoneal infection.

One of the mechanisms of neutrophils to trap and kill pathogens is through the formation of NETs (Brinkmann *et al.*, 2004; Urban *et al.*, 2006; Urban *et al.*, 2009). Previously, our research group observed that neutrophils undergo NETosis in the presence of *T. vaginalis in vitro* (Huynh and Ávila-Muro, 2015). NETs formation involves a series of steps, which include the generation of ROS by the NADPH oxidase (Fuchs *et al.*, 2007; Röhm *et al.*, 2014), and the citrullination of histone H3 that converts arginine residues to citrulline (Wang *et al.*, 2009). This reaction is catalysed by PAD4 (Leshner *et al.*, 2012). When we used metformin and Cl-amidine to inhibit both NADPH oxidase and PAD4, we showed an increased *T. vaginalis* viability up to 43% using metformin, and 86% using Cl-amidine. Throughout those experiments, we demonstrate the importance of NETs participation to control *T. vaginalis* growth. NETs are capable to capture and kill pathogens such as bacteria and fungus (Brinkmann *et al.*, 2004; Urban *et al.*, 2006), because the DNA of NETs is associated with histones and granular proteins with antimicrobial activities (Urban *et al.*, 2009; Papayannopoulos *et al.*, 2010; Metzler *et al.*, 2014). However, recent research has revealed that using DNase I to break NETs in assays *in vitro* does not affect the ability of PMN cells to kill trichomonas in a system free of other immune cells (Rein *et al.*, 1980; Mercer *et al.*, 2018). Hence the mechanisms involved require further investigations.

Previously it has been described that ROS inhibition in PMN cells interacting with *T. vaginalis* improves the ability of the parasite to survive (Rein *et al.*, 1980). These reports suggest that ROS have a direct activity on the parasite and PMN cells undergo apoptosis in the cell–parasite interaction. We eliminate ROS production induced by trichomonas by the inhibition of NADPH oxidase activity (Song *et al.*, 2008). Nevertheless, our results show that the ROS effect on the reduction of *T. vaginalis* viability was indirect probably because NETosis could also be inhibited. Our results, based on new knowledge, suggest that NADPH oxidase-derived ROS trigger the NETosis, and the traps are decorated with antimicrobial peptides promoting parasite death. In spite of previous reports suggesting that NETosis has a little contribution on PMN cells on the parasite elimination *in vitro* (Mercer *et al.*, 2018), we believe that it is very important to consider the participation of other immune cells or antimicrobial peptides present in the NETs during the *in vivo* infection, as it happens in our model of trichomoniasis. In consequence, it is necessary to elucidate how NETs-antimicrobial peptides contribute to kill the parasite or how NETs catch the parasite while other cells attack and destroy it.

To our knowledge, this is the first report about the relative and total count of immune cells arriving at the early infection of *T. vaginalis* in an *in vivo* model, which has allowed us to explore the role that those cell populations play, both individually and as a team during trichomoniasis. Likewise, the implementation of this *in vivo* model will allow us to investigate the strategies used by the immune cells on the whole organism for parasite clearance.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182021001311>

**Acknowledgements.** The authors thank Dr Felipe Padilla-Vaca and Dr Fernando Anaya-Velazquez for their kind donation of *T. vaginalis* GT-21 strain.

**Author contributions.** Cuéllar-Mata and Ávila conceived, designed the study and reviewed the manuscript. Rangel-Mata conducted data gathering and wrote the manuscript. Arriaga-Pizano established the antibodies panel for flow cytometry assays and facilitated the FlowCytometer; Reyes-Martínez and Brunck performed the data analysis. Olmos-Ortiz preserved parasite cultures and reactants.

**Financial support.** Rangel-Mata was supported by a scholarship from CONACyT, México. The project was supported by CONACyT (CB-2012-01 182671) and the Universidad de Guanajuato (CIUG 2016-2017).

**Conflict of interest.** None.

**Ethical standards.** All animals were handled according to the Mexican norm NOM-062-1999. Permission for this research was granted by the Ethics Committee of the University of Guanajuato.

## References

- Abraham MC, Desjardins M, Filion LG and Garber GE (1996) Inducible immunity to *Trichomonas vaginalis* in a mouse model of vaginal infection. *Infection and Immunity* **64**, 3571–3575.
- Alderete JF and Garza GE (1985) Specific nature of *Trichomonas vaginalis* parasitism of host cell surfaces. *Infection and Immunity* **50**, 701–708.
- Arias-Negrete S, Jiménez-Romero LA, Solís-Martínez MO, Ramírez-Emiliano J, Avila EE and Cuéllar-Mata P (2004) Indirect determination of nitric oxide production by reduction of nitrate with a freeze-thawing-resistant nitrate reductase from *Escherichia coli* MC1061. *Analytical Biochemistry* **328**, 14–21.
- Barajas-Mendiola MA, Castellano LE, Barrios-Rodiles M, Deveze-Alvarez MA, Avila EE and Cuéllar-Mata P (2019) Reduced *Trichomonas vaginalis* viability in mice pretreated with parasite DNA. *Parasitology* **146**, 1636–1645.
- Bogdan C (2015) Nitric oxide synthase in innate and adaptive immunity: an update. *Trends in Immunology* **36**, 161–178.
- Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y and Zychlinsky A (2004) Neutrophil extracellular traps kill bacteria. *Science (New York, N.Y.)* **303**, 1532–1535.
- Candeias LP, Patel KB, Stratford MR and Wardman P (1993) Free hydroxyl radicals are formed on reaction between the neutrophil-derived species superoxide anion and hypochlorous acid. *FEBS Letters* **333**, 151–153.
- Carestia A, Frechtel G, Cerrone G, Linari MA, Gonzalez CD, Casais P and Schattner M (2016) NETosis before and after hyperglycemic control in type 2 diabetes mellitus patients. *PLoS ONE* **11**, e0168647.
- Cheng WH, Huang KY, Huang PJ, Hsu JH, Fang YK, Chiu CH and Tang P (2015) Nitric oxide maintains cell survival of *Trichomonas vaginalis* upon iron depletion. *Parasites & Vectors* **8**, 393.
- Critchlow CW, Wolner-Hanssen P, Eschenbach DA, Kiviat NB, Koutsky LA, Stevens CE and Holmes KK (1995) Determinants of cervical ectopia and of cervicitis: age, oral contraception, specific cervical infection, smoking, and douching. *American Journal Obstetrics & Gynecology* **173**, 534–543.
- Cusumano ZT, Watson ME and Caparon MG (2014) *Streptococcus pyogenes* arginine and citrulline catabolism promotes infection and modulates innate immunity. *Infection and Immunity* **82**, 233–242.
- Edwards JN and Morris HB (1985) Langerhans' cells and lymphocyte subsets in the female genital tract. *British Journal of Obstetrics and Gynaecology* **92**, 974–982.
- Fichorova RN (2009) Impact of *T. vaginalis* infection on innate immune responses and reproductive outcome. *Journal of Reproductive Immunology* **83**, 185–189.
- Fraga J, Rojas L, Sariego I, Fernández-Calienes A and Nuñez FA (2012) Species typing of Cuban *Trichomonas vaginalis* virus by RT-PCR, and association of TVV-2 with high parasite adhesion levels and high pathogenicity in patients. *Archives of Virology* **157**, 1789–1795.
- Fuchs TA, Abed U, Goosmann C, Hurwitz R, Schulze I, Wahn V, Weinrauch Y, Brinkmann V and Zychlinsky A (2007) Novel cell death program leads to neutrophil extracellular traps. *Journal of Cell Biology* **176**, 231–241.
- Gasper DJ, Tejera MM and Suresh M (2014) CD4 T-cell memory generation and maintenance. *Critical Reviews in Immunology* **34**, 121–146.
- Gomez-Barrio A, Nogal-Ruiz J.J., Montero-Pereira D, Rodríguez-Gallego E, Romero-Fernández E and Escario J.A. (2002) Biological variability in clinical isolates of *Trichomonas vaginalis*. *Memorias do Instituto Oswaldo Cruz* **97**, 893–896.
- Goodman RP, Freret TS, Kula T, Geller AM, Talkington MW, Tang-Fernandez V, Suciú O, Demidenko AA, Ghabrial SA, Beach DH, Singh BN, Fichorova RN and Nibert ML (2011) Clinical isolates of *Trichomonas vaginalis* concurrently infected by strains of up to four

- Trichomonas* virus species (Family Totiviridae). *Journal of Virology* **85**, 4258–4270.
- Green SJ, Mellouk S, Hoffman SL, Meltzer MS and Nacy CA** (1990) Cellular mechanisms of nonspecific immunity to intracellular infection: cytokine-induced synthesis of toxic nitrogen oxides from L-arginine by macrophages and hepatocytes. *Immunology Letters* **25**, 15–19.
- Han IH, Goo SY, Park SJ, Hwang SJ, Kim YS, Yang MS, Ahn MH and Ryu JS** (2009) Proinflammatory cytokine and nitric oxide production by human macrophages stimulated with *Trichomonas vaginalis*. *Korean Journal of Parasitology* **47**, 205–212.
- Hänninen A, Maksimov M, Alam C, Morgan DJ and Jalkanen S** (2011) Ly6C supports preferential homing of central memory CD8<sup>+</sup> T cells into lymph nodes. *European Journal of Immunology* **41**, 634–644.
- Henderson SC, Kamdar MM and Bamezai A** (2002) Ly-6A.2 expression regulates antigen-specific CD4<sup>+</sup> T cell proliferation and cytokine production. *Journal of Immunology* **168**, 118–126.
- Huynh AV and Ávila-Muro EE** (2015) Efecto del anticuerpo contra el receptor TLR4 en la formación de trampas extracelulares de neutrófilos inducidas por el parásito *T. vaginalis*. *Jovenes en la Ciencia* **1**, 245–249.
- James SL and Glaven J** (1989) Macrophage cytotoxicity against schistosomula of *Schistosoma mansoni* involves arginine-dependent production of reactive nitrogen intermediates. *Journal of Immunology* **143**, 4208–4212.
- James SL and Hibbs JB** (1990) The role of nitrogen oxides as effector molecules of parasite killing. *Parasitology Today* **6**, 303–305.
- Kharsany AB, Hoosen AA, Moodley J, Bagaratee J and Gouws E** (1993) The association between sexually transmitted pathogens and cervical intra-epithelial neoplasia in a developing community. *Genitourinary Medicine* **69**, 357–360.
- Kissinger P** (2015) *Trichomonas vaginalis*: a review of epidemiologic, clinical and treatment issues. *BMC Infectious Diseases* **15**, 307.
- Kissinger P, Amedee A, Clark RA, Dumestre J, Theall KP, Myers L, Hagensee ME, Farley TA and Martin DH** (2009) *Trichomonas vaginalis* treatment reduces vaginal HIV-1 shedding. *Sexually Transmitted Diseases* **36**, 11–16.
- Knight JS, Zhao W, Luo W, Subramanian V, O'Dell AA, Yalavarthi S, Hodgins JB, Eitzman DT, Thompson PR and Kaplan MJ** (2013) Peptidylarginine deiminase inhibition is immunomodulatory and vasculoprotective in murine lupus. *The Journal of Clinical Investigation* **123**, 2981–2993.
- Kusunoki Y, Nakazawa D, Shida H, Hattanda F, Miyoshi A, Masuda S, Nishio S, Tomaru U, Atsumi T and Ishizu A** (2016) Peptidylarginine deiminase inhibitor suppresses neutrophil extracellular trap formation and MPO-ANCA production. *Frontiers in Immunology* **7**, 227.
- Lazenby GB, Soper DE and Nolte FS** (2013) Correlation of leukorrhea and *Trichomonas vaginalis* infection. *Journal of Clinical Microbiology* **51**, 2323–2327.
- Leshner M, Wang S, Lewis C, Zheng H, Chen XA, Santy L and Wang Y** (2012) PAD4 mediated histone hypercitrullination induces heterochromatin decondensation and chromatin unfolding to form neutrophil extracellular trap-like structures. *Frontiers in Immunology* **3**, 307.
- Levine WC, Pope V, Bhoonkar A, Tambe P, Lewis JS, Zaidi AA, Farshy CE, Mitchell S and Talkington DF** (1998) Increase in endocervical CD4 lymphocytes among women with nonulcerative sexually transmitted diseases. *The Journal of Infectious Diseases* **177**, 167–174.
- Lin Y, Roberts TJ, Sriram V, Cho S and Brutkiewicz RR** (2003) Myeloid marker expression on antiviral CD8<sup>+</sup> T cells following an acute virus infection. *European Journal of Immunology* **33**, 2736–2743.
- Luo Y, Arita K, Bhatia M, Knuckley B, Lee YH, Stallcup MR, Sato M and Thompson PR** (2006) Inhibitors and inactivators of protein arginine deiminase 4: functional and structural characterization. *Biochemistry* **45**, 11727–11736.
- MacLennan IC, Gulbranson-Judge A, Toellner KM, Casamayor-Palleja M, Chan E, Sze DM, Luther SA and Orbea HA** (1997) The changing preference of T and B cells for partners as T-dependent antibody responses develop. *Immunology Reviews* **156**, 53–66.
- Malla N, Valadkhani Z, Harjai K, Sharma S and Gupta I** (2004) Reactive nitrogen intermediates in experimental trichomoniasis induced with isolates from symptomatic and asymptomatic women. *Parasitology Research* **94**, 101–105.
- Marcel AM, Rojas L, Rodríguez SS, López Y and Ramos IS** (2001) [Application of an alternative method to counting in Neubauer chamber to determine *Trichomonas vaginalis* concentrations]. *Revista Cubana de Medicina Tropical* **53**, 180–188.
- Marshall HD, Chandele A, Jung YW, Meng H, Poholek AC, Parish IA, Rutishauser R, Cui W, Kleinstein SH, Craft J and Kaech SM** (2011) Differential expression of Ly6C and T-bet distinguish effector and memory Th1 CD4(+) cell properties during viral infection. *Immunity* **35**, 633–646.
- Mason PR and Gwanzura L** (1990) Reduced lymphocyte responses to mitogens in natural and experimental trichomoniasis. *Infection and Immunity* **58**, 3553–3557.
- Matsuda JL, Zhang Q, Ndonge R, Richardson SK, Howell AR and Gapin L** (2006) T-bet concomitantly controls migration, survival, and effector functions during the development of Valpha14i NKT cells. *Blood* **107**, 2797–2805.
- McClelland RS, Sangare L, Hassan WM, Lavreys L, Mandaliya K, Kiarie J, Ndinya-Achola J, Jaoko W and Baeten JM** (2007) Infection with *Trichomonas vaginalis* increases the risk of HIV-1 acquisition. *The Journal of Infectious Diseases* **195**, 698–702.
- Menezes CB and Tasca T** (2016) Trichomoniasis immunity and the involvement of the purinergic signaling. *Biomedical Journal* **39**, 234–243.
- Mercer F, Ng SH, Brown TM, Boatman G and Johnson PJ** (2018) Neutrophils kill the parasite *Trichomonas vaginalis* using trogocytosis. *PLoS Biology* **16**, e2003885.
- Metzler KD, Goosmann C, Lubojemska A, Zychlinsky A and Papayannopoulos V** (2014) A myeloperoxidase-containing complex regulates neutrophil elastase release and actin dynamics during NETosis. *Cell Reports* **8**, 883–896.
- Misharin AV, Morales-Nebreda L, Mutlu GM, Budinger GR and Perlman H** (2013) Flow cytometric analysis of macrophages and dendritic cell subsets in the mouse lung. *American Journal of Respiratory Cell Molecular Biology* **49**, 503–510.
- Misharin AV, Cuda CM, Saber R, Turner JD, Gierut AK, Haines GK, Berdnikovs S, Filer A, Clark AR, Buckley CD, Mutlu GM, Budinger GR and Perlman H** (2014) Nonclassical Ly6C(-) monocytes drive the development of inflammatory arthritis in mice. *Cell Reports* **9**, 591–604.
- Morias Y, Abels C, Laoui D, Van Overmeire E, Williams M, Schoupe E, Tacke F, deVries CJ, De Baetselier P and Beschin A** (2015) Ly6C-monocytes regulate parasite-induced liver inflammation by inducing the differentiation of pathogenic Ly6C<sup>+</sup> monocytes into macrophages. *PLoS Pathogens* **11**, e1004873.
- Nahrendorf M, Swirski FK, Aikawa E, Stangenberg L, Wurdinger T, Figueiredo JL, Libby P, Weissleder R and Pittet MJ** (2007) The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. *Journal of Experimental Medicine* **204**, 3037–3047.
- Nam YH, Min A, Kim SH, Lee YA, Kim KA, Song KJ and Shin MH** (2012) Leukotriene B(4) receptors BLT1 and BLT2 are involved in interleukin-8 production in human neutrophils induced by *Trichomonas vaginalis*-derived secretory products. *Inflammation Research* **61**, 97–102.
- Nemati M, Malla N, Yadav M, Khorramdelazad H and Jafarzadeh A** (2018) Humoral and T cell-mediated immune response against trichomoniasis. *Parasite Immunology* **40**, 1–11. doi: 10.1111/pim.12510
- Nogal Ruiz JJ, Escario JA, Martinez Diaz RA and Gomez Barrio A** (1997) Evaluation of a murine model of experimental trichomoniasis. *Parasite* **4**, 127–132.
- Olmos-Ortiz LM, Barajas-Mendiola MA, Barrios-Rodiles M, Castellano LE, Arias-Negrete S, Avila EE and Cuéllar-Mata P** (2017) *Trichomonas vaginalis* exosome-like vesicles modify the cytokine profile and reduce inflammation in parasite-infected mice. *Parasite Immunology* **39**, 1–10. doi: 10.1111/pim.12426
- Papayannopoulos V, Metzler KD, Hakkim A and Zychlinsky A** (2010) Neutrophil elastase and myeloperoxidase regulate the formation of neutrophil extracellular traps. *Journal of Cell Biology* **191**, 677–691.
- Park GC, Ryu JS and Min DY** (1997) The role of nitric oxide as an effector of macrophage-mediated cytotoxicity against *Trichomonas vaginalis*. *The Korean Journal of Parasitology* **35**, 189–195.
- Petrin D, Delgaty K, Bhatt R and Garber G** (1998) Clinical and microbiological aspects of *Trichomonas vaginalis*. *Clinical Microbiology Reviews* **11**, 300–317.
- Reighard SD, Sweet RL, Vicetti Miguel C, Vicetti Miguel RD, Chivukula M, Krishnamurti U and Cherpes TL** (2011) Endometrial leukocyte subpopulations associated with *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis* genital tract infection. *American Journal of Obstetrics & Gynecology* **205**, 324.e321–327.

- Rein MF, Sullivan JA and Mandell GL (1980) Trichomonocidal activity of human polymorphonuclear neutrophils: killing by disruption and fragmentation. *The Journal of Infection Diseases* **142**, 575–585.
- Röhm M, Grimm MJ, D'Auria AC, Almyroudis NG, Segal BH and Urban CF (2014) NADPH oxidase promotes neutrophil extracellular trap formation in pulmonary aspergillosis. *Infection and Immunity* **82**, 1766–1777.
- Ryu JS and Lloyd D (1995) Cell cytotoxicity of sodium nitrite, sodium nitroprusside and Roussin's black salt against *Trichomonas vaginalis*. *FEMS Microbiology Letters* **130**, 183–187.
- Ryu JS, Kang JH, Jung SY, Shin MH, Kim JM, Park H and Min DY (2004) Production of interleukin-8 by human neutrophils stimulated with *Trichomonas vaginalis*. *Infection and Immunity* **72**, 1326–1332.
- Shaio MF, Lin PR, Lee CS, Hou SC, Tang P and Yang KD (1992) A novel neutrophil-activating factor released by *Trichomonas vaginalis*. *Infection and Immunity* **60**, 4475–4482.
- Shaio MF, Lin PR, Liu JY and Tang KD (1994) Monocyte-derived interleukin-8 involved in the recruitment of neutrophils induced by *Trichomonas vaginalis* infection. *The Journal of Infection Diseases* **170**, 1638–1640.
- Shaio MF, Lin PR, Liu JY and Yang KD (1995) Generation of interleukin-8 from human monocytes in response to *Trichomonas vaginalis* stimulation. *Infection and Immunity* **63**, 3864–3870.
- Smith JD and Garber GE (2015) *Trichomonas vaginalis* infection induces vaginal CD4+ T-cell infiltration in a mouse model: a vaccine strategy to reduce vaginal infection and HIV transmission. *The Journal of Infection Diseases* **212**, 285–293.
- Song HO, Shin MH, Ahn MH, Min DY, Kim YS and Ryu JS (2008) *Trichomonas vaginalis*: reactive oxygen species mediates caspase-3 dependent apoptosis of human neutrophils. *Experimental Parasitology* **118**, 59–65.
- Song HO, Lim YS, Moon SJ, Ahn MH and Ryu JS (2010) Delayed human neutrophil apoptosis by *Trichomonas vaginalis* lysate. *The Korean Journal of Parasitology* **48**, 1–7.
- Stadelmann B, Hanevik K, Andersson MK, Bruserud O and Svärd SG (2013) The role of arginine and arginine-metabolizing enzymes during Giardia – host cell interactions in vitro. *BMC Microbiology* **13**, 256.
- Swygard H, Seña AC, Hobbs MM and Cohen MS (2004) Trichomoniasis: clinical manifestations, diagnosis and management. *Sexually Transmitted Infections* **80**, 91–95.
- Terrazas C, Varikuti S, Oghumu S, Steinkamp HM, Ardic N, Kimble J, Nakhasi H and Satoskar AR (2017) Ly6C. *Scientific Report* **7**, 14693.
- Twu O, de Miguel N, Lustig G, Stevens GC, Vashisht AA, Wohlschlegel JA and Johnson PJ (2013) *Trichomonas vaginalis* exosomes deliver cargo to host cells and mediate host:parasite interactions. *PLoS Pathogens* **9**, e1003482.
- Urban CF, Reichard U, Brinkmann V and Zychlinsky A (2006) Neutrophil extracellular traps capture and kill *Candida albicans* yeast and hyphal forms. *Cell Microbiology* **8**, 668–676.
- Urban CF, Ermert D, Schmid M, Abu-Abed U, Goosmann C, Nacken W, Brinkmann V, Jungblut PR and Zychlinsky A (2009) Neutrophil extracellular traps contain calprotectin, a cytosolic protein complex involved in host defense against *Candida albicans*. *PLoS Pathogens* **5**, e1000639.
- Walunas TL, Bruce DS, Dustin L, Loh DY and Bluestone JA (1995) Ly-6C is a marker of memory CD8+ T cells. *The Journal of Immunology* **155**, 1873–1883.
- Wang Y, Li M, Stadler S, Correll S, Li P, Wang D, Hayama R, Leonelli L, Han H, Grigoryev SA, Allis CD and Coonrod SA (2009) Histone hyperacetylation mediates chromatin decondensation and neutrophil extracellular trap formation. *Journal of Cell Biology* **184**, 205–213.
- Wang H, Li T, Chen S, Gu Y and Ye S (2015) Neutrophil extracellular trap mitochondrial DNA and its autoantibody in systemic lupus erythematosus and a proof-of-concept trial of metformin. *Arthritis & Rheumatology* **67**, 3190–3200.
- Yano A, Aosai F, Yui K, Kojima S and Kawana T (1983) Antigen-specific proliferation responses of peripheral blood lymphocytes to *Trichomonas vaginalis* antigen in patients with *Trichomonas vaginitis*. *Journal of Clinical Microbiology* **17**, 175–180.