

This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

doi:10.1017/etds.2017.42

Badly approximable points on self-affine sponges and the lower Assouad dimension

TUSHAR DAS[†], LIOR FISHMAN[‡], DAVID SIMMONS[§] and
MARIUSZ URBANŃSKI[‡]

[†] *University of Wisconsin – La Crosse, Department of Mathematics & Statistics,
1725 State Street, La Crosse, WI 54601, USA
(e-mail: tdas@uwlax.edu)*

[‡] *University of North Texas, Department of Mathematics, 1155 Union Circle #311430,
Denton, TX 76203-5017, USA
(e-mail: lior.fishman@unt.edu, urbanski@unt.edu)*

[§] *University of York, Department of Mathematics, Heslington, York YO10 5DD, UK
(e-mail: David.Simmons@york.ac.uk)*

(Received 23 August 2016 and accepted in revised form 20 April 2017)

Abstract. We highlight a connection between Diophantine approximation and the lower Assouad dimension by using information about the latter to show that the Hausdorff dimension of the set of badly approximable points that lie in certain non-conformal fractals, known as self-affine sponges, is bounded below by the dynamical dimension of these fractals. For self-affine sponges with equal Hausdorff and dynamical dimensions, the set of badly approximable points has full Hausdorff dimension in the sponge. Our results, which are the first to advance beyond the conformal setting, encompass both the case of Sierpiński sponges/carpets (also known as Bedford–McMullen sponges/carpets) and the case of Barański carpets. We use the fact that the lower Assouad dimension of a hyperplane diffuse set constitutes a lower bound for the Hausdorff dimension of the set of badly approximable points in that set.

Fix $d \in \mathbb{N}$. Dirichlet’s theorem in Diophantine approximation states that, for all $\mathbf{x} \in \mathbb{R}^d$, there exist infinitely many rational points $\mathbf{p}/q \in \mathbb{Q}^d$ such that

$$\left\| \mathbf{x} - \frac{\mathbf{p}}{q} \right\| < \frac{1}{q^{1+1/d}}.$$

A point $\mathbf{x} \in \mathbb{R}^d$ is said to be *badly approximable* if this inequality cannot be improved by more than a constant, i.e. if there exists a constant $c > 0$ such that, for any rational

point $\mathbf{p}/q \in \mathbb{Q}^d$, we have

$$\left\| \mathbf{x} - \frac{\mathbf{p}}{q} \right\| \geq \frac{c}{q^{1+1/d}}.$$

We denote the set of all badly approximable points in \mathbb{R}^d by BA_d . Dirichlet's theorem shows that in some sense, out of all the points in \mathbb{R}^d , badly approximable points are the hardest to approximate by rationals. It is well known that BA_d is a Lebesgue nullset of full Hausdorff dimension in \mathbb{R}^d , see e.g. [25, Ch. III].

For more than a decade now, as part of the burgeoning study of Diophantine properties of fractal sets and measures [5, 9, 10, 13, 19], there has been a growing interest in computing the Hausdorff dimension of the intersection of BA_d with various fractal sets. Since BA_d has full dimension, one expects its intersection with any fractal set $J \subseteq \mathbb{R}^d$ to have the same dimension as J , and this can be proven for certain broad classes of fractal sets J , see e.g. [4, 6, 12, 20].

However, progress so far has been limited to the class of fractals defined by conformal dynamical systems, and it has been a natural challenge to understand what happens beyond this case. Non-conformal dynamical systems (where the system is expanding but may have different rates of expansion in different directions) are often much more complicated than conformal ones, which can often be thought of as essentially the same as one-dimensional systems. For instance, the Hausdorff dimension of any conformal expanding repeller can be computed via Bowen's formula (e.g. [23, Corollary 9.1.7]), but it is far more difficult to compute the Hausdorff dimension of even relatively simple non-conformal fractals, such as the limit sets/measures of affine iterated function systems (IFSs) satisfying the open set condition. To make progress one generally has to assume either some randomness in the contractions defining the IFS, as in [11, 16], or some special relations between these contractions, as in [1, 7]. An exception to this is a recent theorem of Bárány and Käenmäki [2], who showed that every self-affine measure on the plane is exact dimensional.

In this paper, we will concentrate on the latter situation, considering the class of *self-affine sponges*, and in particular analyzing the Hausdorff dimension of the intersection of a self-affine sponge with the set of badly approximable points. The class of self-affine sponges is the generalization to higher dimensions of the class of *self-affine carpets*, which consists of subsets of \mathbb{R}^2 defined according to a certain recursive construction where each rectangle in the construction is replaced by the union of several rectangles contained in that rectangle (see Definition 1.1 below). The Hausdorff dimension of certain self-affine carpets was computed independently by Bedford [3] and McMullen [22], and their results were extended by several authors [1, 7, 18, 21].

Given a self-affine sponge, we would like to know what the Hausdorff dimension of its intersection with BA_d is. There are two subtleties that make this question more difficult to answer than in the conformal case. One involves the question of what hypotheses are sufficient to deduce that a self-affine sponge intersects BA_d non-trivially. In the case of self-conformal sets, the answer has always turned out to be an irreducibility assumption: in the most general case, that the set in question is not contained in any real-analytic manifold of dimension strictly less than d (see [6]). This assumption is natural because of a well-known obstruction: any point contained in a rational affine hyperplane cannot be badly

approximable, and thus any set that intersects BA_d cannot be contained in a rational affine hyperplane. Strengthening this requirement from rational hyperplanes to all hyperplanes, and then from hyperplanes to real-analytic manifolds, is natural from a geometric point of view. However, in the case of self-affine sponges the irreducibility assumption needs to be stronger than the condition of not being contained in a manifold; in the next section we will say precisely what assumption is needed (see Definition 1.2). Our irreducibility assumption is satisfied in ‘most’ examples, and we can show that standard techniques (i.e. Schmidt’s game and hyperplane diffuseness) must fail for sponges that are not irreducible in our sense (see Proposition 3.4).

The other subtlety is that self-affine sponges may have no natural measure of full dimension, as recently discovered by two of the authors [7]. For such sponges, our techniques cannot prove that the intersection of the sponge with BA_d has full dimension in the sponge, but only that the dimension of this intersection is bounded below by the *dynamical dimension* of the sponge, i.e. the supremum of the dimensions of the invariant measures. Regarding this difficulty, we leave open the possibility that it may still be possible to prove the full Hausdorff dimension of the sponge’s intersection with BA_d using Schmidt’s game and hyperplane diffuseness, but new ideas would be needed. This problem as well as a few other open problems are listed at the end of the paper.

1. Main results

We use the same notation to describe self-affine sponges as in [7].

Definition 1.1. [7, Definitions 2.1 and 2.2] Fix $d \geq 1$, and let $D = \{1, \dots, d\}$. For each $i \in D$, let A_i be a finite index set, and let $\Phi_i = (\phi_{i,a})_{a \in A_i}$ be a finite collection of contracting similarities of $[0, 1]$, called the *base iterated function system (IFS) in coordinate i* . Let $A = \prod_{i \in D} A_i$, and for each $\mathbf{a} = (a_1, \dots, a_d) \in A$, consider the contracting affine map $\phi_{\mathbf{a}} : [0, 1]^d \rightarrow [0, 1]^d$ defined by the formula

$$\phi_{\mathbf{a}}(x_1, \dots, x_d) = (\phi_{\mathbf{a},1}(x_1), \dots, \phi_{\mathbf{a},d}(x_d)),$$

where $\phi_{\mathbf{a},i}$ is shorthand for ϕ_{i,a_i} in the formula above, as well as elsewhere. Geometrically, $\phi_{\mathbf{a}}$ can be thought of as corresponding to the rectangle to which it sends $[0, 1]^d$:

$$\phi_{\mathbf{a}}([0, 1]^d) = \prod_{i \in D} \phi_{\mathbf{a},i}([0, 1]) \subseteq [0, 1]^d.$$

Given $E \subseteq A$, we call the collection $\Phi \stackrel{\text{def}}{=} (\phi_{\mathbf{a}})_{\mathbf{a} \in E}$ a *diagonal IFS*. It is a special case of the more general notion of an *affine IFS*. The *coding map* of Φ is the map $\pi : E^{\mathbb{N}} \rightarrow [0, 1]^d$ defined by the formula

$$\pi(\omega) = \lim_{n \rightarrow \infty} \phi_{\omega \upharpoonright n}(\mathbf{0}),$$

where $\phi_{\omega \upharpoonright n} \stackrel{\text{def}}{=} \phi_{\omega_1} \circ \dots \circ \phi_{\omega_n}$. Finally, the *limit set* of Φ is the set $\Lambda_{\Phi} \stackrel{\text{def}}{=} \pi(E^{\mathbb{N}})$. We call the limit set of a diagonal IFS a *self-affine sponge*. If $d = 2$, the limit set is also called a *self-affine carpet*.

The sponge Λ_{Φ} is called *Barański* (respectively *strongly Barański*) if the base IFSs all satisfy the open set condition (respectively the strong separation condition) with respect to




FIGURE 1. Generating templates for the IFSs $\Phi^{(1)}$ and $\Phi^{(2)}$ considered in Example 1.3. The left-hand picture is irreducible but the right-hand picture is reducible. Both IFSs have distinguishable coordinates and are Barański but are not strongly Barański.

the interval $\mathbb{I} = (0, 1)$ (respectively $\mathbb{I} = [0, 1]$), i.e. if for all $i \in D$ the collection

$$(\phi_{i,a}(\mathbb{I}))_{a \in A_i}$$

is disjoint.

We now define the notion of irreducibility that we need to state our theorem.

Definition 1.2. Define a partial order \leq on D by writing $i \leq j$ if $|\phi'_{\mathbf{a},i}| \geq |\phi'_{\mathbf{a},j}|$ for all $\mathbf{a} \in E$. In other words, $i \leq j$ if all contractions of Φ contract at least as fast in coordinate j as in coordinate i . The sponge Λ_Φ is said to be *irreducible* if, for all $i \in D$, there exist $\mathbf{a}, \mathbf{b} \in E$ such that $a_i \neq b_i$ but $\{j \in D : a_j \neq b_j\} \subseteq \{j \in D : j \geq i\}$. In other words, Φ is irreducible if, for every coordinate, Φ contains two contractions that can be distinguished in that coordinate but not in any coordinate that allows for slower contraction than in the original coordinate.

The sponge Λ_Φ is said to have *distinguishable coordinates* if, for all $i, j \in D$ with $i \neq j$, there exists $\mathbf{a} \in E$ such that $|\phi'_{\mathbf{a},i}| \neq |\phi'_{\mathbf{a},j}|$. Note that in two dimensions a carpet has distinguishable coordinates if and only if its IFS does not consist entirely of similarities.

Example 1.3. Let $d = 2$, $m_1 = 3$, and $m_2 = 2$, and consider the base IFSs

$$\Phi_i = (\phi_{i,a})_{0 \leq a \leq m_i - 1}, \quad \phi_{i,a}(x) = \frac{a + x}{m_i}.$$

The product IFS consists of the affine contractions

$$\phi_{\mathbf{a}}(x_1, x_2) = \left(\frac{a_1 + x_1}{3}, \frac{a_2 + x_2}{2} \right), \quad \mathbf{a} \in A = \{0, 1, 2\} \times \{0, 1\}.$$

Let $E^{(1)} = \{(0, 0), (1, 1), (2, 0)\} \subseteq A$ and $E^{(2)} = \{(0, 0), (2, 1)\} \subseteq A$, and consider the diagonal IFSs $\Phi^{(k)} = (\phi_{\mathbf{a}})_{\mathbf{a} \in E^{(k)}} (k = 1, 2)$. (See Figure 1.) The IFS $\Phi^{(1)}$ is irreducible, since for $i = 1$ we can take $\mathbf{a} = (0, 0)$ and $\mathbf{b} = (2, 0)$ and for $i = 2$ we can take $\mathbf{a} = (0, 0)$ and $\mathbf{b} = (1, 1)$. On the other hand, the IFS $\Phi^{(2)}$ is reducible, since if $i = 1$, then there do not exist $\mathbf{a}, \mathbf{b} \in E^{(2)}$ such that $a_i \neq b_i$ and $\{j \in D : a_j \neq b_j\} \subseteq \{j \in D : j \geq i\}$. Both IFSs have distinguishable coordinates.

We note that although $\Phi^{(2)}$ is reducible, its limit set is not contained in any line or smooth curve in \mathbb{R}^2 . This contrasts with the case of limit sets of conformal IFSs, where a set is defined to be irreducible if it is not contained in any real-analytic manifold of dimension

strictly smaller than the ambient dimension. The reason that we call $\Phi^{(2)}$ reducible is that its limit set does not have any hyperplane diffuse subsets, meaning that Schmidt's game cannot be used to deduce lower bounds on the dimension of its intersection with BA_d ; see §2 and Proposition 3.4 for details.

The last notion we need to define before we can state our theorem is the notion of the dynamical dimension of a self-affine sponge.

Definition 1.4. [7, Definition 2.6] The *dynamical dimension* of a self-affine sponge Λ_Φ is the number

$$\dim_D(\Phi) \stackrel{\text{def}}{=} \sup_{\mu} \{\dim_H(\pi_*[\mu])\},$$

where the supremum is taken over all probability measures μ on $E^{\mathbb{N}}$ that are invariant under the shift map[†]. Here, $\pi_*[\mu]$ denotes the pushforward of μ under the coding map π , and the *Hausdorff dimension* of $\pi_*[\mu]$ is the infimum of the Hausdorff dimensions of sets that have full measure under $\pi_*[\mu]$.

The dynamical dimension of a self-affine sponge is always bounded above by the Hausdorff dimension of the sponge. Equality holds in the case of Sierpiński sponges (in which the coordinatewise rates of contraction are the same for all contractions in the IFS) [18] and also in the case of Barański carpets (i.e. two-dimensional Barański sponges) [1]. In general, the dynamical dimension may be strictly less than the Hausdorff dimension, see [7]; this is true even for three-dimensional Barański sponges satisfying the coordinate ordering condition (see Definition 2.8 below).

We are now ready to state our main result.

THEOREM 1.5. *Let $\Lambda_\Phi \subseteq [0, 1]^d$ be an irreducible Barański sponge with distinguishable coordinates. Then*

$$\dim_H(\Lambda_\Phi \cap \text{BA}_d) \geq \dim_D(\Lambda_\Phi).$$

In particular, if $d = 2$ or if Λ_Φ is a Sierpiński sponge, then $\Lambda_\Phi \cap \text{BA}_d$ has full Hausdorff dimension in Λ_Φ .

The idea of the proof can be described succinctly as follows. Let $\nu = \pi_*[\mu]$ be the image under the coding map of an ergodic shift-invariant probability measure μ on $E^{\mathbb{N}}$. Let N be a large number, and let $F \subseteq E^N$ be a subset consisting of ' μ -typical' words. Then the limit set of the IFS Ψ_F corresponding to F can be shown to intersect BA_d in a set of dimension close to the dimension of ν . Roughly, this is because the elements of Ψ_F are all relatively 'homogeneous' and so the lower Assouad dimension of the limit set of Ψ_F , which is a lower bound for the dimension of its intersection with BA_d (see Proposition 2.5), is close to the Hausdorff dimension of the limit set of Ψ_F , which is in turn close to the Hausdorff dimension of ν (computed using a Ledrappier–Young-type formula [7, (2.13)]). Finally, ν can be chosen so that its Hausdorff dimension is close to the dynamical dimension of Λ_Φ .

[†] By [7, Theorem 2.7 and (2.13)], the dynamical dimension is the same if we take the supremum only over ergodic measures of positive entropy. This means that our definition agrees with the usual one in the literature e.g. [8, (2.14)].

We do not need to deal with all possible ergodic shift-invariant measures; it suffices to consider the smaller class of Bernoulli measures. A *Bernoulli measure* is a measure of the form $\nu_{\mathbf{p}} = \pi_*[\mathbf{p}^{\mathbb{N}}]$, where \mathbf{p} is a probability measure on E . In [7, Theorem 2.7], it was shown that the supremum of the Hausdorff dimensions of the Bernoulli measures is equal to the dynamical dimension, so in the above proof sketch ν can be assumed to be a Bernoulli measure. The function sending a Bernoulli measure to its Hausdorff dimension is continuous [7, Theorem 2.9], so by compactness there exists a (not necessarily unique) Bernoulli measure whose Hausdorff dimension is equal to the dynamical dimension.

If $\nu_{\mathbf{p}}$ is a Bernoulli measure of maximal dimension, then to show that the conclusion of Theorem 1.5 holds it suffices to check that the above proof sketch can be made rigorous for the measure $\nu = \nu_{\mathbf{p}}$. It turns out that the conditions under which this is possible are more general than the hypotheses of Theorem 1.5. To make this statement precise, we introduce a ‘local’ analogue of the partial order \preceq considered in Definition 1.2.

Definition 1.6. Let \mathbf{p} be a probability measure on E , and for each $i \in D$ we define the *Lyapunov exponent of \mathbf{p} in coordinate i* to be the number

$$\chi_i(\mathbf{p}) = - \int \log|\phi'_{\mathbf{a},i}| d\mathbf{p}(\mathbf{a}).$$

Define a partial order $\preceq_{\mathbf{p}}$ on D by writing $i \preceq_{\mathbf{p}} j$ if $\chi_i(\mathbf{p}) \leq \chi_j(\mathbf{p})$. The measure \mathbf{p} is said to be *irreducible* (or equivalently, the sponge Λ_{Φ} is said to be *irreducible with respect to \mathbf{p}*) if for all $i \in D$, there exist $\mathbf{a}, \mathbf{b} \in E$ such that $a_i \neq b_i$ but $\{j \in D : a_j \neq b_j\} \subseteq \{j \in D : j \succeq_{\mathbf{p}} i\}$. Note that $\preceq_{\mathbf{p}}$ is a finer partial order than \preceq , so if a sponge is irreducible in the sense of Definition 1.2 then it is irreducible with respect to every probability measure on E .

The measure \mathbf{p} is said to have *distinct Lyapunov exponents* if the numbers $\chi_i(\mathbf{p})$ ($i \in D$) are all distinct.

We also introduce a ‘local’ analogue of the Barański condition.

Definition 1.7. (See [7, Definition 3.1]) Let Λ_{Φ} be a self-affine sponge, and let $I \subseteq D$ be a coordinate set. Let

$$\Phi_I = (\phi_{I,\mathbf{a}})_{\mathbf{a} \in \pi_I(E)},$$

where $\phi_{I,\mathbf{a}} : [0, 1]^I \rightarrow [0, 1]^I$ is defined by the formula

$$\phi_{I,\mathbf{a}}(\mathbf{x}) = (\phi_{\mathbf{a},i}(x_i))_{i \in I}$$

and $\pi_I : A \rightarrow A_I \stackrel{\text{def}}{=} \prod_{i \in I} A_i$ is the projection map. We call I *good* (respectively *strongly good*) if the collection

$$(\phi_{I,\mathbf{a}}(\mathbb{I}^I))_{\mathbf{a} \in \pi_I(E)}$$

is disjoint, where $\mathbb{I} = (0, 1)$ (respectively $\mathbb{I} = [0, 1]$). Also, a measure \mathbf{p} on E is called *good* (respectively *strongly good*) if for every $x > 0$, the set

$$I(\mathbf{p}, x) = \{i \in D : \chi_i(\mathbf{p}) \leq x\}$$

is good (respectively strongly good).

We can now state a ‘local’ version of Theorem 1.5.

THEOREM 1.8. *Let $\Lambda_\Phi \subseteq [0, 1]^d$ be a self-affine sponge, and let \mathbf{p} be an irreducible good probability measure on E with distinct Lyapunov exponents. Then*

$$\dim_H(\Lambda_\Phi \cap \text{BA}_d) \geq \dim_H(\nu_{\mathbf{p}}).$$

In particular, if $\dim_H(\nu_{\mathbf{p}}) = \dim_H(\Lambda_\Phi)$, then $\Lambda_\Phi \cap \text{BA}_d$ has full Hausdorff dimension in Λ_Φ .

If Λ_Φ is an irreducible Barański sponge with distinguishable coordinates, then every probability measure on E is both irreducible and good, and the set of measures with distinct Lyapunov exponents forms an open dense set. Thus, Theorem 1.8 implies Theorem 1.5.

Outline of the paper. In the next section, we recall some known results about the dimension of intersection of BA_d with fractals and its relation to the lower Assouad dimension, and state a strengthening of Theorem 1.8, namely Theorem 2.7. In §3, we prove some results which suffice to give a useful estimate of the Hausdorff dimension of $\Lambda_{\Psi_F} \cap \text{BA}_d$, where Ψ_F is a ‘homogeneous’ IFS as described in our proof sketch above. In §4 we use this estimate to prove Theorem 2.7.

Convention 1. The symbols \lesssim , \gtrsim , and \asymp will denote coarse multiplicative asymptotics. For example, $A \lesssim B$ means that there exists a constant $C > 0$ (the *implied constant*) such that $A \leq CB$.

2. Schmidt’s game, hyperplane diffuse sets, and the lower Assouad dimension

For $d \geq 2$, the full dimension of BA_d in \mathbb{R}^d was proven by Schmidt [24] using a technique now known as Schmidt’s game. Since then, Schmidt’s game and its variants have been used to prove the full dimension intersection of BA_d with various fractals, as well as various stability properties such as C^1 incompressibility [4]. The modern approach [4] is to first show that BA_d is winning for a variant of Schmidt’s game known as the hyperplane absolute game, and then to show that any set winning for the hyperplane absolute game is also winning for Schmidt’s game played on any fractal satisfying a certain geometric condition called hyperplane diffuseness. The Hausdorff dimension of a set winning for Schmidt’s game played on a fractal can be bounded from below based on the geometry of that fractal; there are currently two known ways of doing this, one based on the dimensions of fully supported doubling Frostman measures on the fractal [20, Theorem 1.1], and the other based on the lower Assouad dimension of the fractal [12, Theorem 3.1]. The two methods give the same bound whenever the fractal in question is Ahlfors regular[†], which is true in most applications that have been considered so far. However, in our case the fractals are not Ahlfors regular, and there is a difference between the two methods. We work with the second method, based on the lower Assouad dimension, as it seems to be more suited to the situation we consider here.

[†] We recall that a measure μ on \mathbb{R}^d is said to be *Ahlfors s -regular* if, for all $\mathbf{x} \in \text{Supp}(\mu)$ and $0 < \rho \leq 1$, we have

$$\mu(B(\mathbf{x}, \rho)) \asymp \rho^\delta.$$

A closed set $K \subseteq \mathbb{R}^d$ is said to be *Ahlfors s -regular* if there exists an Ahlfors s -regular measure μ such that $K = \text{Supp}(\mu)$, and *Ahlfors regular* if it is Ahlfors s -regular for some $s > 0$.

Definition 2.1. Let K be a closed subset of \mathbb{R}^d . For any $0 < \alpha, \beta < 1$, *Schmidt's (α, β) -game* is an infinite game played by two players, Alice and Bob, who take turns choosing balls in \mathbb{R}^d whose centers lie in K , with Bob moving first. The players must choose their moves so as to satisfy the relations

$$B_1 \supseteq A_1 \supseteq B_2 \supseteq \dots$$

and

$$\rho(A_k) = \alpha\rho(B_k) \quad \text{and} \quad \rho(B_{k+1}) = \beta\rho(A_k) \quad \text{for } k \in \mathbb{N},$$

where B_k and A_k denote Bob's and Alice's k th moves, respectively, and where $\rho(B)$ denotes the radius of a ball B . Since the sets B_1, B_2, \dots form a nested sequence of non-empty closed sets whose diameters tend to zero, it follows that the intersection $\bigcap_k B_k$ is a singleton, say $\bigcap_k B_k = \{\mathbf{x}\}$, whose unique member \mathbf{x} lies in K . The point \mathbf{x} is called the *outcome* of the game. A set $S \subseteq K$ is said to be (α, β) -*winning on K* if Alice has a strategy guaranteeing that the outcome lies in S , regardless of the way Bob chooses to play. It is said to be α -*winning on K* if it is (α, β) -winning on K for every $0 < \beta < 1$, and *winning on K* if it is α -winning on K for some $0 < \alpha < 1$.

Definition 2.2. A set $K \subseteq \mathbb{R}^d$ is said to be *hyperplane diffuse* if there exists $\beta > 0$ such that, for all $0 < \rho \leq 1$ and $\mathbf{x} \in K$ and for any (affine) hyperplane $\mathcal{L} \subseteq \mathbb{R}^d$, we have

$$B(\mathbf{x}, \rho) \cap K \setminus \mathcal{N}(\mathcal{L}, \beta\rho) \neq \emptyset,$$

where $\mathcal{N}(\mathcal{L}, \varepsilon)$ denotes the closed ε -thickening of \mathcal{L} , i.e. $\mathcal{N}(\mathcal{L}, \varepsilon) = \{\mathbf{y} \in \mathbb{R}^d : d(\mathbf{y}, \mathcal{L}) \leq \varepsilon\}$.

PROPOSITION 2.3. [4, Theorem 2.5 + Proposition 4.7] *Let $K \subseteq \mathbb{R}^d$ be a closed and hyperplane diffuse set. Then $BA_d \cap K$ is winning on K .*

This result remains true if BA_d is replaced by any *hyperplane absolute winning set*, see [4, p. 4] for the definition. The same applies to all the results of this paper.

To state in a clearer way the lower bound for the Hausdorff dimension of a winning set discovered by the second-named author [12, Theorem 3.1], we recall the definition of the lower Assouad dimension of a set. (The lower Assouad dimension has been given several names in the literature, see [14, p. 6688], but the name 'lower Assouad dimension' seems by far the most natural to us.)

Definition 2.4. Given $\rho > 0$ and $S \subseteq \mathbb{R}^d$, we let $N_\rho(S)$ denote the cardinality of any maximal ρ -separated subset of S . (Choosing a different maximal ρ -separated set will not change $N_\rho(S)$ by more than a constant factor.) The *lower Assouad dimension* of a non-empty closed set $K \subseteq \mathbb{R}^d$, denoted $\underline{\dim}_A(K)$, is the supremum of $s \geq 0$ such that there exists a constant $c > 0$ such that, for all $\mathbf{x} \in K$ and $0 < \beta, \rho \leq 1$, we have

$$N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K) \geq c\beta^{-s}.$$

Equivalently,

$$\begin{aligned} \underline{\dim}_A(K) &= \liminf_{\beta \rightarrow 0} \inf_{0 < \rho \leq 1} \inf_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \\ &= \liminf_{\beta \rightarrow 0} \liminf_{\rho \rightarrow 0} \inf_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \end{aligned}$$

(see Appendix A).

The lower Assouad dimension is the smallest of the standard fractal dimensions. In particular, if K is closed then $\underline{\dim}_A(K) \leq \dim_H(K)$, see e.g. [17, Lemma 2.2]. Note that unlike most notions of dimension, the lower Assouad dimension is not monotone: a subset may have larger lower Assouad dimension than the set it is contained in.

The essential idea of the following result is found in [12, Theorem 3.1]. We include the proof for completeness.

PROPOSITION 2.5. *Let $K \subseteq \mathbb{R}^d$ be closed and let $S \subseteq K$ be winning on K . Then*

$$\dim_H(S) \geq \underline{\dim}_A(K),$$

where $\underline{\dim}_A$ denotes the lower Assouad dimension.

Proof. Let $\delta = \underline{\dim}_A(K)$, and fix $\varepsilon > 0$. Then by definition, there exists a constant $c = c_\varepsilon > 0$ such that, for all $\mathbf{x} \in K$, $0 < \beta \leq 1/4$, and $0 < \rho \leq 1$, we have

$$N_{3\beta\rho}(B(\mathbf{x}, (1 - \beta)\rho) \cap K) \geq N = N(\beta) \stackrel{\text{def}}{=} \lfloor c_\varepsilon \beta^{-(\delta - \varepsilon)} \rfloor.$$

Now let $\alpha > 0$ be chosen so that S is α -winning, and fix $0 < \beta \leq 1/2$. For each ball $A = B(\mathbf{x}, \rho)$, we choose a $3\beta\rho$ -separated sequence $\mathbf{y}^{(1)}(A), \dots, \mathbf{y}^{(N)}(A) \in B(\mathbf{x}, (1 - \beta)\rho) \cap K$, and we let $f_i(A)$ denote the ball centered at $\mathbf{y}^{(i)}(A)$ of radius $\beta\rho$. Then the balls $f_1(A), \dots, f_N(A)$ are contained in A and are separated by distances of at least $\beta\rho$. Moreover, each ball $f_i(A)$ is a legal move for Bob to make in response to Alice playing A as her move.

Now fix a winning strategy for Alice to win the (α, β) -game, and we will consider the family of counter-strategies for Bob such that whenever Alice plays a ball A_k , Bob responds by playing one of the balls $f_1(A_k), \dots, f_N(A_k)$. We fix Bob's initial ball (chosen to have radius less than 1), and for each function $\omega : \mathbb{N} \rightarrow E \stackrel{\text{def}}{=} \{1, \dots, N(\beta)\}$, we consider the counter-strategy in which Bob responds to Alice's k th move A_k by choosing the ball $B_{k+1} = f_{\omega(k)}(A_k)$. We denote the outcome of this counter-strategy by $\pi(\omega)$, so that $\pi : E^\mathbb{N} \rightarrow K$. The separation conditions on the balls $f_1(A), \dots, f_N(A)$ guarantee that

$$d(\pi(\omega), \pi(\tau)) \asymp (\alpha\beta)^{|\omega \wedge \tau|} \quad \text{for all } \omega, \tau \in E^\mathbb{N},$$

where $|\omega \wedge \tau|$ denotes the length of the longest common initial segment of ω and τ . Thus, the uniform Bernoulli measure on $\pi(E^\mathbb{N})$ is Ahlfors $s(\beta)$ -regular, where

$$s(\beta) \stackrel{\text{def}}{=} \frac{\log N(\beta)}{-\log(\alpha\beta)} = \frac{-(\delta - \varepsilon) \log(\beta) + O(1)}{-\log(\alpha\beta)} \xrightarrow{\beta \rightarrow 0} \delta - \varepsilon.$$

It follows that $\dim_H(\pi(E^\mathbb{N})) \geq s(\beta)$. Now, since each element of $\pi(E^\mathbb{N})$ is the outcome of a game where Alice played her winning strategy, we have $\pi(E^\mathbb{N}) \subseteq S$ and thus $\dim_H(S) \geq s(\beta) \rightarrow \delta - \varepsilon$. Since ε was arbitrary, we have $\dim_H(S) \geq \delta$. □

Combining Propositions 2.3 and 2.5 gives the following corollary.

COROLLARY 2.6. *Let $K \subseteq \mathbb{R}^d$ be a closed and hyperplane diffuse set. Then*

$$\dim_H(\text{BA}_d \cap K) \geq \underline{\dim}_A(K).$$

With this result in mind, we can see how the following theorem is a strengthening of Theorem 1.8.




FIGURE 2. Generating templates for two carpets satisfying the coordinate ordering condition. The picture on the right also satisfies the disjointness condition, making it a Lalley–Gatzouras carpet.

THEOREM 2.7. *Let $\Lambda_\Phi \subseteq [0, 1]^d$ be a self-affine sponge, and let \mathbf{p} be an irreducible good probability measure on E with distinct Lyapunov exponents. Then there exists a sequence of strongly Lalley–Gatzouras sponges $\Lambda_{\Psi_N} \subseteq \Lambda_\Phi$ that are hyperplane diffuse and satisfy*

$$\underline{\dim}_A(\Lambda_{\Psi_N}) \rightarrow \dim_H(\nu_{\mathbf{p}}).$$

Here, following [7], we use the term ‘Lalley–Gatzouras’ to refer to a certain class of sponges that includes the carpets considered by Lalley and Gatzouras.

Definition 2.8. (See [7, Definition 3.6]) A self-affine sponge Λ_Φ is *Lalley–Gatzouras* (respectively *strongly Lalley–Gatzouras*) if there exists a permutation σ of D such that both of the following hold (see Figure 2).

- (Coordinate ordering condition) For all $\mathbf{a} \in E$, we have

$$|\phi'_{\mathbf{a},\sigma(1)}| > |\phi'_{\mathbf{a},\sigma(2)}| > \dots > |\phi'_{\mathbf{a},\sigma(d)}|.$$

- (Disjointness condition) The coordinate sets $\sigma(I_{\leq i})$ ($i = 1, \dots, d$) are all good (respectively strongly good), where $I_{\leq i} \stackrel{\text{def}}{=} \{1, \dots, i\}$.

3. Some results on Lalley–Gatzouras sponges

In this section, we give a necessary and sufficient condition for the hyperplane diffuseness of a strongly Lalley–Gatzouras sponge Λ_Φ (which will be fixed throughout the section), as well as a formula for the lower Assouad dimension of Λ_Φ . For conceptual completeness we also state the formula for the upper Assouad dimension of Λ_Φ , which is defined in the following.

Definition 3.1. The *upper Assouad dimension* of a non-empty closed set $K \subseteq \mathbb{R}^d$, denoted $\overline{\dim}_A(K)$, is the infimum of $s \geq 0$ such that there exists $C > 0$ such that, for all $\mathbf{x} \in K$ and $0 < \beta, \rho \leq 1$, we have

$$N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K) \leq C\beta^{-s},$$

where $N_{\beta\rho}$ is as in Definition 2.4. Equivalently,

$$\begin{aligned} \overline{\dim}_A(K) &= \limsup_{\beta \rightarrow 0} \sup_{0 < \rho \leq 1} \sup_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \\ &= \limsup_{\beta \rightarrow 0} \limsup_{\rho \rightarrow 0} \sup_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \end{aligned}$$

(see Appendix A).

Definition 3.2. The sponge Λ_Φ is called *uniformly irreducible* if, for all $i \in D$ and $\mathbf{a} \in E$, there exists $\mathbf{b} \in E$ such that $b_i \neq a_i$ but $\{j \in D : a_j \neq b_j\} \subseteq \{j \in D : j \geq i\}$, where the partial order \leq is the same as in Definition 1.2.

For example, the irreducible Barański sponge $\Lambda_{\Phi^{(1)}}$ appearing in Example 1.3 is not uniformly irreducible, since if $i = 1$ and $\mathbf{a} = (1, 1)$ then there is no $\mathbf{b} \in E^{(1)}$ such that $b_i \neq a_i$ but $\{j \in D : a_j \neq b_j\} \subseteq \{j \in D : j \geq i\}$. On the other hand, if $E^{(3)} = \{(0, 0), (1, 1), (2, 0), (2, 1)\}$, then the corresponding Barański sponge $\Lambda_{\Phi^{(3)}}$ is uniformly irreducible.

In the remainder of this section, we assume without loss of generality that the permutation σ appearing in Definition 2.8 is trivial, i.e. that the orders \leq and \leq on D are equivalent. (Note that this is not true for the sponges of Example 1.3.)

PROPOSITION 3.3. *The sponge Λ_Φ is hyperplane diffuse if and only if it is uniformly irreducible.*

We will use the backwards direction of this proposition in the proof of Theorem 2.7; we include the proof of the forwards direction for completeness.

Proof of backwards direction. Fix $\omega \in E^{\mathbb{N}}$ and $0 < \rho \leq 1$, and let $\mathbf{x} = \pi(\omega)$; we will prove that

$$\Lambda_\Phi \cap B(\mathbf{x}, \rho) \setminus \mathcal{N}(\mathcal{L}, \beta\rho) \neq \emptyset \tag{3.1}$$

for any hyperplane \mathcal{L} , where $\beta > 0$ is an appropriate constant. For each $i \in D$, let $N_i \in \mathbb{N}$ be the smallest number such that

$$\prod_{n=1}^{N_i} |\phi'_{\omega_n, i}| \leq \rho, \tag{3.2}$$

and note that by the coordinate ordering condition, we have $N_1 \geq N_2 \geq \dots \geq N_d$. Let

$$\begin{aligned} [\mathbf{a}]_i &= \{\mathbf{b} \in E : a_i = b_i\}, \\ [\omega \upharpoonright N]_i &\stackrel{\text{def}}{=} \{\tau \in E^{\mathbb{N}} : \tau_n \in [\omega_n]_i \ \forall n \leq N\}, \\ B_\omega(N_1, \dots, N_d) &\stackrel{\text{def}}{=} \bigcap_{i \in D} [\omega \upharpoonright N_i]_i. \end{aligned} \tag{3.3}$$

Then, by (3.2),

$$B_\omega(N_1, \dots, N_d) \subseteq \pi^{-1}(B(\pi(\omega), \rho)).$$

(Here for convenience we work with the max norm on \mathbb{R}^d .) Now fix $i \in D$ and let $\mathbf{a} = \omega_{N_i+1}$. Let $\mathbf{b} \in E$ be as in Definition 3.2. Define the point $\tau \in E^{\mathbb{N}}$ as follows: let $\tau_n = \omega_n$ for all $n \neq N_i + 1$, and let $\tau_{N_i+1} = \mathbf{b}$. Finally, let $\mathbf{y}^{(i)} = \pi(\tau)$.

Fix $j \in D$. If $a_j = b_j$, then clearly $\tau \in [\omega \upharpoonright N_j]_j$. On the other hand, if $a_j \neq b_j$, then $j \geq i$ and thus $N_j \leq N_i$, which implies $\tau \in [\omega \upharpoonright N_j]_j$. So either way we have $\tau \in [\omega \upharpoonright N_j]_j$, and thus $\tau \in B_\omega(N_1, \dots, N_d)$. Consequently, $\mathbf{y}^{(i)} \in B(\mathbf{x}, \rho)$.

If $j < i$, then $a_j = b_j$ and thus $x_j = y_j^{(i)}$. On the other hand, since $a_i \neq b_i$ and since $I_{\leq i}$ is strongly good, we have $\max_{j \leq i} |y_j^{(i)} - x_j| \asymp \rho$. Combining these facts gives $|y_i^{(i)} - x_i| \asymp \rho$. Since i, j were arbitrary, this means that the matrix $M = (y_j^{(i)} - x_j)_{i,j} = \sum_{i \in D} (\mathbf{e}^{(i)} \cdot (\mathbf{y}^{(i)} - \mathbf{x})^T$ is upper triangular and its diagonal entries are asymptotic to ρ ,

while all of its entries are bounded in magnitude by ρ . Here $\mathbf{e}^{(i)}$ denotes the (column) vector whose i th entry is 1 (and whose other entries are 0). This implies that

$$\|M^{-1}\| \lesssim \rho^{-1}.$$

So if $\mathbf{v} \in \mathbb{R}^d$ is any unit vector, then $\|M\mathbf{v}\| \gtrsim \rho$ and thus there exists $i \in D$ such that $|(\mathbf{y}^{(i)} - \mathbf{x}) \cdot \mathbf{v}| \gtrsim \rho$. It follows that there exists a constant $\beta > 0$ (independent of $\mathbf{x}, \rho, \mathbf{v}$) such that

$$d(\mathbf{y}^{(i)} - \mathbf{x}, \mathbf{v}^\perp) > 2\beta\rho. \tag{3.4}$$

Now if $\mathcal{L} \subseteq \mathbb{R}^d$ is a hyperplane, then we can write $\mathcal{L} = \mathbf{p} + \mathbf{v}^\perp$ for some $\mathbf{p} \in \mathbb{R}^d$ and some unit vector \mathbf{v} , and then (3.4) implies that

$$\max(d(\mathbf{x}, \mathcal{L}), d(\mathbf{y}^{(i)}, \mathcal{L})) > \beta\rho.$$

Since $\mathbf{x}, \mathbf{y}^{(i)} \in \Lambda_\Phi \cap B(\mathbf{x}, \rho)$, this demonstrates (3.1), completing the proof. □

Proof of forwards direction. By contradiction suppose that Λ_Φ is not uniformly irreducible. Then there exist $i \in D$ and $\mathbf{a} \in E$ such that, for all $\mathbf{b} \in E$ satisfying $b_i \neq a_i$, we have $\{j \in D : a_j \neq b_j\} \not\subseteq \{j \in D : j \geq i\}$. Let $\omega = \mathbf{a}^\infty$, $\mathbf{x} = \pi(\omega)$, and $\mathcal{L} = \mathbf{x} + \sum_{j \neq i} \mathbb{R}\mathbf{e}^{(j)}$. We claim that, for all $0 < \rho \leq 1$,

$$\Lambda_\Phi \cap B(\mathbf{x}, \rho) \subseteq \mathcal{N}(\mathcal{L}, C\rho^\alpha), \tag{3.5}$$

where $C > 0$ and $\alpha > 1$ are constants. This implies that Λ_Φ is not hyperplane diffuse.

Indeed, fix $0 < \rho \leq 1$, and for each $j \in D$, let $N_j \in \mathbb{N}$ be the largest number such that

$$\prod_{n=1}^{N_j} |\phi'_{\omega_n, j}| = |\phi'_{\mathbf{a}, j}|^{N_j} \geq \varepsilon^{-1}\rho,$$

where

$$\varepsilon = \min_{\substack{I=I \leq j \\ j \in D}} \min_{\substack{\mathbf{a}, \mathbf{b} \in \pi_I(E) \\ \text{distinct}}} d(\phi_{I, \mathbf{a}}([0, 1]^I), \phi_{I, \mathbf{b}}([0, 1]^I)). \tag{3.6}$$

Since Λ_Φ satisfies the disjointness condition, we have $\varepsilon > 0$. As before we have $N_1 \geq N_2 \geq \dots \geq N_d$. We let the notation $[\mathbf{a}]_i$, $[\omega \upharpoonright N]_i$, and $B_\omega(N_1, \dots, N_d)$ be as in the previous proof, but this time our definition of N_j implies that

$$\pi^{-1}(B(\pi(\omega), \rho)) \subseteq B_\omega(N_1, \dots, N_d).$$

Fix $\tau \in \pi^{-1}(B(\pi(\omega), \rho))$, and we will estimate $d(\mathbf{y}, \mathcal{L})$, where $\mathbf{y} = \pi(\tau)$. Fix $n \leq N_{i-1}$ (with the convention that $N_0 = \infty$), and let $\mathbf{b} = \tau_n$. Since $\tau \in B_\omega(N_1, \dots, N_d)$, we have $b_j = \tau_{n, j} = \omega_{n, j} = a_j$ for all $j < i$. By the definition of i , this implies that $b_i = a_i$, and thus $\tau_{n, i} = \omega_{n, i}$ for all $n \leq N_{i-1}$. It follows that

$$d(\mathbf{y}, \mathcal{L}) = |y_i - x_i| \leq \prod_{n=1}^{N_{i-1}} |\phi'_{\omega_n, i}| = |\phi'_{\mathbf{a}, i}|^{N_{i-1}}.$$

If $i = 1$, then we have shown that $d(\mathbf{y}, \mathcal{L}) = 0$. Suppose $i > 1$. Since Λ_Φ satisfies the coordinate ordering condition, we have

$$\alpha \stackrel{\text{def}}{=} \frac{\log|\phi'_{\mathbf{a}, i}|}{\log|\phi'_{\mathbf{a}, i-1}|} > 1.$$

On the other hand, the definition of N_{i-1} implies that $|\phi'_{\mathbf{a},i-1}|^{N_{i-1}+1} < \varepsilon^{-1}\rho$, so

$$d(\mathbf{y}, \mathcal{L}) \leq |\phi'_{\mathbf{a},i-1}|^{N_{i-1}\alpha} \lesssim \rho^\alpha,$$

demonstrating (3.5). □

If Λ_Φ is irreducible but not uniformly irreducible, then in the next section we will show that Λ_Φ contains uniformly irreducible sponges. So, in this case, even though Λ_Φ is not hyperplane diffuse it contains hyperplane diffuse subsets. On the other hand, we have the following.

PROPOSITION 3.4. *If Λ_Φ is reducible, then it contains no hyperplane diffuse subsets.*

In this case, the techniques of §2 cannot possibly be used to prove that $\text{BA}_d \cap \Lambda_\Phi$ is large, because these techniques rely on finding hyperplane diffuse subsets of Λ_Φ (with sufficiently large lower Assouad dimension).

Proof. Since Λ_Φ is reducible, there exists $i \in D$ such that, for all $\mathbf{a}, \mathbf{b} \in E$ such that $a_i \neq b_i$, there exists $j < i$ such that $a_j \neq b_j$. Fix $\mathbf{x} = \pi(\omega) \in \Lambda_\Phi$, and let $\mathcal{L} = \mathbf{x} + \sum_{j \neq i} \mathbb{R}\mathbf{e}^{(j)}$. Repeating the second paragraph of the proof of the forwards direction of Proposition 3.3 shows that (3.5) holds for all $0 < \rho \leq 1$. (One minor change is needed: after fixing $n \leq N_{i-1}$, we let $\mathbf{a} = \omega_n$. Since the condition on i now holds for all $\mathbf{a}, \mathbf{b} \in E$, the subsequent argument is still valid.) Since $\mathbf{x} \in \Lambda_\Phi$ was arbitrary, (3.5) implies that no subset of Λ_Φ is hyperplane diffuse. □

Next, we compute the upper and lower Assouad dimensions of Λ_Φ . For each $i \in D$, let $\pi_i = \pi_{I_{<i}} = \pi_{I_{\leq i-1}}$, and for each $\mathbf{a} \in \pi_i(E)$, consider the ‘fiber IFS’

$$\Phi_{i,\mathbf{a}} = (\phi_{i,b})_{b \in E_{i,\mathbf{a}}} \quad \text{where } E_{i,\mathbf{a}} = \{b \in A_i : (\mathbf{a}, b) \in \pi_{i+1}(E)\}. \tag{3.7}$$

Note that $\pi_1(E) = \{\emptyset\}$ and $\pi_{d+1}(E) = E$. We let $\dim(\Phi_{i,\mathbf{a}})$ denote the dimension of the limit set of the IFS $\Phi_{i,\mathbf{a}}$. (Since the limit set is Ahlfors regular, it does not matter what notion of fractal dimension we use.)

THEOREM 3.5. *Recall that Λ_Φ denotes a strongly Lalley–Gatzouras sponge such that the orders \preceq and \leq on D are equivalent. For each $i \in D$ let*

$$\underline{\delta}_i = \min_{\mathbf{a} \in \pi_i(E)} \dim(\Phi_{i,\mathbf{a}}),$$

$$\overline{\delta}_i = \max_{\mathbf{a} \in \pi_i(E)} \dim(\Phi_{i,\mathbf{a}}).$$

Then

$$\underline{\dim}_A(\Lambda_\Phi) = \sum_{i \in D} \underline{\delta}_i, \tag{3.8}$$

$$\overline{\dim}_A(\Lambda_\Phi) = \sum_{i \in D} \overline{\delta}_i. \tag{3.9}$$

The case $d = 2$ of Theorem 3.5 (i.e. carpets) was proven recently by Fraser [14, Theorems 2.12 and 2.13], and the case of Sierpiński sponges by Fraser and Howroyd [15]. The case of general Barański sponges appears to be subtler.

We will prove only (3.8), since that is the equation that we will need in the proof of Theorem 2.7. (To be precise, we only need the \geq direction of (3.8).) The proof of (3.9) is similar.

Proof of \geq direction. Fix $\omega \in E^{\mathbb{N}}$ and $0 < \beta, \rho \leq 1$, and let $\mathbf{x} = \pi(\omega)$. For each $i \in D$, let $N_i \in \mathbb{N}$ be the smallest number such that (3.2) holds. Fix $i \in D$, and consider the space

$$X_i \stackrel{\text{def}}{=} \prod_{n=1}^{N_i} \{\omega_{n,i}\} \times \prod_{n=N_i+1}^{N_{i-1}} E_{i,\pi_i(\omega_n)}.$$

Here we use the convention that $N_0 = \infty$. Let r_i be the map

$$r_i([\tau \upharpoonright N]) = \prod_{n=1}^N |\phi'_{\tau_n,i}|,$$

i.e. up to a constant, r_i sends a cylinder in X_i to the diameter of the i th coordinate of its image under the coding map. Here τ denotes any element of $E^{\mathbb{N}}$.

CLAIM 3.6. *If $[\tau \upharpoonright N_{i-1}]$ is a maximal-length cylinder of X_i , then*

$$r_i([\tau \upharpoonright N_{i-1}]) \lesssim \rho^{1+\delta} \tag{3.10}$$

where $\delta > 0$ is a constant.

Proof. We have

$$r_i([\tau \upharpoonright N_{i-1}]) = \left(\prod_{n=1}^{N_i} |\phi'_{\omega_n,i}| \right) \left(\prod_{n=N_i+1}^{N_{i-1}} |\phi'_{\tau_n,i}| \right) \leq \lambda_+^{N_{i-1}-N_i} \rho,$$

where $\lambda_+ = \max_{i \in D} \max_{b \in A_i} |\phi'_{i,b}| < 1$. Also,

$$r_i([\omega \upharpoonright N_{i-1}]) = \left(\prod_{n=1}^{N_i} |\phi'_{\omega_n,i}| \right) \left(\prod_{n=N_i+1}^{N_{i-1}} |\phi'_{\omega_n,i}| \right) \geq \lambda_-^{N_{i-1}-N_i} \rho,$$

where $\lambda_- = \min_{i \in D} \min_{b \in A_i} |\phi'_{i,b}| > 0$. On the other hand, if

$$\alpha \stackrel{\text{def}}{=} \max_{i \in D} \max_{\mathbf{a} \in E} \frac{\log |\phi'_{\mathbf{a},i}|}{\log |\phi'_{\mathbf{a},i-1}|} > 1,$$

then

$$r_i([\omega \upharpoonright N_{i-1}]) = \prod_{n=1}^{N_{i-1}} |\phi'_{\omega_n,i}| \leq \prod_{n=1}^{N_{i-1}} |\phi'_{\omega_n,i-1}|^\alpha \asymp \rho^\alpha.$$

Combining these inequalities gives

$$\lambda_-^{N_{i-1}-N_i} \lesssim \rho^{\alpha-1}$$

and thus

$$r_i([\tau \upharpoonright N_{i-1}]) \lesssim \rho^{1+(\alpha-1) \log(\lambda_+)/\log(\lambda_-)}. \tag{3.11} \quad \triangleleft$$

In the remainder of the proof we assume that $\rho \leq (C^{-1}\beta)^{1/\delta}$, where C is the implied constant of (3.10). Then

$$r_i([\tau \upharpoonright N_{i-1}]) \leq \beta\rho$$

for all maximal-length cylinders $[\tau \upharpoonright N_{i-1}]$ in X_i . It follows that the collection

$$\mathcal{P}_i = \{[\tau \upharpoonright N] : r_i([\tau \upharpoonright N]) \leq \beta\rho < r_i([\tau \upharpoonright N - 1])\}$$

is a partition of X_i .

Now let $s_i = \delta_i$. By the definition of δ_i , we have

$$\sum_{b \in E_{i,a}} |\phi'_{i,b}|^{s_i} \geq 1 \quad \text{for all } \mathbf{a} \in \pi_i(E),$$

and thus the map $r_i^{s_i}$ is subadditive on cylinders of length at least N_i . So

$$\rho^{s_i} \asymp r_i^{s_i}([\omega \upharpoonright N_i]) \leq \sum_{P \in \mathcal{P}_i} r_i^{s_i}(P) \asymp (\beta\rho)^{s_i} \#(\mathcal{P}_i),$$

i.e. $\#(\mathcal{P}_i) \gtrsim \beta^{-s_i}$. Now let $\mathcal{P} = \prod_{i=1}^d \mathcal{P}_i$, and let $\mathbb{P}(E^{\mathbb{N}})$ denote the power set of $E^{\mathbb{N}}$. Define the map $\iota : \mathcal{P} \rightarrow \mathbb{P}(E^{\mathbb{N}})$ as follows:

$$\iota([\tau_1 \upharpoonright M_1], \dots, [\tau_d \upharpoonright M_d]) = \bigcap_{i \in D} [\tau_i \upharpoonright M_i]_i \subseteq B_\omega(N_1, \dots, N_d). \tag{3.11}$$

Then the sets $\pi(\iota(\mathbf{P}))$ ($\mathbf{P} \in \mathcal{P}$) are contained in $B(\mathbf{x}, \rho)$ and separated by distances $\gtrsim \beta\rho$. Thus

$$N_{\beta\rho}(B(\mathbf{x}, \rho) \cap \Lambda_\Phi) \gtrsim \#(\mathcal{P}) = \prod_{i=1}^d \#(\mathcal{P}_i) \gtrsim \prod_{i=1}^d \beta^{-s_i} = \beta \wedge \left(- \sum_{i=1}^d \delta_i\right),$$

assuming that $\rho \leq (C^{-1}\beta)^{1/\delta}$. Here $\beta \wedge s$ denotes β raised to the power of s . Taking the infimum over $\mathbf{x} \in K$, the liminf as $\rho \rightarrow 0$, and then the liminf as $\beta \rightarrow 0$ completes the proof. □

Proof of \leq direction. For each $i \in D$, let $\mathbf{a}^{(i)} \in E$ be chosen so that

$$\dim(\Phi_{i,\pi_i(\mathbf{a}^{(i)})}) = \delta_i. \tag{3.12}$$

Fix $0 < \rho \leq 1$, and define the sequence N_1, \dots, N_d by backwards recursion: if N_{i+1}, \dots, N_d are defined, then let N_i be the smallest integer such that

$$\prod_{j=i}^d |\phi'_{\mathbf{a}^{(j)},i}|^{N_j - N_{j+1}} \leq \rho.$$

Then let $\omega \in E^{\mathbb{N}}$ be the infinite word defined by the formula

$$\omega_n = \mathbf{a}^{(i)} \text{ for all } i \in D \text{ for all } n = N_i + 1, \dots, N_{i-1}. \tag{3.13}$$

Note that, for each $i \in D$, N_i is the smallest integer that satisfies (3.2), i.e. N_i has the same value in this proof as it did in the proof of the \geq direction. Fix $i \in D$, and let X_i and r_i be as in the proof of the \geq direction. By (3.13) and (3.12), r_i is additive on cylinders rather

than merely being subadditive. Moreover, since the sets $\pi(\iota(\mathbf{P}))$ ($\mathbf{P} \in \mathcal{P}$) defined by (3.11) have diameter $\lesssim \beta\rho$ and form a cover of $B(\mathbf{x}, \lambda_{-\varepsilon}\rho) \cap \Lambda_\Phi$ (here ε is as in (3.6)), we have \asymp in the last calculation rather than just \gtrsim :

$$N_{\beta\rho}(B(\mathbf{x}, \lambda_{-\varepsilon}\rho) \cap \Lambda_\Phi) \asymp \beta \wedge \left(- \sum_{i=1}^d \delta_i \right). \tag{3.14}$$

The quantifiers on this statement are: for all ρ , there exists \mathbf{x} such that (3.14) holds for all $\beta \geq C\rho^\delta$, where C is the implied constant of (3.10) and δ is as in Claim 3.6. By varying ρ we can make β arbitrarily small while still retaining (3.14). This completes the proof. \square

4. Proof of the main theorems

In this section, we prove Theorem 2.7, thus indirectly proving Theorems 1.5 and 1.8, which are consequences of Theorem 2.7.

We recall that in Theorem 2.7, \mathbf{p} is an irreducible good measure with distinct Lyapunov exponents. Without loss of generality, we suppose that the orders $\leq_{\mathbf{p}}$ and \leq on D are equivalent, i.e. that $\chi_1(\mathbf{p}) < \chi_2(\mathbf{p}) < \dots < \chi_d(\mathbf{p})$.

By perturbing the measure \mathbf{p} , we may assume that $\mathbf{p}(\mathbf{a}) > 0$ for all $\mathbf{a} \in E$. Since Λ_Φ is irreducible with respect to \mathbf{p} , this implies that

$$h_{\mathbf{p}}(I_{\leq i} \uparrow I_{< i}) \stackrel{\text{def}}{=} \int \log \frac{\mathbf{p}([\mathbf{a}]_{I_{< i}})}{\mathbf{p}([\mathbf{a}]_{I_{\leq i}})} d\mathbf{p}(\mathbf{a}) > 0 \tag{4.1}$$

for all $i \in D$. Here $[\mathbf{a}]_I = \bigcap_{j \in I} [\mathbf{a}]_j = \{\mathbf{b} \in E : b_j = a_j \ \forall j \in I\}$.

Now fix $\varepsilon > 0$ and $N \in \mathbb{N}$, and let $S = S_N \subseteq E^N$ be the set of all words $\omega \in E^N$ satisfying

$$(1 - \varepsilon)N\chi_i(\mathbf{p}) \leq -\log|\phi'_{\omega,i}| = \sum_{j=1}^N -\log|\phi_{\omega_j,i}| \leq (1 + \varepsilon)N\chi_i(\mathbf{p}) \tag{4.2}$$

and

$$\log \frac{\mu([\omega]_{I_{< i}})}{\mu([\omega]_{I_{\leq i}})} = \sum_{j=1}^N \log \frac{\mathbf{p}([\omega_j]_{I_{< i}})}{\mathbf{p}([\omega_j]_{I_{\leq i}})} \geq (1 - \varepsilon)Nh_{\mathbf{p}}(I_{\leq i} \uparrow I_{< i}), \tag{4.3}$$

where $\mu = \mu_N = \mathbf{p}^N$. Here the notation is slightly different from in (3.3):

$$[\omega]_I = \{\tau \in E^N : \tau_n \in [\omega_n]_I \ \forall n \leq N\}.$$

By the law of large numbers, we have $\lim_{N \rightarrow \infty} \mu_N(S_N) = 1$, so if N is sufficiently large then $\mu(S) \geq 1 - \varepsilon$.

Now define a sequence of sets $T_d \supseteq T_{d-1} \supseteq \dots \supseteq T_0$ as follows: $T_d = S$, and if T_i is defined then let

$$T_{i-1} = \{\omega \in T_i : \mu(T_i \cap [\omega]_{I_{< i}}) \geq \varepsilon\mu([\omega]_{I_{< i}})\}.$$

Letting

$$\mathcal{I}_i = \{[\mathbf{a}]_{I_{< i}} : \mathbf{a} \in E\} = \{\pi_i^{-1}(\mathbf{a}) : \mathbf{a} \in \pi_i(E)\},$$

we have

$$\mu(T_i \setminus T_{i-1}) = \sum_{\substack{P \in \mathcal{I}_i \\ \mu(T_i \cap P) < \varepsilon\mu(P)}} \mu(T_i \cap P) \leq \sum_{\substack{P \in \mathcal{I}_i \\ \mu(T_i \cap P) < \varepsilon\mu(P)}} \varepsilon\mu(P) \leq \varepsilon\mu(E^N) = \varepsilon,$$

so $\mu(T_0) \geq 1 - (d + 1)\varepsilon$. If ε is small enough then $\mu(T_0) > 0$, and $T_0 \neq \emptyset$.

CLAIM 4.1. For all $\omega \in T_0$, we have $T_0 \cap [\omega]_{I_{<i}} = T_i \cap [\omega]_{I_{<i}}$.

Proof. Fix $\tau \in T_i \cap [\omega]_{I_{<i}}$; we will prove by backwards induction that $\tau \in T_j$ for all $j = i, i - 1, \dots, 0$. Fix $j \leq i$, and suppose that $\tau \in T_j$. Since $\tau \in [\omega]_{I_{<i}} \subseteq [\omega]_{I_{<j}}$, we have $[\tau]_{I_{<j}} = [\omega]_{I_{<j}}$. Since $\omega \in T_0 \subseteq T_{j-1}$, this shows that $\mu(T_i \cap [\tau]_{I_{<i}}) \geq \varepsilon \mu([\tau]_{I_{<i}})$, and thus $\tau \in T_{j-1}$. \triangleleft

Note that in this claim it was crucial that we defined the sequence $(T_i)_0^d$ by backward recursion rather than by forward recursion, since we needed the fact that $[\omega]_{I_{<i}} \subseteq [\omega]_{I_{<j}}$ for all $j \leq i$. Combining Claim 4.1 with the definition of $(T_i)_0^d$ yields

$$\mu(T_0 \cap [\omega]_{I_{<i}}) \geq \varepsilon \mu([\omega]_{I_{<i}}) \quad \text{for all } i \in D \text{ for all } \omega \in T_0. \tag{4.4}$$

Now let $\tau \in E^* = \bigcup_{n=0}^\infty E^n$ be a word of fixed length (independent of N) such that

$$\phi_\tau([0, 1]^d) \subseteq (0, 1)^d, \tag{4.5}$$

and consider the diagonal IFS $\Psi = \Psi_N = (\psi_\omega)_{\omega \in T_0}$, where for each $\omega \in T_0$ we write

$$\psi_\omega = \phi_\omega \circ \phi_\tau.$$

Clearly, $\Lambda_\Psi \subseteq \Lambda_\Phi$. To bound $\underline{\dim}_A(\Lambda_\Psi)$ from below using Theorem 3.5, we fix $i \in D$ and $\mathbf{a} \in \pi_i(T_0)$. We have

$$\dim(\Psi_{i,\mathbf{a}}) \geq \frac{\log \#(E_{i,\mathbf{a}})}{\max_{b \in E_{i,\mathbf{a}}} (-\log |\psi'_{i,b}|)}$$

where the ‘fiber IFS’ $\Psi_{i,\mathbf{a}} = (\psi_b)_{b \in E_{i,\mathbf{a}}}$ is as in (3.7). Now, by (4.2),

$$\max_{b \in E_{i,\mathbf{a}}} (-\log |\psi'_{i,b}|) \leq (1 + \varepsilon)N\chi_i(\mathbf{p}) + (-\log |\phi'_{\tau,i}|),$$

and on the other hand

$$\begin{aligned} \#(E_{i,\mathbf{a}}) &\geq \frac{\mu(T_0 \cap \pi_i^{-1}(\mathbf{a}))}{\max_{b \in E_{i,\mathbf{a}}} \mu(T_0 \cap \pi_{i+1}^{-1}(\mathbf{a}, b))} \\ &\geq \frac{\varepsilon \mu(\pi_i^{-1}(\mathbf{a}))}{\max_{b \in E_{i,\mathbf{a}}} \mu(\pi_{i+1}^{-1}(\mathbf{a}, b))} \quad \text{(by (4.4))} \\ &\geq \varepsilon \exp((1 - \varepsilon)N\mathbf{h}_\mathbf{p}(I_{\leq i} \uparrow I_{<i})). \quad \text{(by (4.3))} \end{aligned}$$

So

$$\dim(\Psi_{i,\mathbf{a}}) \geq \delta_i(N, \varepsilon) \stackrel{\text{def}}{=} \frac{(1 - \varepsilon)N\mathbf{h}_\mathbf{p}(I_{\leq i} \uparrow I_{<i}) + \log(\varepsilon)}{(1 + \varepsilon)N\chi_i(\mathbf{p}) - \log |\phi'_{\tau,i}|} \tag{4.6}$$

and thus, by Theorem 3.5,

$$\underline{\dim}_A(\Psi_N) \geq \sum_{i \in D} \delta_i(N, \varepsilon) \xrightarrow{N \rightarrow \infty} \frac{1 - \varepsilon}{1 + \varepsilon} \sum_{i \in D} \frac{\mathbf{h}_\mathbf{p}(I_{\leq i} \uparrow I_{<i})}{\chi_i(\mathbf{p})} \xrightarrow{\varepsilon \rightarrow 0} \sum_{i \in D} \frac{\mathbf{h}_\mathbf{p}(I_{\leq i} \uparrow I_{<i})}{\chi_i(\mathbf{p})}.$$

The right-hand side is equal to $\dim_H(\nu_\mathbf{p})$ by the Ledrappier–Young formula (e.g. [7, Proposition 2.16]). So to complete the proof, we need to show that the sponge Λ_{Ψ_N} is strongly Lalley–Gatzouras and hyperplane diffuse for all N sufficiently large. The coordinate ordering condition follows from (4.2) and the fact that \mathbf{p} has distinct Lyapunov

exponents. Since \mathbf{p} is good, so is Λ_{Ψ_N} , and (4.5) implies that Λ_{Ψ_N} is in fact strongly good. Finally, combining (4.1) with the calculation preceding (4.6) shows that

$$\min_{i \in D} \min_{\mathbf{a} \in \pi_i(E)} \#(E_{i,\mathbf{a}}) \geq 2$$

for all N sufficiently large, and it is easy to check that this is equivalent to Ψ_N being uniformly irreducible. So, by Proposition 3.3, Λ_{Ψ_N} is hyperplane diffuse for all N sufficiently large.

5. Open questions

We conclude with a list of open questions.

Question 5.1. Let $\Lambda_\Phi \subseteq [0, 1]^d$ be an irreducible Barański sponge with distinguishable coordinates, such that $\dim_D(\Lambda_\Phi) < \dim_H(\Lambda_\Phi)$ (see [7]). Does $\Lambda_\Phi \cap \text{BA}_d$ necessarily have full Hausdorff dimension in Λ_Φ (as opposed to just full dynamical dimension as guaranteed by Theorem 1.5)? Alternatively, can it be shown that it is impossible to prove this using the techniques of this paper, by showing that Λ_Φ does not necessarily have hyperplane diffuse subsets of sufficiently large lower Assouad dimension?

Question 5.2. Can Schmidt’s game be used to show that BA_d has full dimension in some fractal defined by a dynamical system which is both non-conformal and nonlinear?

Question 5.3. Is there any fractal Λ defined by a smooth dynamical system (e.g. the limit set of a C^1 IFS) such that $\Lambda \cap \text{BA}_d$ is non-empty but does not have full dimension in Λ ?

Acknowledgements. The first-named author was supported in part by a 2016-2017 Faculty Research Grant from the University of Wisconsin–La Crosse. The second-named author was supported in part by the Simons Foundation grant #245708. The third-named author was supported in part by the EPSRC Programme grant EP/J018260/1. The fourth-named author was supported in part by the NSF grant DMS-1361677. The authors thank the anonymous referee for helpful comments.

A. Appendix. Equivalent formulas for the upper and lower Assouad dimensions

In this paper, we have used the formulas

$$\begin{aligned} \underline{\dim}_A(K) &= \underline{\delta}_1 \stackrel{\text{def}}{=} \liminf_{\beta \rightarrow 0} \inf_{0 < \rho \leq 1} \inf_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \\ &= \underline{\delta}_2 \stackrel{\text{def}}{=} \liminf_{\beta \rightarrow 0} \liminf_{\rho \rightarrow 0} \inf_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)}, \\ \overline{\dim}_A(K) &= \overline{\delta}_1 \stackrel{\text{def}}{=} \limsup_{\beta \rightarrow 0} \sup_{0 < \rho \leq 1} \sup_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \\ &= \overline{\delta}_2 \stackrel{\text{def}}{=} \limsup_{\beta \rightarrow 0} \limsup_{\rho \rightarrow 0} \sup_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta)} \end{aligned}$$

for the lower and upper Assouad dimension, respectively. It is clear from the definitions that $\underline{\dim}_A(K) = \underline{\delta}_1$ and $\overline{\dim}_A(K) = \overline{\delta}_1$, but it is less clear that $\underline{\delta}_1 = \underline{\delta}_2$ and $\overline{\delta}_1 = \overline{\delta}_2$, so we

prove this now. For brevity, we only prove the equality $\underline{\delta}_1 = \underline{\delta}_2$, as the proof of the equality $\overline{\delta}_1 = \overline{\delta}_2$ is similar.

Obviously $\underline{\delta}_1 \leq \underline{\delta}_2$, so we fix $s < \underline{\delta}_2$, and we will show that $\underline{\delta}_1 \geq s$. Choose $0 < \beta \leq 1$ small enough so that

$$\liminf_{\rho \rightarrow 0} \inf_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta/4)} > s,$$

and then choose $\rho_0 > 0$ small enough so that

$$\inf_{\mathbf{x} \in K} \frac{\log N_{\beta\rho}(B(\mathbf{x}, \rho) \cap K)}{-\log(\beta/4)} \geq s \quad \text{for all } 0 < \rho \leq \rho_0.$$

Then every ball $B(\mathbf{x}, \rho)$ centered at a point in K of radius $\leq \rho_0$ contains a $\beta\rho$ -separated set of cardinality at least $(\beta/4)^{-s}$. Now the balls of radius $\beta\rho/2$ centered at the points of this set are disjoint and contained in $B(\mathbf{x}, 2\rho)$. Letting $\kappa = 2\rho$, we see that every ball $B(\mathbf{x}, \kappa)$ with $\mathbf{x} \in K$ and $\kappa \leq 2\rho_0$ contains at least $(\beta/4)^{-s}$ disjoint balls of radius $(\beta/4)\kappa$. Iterating, every such ball contains at least $(\beta/4)^{-ns}$ disjoint balls of radius $(\beta/4)^n\kappa$. Now fix $\mathbf{x} \in K$ and $0 < r \leq R \leq 1$. Let $r' = \min(\rho_0, r)$ and $R' = \min(\rho_0, R)$, and let $n \geq 0$ be chosen so that $(\beta/4)^n \geq r'/R'$ but $(\beta/4)^{n+1} < r'/R'$. Then by the above argument, $B(\mathbf{x}, R') \subseteq B(\mathbf{x}, R)$ contains at least $(\beta/4)^{-ns} \asymp (r'/R')^{-s}$ disjoint balls of radius $(\beta/4)^n R' \geq r'$. If $r' = r$, this shows that $N_r(B(\mathbf{x}, R)) \gtrsim (R/r)^s$, and if $r' < r$, then $r > \rho_0$ and thus $N_r(B(\mathbf{x}, R)) \geq 1 \asymp (R/r)^s$. By the definition of the lower Assouad dimension, this implies that $\underline{\delta}_1 = \underline{\dim}_A(K) \geq s$.

REFERENCES

- [1] K. Barański. Hausdorff dimension of the limit sets of some planar geometric constructions. *Adv. Math.* **210**(1) (2007), 215–245.
- [2] B. Bárány and A. Käenmäki. Ledrappier–Young formula and exact dimensionality of self-affine measures. *Preprint*, 2015, arXiv:1511.05792.
- [3] T. Bedford. Crinkly curves, Markov partitions and box dimensions in self-similar sets. *PhD Thesis*, The University of Warwick, 1984.
- [4] R. Broderick, L. Fishman, D. Kleinbock, A. Reich and B. Weiss. The set of badly approximable vectors is strongly C^1 incompressible. *Math. Proc. Cambridge Philos. Soc.* **153**(02) (2012), 319–339.
- [5] T. Das, L. Fishman, D. Simmons and M. Urbański. Extremality and dynamically defined measures, part I: Diophantine properties of quasi-decaying measures. *Selecta Math. (N.S.)*, to appear. *Preprint*, 2015, arXiv:1504.04778.
- [6] T. Das, L. Fishman, D. Simmons and M. Urbański. Badly approximable vectors and fractals defined by conformal dynamical systems. *Math. Res. Lett.*, to appear. *Preprint*, 2016, arXiv:1603.01467.
- [7] T. Das and D. Simmons. The Hausdorff and dynamical dimensions of self-affine sponges: a dimension gap result. *Invent. Math.*, to appear. *Preprint*, 2016, arXiv:1604.08166.
- [8] M. Denker and M. Urbański. On Sullivan’s conformal measures for rational maps of the Riemann sphere. *Nonlinearity* **4**(2) (1991), 365–384.
- [9] M. Einsiedler, L. Fishman and U. Shapira. Diophantine approximations on fractals. *Geom. Funct. Anal.* **21**(1) (2011), 14–35.
- [10] M. Einsiedler and J. Tseng. Badly approximable systems of affine forms, fractals, and Schmidt games. *J. Reine Angew. Math.* **660** (2011), 83–97.
- [11] K. Falconer. Random fractals. *Math. Proc. Cambridge Philos. Soc.* **100**(3) (1986), 559–582.
- [12] L. Fishman. Schmidt’s game on fractals. *Israel J. Math.* **171**(1) (2009), 77–92.
- [13] L. Fishman and D. Simmons. Intrinsic approximation for fractals defined by rational iterated function systems – Mahler’s research suggestion. *Proc. Lond. Math. Soc. (3)* **109**(1) (2014), 189–212.

- [14] J. M. Fraser. Assouad type dimensions and homogeneity of fractals. *Trans. Amer. Math. Soc.* **366**(12) (2014), 6687–6733.
- [15] J. M. Fraser and D. Howroyd. Assouad type dimensions for self-affine sponges. *Preprint*, 2015, arXiv: 1508.03393.
- [16] A. Käenmäki. On natural invariant measures on generalised iterated function systems. *Ann. Acad. Sci. Fenn. Math.* **29**(2) (2004), 419–458.
- [17] A. Käenmäki, J. Lehrbäck and M. Vuorinen. Dimensions, Whitney covers, and tubular neighborhoods. *Indiana Univ. Math. J.* **62**(6) (2013), 1861–1889.
- [18] R. Kenyon and Y. Peres. Measures of full dimension on affine-invariant sets. *Ergod. Th. & Dynam. Sys.* **16**(2) (1996), 307–323.
- [19] D. Kleinbock, E. Lindenstrauss and B. Weiss. On fractal measures and Diophantine approximation. *Selecta Math.* **10** (2004), 479–523.
- [20] D. Kleinbock and B. Weiss. Badly approximable vectors on fractals. *Israel J. Math.* **149** (2005), 137–170.
- [21] S. P. Lalley and D. Gatzouras. Hausdorff and box dimensions of certain self-affine fractals. *Indiana Univ. Math. J.* **41**(2) (1992), 533–568.
- [22] C. McMullen. The Hausdorff dimension of general Sierpinski carpets. *Nagoya Math. J.* **96** (1984), 1–9.
- [23] F. Przytycki and M. Urbański. *Conformal Fractals: Ergodic Theory Methods (London Mathematical Society Lecture Note Series, 371)*. Cambridge University Press, Cambridge, 2010.
- [24] W. M. Schmidt. Badly approximable systems of linear forms. *J. Number Theory* **1** (1969), 139–154.
- [25] W. M. Schmidt. *Diophantine Approximation (Lecture Notes in Mathematics, 785)*. Springer, Berlin, 1980.