

# Parasitic infection alters the physiological response of a marine gastropod to ocean acidification

C. D. MACLEOD\* and R. POULIN

Department of Zoology, University of Otago, PO Box 56, Dunedin 9054, New Zealand

(Received 24 February 2016; revised 12 April 2016; accepted 21 April 2016; first published online 25 May 2016)

## SUMMARY

Increased hydrogen ion concentration and decreased carbonate ion concentration in seawater are the most physiologically relevant consequences of ocean acidification (OA). Changes to either chemical species may increase the metabolic cost of physiological processes in marine organisms, and reduce the energy available for growth, reproduction and survival. Parasitic infection also increases the energetic demands experienced by marine organisms, and may reduce host tolerance to stressors associated with OA. This study assessed the combined metabolic effects of parasitic infection and OA on an intertidal gastropod, *Zeacumantus subcarinatus*. Oxygen consumption rates and tissue glucose content were recorded in snails infected with one of three trematode parasites, and an uninfected control group, maintained in acidified (7.6 and 7.4 pH) or unmodified (8.1 pH) seawater. Exposure to acidified seawater significantly altered the oxygen consumption rates and tissue glucose content of infected and uninfected snails, and there were clear differences in the magnitude of these changes between snails infected with different species of trematode. These results indicate that the combined effects of OA and parasitic infection significantly alter the energy requirements of *Z. subcarinatus*, and that the species of the infecting parasite may play an important role in determining the tolerance of marine gastropods to OA.

Key words: ocean acidification, parasite, gastropod, metabolism, oxygen, glucose.

## INTRODUCTION

Stressors associated with ocean acidification (OA) and parasitic infection have the potential to alter the metabolic demands experienced by many marine organisms. OA changes the relative concentrations of physiologically important charged molecules, such as hydrogen, bicarbonate and carbonate ions (Dickson *et al.* 2007), potentially increasing the energetic costs of intra- and extra-cellular acid–base homeostasis (Pan *et al.* 2015) and the biomineralization of calcium carbonate (Kelly and Hofmann, 2013). This is especially true for marine invertebrates, many of which possess a limited ionoregulatory capacity and biogenically form calcified structures (Pörtner, 2008). Parasitic infection can also increase the energetic demands of host organisms, by absorbing nutrients from, or causing damage to, host tissue, and by inducing behaviours that are energetically costly (Bates *et al.* 2011; Toledo and Fried, 2011).

The process of maintaining pH balance across cellular spaces requires a significant investment of metabolic energy (Pörtner *et al.* 2000), and exposure to an acidified environment has often been correlated with increased respiration (Zhang *et al.* 2014). The simultaneous reduction of the carbonate ion concentration of seawater could also dramatically increase

the energy required to form calcified structures (see review in Parker *et al.* 2013). Consequently, changes to the chemical speciation of seawater caused by OA may have wide-ranging physiological effects on many marine species, which may manifest themselves as changes in energy usage.

By definition, parasitic infection has a negative effect on host organisms, and, like changing abiotic environmental conditions, can alter the energy demands of the host. Parasites can withdraw energy from hosts directly, through the absorption of nutrients or ingestion of tissue (Sorensen and Minchella, 2001), and indirectly, by initiating behaviours that increase the host's metabolic demands (Bates *et al.* 2011; Macnab and Barber, 2012). Accordingly, parasites additively increase the energy requirements of infected organisms, as the host must generate sufficient energy to meet its own needs plus the needs of the parasite, invest metabolic energy into tissue repair, and/or compensate for parasite-mediated increases in energetically costly behaviour.

The energy status, and rates of energy consumption, of living organisms are commonly quantified by measuring oxygen consumption rates (e.g. Melatunan *et al.* 2011) and the concentration of free glucose in body tissues and fluids (e.g. Berthelin *et al.* 2000). The oxygen consumption rates of marine gastropods typically increase in response to moderate abiotic stress, as a result of compensatory or regulatory processes requiring more metabolic energy, and decrease in response to extreme abiotic stress as their compensatory ability is overwhelmed,

\* Corresponding author. Department of Zoology, University of Otago, PO Box 56, Dunedin 9054, New Zealand. Tel: +64 3 479 7964. Fax: +64 3 479 7584. E-mail: [colin.macleod@postgrad.otago.ac.nz](mailto:colin.macleod@postgrad.otago.ac.nz)

i.e. metabolic depression (Pörtner and Farrell, 2008). Changes in tissue glucose concentration can be caused by an imbalance between glycogen breakdown and the rate of glucose consumption in the glycolytic process. If glycogen breakdown outpaces glucose consumption, tissue glucose levels will rise; conversely, if glucose consumption outpaces glycogen breakdown, tissue glucose will fall (Martínez-Quintana and Yepiz-Plascencia, 2012). Elevated rates of glucose consumption can eventually lead to mortality, as energy stores are depleted at a greater rate than they are replenished.

The results of previous studies of marine gastropods exposed to simulated OA conditions are variable: increased rates of oxygen consumption at 6.63 and 7.8 pH (Zhang *et al.* 2014); decreased rates of oxygen consumption at 7.6 pH (Melatunan *et al.* 2011); reduced heart rates at 7.6 pH (Ellis *et al.* 2009); increased ATP concentration at 7.6 pH (Melatunan *et al.* 2011); and increased mortality between 7.4 and 7.98 pH (Kimura *et al.* 2011; Zhang *et al.* 2014). Variability in the oxygen consumption rates of different species exposed to the same pH suggests that the metabolic tolerance of marine gastropods is highly species-specific. The few studies that have exposed more than one species of marine gastropod to acidified seawater (Coleman *et al.* 2014; Zhang *et al.* 2014), or conspecifics taken from different populations (Lardies *et al.* 2014), concluded that abiotic characteristics of the gastropods' habitat were the major predictors of tolerance to stressors associated with OA. Although caused by very different physiological mechanisms, the effects of trematode infection and OA on marine gastropods are relatively similar. Trematode infection can increase or decrease oxygen consumption rates, elevate heart rates and positively or negatively affect the survival of infected gastropods (see the review in Galaktionov and Dobrovolskij, 2003). In addition, trematode infection has been linked with changes in haemolymph glucose levels and increases in the rate of glycolysis; these effects appear to be specific to host and parasite species (Fried and Graczyk, 1997).

The aim of this study was to describe the combined effects of trematode infection and exposure to acidified seawater on the metabolic status of the intertidal mud snail *Zeacumantus subcarinatus*, and to identify species-specific effects of the infecting parasite. Accordingly, *Z. subcarinatus* snails were categorized as infected with one of three trematode species (*Maritrema novaezealandensis*, *Acanthoparyphium* sp. or *Philophthalmus* sp.), or as uninfected, and exposed to acidified (7.6 and 7.4 pH) or unmodified (8.1 pH) seawater. The pH treatments chosen for this experiment correspond to the current average seawater pH measured at the collection site of the experimental organisms (8.085 pH, see Supplementary Material Table S1 and MacLeod, 2015), which was similar to the global average of 8.1 pH, and predicted

average conditions for the year 2100 (7.6 pH) and 2300 (7.4 pH) reported in the Intergovernmental Panel on Climate Change (IPCC, 2014).

## MATERIALS AND METHODS

### *Trematode–snail system*

The herbivorous New Zealand mud snail, *Z. subcarinatus*, is widely distributed along the New Zealand coast, occupies a wide range of habitats (sand, rock and mud substrates) from the sublittoral zone to the littoral fringe, and is a first intermediate host, i.e. the site of asexual reproduction, for many species of trematode parasites (Fredensborg *et al.* 2005). Trematode infection of *Z. subcarinatus* causes complete sterilization, as parasite tissue displaces the host's reproductive organs. At the collection site used in this experiment [Lower Portobello Bay (LPB), South Island, New Zealand] *Z. subcarinatus* is infected by eight trematode parasites (Leung *et al.* 2009), although in the following experiments we focus on the three most common species: *M. novaezealandensis* (60% prevalence), *Philophthalmus* sp. (10% prevalence) and *Acanthoparyphium* sp. (3% prevalence). All three trematode species produce free-swimming larvae (cercariae) within the snail host, which periodically emerge in large numbers to actively infect the next host in the parasite's life cycle (*Acanthoparyphium* sp. and *M. novaezealandensis*) or form a cyst on the surface of a transport host (*Philophthalmus* sp.) (Martorelli *et al.* 2004, 2008).

### *Snail collection and morphology, and parasite identification*

Approximately 2000 *Z. subcarinatus* snails were collected at LPB in July 2013 and subsequently screened for trematode infection by exposing snails to physical conditions that trigger cercarial emergence: warmed seawater (25 °C) and constant light. Trematode species were identified by inspecting cercariae under a dissecting microscope and comparing cercarial morphology with published descriptions of all parasite species (Martorelli *et al.* 2008). Snails that were positively identified as infected with a parasite of interest were maintained at room temperature (approximately 18–20 °C) for one week before being screened a second time, thus reducing the probability of selecting snails that were infected by two parasite species. All snails selected for the experiment were then marked with individual identification labels (Bee Works, Orillia, Canada), maintained at room temperature in aerated seawater (approximately pH 8.1, 20 °C), and fed sea lettuce (*Ulva* spp.) *ad libitum*. Prior to the experiments, the wet weight of each snail was recorded ( $\pm 0.001$  g), and the shell length measured using digital photographs (64 $\times$  magnification) and image analysis software (ImageJ). Snail length was used

Table 1. Mean values ( $\pm$ S.D.) of all measured and calculated parameters used to characterize the carbonate chemistry of unmodified and acidified seawater

Treatment	pH <sub>T</sub> (measured)	Temp. (°C)	Salinity	Alkalinity ( $\mu\text{mol kg}^{-1}$ )	DIC ( $\mu\text{mol kg}^{-1}$ )	pH <sub>T</sub> (calculated)	pCO <sub>2</sub> (calculated)	$\Omega$ Aragonite
pH 8.1	8.09 $\pm$ 0.03	12.5 $\pm$ 0.3	31.7 $\pm$ 0.6	2361 $\pm$ 10	2138 $\pm$ 11	8.12 $\pm$ 0.03	365 $\pm$ 30	2.52 $\pm$ 0.2
pH 7.6	7.60 $\pm$ 0.03	12.6 $\pm$ 0.6	31.9 $\pm$ 0.6	2389 $\pm$ 7	2351 $\pm$ 16	7.64 $\pm$ 0.04	1304 $\pm$ 115	0.94 $\pm$ 0.1
pH 7.4	7.40 $\pm$ 0.03	12.6 $\pm$ 0.5	31.3 $\pm$ 0.6	2375 $\pm$ 12	2397 $\pm$ 13	7.45 $\pm$ 0.04	1980 $\pm$ 110	0.62 $\pm$ 0.1

Note: pH was measured on the total hydrogen ion scale.

Temp., temperature; DIC, dissolved inorganic carbon;  $\Omega$ , saturation state.

throughout the experiment as a proxy for age, as snails taken from the same population typically have similar growth rates (Graham, 2003).

#### OA simulation system

In order to expose snails to acidified seawater, a modular OA simulation system was designed (MacLeod *et al.* 2015). Three seawater aquaria were constructed, each consisting of a 120 L culture tank (Stowers Containment Solutions, Christchurch, New Zealand) [870 mm (L)  $\times$  600 mm (W)  $\times$  295 mm (H)], an Aquis700 pump and filtration unit (Aqua One<sup>®</sup>, Nelson, New Zealand), a Hailea HC-150A refrigeration unit (Hailea, Guangdong, China) and a TUNZE<sup>™</sup> pH/CO<sub>2</sub> regulation unit (TUNZE<sup>™</sup>, Penzberg, Germany). The pH, measured on the total hydrogen ion scale, was adjusted with 100% CO<sub>2</sub> gas and monitored potentiometrically with glass electrodes calibrated with saltwater buffers [2-amino-2-hydroxy-1,3-propanediol (Tris) and 2-aminopyridine (AMP)]. Temperature was actively controlled using the flow-through chiller unit, while total alkalinity ( $A_T$ ) and salinity were passively controlled by the regular replacement of unmodified seawater (20 L/48 h); light levels were also standardized across all culture tanks. Seawater in the three culture tanks was maintained at 12.5 °C, 32 salinity and at one of the three pH treatment levels: pH 7.4, 7.6 and 8.1. Each culture tank was also aerated with ambient air by an AquaOne 9500 aquarium bubbler (Aqua One<sup>®</sup>, Nelson, New Zealand), and oxygen saturation, measured daily with a YSI ProODO (YSI, Yellow Springs, OH, USA) was maintained at similar levels to those found in the habitat of *Z. subcarinatus* (approximately 98%, MacLeod, 2015). We also validated the potentiometric regulation of pH by measuring  $A_T$  and dissolved inorganic carbon (DIC) in seawater samples taken from each culture tank, and used that data to calculate pH with the software package SWCO2 (Hunter, 2007) (Table 1).

#### Experimental design

The combined effects of parasitic infection and exposure to acidified seawater on *Z. subcarinatus* individuals

were investigated by exposing snails from four infection categories, i.e. uninfected snails and those infected with *M. novaezealandensis*, *Philophthalmus* sp. or *Acanthoparyphium* sp., to all pH treatments. In each pH treatment, approximately 30 snails from each infection category were distributed evenly between five nylon mesh chambers, i.e. 20 chambers and 120 snails per treatment, and the chambers submerged in unmodified or acidified seawater for a period of 90 days. To minimize the effects of abrupt changes in seawater pH, snails in the acidified treatments were gradually acclimated to reduced pH over a 2-week period. Average snail length was 14.223  $\pm$  1.52 (S.D.) mm (max. 19.64 mm, min. 10.38 mm) and average wet weight was 0.233  $\pm$  0.06 (S.D.) g (max. 0.503 g, min. 0.115 g). Preparatory analysis showed no significant differences in these parameters between groups (length –  $F_{11,228} = 1.831$ ,  $P = 0.205$ , weight –  $F_{11,228} = 0.965$ ,  $P = 0.479$ ). Throughout the 90-day period, snails were provided with a constant supply of sea lettuce (*Ulva* spp.). To account for any unrecorded and unwanted variation in the performance of a particular culture tank and associated apparatus, i.e. tank effect, the pH assigned to each culture tank was changed at 30 and 60 days and snails transferred between tanks. Consequently, all snails experienced constant pH conditions and spent equal amounts of time in each culture tank. In addition to transferring snails between culture tanks, the position of each chamber was changed within each tank every four days, so that all chambers spent an equal period of time at each of the 20 positions available in the culture tanks. The periodic movement of snail chambers was carried out so that no snails spent an unequal amount of time in close proximity to the CO<sub>2</sub> inflow point, although frequent testing of seawater pH did not identify any pH gradient in the culture tanks.

During the 90-day period, the oxygen consumption of 20 snails from each infection category in each pH treatment was measured three times at 4-week intervals, i.e. three separate trials of 240 snails. Oxygen trials were completed before snails were transferred between tanks. Consequently, oxygen consumption data represent an average value for each snail using data from all trials, thus compensating for a potential tank effect. At the end of the 90-day period, all snails

were euthanized, parasite tissue was removed, and the remaining tissue was frozen ( $-80^{\circ}\text{C}$ ) for glucose analysis.

#### Oxygen consumption

In each trial, the oxygen consumption of 20 snails was measured over 4 h, normalized for snail mass (g), and converted into the rate of oxygen consumed in micromoles per hour ( $\mu\text{mO}_2 \text{ h}^{-1} \text{ g}^{-1}$ ). The snails were placed in individual plastic 50 mL screw-top containers which were then completely filled with acidified or unmodified seawater by fully submerging container and lid in the culture tank associated with each snail; the oxygen concentration of source water was also measured prior to filling containers. A blank container, i.e. one containing only acidified or unmodified seawater, was also used for each pH/infection category combination, i.e. 12 blank containers per trial. The containers were then closed, sealed with Parafilm, and fully submerged in the appropriate culture tank to maintain constant temperature. After 4 h, the oxygen content of seawater ( $\pm 0.1 \text{ mg L}^{-1}$ ) in each container was measured with an optical oxygen sensor (YSI-proODO), and the total oxygen consumed by each snail was calculated using the formula:

$$\text{Oxygen consumed} = (\text{O}_{2(i)} - \text{O}_{2(b)}) - (\text{O}_{2(i)} - \text{O}_{2(s)}) \quad (1)$$

where  $\text{O}_{2(i)}$  is the initial oxygen concentration of the source water,  $\text{O}_{2(b)}$  is the final oxygen concentration of seawater in the blank container and  $\text{O}_{2(s)}$  is the final oxygen concentration of seawater in the containers that held snails. The oxygen concentration of the seawater in the blank container was used to account for changes to the oxygen content of source water caused by planktonic photosynthesis and/or respiration, which would also occur in containers holding snails. The oxygen concentration in the 50 mL containers did not fall below 85% during the 4 h trials, and we assumed that this minimal decrease in available oxygen would not alter the respiration rate of the snails.

#### Glucose concentration

The free glucose concentration of snail muscle tissue was measured in approximately 20 snails per infection category per pH treatment using a GAGO-20 colorimetric assay kit (Sigma-Aldrich, St. Louis, MO, USA). Assay reagents in this kit react with D-glucose to form a pink colour, the intensity of which is proportional to the concentration of D-glucose in a sample fluid. D-glucose is the most common monosaccharide in living organisms, and is integral to the glycolytic pathway of gastropods (Martínez-Quintana and Yepiz-Plascencia, 2012). Free glucose was extracted from pre-weighed snail tissue ( $\pm 0.001 \text{ g}$ ) by immersing the tissue in 500  $\mu\text{L}$  of de-ionized water, heating to  $60^{\circ}\text{C}$  for 1 h, and

grinding to an opaque slurry with a small pestle. The snail solution was then centrifuged at 14 000 rpm for 5 min and 200  $\mu\text{L}$  of clear supernatant removed and stored at  $-20^{\circ}\text{C}$  until assays were conducted.

For each assay, 40  $\mu\text{L}$  of supernatant was mixed with 80  $\mu\text{L}$  of assay reagent in one well of a 96-well plastic culture plate and incubated at  $37^{\circ}\text{C}$  for 30 min. After 30 min, 80  $\mu\text{L}$  of 12N sulphuric acid was added to each well to stop the colorimetric reaction. Duplicate assays were completed for each snail and used to calculate an average value of glucose concentration for each individual. As a precaution against sample loss or human error, samples from each infection category/pH combination were divided such that no entire group was processed in a single incubation.

The colour intensity of all samples was measured using a multi-mode Fluostar Omega microplate reader (BMG Labtech GmbH, Ortenberg, Germany), which exposed each well to 540 nm light and measured the absorbance of each sample. A single well filled with de-ionized water was included in each 96-well plate to provide a negative control, allowing the absorbance of de-ionized water to be subtracted from the absorbance of the samples. Per the protocol for the assay kit, each plate also contained a series of standard solutions of D-glucose: 5, 10, 20, 30 and 40  $\mu\text{g}$  glucose  $\text{mL}^{-1}$ . The absorbance readings for these standards were used to create an absorbance concentration curve that was then used, in conjunction with the initial weight of the tissue sample, to convert sample absorbance to micrograms of glucose per gram of snail tissue.

#### Statistical analysis

Oxygen consumption data were analysed using three different model structures. The first model analysed the combined oxygen data from all three trials with a linear mixed-effect structure, and used oxygen consumption as the response variable, pH, infection category and initial shell length as fixed effects, and 'Snail ID', 'Chamber ID' and 'Trial' as random effects. To further quantify the effect of infection, this model was also run with two modifications: data from all infected snails were pooled to compare only 'infected' *vs* 'uninfected' snails; and infection status was removed from the analysis, thus pooling data from all snails. The second model structure analysed each pH treatment (three groups) and infection category (four groups) separately, i.e. seven separate models each using a different subset of the overall data. This model type also had a linear mixed-effect structure and used oxygen consumption as the response variable, infection category/pH (depending on data subset) and initial shell length as fixed effects, and 'Snail ID', 'Chamber ID' and 'Trial' as random effects. These models were used to identify significant




Fig. 1. Mean oxygen consumption rates ( $\pm$ S.E.M.) of snails exposed to acidified and unmodified seawater for 90 days: (A) data grouped by pH treatment and (B) data grouped by infection category. Lowercase letters in bold indicate significant differences between infection categories within pH treatments (A), and between pH treatments within infection categories (B).  $n = 20$  in each pH/infection category combination.

differences within the 'infection category' and 'pH' groups (Fig. 1 and Supplementary Material Tables S2 and S3). The third model used a linear structure to analyse the consumption of oxygen in each trial separately, with oxygen consumption as a response variable, and pH and initial shell length as fixed effects (Supplementary Material Table S4).

Glucose concentration was analysed using two linear mixed-effect models. The first provided an overall analysis, using glucose concentration as the response variable, pH, infection category, and initial shell length as fixed effects, and 'Chamber ID' as a random effect. As with the oxygen analysis, this model was also run with two modifications: data from all infected snails were pooled to compare only 'infected' vs 'uninfected' snails; and infection status was removed from the analysis, thus pooling data from all snails. The second model analysed each pH treatment (three groups) and infection category

separately (four groups), i.e. seven separate models each using a different subset of the overall data. This model type also had a linear mixed-effect structure and used glucose concentration as the response variable, infection category/pH (depending on data subset) and initial shell length as fixed effects, and 'Chamber ID' as a random effect. As with oxygen consumption data, these models were used to identify significant differences in glucose concentration within the 'infection category' and 'pH' groups (Fig. 2 and Supplementary Material Tables S5 and S6).

The random effects 'Snail ID' and 'Trial' were used in the oxygen data analysis to compensate for repeated measurements of each snail and for pooling data from each trial, respectively. The random effect 'Chamber ID' was used in both oxygen and glucose trials to quantify the repeatability of measurements of groups of snails maintained in the same tank. Repeatability can be estimated by




Fig. 2. Mean tissue glucose concentration ( $\pm$ S.E.M.) of snails exposed to acidified and unmodified seawater for 90 days: (A) data grouped by pH treatment and (B) data grouped by infection category. Lowercase letters in bold indicate significant differences between infection categories within pH treatments (A), and between pH treatments within infection categories (B). Sample sizes: uninfected, 8.1 pH = 14, 7.6 pH = 18, 7.4 pH = 20; *Philophthalmus* sp., 8.1 pH = 18, 7.6 pH = 19, 7.4 pH = 18; *M. novaezealandensis*, 8.1 pH = 19, 7.6 pH = 17, 7.4 pH = 13; *Acanthoparyphium* sp., 8.1 pH = 18, 7.6 pH = 20, 7.4 pH = 18.

calculating the intra-class correlation (ICC) between groups, in this case snails within a particular chamber, by finding the per cent variance attributed to the grouping factor (Chamber ID). An ICC score of 0% indicates no repeatability of measurements between groups and a score of 100% indicates identical measurements, i.e. pseudoreplication. Calculating ICC scores allowed us to assess the independence, or lack thereof, of data points taken from multiple chambers:

$$ICC = \left( \frac{(\text{between} - \text{chamber variance})}{(\text{between} - \text{chamber variance}) + (\text{within} - \text{chamber variance})} \right) \times 100$$

An *a posteriori* correlation analysis was also conducted on oxygen consumption rates and tissue glucose concentration, after it became clear that there was a weak negative relationship between these parameters. This relationship was analysed using a simple linear model that used end point oxygen data and glucose data.

In all analyses, model selection was performed using comparative AIC values and manual simplification of model parameters. Where necessary, oxygen consumption data was normalized using the *powerTransform* function in the package *car* (Fox *et al.* 2014). All models used for oxygen and glucose analysis were designed using the function *lmer* in the package *lme4* (Bates *et al.* 2014) using R version 3.1.0 (R Development Core Team, 2014). Fixed

Table 2. Output for the linear mixed-effect analysis of oxygen consumption rates in snails maintained for 90 days in acidified or unmodified seawater ( $n = 240$ ). ‘Length’ represents the length of snail shells at the beginning of the 90-day trial. Significant  $P$ -values are shown in bold

	df	MS	$F$	$P$
pH	2	0.715	22.94	<b>&lt;0.001</b>
IC	3	0.053	1.70	0.170
Length	1	1.229	39.40	<b>&lt;0.001</b>
pH*IC	6	0.206	6.60	<b>&lt;0.001</b>
IC*Length	3	0.135	4.32	<b>0.005</b>

IC, infection category.

effects were considered significant if  $P$  values were less than or equal to 0.05.

RESULTS

Oxygen consumption

The oxygen consumption of *Z. subcarinatus* was significantly affected by pH, initial shell length and the interaction of infection category with both pH and initial shell length, but not by infection category alone (Table 2). Both modifications of the overall analysis, i.e. pooling infected snails and pooling all snails, found similar results (Supplementary Material Table S7). Despite the non-significant effect of infection category in the overall analysis, the individual analysis of each pH treatment and infection category found significant differences in oxygen consumption rates in both data subsets (Fig. 1). At 8.1 pH, the rate of oxygen consumption was higher in infected snails relative to uninfected individuals, although these differences were only significant in snails infected with *Acanthoparyphium* sp. and *Philophthalmus* sp. This trend was reversed in the 7.6 pH treatment, where uninfected snails consumed more oxygen per hour than all infected categories. In the same treatment, the oxygen consumption rate of *M. novaezealandensis*-infected snails was significantly lower than uninfected and *Philophthalmus* sp.-infected snails. The relative oxygen consumption rates of infection categories changed again in the 7.4 pH treatment, where there was no statistical difference between uninfected and *Philophthalmus* sp.-infected snails, while *Acanthoparyphium* sp.- and *M. novaezealandensis*-infected snails consumed oxygen at a significantly higher rate than the other two groups. Snails in all infection categories exhibited higher rates of oxygen consumption at 7.6 pH relative to 8.1 pH, and these differences were significant for all infection categories except *Philophthalmus* sp.-infected snails (Fig. 1B). At 7.4 pH, oxygen consumption rates were more varied relative to the 7.6 pH treatment: uninfected and *Philophthalmus* sp.-infected snails exhibited significantly reduced consumption rates,

Table 3. Output for the linear mixed-effect analysis of oxygen consumption rates in each infection category and pH treatment ( $n = 240$ ). ‘Length’ represents the length of snail shells at the beginning of the 90-day trial. Significant  $P$ -values are shown in bold

	df	MS	$F$	$P$
IC				
Uninfected (60)				
pH	2	0.52431	25.1871	<b>&lt;0.001</b>
Length	1	0.02371	1.1392	0.287
<i>M. novaezealandensis</i> (60)				
pH	2	0.25776	14.805	<b>&lt;0.001</b>
Length	1	0.25776	14.805	<b>&lt;0.001</b>
<i>Philophthalmus</i> sp. (60)				
pH	2	0.141265	3.7988	<b>0.0242</b>
Length	1	0.038169	1.0264	0.3604
<i>Acanthoparyphium</i> sp. (60)				
pH	2	0.033858	3.7914	<b>0.0244</b>
Length	1	0.147470	16.5136	<b>&lt;0.001</b>
pH				
8.1 pH (80)				
IC	3	0.30399	6.9054	<b>0.0093</b>
Length	1	0.11777	2.6753	0.1036
7.6 pH (80)				
IC	3	0.059786	2.9686	<b>0.0326</b>
Length	1	0.223457	11.0954	<b>&lt;0.001</b>
7.4 pH (80)				
IC	3	0.027802	4.2092	<b>0.006</b>
Length	1	0.145793	22.0734	<b>&lt;0.001</b>

IC, infection category.

*M. novaezealandensis*-infected snails showed significantly increased consumption rates, and *Acanthoparyphium* sp.-infected snails showed no difference between the two pH treatments. These observations were supported by the overall analysis which found that pH significantly altered rates of oxygen consumption (Table 2), although only one out of three trials found a significant effect of pH on the oxygen consumption rate of *Acanthoparyphium* sp.-infected snails (Supplementary Material Table S4).

ICC scores generated by the random effect ‘Chamber ID’ showed that the repeatability of oxygen consumption data was 11% overall and between 5 and 22% in the data subsets (Supplementary Material Table S8), indicating that repeatability was low and pseudoreplication not a confounding factor in these data.

Initial shell length, the variable used as a proxy for age, significantly affected oxygen consumption rates in the overall analysis (Table 2). However, in the individual analysis of each infection category, we see that initial shell length was only a significant factor for *M. novaezealandensis*- and *Acanthoparyphium* sp.-infected snails (Table 3). Subsequent analysis of the relationship between shell length/age and oxygen consumption showed that infected snails exhibited a negative correlation between length and oxygen consumption (average  $R^2$  value, 0.16), while uninfected snails exhibited a positive correlation (average  $R^2$  value, 0.01).

Table 4. Output for the linear mixed-effect analysis of tissue glucose concentration in snails maintained for 90 days in acidified or unmodified seawater ( $n = 214$ ). 'Length' represents the length of snail shells at the beginning of the 90-day trial. Significant  $P$ -values are shown in bold

	df	MS	$F$	$P$
pH	2	0.015	5.22	<b>0.006</b>
IC	3	0.045	15.45	<b>&lt;0.001</b>
Length	1	0.007	2.37	0.125
pH*IC	6	0.020	6.91	<b>&lt;0.001</b>

IC, infection category.

#### Free glucose concentration

The free glucose concentration in *Z. subcarinatus* tissue was significantly affected by pH, infection category, and the interaction of these variables (Table 4), although the response of each infection category to acidified seawater was quite different (Fig. 2B). Modifications of the overall analysis found that pooling all infected snails also showed a significant effect of infection status, i.e. infected *vs* uninfected, while pooling all snails regardless of infection category removed the significant effect of pH (Supplementary Material Table S9). The individual analysis of each infection category showed that the glucose concentration of *Philophthalmus* sp.-infected snails was not significantly affected by pH, but was significantly affected by initial shell length (Table 5). Within infection categories, uninfected and *Acanthoparyphium* sp.-infected snails exhibited significantly lower glucose concentrations in the 7.6 pH treatment relative to the control (8.1 pH) treatment, while *Philophthalmus* sp.- and *M. novaezealandensis*-infected snails showed no statistical differences between the two treatments. Uninfected, *Acanthoparyphium* sp.-infected and *Philophthalmus* sp.-infected snails exhibited higher glucose concentrations at 7.4 pH relative to the 7.6 pH treatment, although relative to the control (8.1 pH) treatment, glucose concentrations were higher (*Philophthalmus* sp.), lower (*Acanthoparyphium* sp.) or not significantly different (uninfected). Snails infected with *M. novaezealandensis* exhibited significantly lower glucose concentrations at 7.4 pH relative to 8.1 and 7.6 pH treatments. There were significant differences in free glucose concentration between infection categories in all pH treatments (Fig. 2A): at 8.1 pH, *Acanthoparyphium* sp.-infected snails exhibited significantly higher glucose concentrations than *M. novaezealandensis*-infected snails, and both infection categories exhibited significantly higher glucose concentrations than uninfected and *Philophthalmus* sp.-infected snails; at 7.6 pH, all infected snails exhibited significantly higher glucose concentrations

Table 5. Output for the linear mixed-effect analysis of tissue glucose concentration of snails in each infection category and pH treatment. Sample sizes for each data subset are given in parentheses. 'Length' represents the length of snail shells at the beginning of the 90-day trial. Significant  $P$ -values are shown in bold

	df	MS	$F$	$P$
IC				
Uninfected (52)				
pH	2	0.028	7.59	<b>0.001</b>
Length	1	0.001	0.33	0.567
<i>M. novaezealandensis</i> (48)				
pH	2	0.0008	8.74	<b>&lt;0.001</b>
Length	1	0.00005	0.05	0.815
<i>Philophthalmus</i> sp. (55)				
pH	2	0.048	1.85	0.167
Length	1	0.161	6.26	<b>0.015</b>
<i>Acanthoparyphium</i> sp. (56)				
pH	2	1.566	6.54	<b>0.003</b>
Length	1	0.177	0.74	0.393
pH				
8.1 pH (71)				
IC	3	0.020	9.75	<b>&lt;0.001</b>
7.6 pH (71)				
Length	1	0.005	2.46	0.121
IC	3	0.041	17.11	<b>&lt;0.001</b>
7.4 pH (69)				
Length	1	0.00003	0.01	0.913
IC	3	0.043	2.77	<b>0.049</b>
Length	1	0.040	2.60	0.112

IC, infection category.

than uninfected snails, and *M. novaezealandensis*-infected snails exhibited significantly higher glucose concentrations than *Acanthoparyphium* sp.-infected snails; and at 7.4 pH, *Philophthalmus* sp.-infected snails exhibited significantly higher glucose concentrations than uninfected and *M. novaezealandensis*-infected snails, and uninfected and *Acanthoparyphium* sp.-infected snails exhibited significantly higher glucose concentrations than *M. novaezealandensis*-infected snails. ICC scores generated by the random effect 'Chamber ID' showed that the repeatability of glucose data was 13% overall and between 1 and 13% in the data subsets (Supplementary Material Table S8), indicating that repeatability was low and pseudoreplication not a confounding factor in these data. The linear model used to analyse the relationship between oxygen consumption rates recorded in the final trial and tissue glucose concentration showed that there was a weak but significant negative relationship ( $t = 4.003$ ,  $P < 0.001$ ,  $R^2 = 0.10$ ), i.e. as oxygen consumption increased, tissue glucose concentration decreased.

#### DISCUSSION

The goal of this study was to assess changes to the metabolic status of the mud snail *Z. subcarinatus*



caused by the combined stressors of OA and parasitic infection. The metabolic status of *Z. subcarinatus* was quantified by recording oxygen consumption rates and tissue glucose content during a 90-day exposure to unmodified (8.1 pH) or acidified (7.6 and 7.4 pH) seawater. In light of the negative correlation found between endpoint oxygen consumption rates and tissue glucose content, exposure to 7.6 pH seawater appeared to increase the metabolic demands experienced by both uninfected and infected individuals, although the relationship between these parameters was less clear in infected snails. Upon exposure to 7.4 pH seawater, uninfected snails exhibited a decrease in oxygen consumption rates relative to the 7.6 pH treatment, while the response of infected individuals was again less clear and also exhibited a high degree of variability depending on the species of infecting parasite. These results suggest that *Z. subcarinatus* will be negatively affected by the reductions in seawater pH predicted to occur over the next 100–300 years (IPCC, 2014), and that the species of infecting parasite will modify this effect. Given the high degree of variability found in the oxygen consumption rates and tissue glucose content of snails infected with different species of parasite, we will first discuss the effects of acidified seawater on these parameters in uninfected snails and then address the modifications to this response caused by trematode infection.

Relative to snails maintained in control (8.1 pH) conditions, individuals exposed to 7.6 pH seawater exhibited significantly increased oxygen consumption rates and decreased tissue glucose content. These results suggest that *Z. subcarinatus* experiences higher metabolic costs at 7.6 pH due to the increased ionoregulatory activity required to maintain acid–base homeostasis and the biomineralization of calcified structures. As *Z. subcarinatus* snails are occasionally exposed to 7.6 pH seawater in their habitat (MacLeod, 2015), it is unsurprising that this species can tolerate this degree of acidification, albeit at a metabolic cost. The ability of *Z. subcarinatus* to tolerate exposure to these conditions is also supported by the positive shell growth of individuals maintained in 7.6 pH seawater (MacLeod and Poulin, 2015).

*Zeacumantus subcarinatus* is not exposed to seawater pH as low as 7.4 in its habitat, and we predicted that this degree of acidification would cause the snails to enter a state of metabolic depression, as has been found in other species of marine gastropod (Ellis *et al.* 2009; Kimura *et al.* 2011; Melatunan *et al.* 2011; Zhang *et al.* 2014). While the oxygen consumption rates of uninfected snails in the 7.4 pH treatment did decrease relative to those exposed to 7.6 pH seawater, they did not fall below the consumption rates of snails maintained in control (8.1 pH) conditions indicating that the snail did not experience metabolic depression. Similarly, tissue glucose concentration of snails in the 7.4 pH treatment increased relative to those exposed to 7.6 pH seawater, but did not

differ significantly from snails maintained at 8.1 pH. These results indicate that *Z. subcarinatus* is somewhat tolerant of 7.4 pH conditions. However, it is logical that *Z. subcarinatus* would experience some metabolic stress in the more acidic seawater, which suggests that an unknown factor or factors are modifying the metabolic processes of the snails in these conditions. As described above, uninfected snails in the 7.4 pH treatment consume more oxygen relative to those in control (8.1 pH) conditions, but have higher tissue glucose content. Glucose levels in marine gastropods can be altered by a number of factors, such as the quantity and quality of food, stress hormones, hibernation and starvation (Hochachka, 1983; Lacoste *et al.* 2001). Consequently, it is possible that the stress experienced by the snails in the 7.4 pH treatment caused glucose to be released while simultaneously depressing oxygen consumption rates. We must also consider that, despite an acclimation period, the snails in acidified treatments may have been exposed to environmental conditions out with the range found in their natural habitat.

In addition to their own metabolic requirements, infected snails must also supply parasites with sufficient energy to grow and reproduce, leading to a *de facto* increase in the total energetic costs experienced by infected snails. Figure 1A shows that in control (8.1 pH) conditions, the oxygen consumption rates of snails infected with all species of parasite are indeed greater than the consumption rates of uninfected snails, although this increase is only significant in snails infected with either *Acanthoparyphium* sp. or *Philophthalmus* sp. The tissue glucose content of infected snails, however, does not change as we would expect, i.e. reduced tissue glucose content correlates with increased rates of oxygen consumption.

At 8.1 pH, infected snails exhibit increased tissue glucose concentrations relative to uninfected individuals, although, again, this difference is only significant for two of the three infecting parasite species. The lack of correlation between oxygen and glucose in infected individuals is the first indication in these data that parasitic infection alters the concentration of free glucose independently of oxygen consumption rates. At 7.6 pH, snails infected with *M. novaezealandensis* exhibit significantly lower oxygen consumption rates and significantly higher tissue glucose content relative to uninfected individuals, re-establishing the negative relationship between these parameters for this species of parasite. The data on *Acanthoparyphium* sp.- and *Philophthalmus* sp.-infected snails, however, confirm the disconnect between oxygen and glucose, as their rates of oxygen consumption are similar to uninfected individuals while their glucose levels are significantly higher.

There are many aspects of the host–parasite association that may explain the observed differences

between the oxygen consumption and tissue glucose levels of infected and uninfected snails, and between snails infected with different species of parasite. In this paper, we limit discussion to two mechanisms which may explain the observed changes in oxygen consumption and tissue glucose concentration, the effects of parasitic castration and the oxygen uptake and nutrient absorbing capacity of the trematode life stages found within first intermediate gastropod hosts (rediae and sporocysts).

The energy requirements of gastropods are radically altered by trematode infection. Not only does infection withdraw energy from the host organism to meet the metabolic demands of parasite growth and reproduction, but also partially or completely castrates the host, dramatically reducing the energy required by the host for reproduction. This can cause an energy imbalance in which the host generates more metabolic energy than is required by the parasite, and in some cases this results in increased somatic growth rates of infected snails (Hay *et al.* 2005). However, as parasite energetics are likely to be species-specific, the energy available to host organisms may vary between individuals infected with different species of parasite. This variability in the energy available to host organisms may explain the differences in oxygen consumption and glucose concentration we observed between groups of snails infected with different species of parasite. As all parasite species used in this experiment completely sterilize host snails, differences in the metabolic status of snails infected with different species of trematode, but maintained at the same pH, may be solely due to species-specific energy requirements of the infecting parasite.

Rediae and sporocysts asexually produce cercariae, a free-swimming life stage of trematode parasites, within the reproductive organs of the first intermediate gastropod host. Despite their similar function, sporocysts and rediae are morphologically distinct: sporocysts have a simple sac-like structure, while rediae have a primitive gut, a muscular pharynx and an oral sucker (Galaktionov and Dobrovolskij, 2003). Both reproductive morphs have the ability to regulate oxygen (Vernberg, 1963; Popiel and James, 1976) and nutrient consumption rates (Richards, 1970; Richards *et al.* 1972) in response to changing environmental conditions, and absorb exogenous nutrients, such as glucose, through microvilli on their external surface (McDaniel and Dixon, 1967; Pojmanska and Machaj, 1991); rediae also use their muscular pharynx to actively consume host tissue (Cheng, 1963). Rediae and sporocysts are also capable of increasing the concentration of free glucose in host tissue by releasing a digestive enzyme that breaks down host glycogen stores into glucose (Cheng, 1963) or by inducing the host to break down its own glycogen (Cheng and Snyder, 1963). The rediae and sporocysts of many trematode

species are also facultative anaerobes, meaning they can switch between aerobic and anaerobic respiration in response to the changing internal conditions of their host (Van Hellemond *et al.* 1997).

Clearly, the oxygen absorbed by sporocysts and rediae will not be available to host snails, causing a compensatory increase in oxygen consumption by infected individuals. This is the most likely explanation for the higher oxygen consumption rates of infected snails maintained at 8.1 pH. In addition, at 8.1 and 7.6 pH, snails infected with rediae-producing parasites (*Philophthalmus* sp. and *Acanthoparyphium* sp.) exhibit increased rates of oxygen consumption relative to snails infected with sporocyst-producing parasites (*M. novaezealandensis*). The higher oxygen consumption rates of *Acanthoparyphium* sp.- and *Philophthalmus* sp.-infected snails could indicate that rediae have greater oxygen requirements than sporocysts, as the former are more active. Alternatively, increased oxygen consumption could be the metabolic cost of tissue repair in snails infected with rediae-producing parasites, due to the destructive feeding strategy of this reproductive morph.

As both rediae and sporocysts can directly or indirectly increase the concentration of glucose in host tissue, and regulate the rate at which they absorb nutrients from the host, the observed deviation from the negative relationship between oxygen and glucose in infected snails may be caused by parasite-induced breakdown of glycogen and/or species-specific nutrient requirements. If parasite-induced glycogen breakdown occurs at a greater rate than glucose consumption, the resultant nutrient accumulation could explain the elevated glucose levels recorded in infected snails.

#### Concluding remarks

The data presented here support our prediction that parasitic infection may alter the effects of OA on marine organisms. By incorporating multiple parasite species into this study, we have also shown that the species-specific effects of the infecting parasite can be pronounced and alter the apparent tolerance of host organisms to an acidified environment. *The integration of parasitology into the physiological assessment of marine organisms exposed to simulated OA conditions is urgently required.* A failure to do so may lead to the incorrect identification of some marine species as tolerant or susceptible to acidified seawater, when in fact the physiological response of the organism is modified by parasitic infection. This concern is supported by our analysis of the glucose dataset, which showed that when the infection status of the snails was excluded from the analysis, the significant effect of pH was no longer found. In the context of OA, predictions on the future states of marine ecosystems rely on data describing the relative tolerance of sympatric marine

species to acidified seawater; these predictions will be misleading if parasitic infection is not taken into consideration.

#### SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <http://dx.doi.org/10.1017/S0031182016000913>.

#### ACKNOWLEDGEMENTS

The authors would like to thank Matthew Downes and Tony Stumbo for technical assistance with the analysis of tissue glucose and two anonymous reviewers for comments on an earlier draft of this manuscript.

#### FINANCIAL SUPPORT

This research was supported by a University of Otago Doctoral Scholarship (to C.M.), funding from the Department of Zoology, University of Otago, New Zealand (to C.M. and R.P.) and the Royal Society of New Zealand Hutton Fund (to C.M.).

#### REFERENCES

- Bates, A. E., Leiterer, F., Wiedebach, M. L. and Poulin, R. (2011). Parasitized snails take the heat: a case of host manipulation? *Oecologia* **167**, 613–621.
- Bates, D., Maechler, M., Bolker, B. and Walker, S. (2014). *lme4: Linear mixed-effects models using Eigen and S4*. R package version 1.1-7. <http://CRAN.R-project.org/package=lme4>
- Berthelin, C., Kellner, K. and Mathieu, M. (2000). Histological characterization and glucose incorporation into glycogen of the Pacific oyster *Crassostrea gigas* storage cells. *Marine Biotechnology* **2**, 136–145.
- Cheng, T. C. (1963). Biochemical requirements of larval trematodes. *Annals of the New York Academy of Sciences* **113**, 289–321.
- Cheng, T. C. and Snyder, R. W. (1963). Studies on host-parasite relationships between larval trematodes and their hosts. IV. A histochemical determination of glucose and its role in the metabolism of molluscan host and parasite. *Transactions of the American Microscopical Society* **82**, 343–346.
- Coleman, D., Byrne, M. and Davis, A. (2014). Molluscs on acid: gastropod shell repair and strength in acidifying oceans. *Marine Ecology Progress Series* **509**, 203–211.
- Dickson, A. G., Sabine, C. L. and Christian, J. R. (2007). Guide to best practices for ocean CO<sub>2</sub> measurements. *PICES Special Publication 3* **191**, 1–176.
- Ellis, R., Bersey, J., Rundle, S., Hall-Spencer, J. and Spicer, J. (2009). Subtle but significant effects of CO<sub>2</sub> acidified seawater on embryos of the intertidal snail, *Littorina obtusata*. *Aquatic Biology* **5**, 41–48.
- Fox, J., Weisburg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., Firth, D., Friendly, M., Gorjanc, G., Graves, S., Heiburger, R., Laboissiere, R., Monette, G., Murdoch, D., Nilsson, H., Ripley, B., Venables, W. and Zeileis, A. (2014). *Companion to applied regression*. <https://r-forge.r-project.org/projects/car/>
- Fredensborg, B. L., Mouritsen, K. N. and Poulin, R. (2005). Impact of trematodes on host survival and population density in the intertidal gastropod *Zeacumantus subcarinatus*. *Marine Ecology Progress Series* **290**, 109–117.
- Fried, B. and Graczyk, T. K. (1997). *Advances in Trematode Biology*. CRC Press, Boca Raton, FL.
- Galaktionov, K. V. and Dobrovolskij, A. A. (2003). *The Biology of Trematodes*. Kluwer, Dordrecht.
- Graham, A. L. (2003). Effects of snail size and age on the prevalence and intensity of avian schistosome infection: relating laboratory to field studies. *Journal of Parasitology* **89**, 458–463. doi: [http://dx.doi.org/10.1645/0022-3395\(2003\)089\[0458:EOSSAA\]2.0.CO;2](http://dx.doi.org/10.1645/0022-3395(2003)089[0458:EOSSAA]2.0.CO;2)
- Hay, K. B., Fredensborg, B. L. and Poulin, R. (2005). Trematode-induced alterations in shell shape of the mud snail *Zeacumantus subcarinatus* (Prosobranchia: Batillariidae). *Journal of the Marine Biological Association of the United Kingdom* **85**, 989–992. doi: <http://dx.doi.org/10.1017/S0025315405012002>
- Hochachka, P. W. (1983). *Mollusca: Metabolic Biochemistry and Molecular Biomechanics*. Academic Press, London.
- Hunter, K. A. (2007). *SWCO<sub>2</sub> Seawater CO<sub>2</sub> Equilibrium Calculations*, University of Otago, New Zealand. [http://neon.otago.ac.nz/research/mfc/people/keith\\_hunter/software/swco2/](http://neon.otago.ac.nz/research/mfc/people/keith_hunter/software/swco2/)
- IPCC (2014). Summary for policymakers. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R. and White, L. L.), pp. 1–32. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Kelly, M. W. and Hofmann, G. E. (2013). Adaptation and the physiology of ocean acidification. *Functional Ecology* **27**, 980–990.
- Kimura, R., Takami, H., Ono, T., Onitsuka, T. and Nojiri, Y. (2011). Effects of elevated pCO<sub>2</sub> on the early development of the commercially important gastropod, Ezo abalone *Haliotis discus hannai*: effects of high pCO<sub>2</sub> on larval Ezo abalone. *Fisheries Oceanography* **20**, 357–366.
- Lacoste, A., Jalabert, F., Malham, S. K., Cuffe, A. and Poulet, S. A. (2001). Stress and stress-induced neuroendocrine changes increase the susceptibility of juvenile oysters (*Crassostrea gigas*) to *Vibrio splendidus*. *Applied and Environmental Microbiology* **67**, 2304–2309.
- Lardies, M. A., Arias, M. B., Poupin, M. J., Manriquez, P. H., Torres, R., Vargas, C. A., Navarro, J. M. and Lagos, N. A. (2014). Differential response to ocean acidification in physiological traits of *Concholepas concholepas* populations. *Journal of Sea Research* **90**, 127–134.
- Leung, T. L. F., Donald, K. M., Keeney, D. B., Koehler, A. V., Peoples, R. C. and Poulin, R. (2009). Trematode parasites of Otago Harbour (New Zealand) soft-sediment intertidal ecosystems: life cycles, ecological roles and DNA barcodes. *New Zealand Journal of Marine and Freshwater Research* **43**, 857–865.
- MacLeod, C. D. (2015). *The effects of ocean acidification on host-parasite associations*. PhD thesis. University of Otago, New Zealand.
- MacLeod, C. D. and Poulin, R. (2015). Interactive effects of parasitic infection and ocean acidification on the calcification of a marine gastropod. *Marine Ecology – Progress Series* **537**, 137–150.
- MacLeod, C. D., Doyle, H. L. and Currie, K. I. (2015). Technical note: maximising accuracy and minimising cost of a potentiometrically regulated ocean acidification simulation system. *Biogeosciences* **12**, 713–721.
- Macnab, V. and Barber, I. (2012). Some (worms) like it hot: fish parasites grow faster in warmer water, and alter host thermal preferences. *Global Change Biology* **18**, 1540–1548.
- Martínez-Quintana, J. A. and Yepiz-Plascencia, G. (2012). Glucose and other hexoses transporters in marine invertebrates: a mini review. *Electronic Journal of Biotechnology* **15**, 1–12.
- Martorelli, S. R., Fredensborg, B. L., Mouritsen, K. N. and Poulin, R. (2004). Description and proposed life cycle of *Maritrema novaezealandensis* n. sp. (Microphallidae) parasitic in red-billed gulls, *Larus novaehollandiae* scopulinus, from Otago Harbor, South Island, New Zealand. *Journal of Parasitology* **90**, 272–277.
- Martorelli, S. R., Fredensborg, B. L., Leung, T. L. F. and Poulin, R. (2008). Four trematode cercariae from the New Zealand intertidal snail *Zeacumantus subcarinatus* (Batillariidae). *New Zealand Journal of Zoology* **35**, 73–84.
- McDaniel, J. S. and Dixon, K. E. (1967). Utilization of exogenous glucose by the rediae of *Parorchis acanthus* (Digenea: Philophthalmidae) and *Cryptocotyle lingua* (Digenea: Heterophyidae). *Biological Bulletin* **133**, 591–599.
- Melatanun, S., Calosi, P., Rundle, S. D., Moody, A. J. and Widdicombe, S. (2011). Exposure to elevated temperature and pCO<sub>2</sub> reduces respiration rate and energy status in the periwinkle *Littorina littorea*. *Physiological and Biochemical Zoology* **84**, 583–594.
- Pan, T.-C. F., Applebaum, S. L. and Manahan, D. T. (2015). Experimental ocean acidification alters the allocation of metabolic energy. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 4696–4701.
- Parker, L., Ross, P., O'Connor, W., Pörtner, H., Scanes, E. and Wright, J. (2013). Predicting the response of molluscs to the impact of ocean acidification. *Biology* **2**, 651–692.
- Pojmanska, T. and Machaj, K. (1991). Differentiation of the ultrastructure of the body wall of the sporocyst of *Leucochloridium paradoxum*. *International Journal for Parasitology* **21**, 651–659.
- Popiel, I. and James, B. L. (1976). The effect of glycogen and glucose on oxygen consumption in the daughter sporocysts of *Cercaria linearis* stunkard, 1932 and *Cercaria stunkardi* palombi, 1934 (Digenea: Opecoelidae). *Zeitschrift für Parasitenkunde* **51**, 71–77.

- Pörtner, H.** (2008). Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Marine Ecology Progress Series* **373**, 203–217.
- Pörtner, H. O. and Farrell, A. P.** (2008). Physiology and climate change. *Science* **322**, 690–692.
- Pörtner, H. O., Bock, C. and Reipschlager, A.** (2000). Modulation of the cost of pHi regulation during metabolic depression: a (31) P-NMR study in invertebrate (*Sipunculus nudus*) isolated muscle. *Journal of Experimental Biology* **203**, 2417–2428.
- R Development Core Team** (2014). *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria, ISBN 3-900051-07-0.
- Richards, R. J.** (1970). Variations in the oxygen uptake, reduced weight and metabolic rate of starving sporocysts of *Microphallus pygmaeus* (Levinsen, 1881)(Trematoda: Microphallidae). *Journal of Helminthology* **44**, 75–88.
- Richards, R. J., Pascoe, D. and James, B. L.** (1972). Variations in the metabolism of the daughter sporocysts of *Microphallus pygmaeus* in a chemically defined medium. *Journal of Helminthology* **46**, 107–116.
- Sorensen, R. E. and Minchella, D. J.** (2001). Snail–trematode life history interactions: past trends and future directions. *Parasitology* **123**, S3–S18.
- Toledo, R. and Fried, B.** eds. (2011). *Biomphalaria Snails and Larval Trematodes*. Springer New York, New York, NY.
- Van Hellemond, J. J., Van Remoortere, A. and Tielens, A. G. M.** (1997). *Schistosoma mansoni* sporocysts contain rhodoquinone and produce succinate by fumarate reduction. *Parasitology* **115**, 177–182.
- Vernberg, W. B.** (1963). Respiration of digenetic trematodes. *Annals of the New York Academy of Sciences* **113**, 261–271.
- Zhang, H., Cheung, S. G. and Shin, P. K. S.** (2014). The larvae of congeneric gastropods showed differential responses to the combined effects of ocean acidification, temperature and salinity. *Marine Pollution Bulletin* **79**, 39–46.