

Frozen–thawed ampullary cell monolayer improves bovine embryo *in vitro* development and quality

Research Article

Cite this article: Asaadi A *et al.* (2019) Frozen–thawed ampullary cell monolayer improves bovine embryo *in vitro* development and quality. *Zygote* 27: 337–346. doi: [10.1017/S0967199419000388](https://doi.org/10.1017/S0967199419000388)

Received: 22 November 2018
Revised: 28 April 2019
Accepted: 11 June 2019
First published online: 13 August 2019

Keywords:

Ampulla; Embryo production; Gene expression; Oocyte maturation

Address for correspondence:

Mojtaba Kafi, Department of Animal Reproduction, School of Veterinary Medicine, Shiraz University, Shiraz, Iran.
E-mail: kafi@shirazu.ac.ir

Anise Asaadi¹, Mojtaba Kafi¹ , Hadi Atashi², Mehdi Azari¹ and Miel Hostens³

¹Department of Animal Reproduction, School of Veterinary Medicine, Shiraz University, Shiraz, Iran; ²Department of Animal Science, School of Agriculture, Shiraz University, Shiraz, Iran and ³Department of Reproduction, Obstetrics and Herd Health, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium

Summary

The aim of this study was to evaluate the effects of different timing for frozen–thawed bovine ampullary epithelial cell (BAEC) and bovine oviductal epithelial cell (BOEC) co-culture on the development and quality of bovine embryos produced *in vitro*. Embryo development was assessed by day 8 blastocyst yield, whereas embryo quality was determined using blastocyst differential cell count, cryotolerance and the expression of selected genes related to embryo quality. The results showed that the presence of BAECs during the last 6 h of *in vitro* maturation (IVM) increased blastocyst yield and survival of the vitrified–warmed blastocysts. In addition, embryos produced in the presence of BAECs during the last 6 h of IVM or in the presence of BOECs during the first 4 days of *in vitro* culture (IVC) showed a greater number of trophectoderm cells and a greater inner cell mass. In terms of gene expression, *IFN-T* was downregulated and *PLAC8*, *AQP3* and *ATPIA1* were upregulated in the presence of the BAECs during the last 6 h of the IVM and/or in the presence of BOECs during the first 4 days of IVC. In conclusion, co-culturing bovine oocytes with a frozen–thawed ampullary cell monolayer during the last 6 h of maturation increased blastocyst yield and quality.

Introduction

In vitro embryo production (IVP) is a reproductive technology used to increase the number of offspring from genetically elite animals and to study early embryo development in different species of mammals (Lopera-Vásquez *et al.*, 2016). Although researchers have had considerable success in increasing IVP efficiency, the quality of the produced embryos remains lower than those produced *in vivo* (Lonergan and Fair, 2008; Rizos *et al.*, 2008; Pontes *et al.*, 2009). Numerous studies have mimicked *in vivo* conditions with the aim of discovering the optimum *in vitro* embryo culture conditions. The oviduct is a convoluted tube that has an epithelial layer lined with a simple cuboidal-to-columnar epithelium covered by both ciliated and secretory cells (Steinhauer *et al.*, 2004; Ferraz *et al.*, 2017). The oviduct's secretory epithelial cells produce the proteins and microelements needed for normal gamete and embryo development (Rodríguez-Martínez, 2007; Mondéjar *et al.*, 2012; Cordova *et al.*, 2014; Schmaltz-Panneau *et al.*, 2014). The oviduct also acts as an environment for sperm capacitation (Rodríguez-Martínez, 2007), sperm storage, acrosome reaction and reducing the possibility of polyspermy (Ghersevich *et al.*, 2015).

When a follicle ruptures, its cumulus–oocyte complex (COC) ovulates into the infundibulum part of the oviduct. The role of the oviduct in transporting the oocyte to the fertilization site has been attributed to cilia beating and smooth muscle activity, which oestrogen can increase and which progesterone can decrease (Coy *et al.*, 2012). When the COC is transported into the ampulla, it remains attached to the epithelium for a variable amount of time depending on the species. Digital video microscope studies have shown that unfertilized porcine and bovine oocytes reached the ampullary–isthmus region within 30–45 min and 3–6 h post ovulation, respectively (Hunter, 1974; Tanghe *et al.*, 2002; Coy *et al.*, 2012). *In vitro* exposure of oocytes to oviductal cells or oviductal fluid increased sperm–zona binding, zona penetration and fertilization rates in mares, cows and pigs (Mugnier *et al.*, 2009; Coy *et al.*, 2010). In equine, oocyte exposure to osteopontin, a protein secreted by oviductal epithelium cells, increased *in vitro* blastocyst yield (Mugnier *et al.*, 2009). In mice, pre-treatment of COCs with *S100A11*, a protein detected at the oestrous stage of the oestrous cycle in the ampullary region of the oviductal epithelium, increased *in vitro* blastocyst yield (Hanaue *et al.*, 2011). Secretions of different parts of the oviduct at different stages of the oestrous cycle have various effects on gamete function and embryo development (Way *et al.*, 1997; Lopera-Vásquez *et al.*, 2017). Embryo mRNA synthesis starts in the oviduct at the 8- to 16-cell stage (Graf *et al.*, 2014). During early embryo development in the oviduct, epigenetic modifications including DNA methylation or histone changes are associated with embryonic implantation, maintenance of pregnancy and the health of the

progeny (Fazeli and Pewsey, 2008; Gad *et al.*, 2012). As *in vitro* culture conditions have not been completely optimized for normal early embryo development, many researchers have attempted to improve the quality of *in vitro* embryos by co-culturing gametes and embryos with oviductal epithelial cells (Kattal *et al.*, 2008; Mugnier *et al.*, 2009). Final *in vivo* maturation of gametes and early embryo development occurs in the oviduct, therefore there is a vital need to obtain deeper knowledge of the oviduct microenvironment that would help researchers and practitioners to improve the efficiency of assisted reproduction techniques (Avilés *et al.*, 2015). Fresh co-culture has been shown to be methodologically complex, lack of reproducibility and have biosanitary risks (Lopera-Vásquez *et al.*, 2016). One alternative to decrease the variability of co-culture systems could be to use stored frozen oviductal epithelial cells (Lopera-Vásquez *et al.*, 2016). The aim of this study was to evaluate the effects of different timing of frozen-thawed bovine ampullary epithelial cell (BAECs) and bovine oviductal epithelial cell (BOECs) co-cultures on the development and quality of bovine embryos produced *in vitro*. Embryo development was assessed as day 8 blastocyst yield, and embryo quality was evaluated as blastocyst differential cell number, cryotolerance and expression of selected genes related to implantation (*PLAC8* and *IFN-T*), apoptosis (*BAX* and *TP53*) and cryotolerance (*AQP3* and *ATP1A1*).

Materials and Methods

All the chemicals used in the study were purchased from Sigma (St. Louis, MO, USA) unless otherwise stated.

BOEC isolation and freezing

Oviducts and ovaries from slaughtered cows at the early luteal phase of the oestrus cycle (based on corpus luteum morphology) were transported to the laboratory on ice. The connective tissue around the oviducts was removed, and oviducts were washed twice in normal saline solution and 70% ethanol solution. Then, oviducts were trimmed completely, washed with phosphate-buffered saline (PBS) containing gentamicin and kept in TCM-199 medium (Sigma, m4530) until further utilization. Next, the oviducts were divided into two groups: the first group was kept intact, while in the second group, the ampulla of the oviduct was separated at the ampullary-isthmus junction (identified by where oviduct diameter first exhibited a marked reduction in size). The mucosa of the oviductal region (complete oviduct or the ampulla) was expelled by squeezing with a sterile glass slide. Then, the oviductal and ampullary epithelial cells were collected and transferred to washing medium (TCM-199 supplemented with gentamicin 40 µg/ml; Sigma G1272) and amphotericin B (1 µl/ml; Biowest, L0009-100). Oviductal epithelial cells were washed twice by centrifuging at 400 g for 5 min in the washing medium. Next, trypsin-EDTA enzyme was added to the samples for around 3 min at 37°C to obtain single cells. The enzyme was neutralized using a washing medium supplemented with 10% fetal bovine serum (FBS) (Gibco, 10270). Retrieved fresh epithelial single cells were counted using a haemocytometer and methylene blue staining. The quality of the fresh oviductal epithelial cells was evaluated by seeding 1×10^6 cells/ml in the culture medium (TCM-199, 15% fetal calf serum (FCS) and 40 µg/ml gentamicin). The remaining fresh cells (1×10^7 cell/ml concentration) were frozen in FCS-containing cryoprotectant (10% dimethyl sulfoxide (DMSO)) using a slow freezing procedure described by Freshney (2015) with slight modifications. Briefly, the cells with cryoprotectant were placed in a -20°C freezer for around 4 h, then transferred to a -80°C freezer for

at least one night. The frozen cells were stored in liquid nitrogen until cell culturing. All samples of fresh oviductal cells were cultured to examine the quality of their monolayers. Only those cells with an acceptable quality of monolayer were kept frozen in the cell bank for subsequent oocyte and embryo co-culturing.

BOEC thawing and seeding

At 5 days before using the cell monolayer in the embryo culture, frozen cells obtained from a mixture of oviducts and belonging to one pooled culture were thawed in 37°C sterile water, added to culture medium (TCM-199, 15% FCS and 40 µg/ml gentamicin) and centrifuged at 400 g for 5 min to remove DMSO. Next, the cells were re-suspended in warm culture medium and seeded into four-well NUNC dishes at a concentration of 1×10^6 cell/ml. Medium was refreshed every 48 h during cell culture. At 1 day before co-culture of oocytes or embryos with epithelial cells, the culture medium was replaced with either maturation medium (for co-culture of oocytes with ampullary epithelial cells) or synthetic oviductal fluid (SOF) medium (for co-culture of embryos with whole oviductal epithelial cells).

Comparing the monolayer of fresh and frozen-thawed BOEC

Consistency of growth of the fresh and frozen-thawed BOECs was evaluated by culturing both types of BOECs under similar conditions. Cells were monitored using an inverted phase-contrast light microscope (Hund Wetzlar Wilovert 30, Germany) on days 1, 3, 5 and 7 to ensure that they were healthy and growing as expected.

In vitro embryo production

Oocyte collection and *in vitro* maturation

Bovine ovaries were transported from the local slaughterhouse (Shiraz, Iran) to the laboratory in saline solution kept at 33–35°C. Immature COCs were collected by aspirating 2–8 mm diameter follicles with a syringe and 18G needle. Class I and class II COCs were selected, washed in warm equilibrated wash medium (HEPES-buffered TCM-199 (M4530), 10% FCS and 50 µg/ml gentamicin (G1272)) three times (Lopera-Vásquez *et al.*, 2016) and then divided into four groups of 40–45 COCs. COCs were matured in a 500 µl maturation medium with or without bovine ampullary oviductal epithelial cell (BAOEC) monolayers under a 90% humidified atmosphere of 5.5% CO₂ in air at 38.5°C in four-well culture dishes (NUNC, Denmark) for 22–24 h. The oocyte maturation medium contained TCM-199 supplemented with 10% FCS, 5 IU/ml of highly purified human chorionic gonadotropin (hCG) (Karma, Pharmatech GmbH, Germany), 0.1 IU/ml of recombinant human follicle-stimulating hormone (FSH) (Follitrope, LG Life Sciences, South Korea) and 10 ng/ml of epidermal growth factor (E4127).

Sperm preparation and *in vitro* fertilization

Sperm separation and selection was performed using the swim-up procedure described by Brackett and Oliphant (1975). Briefly, frozen-thawed semen straws from the same bull with proven field fertility were placed at the bottom of a 15-ml Falcon tube and incubated in warm sperm capacitation medium (modified calcium-free Tyrode's medium supplemented with 1.12 mg/ml Na lactate, 2 mg/ml Na bicarbonate, 0.1 mg/ml caffeine, 1 mg/ml Na pyruvate, 50 µg/ml gentamicin and 6 mg/ml BSA) for 40–50 min. Next, the upper medium was collected and centrifuged at 200 g for 10 min. The sperm pellet was isolated and re-suspended in warm fertilization medium (modified Tyrode's medium supplemented with 1.12

mg/ml Na lactate, 2.2 mg/ml Na bicarbonate, 0.27 mg/ml caffeine, 1 mg/ml Na pyruvate, 10 µg/µl heparin, 50 µg/µl hypotaurine, 50 µg/ml gentamicin and 6 mg/ml BSA). Matured COCs were washed three times in fertilization medium and transferred into four-well dishes containing 450 µl of IVF medium. Sperm concentration was determined, and a concentration of 1×10^6 sperm/ml was used for *in vitro* insemination (Azari *et al.*, 2017).

In vitro culture of the presumptive zygote

Presumptive zygotes were denuded around 18–20 h post insemination (p.i.) by repeated pipetting. Cumulus cells and excess spermatozoa were deleted by washing in modified SOF (mSOF). Next, the cleaned zygotes were placed into four-well dishes (NUNC, Denmark) containing SOF medium supplemented with 5% bovine calf serum (BCS), 4 mg/ml fatty acid-free BSA (Sigma, A3675) and 50 µg/ml gentamicin with or without a BOECs monolayer. The culture medium was refreshed every 48 h, and cultures were incubated under 90% humidified conditions at 38.5°C in an atmosphere of 5% CO₂ in air.

Assessment of embryo development and quality

Embryo development

Embryo development was determined by cleavage rate (defined as the total number of cleaved zygotes/total number of fertilized oocytes) and blastocyst yield (defined as the total number of blastocysts/total number of cleaved zygotes). Cleavage rate was recorded on day 2 (48 h p.i.), and the blastocyst yield was recorded on days 7 and 8 (p.i. under a stereomicroscope).

Embryo quality

Quality of the expanded blastocysts at days 7 and 8 p.i. was evaluated by cryotolerance assessment, differential cell count and expression analysis of the selected genes.

Blastocyst vitrification. Blastocyst cryotolerance was assessed by vitrification-warming of early- and mid-stage blastocysts using the procedure described by Rizos *et al.* (2002). Blastocyst vitrification was performed in two steps: first, the blastocysts were washed in a holding medium (HM) (TCM-199–M4530 supplemented with 20% (v/v) FCS). Blastocysts were then transferred to HM1 medium (HM containing 7.5% ethylene glycol and 7.5% DMSO) for about 3 min. Second, the blastocysts were transferred to HM2 medium (HM supplemented with 16.5% ethylene glycol, 16.5% DMSO and 0.5 M sucrose) for up to 30 s. Then, blastocysts were placed on a cryotop and quickly floated in liquid nitrogen. In the thawing process, the vitrified blastocysts were warmed twice, initially in HM medium (supplemented with 0.25 M sucrose) and then in HM medium (supplemented with 0.15 M sucrose). Next, blastocysts were cultured in SOF medium with 5% FCS in four-well NUNC dishes. The survival rate of the vitrified-warmed blastocysts was investigated as the re-expansion of the blastocoel and its maintenance for 4 h, 24 h and 48 h post culture.

Differential staining of blastocysts. Differential staining of the inner cell mass (ICM) and trophectoderm (TE) was performed using the procedure described by Thouas *et al.* (2001) with slight modifications. Briefly, the expanded blastocysts were incubated in solution I (5 mg/ml BSA–TCM-199 with 1% Triton X-100 (T8787) and 100 µg/ml propidium iodide (P4170)) for up to 30 s in the dark to stain the TE. Next, blastocysts were immediately transferred to solution II (100% ethanol with 25 µg/ml bisbenzimidazole H33342 (B2261)) and kept for 30 min in the dark to complete the fixation and the ICM staining processes. Stained blastocysts were

transferred to a glycerol droplet and mounted onto a glass microscope slide. Cell counting was carried out using a fluorescence microscope (Olympus BX61, Japan).

Gene expression analysis. Expression levels for six target genes (two apoptosis-related genes (*BAX* and *TP53*), two cryotolerance-related genes (*AQP3* and *ATPIA1*) and two implantation-related genes (*PLAC8* and *IFN-T*) along with two housekeeping genes (reference genes (*GAPDH* and *YWHAZ*)) were assessed using quantitative RT-PCR (qPCR). The names, primers (designed using PerlPrimer software; Marshall, 2004), GenBank accession numbers and the expected product sizes of the genes are presented in Table 1. Total RNA was isolated using an RNeasy Micro Kit (catalogue no. 74004, Qiagen, Hilden, Germany) according to the manufacturer's instructions. Total RNA concentrations were determined using a NanoDrop 2000 spectrophotometer (absorbance at 260 nm wavelength). All samples were treated with DNase and RQ1 DNase (Promega, France) to remove possible DNA contamination. Sample purity was assessed based on the A260/A280 nm ratio, with ratios expected to fall between 1.8 and 2.0. Immediately after RNA extraction, reverse-transcription (RT) was performed using a RevertAid H Minus First Strand cDNA Synthesis Kit (K1632, Fermentas, Germany). All mRNA transcripts for the mentioned genes were quantified using qPCR, with three replicates for each group. The PCR mix in each well contained 5 µl of Power SYBR® Green PCR Master Mix (Applied Biosystems, UK), 1 µl of 5 pM forward and reverse primers, 2 µl of cDNA (12.5 ng/ µl) and 11 µl dH₂O in a final volume of 20 µl. PCR reactions were carried out in the Applied Biosystems StepOne and StepOnePlus Real-Time PCR system using the following program: Stage 1: 95°C for 10 min. Stage 2: 95°C for 15 s and 60°C for 1 min for in total, 40 cycles. Stage 3: 95°C for 15 s, 60°C for 15 s and 95°C for 15 s. The last heating step in Stage 3 was performed with a ramp rate of 2% to enable the generation of a dissociation curve of the products. Each reaction was run in duplicate, and the mean value of each duplicate was used for the further calculations. The expression data were normalized using the geometric mean of the internal reference genes (*GAPDH* and *YWHAZ*). Then, $2^{-\Delta\text{CT}}$ (geometric mean of housekeeping genes) values were calculated for each gene.

Experimental design

A schematic diagram of the experimental design for this study is shown in Fig. 1. The study was carried out using the treatments described below.

Control group

The oocytes were placed into cell-free maturation medium for 24 h (IVM). Next, the matured oocytes were transferred into dishes containing IVF medium and capacitated sperms. At 18–20 h p.i., the presumptive zygotes were denuded and transferred to the plates containing SOF medium for 8 days consecutively (IVC).

AM group

Oocytes were cultured in cell-free maturation medium for 18 h. For the last 6 h *in vitro* maturation they were transferred to medium over a BAOECs monolayer (mimicking the *in vivo* ampulla conditions). IVF and IVC steps were performed as described for the control group.

OV group

IVM and IVF steps were performed as described for the control group. The denuded presumptive zygotes were placed in NUNC

Table 1. Details of primers used for qRT-PCR

Gene		Primer sequences (5'→3')	GenBank accession no.	Product size (bp)
BAX	Forward	TTTGCTTCAGGGTTTCATCCAG;	NM_173894.1	146
	Reverse	TGTTACTGTCCAATTCATCTCCGA		
TP53	Forward;	TCCACAGCCAAGAAGAAACCAC	NM_174201.2	125
	Reverse	TTCCATCCAGAGCATCCTTCAG		
AQP3	Forward;	CTCTGGACACTTGACATGG	NM_001079794.1	177
	Reverse	GAGTTGAAGCCATTGAGGT		
ATP1A1	Forward;	GCTTCCAGTTTGACACCGAC	NM_001076798.1	148
	Reverse	GTGACCATGATGACCTTGATTCC		
PLAC8	Forward;	GTGGTCATTGTAACCTCAGCCT	NM_001076987.2	194
	Reverse	CTCATTGCGACACTTGTTC		
IFNT2	Forward;	GATAATGAGTACCGTCTTCCC	NM_001015511.3	198
	Reverse	AGGTAACAACCCAGAGATCG		
GAPDH	Forward;	TTCAACGGCAGTCAAGG	NM_001034034.2	119
	Reverse	ACATACTCAGCACCAGCATCA		
YWHAZ	Forward;	GCTGGTGATGACAAGAAAGG	NM_174814.2	104
	Reverse	AGTCTGATAGGATGTGTTGGT		




Figure 1. Control: IVM, IVF, and IVC was performed using a cell-free culture medium. AM: IVM was performed using TCM-199 for 18 h and then transferred into a medium containing a monolayer of frozen-thawed bovine ampullary cells for the remaining 6 h of culture. OV: the presumptive zygotes were co-cultured using a monolayer of bovine frozen-thawed oviductal cells for 4 days and then with a cell-free medium for the last 4 days of culture. AMOV: oocytes were cultured similarly to those in the AM group, and the presumptive zygotes were co-cultured similarly to those in the OV group.

plates containing a bovine total oviductal epithelial cell (BTOECs) monolayer for 4 days p.i. Then, embryos were transferred to wells containing cell-free medium for the last 4 days of culture (days 5–8 p.i.).

AMOV group

Oocytes were cultured in maturation medium for 18 h and then transferred to medium containing a BAECs monolayer for the last 6 h of *in vitro* maturation. Next, the matured oocytes were

transferred to dishes containing *in vitro* fertilization medium and the capacitated sperms. The presumptive zygotes were denuded and placed into four-well NUNC dishes containing a BTOEC monolayer for 4 days p.i. and transferred into wells containing cell-free medium for the last 4 days of culture (days 5–8 p.i.). The culture media were refreshed every 48 h for all treatments.

The overall cleavage rate was recorded at 48 h p.i.; blastocyst development was recorded on days 7 and 8 p.i. A representative number of day 7–8 blastocysts from each group (treatment) were

Table 2. Effect of the experimental treatments on the *in vitro* embryo development ($n = 8$ replicates)

Treatments ¹	Oocytes fertilized (n)	Cleavage rate ²	Blastocyst yield ³
Control	340	271 (79.62 ± 0.83)	90 (33.09 ± 0.95) ^b
AM	339	275 (81.09 ± 0.84)	108 (38.90 ± 2.61) ^a
OV	354	280 (79.10 ± 0.80)	68 (24.21 ± 1.18) ^c
AMOV	349	281 (80.52 ± 0.74)	74 (26.30 ± 1.66) ^c

¹Control: IVM, IVF, and IVC was performed using a cell-free culture medium. AM: IVM was performed using TCM-199 for 18 h and then transferred into a medium containing a monolayer of frozen-thawed bovine ampullary cells for the remaining 6 h of culture. OV: the presumptive zygotes were co-cultured using a monolayer of bovine frozen-thawed oviductal cells for 4 days and then with a cell-free medium for the last 4 days of culture. AMOV: oocytes were cultured similarly to those in the AM group, and the presumptive zygotes were co-cultured similarly to those in the OV group (mean ± standard error of the mean (SEM)).

²Number of cleaved zygotes on day 2 out of total number of fertilized oocytes.

³Number of blastocysts (day 8) out of number of cleaved zygotes.

^{a,b,c}Values in the same column with different superscripts differ significantly ($P < 0.05$).




Figure 2. BOEC morphology. Fresh BOECs monolayer on day 3 (a) and day 5 (b). Frozen-thawed BOECs monolayer on day 3 (c) and day 5 (d); $\times 40$ magnification.

either vitrified-warmed for the survival rate analysis at 4 h, 24 h and 48 h post warming, fixed for the differential cell count or frozen in liquid nitrogen and stored at -80°C for gene expression analysis. Eight replicates were generated for each treatment.

Statistical analysis

Data on the cleavage rate, blastocyst yield, survival rate of the vitrified-warmed blastocysts, embryo cell number (ICM, TE, ICM/TE ratio) and relative mRNA abundance were analyzed using one-way analysis of variance (ANOVA) in R (R Core Team, 2017). The assumptions of normality and homogeneity of the residual variances were checked at P -value < 0.05 using Shapiro-Wilk and Bartlett tests, respectively. Differences were considered to be statistically significant at $P < 0.05$.

Results

Comparing the monolayer of fresh versus frozen-thawed BOECs

Fresh and frozen-thawed BOEC monolayers showed similar morphology and growing patterns on days 3 and 5 of culture (Fig. 2).

Embryo development

Effects of co-culture conditions on the cleavage rate and blastocyst yield are presented in Table 2. Although various co-culture conditions had no effect on cleavage rate ($P \geq 0.05$), blastocyst yield was affected by the treatments ($P < 0.05$). The highest blastocyst yield was found in the AM group, followed by the control group.

Table 3. Effect of experimental treatments on the survival and 48-h post-culture hatch rate of the vitrified-warmed day 7–8 blastocysts

Treatments ¹	n ²	4-h	24-h	48-h	48-h hatch rate
Control	55	46 (83.73 ± 1.5)	29 (52.57 ± 2.58) ^b	21 (38.30 ± 3.08) ^b	13 (23.16 ± 3.92) ^b
AM	66	59 (89.14 ± 1.53)	50 (74.38 ± 2.98) ^a	42 (62.41 ± 2.81) ^a	33 (48.76 ± 3.29) ^a
OV	36	31 (85.62 ± 1.22)	21 (56.86 ± 3.65) ^b	16 (43.33 ± 4.66) ^b	10 (27.43 ± 2.83) ^b
AMOV	41	36 (88.00 ± 3.71)	30 (73.37 ± 3.69) ^a	25 (60.74 ± 3.83) ^a	19 (45.24 ± 3.73) ^a

¹Control: IVM, IVF, and IVC was performed using a cell-free culture medium. AM: IVM was performed using TCM-199 for 18 h and then transferred into a medium containing a monolayer of frozen–thawed bovine ampullary cells for the remaining 6 h of culture. OV: the presumptive zygotes were co-cultured using a monolayer of bovine frozen–thawed oviductal cells for 4 days and then with a cell-free medium for the last 4 days of culture. AMOV: oocytes were cultured similarly to those in the AM group, and the presumptive zygotes were co-cultured similarly to those in the OV group (mean ± standard error of the mean (SEM)).

²Total number of vitrified-warmed day 8 blastocysts.

^{a,b}Values in the same column with different superscripts differ significantly ($P < 0.05$).

Table 4. Effect of the experimental treatments on the differential cell count measures of day 8 blastocysts

Treatments ¹	n ²	ICM ³	TE ⁴	TOT ⁵	ICM/TOT	TE/TOT
Control	27	35.01 (±0.53) ^b	82.54 (±1.65) ^b	117.55 (±2.01) ^b	29.85 (±0.35) ^b	70.14 (±0.35) ^a
AM	35	46.03 (±1.00) ^a	98.38 (±2.48) ^a	144.42 (±3.39) ^a	31.95 (±0.26) ^a	68.05 (±0.26) ^b
OV	24	47.08 (±0.65) ^a	98.25 (±1.78) ^a	145.33 (±2.29) ^a	32.45 (±0.27) ^a	67.55 (±0.28) ^b
AMOV	25	46.30 (±0.71) ^a	97.96 (±1.84) ^a	144.26 (±2.19) ^a	32.15 (±0.31) ^a	67.58 (±0.31) ^b

¹Control: IVM, IVF, and IVC was performed using a cell-free culture medium. AM: IVM was performed using TCM-199 for 18 h and then transferred into a medium containing a monolayer of frozen–thawed bovine ampullary cells for the remaining 6 h of culture. OV: the presumptive zygotes were co-cultured using a monolayer of bovine frozen–thawed oviductal cells for 4 days and then with a cell-free medium for the last 4 days of culture. AMOV: oocytes were cultured similarly to those in the AM group, and the presumptive zygotes were co-cultured similarly to those in the OV group (mean ± standard error of the mean (SEM)).

²Total number of day 8 blastocysts fixed for the differential cell count.

³Inner cell mass.

⁴Trophectoderm.

⁵Inner cell mass plus trophectoderm.

^{a,b}Values in the same column with different superscripts differ significantly ($P < 0.05$).

However, no difference was found between the OV and AMOV groups (Table 2).

Blastocyst cryotolerance

The effects of co-culture conditions on the survival and hatching rates of vitrified-warmed blastocysts are presented in Table 3. The 4 h post-warming survival of vitrified-warmed blastocysts was not affected by the co-culture conditions ($P \geq 0.05$), whereas co-culture conditions for the AM and AMOV groups resulted in a higher survival rate at 24 and 48 h post warming, as well as a higher 48 h post-warming hatching rate than for the OV and control treatments ($P < 0.05$).

Differential cell count

The effects of experimental treatments on differential cell counts are presented in Table 4. The lowest average number of ICM, TE and total blastomere cells was found in the control group, whereas there was no difference in cell count when using various co-culture conditions in the AM, AMOV and OV groups (Table 4).

Gene expression analysis

Effects of the experimental treatments on the expression levels for six target genes (two apoptosis-related genes (*BAX* and *TP53*), two cryotolerance-related genes (*AQP3* and *ATPIA1*) and two implantation-related genes (*PLAC8* and *IFN-T*)) are presented in Fig. 3. *BAX* was significantly downregulated in the OV group compared with the rest of the groups ($P < 0.05$). The *P53* gene

showed no difference in expression among groups ($P \geq 0.05$). Expression of the *IFN-T* gene was downregulated in all co-culture conditions compared with the control group; its relative expression in the OV group was lower than that in the AM and AMOV groups ($P < 0.05$). Highest expression levels for *PLAC8* and *AQP3* were found in the AMOV group, followed by the AM group; there was no difference in expression between the control and OV groups ($P \geq 0.05$). The *ATPIA1* gene showed the highest expression level for samples in the AMOV group, followed by the OV group; there was no difference between expression levels in the control and AM groups ($P \geq 0.05$).

Discussion

Fresh co-culture is associated with methodological complexity, lack of reproducibility and biosanitary risks (Lopera-Vásquez et al., 2016). This study optimized a procedure to provide a cell bank of fresh BOECs. In this procedure, fresh bovine oviductal epithelial single cells were frozen and successfully thawed, a process that can improve the reproducibility and time efficiency of cell culture. Variation may exist among fresh BOEC lines recovered from the oviduct of different cows. However, the use of frozen–thawed BOECs makes it possible to find the most appropriate cell line for BOECs and use it to consistently produce bovine blastocysts with less laboratory complications and in a more predictable co-culture system. In addition, using the same pooled frozen–thawed oviductal epithelial cells and different experimental replicates can increase reproducibility. To the best of our knowledge, this study is the first to document that a monolayer of frozen–thawed




Figure 3. Relative mRNA expression levels from apoptosis genes (*BAX* and *P53*), implantation genes (*INF-T* and *PLAC8*), and cryotolerance genes (*ATP1A1* and *AQP3*) in day 8 blastocysts cultured in different co-culturing treatments. Data are relative to the geometric mean of internal genes *GAPDH* and *YWHAZ*. ^{a,b,c}Different letters indicate significant differences for each gene ($P < 0.05$). Control: IVM, IVF, and IVC was performed using cell-free culture medium. AM: IVM was performed using TCM-199 for 18 h and then transferred into a medium containing monolayer of frozen-thawed bovine ampullary cells for the remaining 6 h of culture. OV: the presumptive zygotes were co-cultured using monolayer of bovine frozen-thawed oviductal cells for 4 days and then with cell-free medium for the last 4 days of culture. AMOV: The oocytes were cultured similarly to those in the AM group, and the presumptive zygotes were co-cultured similarly to those in the OV group (mean \pm standard error of the mean (SEM)).

BAECs can be used as an alternative method for *in vitro* maturation of bovine oocytes.

Although there has been considerable success in increasing the efficiency of IVP, the quality of the produced embryos still remains lower than that of embryos produced *in vivo* (Lonergan and Fair, 2008; Rizos *et al.*, 2008; Pontes *et al.*, 2009). This situation illustrates that researchers must attempt to create an *in vitro* environment that is relatively similar to conditions during final maturation of the oocyte in the oviduct. Evidence suggests that the quality of the oocyte at the start of IVM is a vital determinant of the proportion of oocytes that will reach the blastocyst stage (Lonergan and Fair, 2016). By mimicking *in vivo* conditions, a better quality of matured oocyte can be obtained, and this can have a positive effect on developmental potential of early embryos throughout IVP (Avilés *et al.*, 2015). Under *in vivo* conditions, bovine COCs are situated in the ampulla part of the oviduct for the first few hours after ovulation to complete the maturation process. The oviduct provides nutrients and enzymes that have anti-oxidant effects, and that are useful for oocyte survival (Leese *et al.*, 2007; Avilés *et al.*, 2010). The main aim of this study was to mimic these *in vivo* conditions to improve the COC maturation process in an IVM procedure. The most important finding of this study was that blastocyst yield and the quality of the produced blastocysts improved when the last

6 h of *in vitro* oocyte maturation was carried out in the presence of a frozen-thawed BAOECs monolayer. Previous researchers have reported that the presence of oviductal cells or fluid from cows in the non-luteal phase of the oestrous cycle had a positive effect on *in vitro* sperm-egg binding and fertilization rates (Lopera-Vásquez *et al.*, 2017). When the oocyte moves forward in the ampulla, it becomes exposed to different proteins and factors such as oviduct-specific glycoprotein 1 and oviductine, which are needed for sperm-zona binding and zona hardening, respectively (Coy *et al.*, 2008; Ghersevich *et al.*, 2015). When the COC moves forward in the ampulla, it is influenced continuously by factors secreted from the ampulla epithelial cells that are vital for the improvement of sperm-zona binding, zona pellucida (ZP) hardening and fertilization rate (Coy *et al.*, 2008; Mugnier *et al.*, 2009; Hanaue *et al.*, 2011; Coy *et al.*, 2012; Ghersevich *et al.*, 2015). There was a higher blastocyst yield in the AM group than in the AMOV group, in which monolayers of frozen/thawed BAOECs and BTOECs were used for oocyte maturation and embryo culture, respectively. This finding could be partly explained by the fact that under *in vivo* conditions, continuous renewal of oviductal fluid creates an optimal microenvironment for embryo development, whereas the accumulation of waste metabolites during co-culture adversely affects early-stage embryo development (Buhi, 2002).

The blastocyst's ability to withstand cryopreservation can be used as an indicator for assessing embryo quality (Rizos *et al.* 2002, 2008; Moore *et al.*, 2007). In the present study, cryotolerance at 4 h post thawing was not affected by the co-culture conditions. However, blastocysts in the AM and AMOV groups showed higher cryotolerance at both 24 and 48 h post thawing compared with the OV and control groups. Previous studies have shown that using BOEC (Lopera-Vásquez *et al.*, 2016) and oviductal fluid (Lopera-Vásquez *et al.*, 2017) in *in vitro* embryo cultures improved the survival rate of vitrified-warmed blastocysts.

It is well documented that TE cells play a crucial role in implantation and placentation (Lopera-Vásquez *et al.*, 2016). The present results showed that the numbers of ICM and TE cells present in all co-culture treatments were higher compared with the numbers in the control group; this finding was in close agreement with results of previous studies (Galli and Lazzari, 1996; Rizos *et al.*, 2001; Lopera-Vásquez *et al.*, 2016). These findings can be partly explained because TE cells are essential for blastocyst re-expansion and preservation after vitrification (Lopera-Vásquez *et al.*, 2016).

Expression patterns of relevant genes are a useful tool for tracking and predicting the clinical outcome of IVF (Lonergan *et al.*, 2003; Jurisicova *et al.*, 2003; Corcoran *et al.*, 2007). This study showed that the expression patterns of genes – including *BAX*, *INF-T*, *PLAC8*, *ATP1A1* and *AQP3* – in day 8 blastocysts were influenced by co-culture conditions. However, the *P53* expression level was not affected. Na/K-ATPase α -1 (*ATP1A1*) is a gene involved in membrane trafficking (Lopera-Vásquez *et al.*, 2017), and its product is essential for blastocyst expansion (Barcroft *et al.*, 2004). Monitoring its expression is a suitable way to assess cryotolerance. The highest expression level for *ATP1A1* was found in the AMOV group and the lowest levels were found in the control and AM groups. Hosseini *et al.* (2015) reported that the number of *ATP1A1* transcripts in high-quality ovine embryos was significantly higher than in those groups with poor-quality embryos. The higher survival rate of the vitrified-warmed blastocysts in the AMOV group could be partly explained by the higher level of *ATP1A1* expression in this group. The highest expression level for *AQP3* was found in the AMOV group, followed by the AM group. One of the well known molecular explanations for assessing cryotolerance ability after vitrification is the water channel aquaporin (*AQP3*) protein, which is involved in membrane permeability regulation (Cordova *et al.*, 2014). This gene codes for a protein involved in facilitating rapid water movement through the membrane and therefore increases the survival rate of the vitrified-warmed blastocysts (Huang *et al.*, 2006; Lopera-Vásquez *et al.*, 2017). Consequently, upregulation of the *AQP3* gene increases the survival rate and dummy expression of *AQP3* in mouse oocytes increased cryotolerance (Edashige *et al.*, 2003). In this study, results for the obtained cryotolerance data correlated with the observed expression level for *AQP3* in the treatments. Cordova *et al.* (2014) reported that there was no difference between the expression levels for *AQP3* in the groups co-cultured with BOEC in the first 4 days of IVC compared with the expression level in the control group.

PLAC8 is related to successful embryo development, more particularly it plays a role in feto-maternal communication and placental development (Galaviz-Hernandez *et al.*, 2003). *PLAC8* and *AQP3* showed similar expression patterns under different treatments. *INF-T* gene products modify ovarian progesterone secretion and stimulate endometrial nutrient secretion, which is required for growth of the conceptus. In addition, *INF-T* gene products regulate the expression of uterine-derived factors, which are responsible for

the early development of the conceptus and placental attachment (Austin *et al.*, 1996). In this study, the lowest expression level for *INF-T* was found in the OV group; this finding partly explains the lowest rate of development of blastocysts in this group.

Apoptosis, a complex programmed cell death process that is activated in embryonic cells in stressful conditions, can be evaluated by examining the expression patterns of the involved genes (e.g. *BAX*) (Betts and King, 2001). *BAX* is a proapoptotic gene (Stallock *et al.*, 2003) and can be used as a potential marker for embryo quality. Melka *et al.* (2010) showed that the expression level of *BAX* in poor-quality embryos was higher than that in high-quality embryos. The present results showed that the expression level of the *BAX* gene in the OV group was lower than that in the other groups. However, the pattern of expression of *BAX* among groups did not explain the observed survival rate of the vitrified-warmed blastocysts. The *P53* gene, also known as *TP53*, codes for a protein that regulates the cell cycle. *P53* plays an important role in conserving stability by preventing genome mutation (Strachan and Read, 2018). However, the present results showed that the treatments did not affect *P53* expression. Similarly, Cordova *et al.* (2014) reported no difference between expression levels of *P53* in the control and in the BOEC co-culture treatments.

Conclusions

This study found that co-culture of bovine oocytes with a monolayer of frozen-thawed bovine ampullary cells during the last 6 h of IVM increased both the blastocyst yield and quality of the produced embryos. The achieved improvement in blastocyst development and embryo quality suggested that using BAOEC co-culture in *in vitro* maturation provides the conditions needed for oocyte maturation and embryo development. Furthermore, bio-banking of fresh oviductal single cells can provide a reliable and reproducible way for IVM/IVF/IVC to effectively mimic *in vivo* conditions.

Acknowledgements. We specifically thank Sheryl Nikpoor, the editor of scientific manuscripts at the Shiraz University, for her meticulous editing of our manuscript. The authors also thank A. Shams, the Chief Director of the Shiraz Abattoir for his kind assistance in collecting the ovaries. Thanks also to Z. Heidary of Royan Institute and H. Jesmani for their technical help in preparing the IVF laboratory and performing RT-PCR.

Financial support. This study was financially supported by Shiraz University (grant no. 97GCU1M1251) and the Centre of Excellence for Research on High Producing Dairy Cows.

Conflicts of interest. The authors declare no conflict of interest.

Ethical standards. The ethical and research committee of the School of Veterinary Medicine, Shiraz University (97GCU1M1251) approved this study.

References

- Austin KJ, Ward SK, Glauca Teixeira M, Dean VC, Moore DW and Hansen TR (1996) Ubiquitin cross-reactive protein is released by the bovine uterus in response to interferon during early pregnancy. *Biol Reprod* **54**, 600–6.
- Avilés M, Gutierrez-Adán A and Coy P (2010) Oviductal secretions: will they be key factors for the future ARTs? *Mol Hum Reprod* **16**, 896–906.
- Avilés M, Coy P and Rizos D (2015) The oviduct: A key organ for the success of early reproductive events. *Anim Front* **5**, 25–31.
- Azari M, Kafi M, Ebrahimi B, Fatehi R and Jamalzadeh M (2017) Oocyte maturation, embryo development and gene expression following two

- different methods of bovine cumulus–oocyte complexes vitrification. *Vet Res Commun* **41**, 49–56.
- Barcroft L, Moseley A, Lingrel J and Watson A** (2004) Deletion of the Na/K-ATPase α 1-subunit gene (*Atpl1a*) does not prevent cavitation of the preimplantation mouse embryo. *Mech Dev* **121**, 417–26.
- Betts D and King W** (2001) Genetic regulation of embryo death and senescence. *Theriogenology* **55**, 171–91.
- Brackett BG and Oliphant G** (1975) Capacitation of rabbit spermatozoa in vitro. *Biol Reprod* **12**, 260–74.
- Buhi WC** (2002) Characterization and biological roles of oviduct-specific, oestrogen-dependent glycoprotein. *Reproduction* **123**, 355–62.
- Corcoran D, Rizos D, Fair T, Evans AC and Lonergan P** (2007) Temporal expression of transcripts related to embryo quality in bovine embryos cultured from the two-cell to blastocyst stage in vitro or in vivo. *Mol Reprod Dev* **74**, 972–7.
- Cordova A, Perreau C, Uzbekova S, Ponsart C, Locatelli Y and Mermillod P** (2014) Development rate and gene expression of IVP bovine embryos cocultured with bovine oviduct epithelial cells at early or late stage of preimplantation development. *Theriogenology* **81**, 1163–73.
- Coy P, Cánovas S, Mondéjar I, Saavedra MD, Romar R, Grullón L, Matás C and Avilés M** (2008) Oviduct-specific glycoprotein and heparin modulate sperm–zona pellucida interaction during fertilization and contribute to the control of polyspermy. *Proc Natl Acad Sci* **105**, 15809–14.
- Coy P, Lloyd R, Romar R, Satake N, Matás C, Gadea J and Holt WV** (2010) Effects of porcine pre-ovulatory oviductal fluid on boar sperm function. *Theriogenology* **74**, 632–42.
- Coy P, Garcia-Vázquez FA, Visconti PE and Avilés M** (2012) Roles of the oviduct in mammalian fertilization. *Reproduction* **144**, 649–60.
- Edashige K, Yamaji Y, Kleinhans F and Kasai M** (2003) Artificial expression of aquaporin-3 improves the survival of mouse oocytes after cryopreservation. *Biol Reprod* **68**, 87–94.
- Fazeli A and Pewsey E** (2008) Maternal communication with gametes and embryos: a complex interactome. *Brief Func Genomics Proteomics* **7**, 111–8.
- Ferraz MA, Henning HH, Stout TA, Vos PL and Gadella BM** (2017) Designing 3-dimensional in vitro oviduct culture systems to study mammalian fertilization and embryo production. *Annals Biomed Eng* **45**, 1731–44.
- Freshney RI** (2015) *Culture of Animal Cells: A Manual of Basic Technique and Specialized Applications*, 7th edition. New Jersey, USA: John Wiley & Sons.
- Gad A, Hoelker M, Besenfelder U, Havlicek V, Cinar U, Rings F, Held E, Dufort I, Sirard M-A and Schellander K** (2012) Molecular mechanisms and pathways involved in bovine embryonic genome activation and their regulation by alternative *in vivo* and *in vitro* culture conditions. *Biol Reprod* **87**, 100.
- Galaviz-Hernandez C, Stagg C, De Ridder G, Tanaka TS, Ko MS, Schlessinger D and Nagaraja R** (2003) Plac8 and Plac9, novel placental-enriched genes identified through microarray analysis. *Gene* **309**, 81–9.
- Galli C and Lazzari G** (1996) Practical aspects of IVM/IVF in cattle. *Anim Reprod Sci* **42**, 371–9.
- Ghersevich S, Massa E and Zumoffen C** (2015) Oviductal secretion and gamete interaction. *Reproduction* **149**, R1–R14.
- Graf A, Krebs S, Heininen-Brown M, Zakhartchenko V, Blum H and Wolf E** (2014) Genome activation in bovine embryos: review of the literature and new insights from RNA sequencing experiments. *Anim Reprod Sci* **149**, 46–58.
- Hanaue M, Miwa N, Uebi T, Fukuda Y, Katagiri Y and Takamatsu K** (2011) Characterization of S100A11, a suppressive factor of fertilization, in the mouse female reproductive tract. *Mol Reprod Dev* **78**, 91–103.
- Hosseini S, Asgari V, Ostadhosseini S, Hajian M, Ghanaei H and Nasr-Esfahani M** (2015) Developmental competence of ovine oocytes after vitrification: differential effects of vitrification steps, embryo production methods, and parental origin of pronuclei. *Theriogenology* **83**, 366–76.
- Huang H-F, He R-H, Sun C-C, Zhang Y, Meng Q-X and Ma Y-Y** (2006) Function of aquaporins in female and male reproductive systems. *Hum Reprod Update* **12**, 785–95.
- Hunter R** (1974) Chronological and cytological details of fertilization and early embryonic development in the domestic pig, *Sus scrofa*. *Anatom Rec* **178**, 169–185.
- Juriscova A, Antenos M, Varmuza S, Tilly JL and Casper RF** (2003) Expression of apoptosis-related genes during human preimplantation embryo development: potential roles for the Harakiri gene product and caspase-3 in blastomere fragmentation. *MHR: Basic Sci Reprod Med* **9**, 133–41.
- Kattal N, Cohen J and Barmat LI** (2008) Role of coculture in human *in vitro* fertilization: a meta-analysis. *Fertil Steril* **90**, 1069–76.
- Leese HJ, Hugentobler SA, Gray SM, Morris DG, Sturmey RG, Whitear S-L and Sreenan JM** (2007) Female reproductive tract fluids: composition, mechanism of formation and potential role in the developmental origins of health and disease. *Reprod Fertil Dev* **20**, 1–8.
- Lonergan P and Fair T** (2008) In vitro-produced bovine embryos—dealing with the warts. *Theriogenology* **69**, 17–22.
- Lonergan P and Fair T** (2016) Maturation of oocytes in vitro. *Ann Rev Anim Biosci* **4**, 255–68.
- Lonergan P, Rizos D, Kanka J, Nemcova L, Mbaye A, Kingston M, Wade M, Duffy P and Boland M** (2003) Temporal sensitivity of bovine embryos to culture environment after fertilization and the implications for blastocyst quality. *Reproduction* **126**, 337–46.
- Lopera-Vázquez R, Hamdi M, Fernandez-Fuertes B, Maillo V, Beltrán-Breña P, Calle A, Redruello A, López-Martín S, Gutierrez-Adán A and Yáñez-Mó M** (2016) Extracellular vesicles from BOEC in *in vitro* embryo development and quality. *PLoS One* **11**, e0148083.
- Lopera-Vázquez R, Hamdi M, Maillo V, Gutierrez-Adán A, Bermejo-Alvarez P, Ramirez MA, Yáñez-Mó M and Rizos D** (2017) Effect of bovine oviductal extracellular vesicles on embryo development and quality *in vitro*. *Reproduction* **153**, 461–70.
- Marshall OJ** (2004) PerlPrimer: cross-platform, graphical primer design for standard, bisulphite and real-time PCR. *Bioinformatics* **20**, 2471–2.
- Melka M, Rings F, Hölker M, Tholen E, Havlicek V, Besenfelder U, Schellander K and Tesfaye D** (2010) Expression of apoptosis regulatory genes and incidence of apoptosis in different morphological quality groups of *in vitro*-produced bovine pre-implantation embryos. *Reprod Domest Anim* **45**, 915–21.
- Mondéjar I, Acuna O, Izquierdo-Rico M, Coy P and Avilés M** (2012) The oviduct: functional genomic and proteomic approach. *Reprod Domest Anim* **47**, 22–9.
- Moore K, Rodríguez-Sallaberry C, Kramer J, Johnson S, Wroclawska E, Goicoa S and Niasari-Naslaji A** (2007) In vitro production of bovine embryos in medium supplemented with a serum replacer: effects on blastocyst development, cryotolerance and survival to term. *Theriogenology* **68**, 1316–25.
- Mugner S, Kervella M, Douet C, Canepa S, Pascal G, Deleuze S, Duchamp G, Monget P and Goudet G** (2009) The secretions of oviduct epithelial cells increase the equine *in vitro* fertilization rate: are osteopontin, atrial natriuretic peptide A and oviductin involved? *Reprod Biol Endocrinol* **7**, 129.
- Pontes J, Nonato-Junior I, Sanches B, Ereno-Junior J, Uvo S, Barreiros T, Oliveira J, Hasler J and Seneda M** (2009) Comparison of embryo yield and pregnancy rate between *in vivo* and *in vitro* methods in the same Nelore (*Bos indicus*) donor cows. *Theriogenology* **71**, 690–7.
- R Core Team** (2017) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rizos D, Ward F, Boland M and Lonergan P** (2001) Effect of culture system on the yield and quality of bovine blastocysts as assessed by survival after vitrification. *Theriogenology* **56**, 1–16.
- Rizos D, Ward F, Duffy P, Boland MP and Lonergan P** (2002) Consequences of bovine oocyte maturation, fertilization or early embryo development *in vitro* versus *in vivo*: implications for blastocyst yield and blastocyst quality. *Mol Reprod Dev* **61**, 234–48.
- Rizos D, Clemente M, Bermejo-Alvarez P, de La Fuente J, Lonergan P and Gutiérrez-Adán A** (2008) Consequences of *in vitro* culture conditions on embryo development and quality. *Reprod Domest Anim* **43**, 44–50.
- Rodríguez-Martínez H** (2007) Role of the oviduct in sperm capacitation. *Theriogenology* **68**, S138–46.
- Schmaltz-Panneau B, Cordova A, Dhorne-Pollet S, Hennequet-Antier C, Uzbekova S, Martinot E, Doret S, Martin P, Mermillod P and Locatelli Y** (2014) Early bovine embryos regulate oviduct epithelial cell gene expression during *in vitro* co-culture. *Anim Reprod Sci* **149**, 103–16.

- Stallock J, Molyneaux K, Schaible K, Knudson CM and Wylie C** (2003) The pro-apoptotic gene Bax is required for the death of ectopic primordial germ cells during their migration in the mouse embryo. *Development* **130**, 6589–97.
- Steinhauer N, Boos A and Günzel-Apel AR** (2004) Morphological changes and proliferative activity in the oviductal epithelium during hormonally defined stages of the oestrous cycle in the bitch. *Reprod Domest Anim* **39**, 110–9.
- Strachan T and Read A** (2018) *Human Molecular Genetics*, 5th edition. Taylor, Francis Group, Boca Roca, USA: Garland Science.
- Tanghe S, Van Soom A, Nauwynck H, Coryn M and de Kruif A** (2002) Minireview: Functions of the cumulus oophorus during oocyte maturation, ovulation, and fertilization. *Mol Reprod Dev* **61**, 414–24.
- Thouas G, Korfiatis N, French A, Jones G and Trounson A** (2001) Simplified technique for differential staining of inner cell mass and trophectoderm cells of mouse and bovine blastocysts. *Reprod Biomed Online* **3**, 25–9.
- Way A, Schuler A and Killian G** (1997) Influence of bovine ampullary and isthmic oviductal fluid on sperm–egg binding and fertilization *in vitro*. *J Reprod Fertil* **109**, 95–101.