

# Chemo-dynamical signatures in simulated Milky Way-like galaxies

Alessandro Spagna<sup>1</sup>, Anna Curir<sup>1</sup>, Marco Giammaria<sup>2</sup>,  
Mario G. Lattanzi<sup>1</sup>, Giuseppe Murante<sup>3</sup> and Paola Re Fiorentin<sup>1</sup>

<sup>1</sup>INAF – Osservatorio Astrofisico di Torino  
via Osservatorio 20, 10025 Pino Torinese, Italy  
email: spagna@oato.inaf.it

<sup>2</sup>Università degli Studi di Torino, Italy

<sup>3</sup>INAF – Osservatorio Astronomico di Trieste, Italy

**Abstract.** We have investigated the chemo-dynamical evolution of a Milky Way-like disk galaxy, AqC4, produced by a cosmological simulation integrating a sub-resolution ISM model. We evidence a global inside-out *and* upside-down disk evolution, that is consistent with a scenario where the “thin disk” stars are formed from the accreted gas close to the galactic plane, while the older “thick disk” stars are originated *in situ* at higher heights. Also, the bar appears the most effective heating mechanism in the inner disk. Finally, no significant metallicity-rotation correlation has been observed, in spite of the presence of a negative [Fe/H] radial gradient.

**Keywords.** Galaxy: abundance, evolution, structure, kinematics and dynamics


In order to study the chemo-dynamical signatures related to the galactic formation processes, we have analyzed a new  $\Lambda$ CDM cosmological simulation, AqC4, carried out with the GADGET-3 TreePM+SPH code, where star formation, chemical evolution and stellar feedback are described using a sub-grid Multi Phase Particle Integrator (MUPPI) model. The main parameters of this simulation are listed in Table 1, while further details on the model are described by Murante *et al.* (2015).

The morphology of AqC4 at redshift  $z = 0$  is well represented by the three main stellar components of a Milky Way-like galaxy (i.e. bulge/spheroid, thin and thick disk). An extensive analysis of the spatial, kinematic, and chemical properties of AqC4 is presented by Giammaria (2017), who confirmed that the 6D ( $\mathbf{x}, \mathbf{v}$ ) stellar distribution is similar to that observed in the Milky Way. The main difference is the presence of a more massive bulge ( $B/T=0.34$ ), which also causes a faster rotation velocity ( $V_\phi \sim 270$  km/s at  $R \simeq 5 - 10$  kpc). Moreover, the metallicity distribution of the stellar halo results a few dex higher than that of the Galactic halo.

However, the overall evolution of AqC4 appears quite consistent with similar studies based on independent simulations (e.g. Minchev *et al.* 2012, Bird *et al.* 2013, Martig *et al.* 2014, Ma *et al.* 2017).

**Table 1.** AqC4 parameters.  $M_{\text{DM}}$ : mass of DM particles;  $M_{\text{gas}}$  initial mass of gas particle;  $\epsilon$  = smoothing parameter;  $M_{\text{vir}}$  and  $R_{\text{vir}}$ : DM mass and virial radius at redshift  $z=0$ ;  $N_{\text{DM}}, N_{\text{gas}}, N_{\text{star}}$ : number of DM particles, gas particles and stellar particles within  $R_{\text{vir}}$  at  $z = 0$ ;  $\Omega_i$ : density parameters;  $H_0$ : Hubble constant.

$M_{\text{DM}}$ [ $M_\odot$ ]	$M_{\text{gas}}$ [ $M_\odot$ ]	$\epsilon$ [kpc]	$M_{\text{vir}}$ [ $M_\odot$ ]	$R_{\text{vir}}$ [kpc]	$N_{\text{DM}}$	$N_{\text{gas}}$	$N_{\text{star}}$	$\Omega_{\text{m}}$	$\Omega_\Lambda$	$\Omega_{\text{b}}$	$H_0$
$2.7 \cdot 10^5$	$5.1 \cdot 10^4$	0.163	$1.49 \cdot 10^{12}$	237	5 518 587	1 348 120	6 919 646	0.25	0.75	0.04	71



**Figure 1.** Disk  $|z|$  distributions with  $R$  between 3-5 kpc and 6-8 kpc.

Figure 1 compares the evolution of the vertical distributions at redshift 0, 1, and 2.16, that clearly evidence an *upside-down* disk evolution in the “solar” annulus,  $6 \text{ kpc} < R < 8 \text{ kpc}$ . Such result supports a formation scenario that firstly generates the ancient “thick disk” stars *in situ*, from an initially turbulent ISM characterized by shorter scale-lengths and higher scale-heights with respect to the younger “thin disk” stars (Haywood *et al.* 2013, Bird *et al.* 2013). Conversely, after about 6 Gyr ( $z = 1$ ), a strong disk thickening is observed in the inner disk,  $3 \text{ kpc} < R < 5 \text{ kpc}$ , possibly due to the heating induced by the central bar (cfr. Grand *et al.* 2016).

Finally, although a typical negative  $[\text{Fe}/\text{H}]$  radial gradient is present in the disk of AqC4, no significant rotation-metallicity relation is present, as observed in our Milky Way (Spagna *et al.* 2010, Lee *et al.* 2011; see also Schönrich & McMillan 2017). This result deserves further investigation as it may be an effect of the specific formation history of this simulation.

## References

- Bird, J. C. *et al.* 2013, *ApJ*, 773, 43  
 Giammaria, M. 2017, *Tesi laurea magistrale in Fisica - Ind. Astrofisica*, Turin University (Italy)  
 Grand, R. J. J., Springel, V., Gómez, F. A. *et al.* 2016, *MNRAS*, 459, 199  
 Haywood, M., Di Matteo, P., Lehnert, M. D. *et al.* 2013, *A&A*, 560, A109  
 Lee, Y. S., Beers, T. C., An, D. *et al.* 2011, *ApJ*, 738, 187  
 Ma, X., Hopkins, P. F., Wetzell, A. R. *et al.* 2017, *MNRAS*, 467, 2430  
 Martig, M., Minchev, I., & Flynn, C. 2014, *MNRAS*, 442, 2474  
 Minchev, I., Famaey, B., Quillen, A. C. *et al.* 2012, *A&A*, 548, A127  
 Murante, G., Monaco, P., Borgani, S., *et al.* 2015, *MNRAS*, 447, 178  
 Schönrich, R. & McMillan, P. J. 2017, *MNRAS*, 467, 1154  
 Spagna, A., Lattanzi, M. G., Re Fiorentin, P., & Smart, R. L. 2010, *A&A*, 510, L4