

Smoke-enhanced seed germination in Mediterranean Lamiaceae

Şükrü Serter Çatav¹, Köksal Küçükakyüz¹, Kenan Akbaş¹ and Çağatay Tavşanoğlu^{2*}

¹Department of Biology, Mugla Sıtkı Koçman University, Kötekli 48000, Mugla, Turkey; ²Fire Ecology and Seed Research Lab., Division of Ecology, Department of Biology, Hacettepe University, Beytepe 06800, Ankara, Turkey

(Received 22 August 2013; accepted after revision 11 April 2014; first published online 22 May 2014)

Abstract

The role of smoke in fire-stimulated germination in the Mediterranean Basin has often been underestimated. A few records on smoke-enhanced germination are present in Mediterranean Lamiaceae species, but there is still a shortage of information to allow generalizations about this family to be made. To test the hypothesis that smoke enhances germination in Mediterranean Lamiaceae species, we performed a germination experiment, including aqueous smoke treatments in various concentrations (1:1, 1:10 and 1:100) on seven eastern Mediterranean Lamiaceae taxa. Six of the studied taxa (*Lavandula stoechas*, *Origanum onites*, *Phlomis bourgaei*, *Stachys cretica* ssp. *smyrnaea*, *Satureja thymbra*, *Teucrium lamiifolium* ssp. *stachyophyllum*) showed significant increments in germination percentage in at least one smoke treatment, as compared to the control. Moreover, *L. stoechas*, *S. thymbra* and *T. lamiifolium* ssp. *stachyophyllum* displayed faster germination in at least one smoke treatment than in the control. Of the species showing significant increments in germination percentages after aqueous smoke application, at least one single concentration of smoke solution did not stimulate germination, except in *L. stoechas* and *S. thymbra* which responded positively to all smoke treatments. Therefore, the concentration of aqueous smoke that improved germination was species-specific. Our results contribute to the current limited knowledge on smoke-enhanced germination in Mediterranean Lamiaceae, and support the idea that smoke is an important germination cue for this family.

Keywords: fire, germination, Lamiaceae, Mediterranean Basin, smoke

*Correspondence
Fax: + 903122992028
Email: ctavsan@hacettepe.edu.tr

Introduction

Plant community dynamics in many ecosystems are largely shaped by wildfires (Brown and Smith, 2000; Bond and Keeley, 2005). Many plants are able to regenerate by mechanisms such as resprouting and seeding after fires, especially in Mediterranean-type ecosystems (Keeley and Zedler, 1978; Pausas and Verdú, 2005). Post-fire regeneration by seeds is driven by fire-induced seed release or germination in Mediterranean plant species (Keeley *et al.*, 2012). The important role of fire-related cues (heat and smoke) in the stimulation of germination in many plants is well known. Heat-stimulated germination, which is a common property among hard-seeded plants, has been shown in fire-prone Mediterranean-type ecosystems (Thanos *et al.*, 1992; Keeley and Bond, 1997; Herranz *et al.*, 1998; Moreira *et al.*, 2010). Smoke-stimulated germination has also been demonstrated in a wide range of species from both fire-prone (Brown, 1993a; Dixon *et al.*, 1995; Keeley and Bond, 1997; Keeley and Fotheringham, 1998; Moreira *et al.*, 2010) and non-fire-prone areas (Pierce *et al.*, 1995; Adkins and Peters, 2001; Daws *et al.*, 2007) of the world.

Lamiaceae is a plant family containing over 240 genera and 6500 species worldwide. Although the family has a worldwide distribution, it shows great diversity in the Mediterranean Basin, Irano-Turanian floristic region and eastern Asia (Takhtajan, 2009), and is represented by many endemic taxa in local floras of this region (e.g. Yeşilyurt and Akaydın, 2012). Many members of the family are of economic importance, including medicinal plants, culinary herbs, fragrance plants and ornamentals (Simpson, 2010). In particular, species of *Lavandula* (lavender), *Mentha* (mint), *Melissa* (lemon balm), *Ocimum* (basil), *Origanum* (oregano), *Rosmarinus* (rosemary), *Salvia* (sage), *Satureja* (savory) and *Thymus* (thyme) are widely used as culinary herbs by local people of circum-Mediterranean areas (Naghbi *et al.*, 2005; Gürdal and Kültür, 2013) and in North America (Small, 2006). At the same time, Lamiaceae taxa are of ecological importance in plant

communities of the Mediterranean area, and can be found in various Mediterranean habitats, such as maquis and phrygana, where fire plays a key role in shaping the structure of plant communities (Verdú and Pausas, 2007; Keeley *et al.*, 2012). Species of Lamiaceae are also found frequently in burned Mediterranean habitats (Kazanis and Arianoutsou, 2004; Kavgacı *et al.*, 2010; Tavşanoğlu and Gürkan, 2014) on account of seedling emergence after fire (Paula *et al.*, 2009). This can be attributed to their enhanced germination ability after fires (Moreira *et al.*, 2010). Indeed, Lamiaceae is one of the main families found in the Northern Hemisphere with numerous species showing smoke-stimulated germination (Keeley *et al.*, 2012).

Although there are records of smoke-stimulated germination in Lamiaceae taxa worldwide, many of these come from outside of the Mediterranean-type ecosystems (Clarke *et al.*, 2000; Tang *et al.*, 2003; Pennacchio *et al.*, 2005; Tierney, 2006; Todorović *et al.*, 2007; Ervin *et al.*, 2010; Schwilk and Zavala, 2012; Tavşanoğlu *et al.*, in prep.). This suggests that the possibly significant role of smoke in the recovery of Lamiaceae species in fire-driven Mediterranean ecosystems might have been overlooked (but see Keeley and Fotheringham, 1998; Moreira *et al.*, 2010). In the Mediterranean Basin, in particular, few studies have tested the germination response of Lamiaceae species to smoke (Crosti *et al.*, 2006; Reyes and Trabaud, 2009; Moreira *et al.*, 2010; Çatav *et al.*, 2012). As a consequence, there is still a shortage of information on the role of smoke in post-fire germination of the members of the Lamiaceae family.

Based on the records of smoke-stimulated germination in many species belonging to the Lamiaceae worldwide, we hypothesized that smoke will enhance germination of Mediterranean Lamiaceae species. To test this hypothesis, we performed a germination experiment with seeds of seven Lamiaceae taxa growing in natural Mediterranean habitats. The percentage and rate of germination of each taxon were assessed in aqueous smoke treatments at different concentrations, and these results were compared to the controls to determine if smoke

treatments resulted in any increment in germination percentage and rate in the studied taxa. By conducting this experiment, we aimed to clarify the role of smoke in enhanced germination of Mediterranean Lamiaceae species.

Materials and methods

Study taxa, study area and seed collection

We collected the seeds of seven Lamiaceae taxa growing in natural habitats of Muğla province, south-western Turkey, the eastern Mediterranean Basin (Table 1). These taxa have a soil seed bank, have obligate or facultative seeder regeneration mode, and are found in post-fire environments in the study area (Tavşanoğlu and Gürkan, 2014). The study area has a Mediterranean climate with wet winters and dry summers. The collections were carried out between July and September 2012, coinciding with the propagule dispersal period of each taxon. Ripe nutlets (seeds) of each taxon were collected from a minimum of ten individuals of the same population, except *Phlomis bourgaei* in which fewer individuals could be found. Seeds were separated from the dry flower parts by hand and stored in paper envelopes under laboratory conditions until the start of the experiment in October 2012. For each taxon, mean seed mass was determined for four replicates of 100 seeds (three replicates in *P. bourgaei* due to the limited number of seeds).

Preparation of aqueous smoke solutions

Since the active compounds of smoke for germination are very stable in aqueous solutions (Van Staden *et al.*, 2000) and aqueous extracts of smoke are equally as effective as airborne smoke in stimulating seed germination (Brown, 1993b; Dixon *et al.*, 1995), we used aqueous smoke solutions as smoke treatment for the experiment. Dry needles of *Pinus brutia*, the pine

Table 1. List of the studied taxa. Growth form (GF), seed collection locality (Loc) and altitude (alt; m a.s.l.), mean (\pm SE) seed size (in mg) and distributional range of the taxa are given. Nomenclature follows Davis (1965–1985)

Taxa	GF	Loc	alt	Seed size	Range
<i>Lavandula stoechas</i> L.	w	36°53'N–28°11'E	22	0.60 \pm 0.01	MB
<i>Origanum onites</i> L.	w	37°00'N–28°21'E	65	0.06 \pm 0.00	EMB
<i>Phlomis bourgaei</i> Boiss.	w	37°09'N–28°22'E	650	6.55 \pm 0.32	EMB (E)
<i>Satureja thymbra</i> L.	w	36°43'N–27°26'E	273	0.56 \pm 0.02	EMB
<i>Stachys cretica</i> L. ssp. <i>smyrnaea</i> Rech.Fil.	p	37°09'N–28°22'E	650	2.64 \pm 0.05	EMB (E)
<i>Teucrium divaricatum</i> Sieber ssp. <i>divaricatum</i>	w	36°43'N–27°27'E	405	1.69 \pm 0.03	EMB
<i>Teucrium lamifolium</i> D'Urv. ssp. <i>stachyophyllum</i> (P.H. Davis) Hedge & Ekim	p	37°00'N–28°21'E	65	0.63 \pm 0.02	EMB

w, woody; p, perennial herb; MB, the whole Mediterranean Basin; EMB, the eastern MB; E, regionally endemic to Turkey and the east Aegean islands.

species dominating the vegetation in most of the study region, were separated into small pieces to use as the plant material needed for preparing aqueous smoke solutions. Four replicates of 5 g of this plant material were heated separately in metallic containers in the oven for 30 min at $193 \pm 1^\circ\text{C}$ (Jäger *et al.*, 1996; Moreira *et al.*, 2010; Çatav *et al.*, 2012). The mouth of each container was tightly covered by an aluminium slab to capture the smoke generated from the burnt plant material. After the treatment, 50 ml of distilled water was added to the plant material in the container and was left for 10 min (Jäger *et al.*, 1996). By this procedure, the active chemicals found in smoke dissolved in the water in the container. The solution was filtered into a bottle to obtain concentrated smoke solution (hereafter, smoke 1:1; pH 4.32). One millilitre of concentrated solution was diluted in 10 and 100 ml of distilled water to prepare lower concentrations of smoke solutions (hereafter, smoke 1:10, pH 4.40, and smoke 1:100, pH 5.03, respectively). Since previous observations revealed that smoke-stimulated germination was independent of the pH value (Brown and Van Staden, 1997), we did not adjust the pH of the different concentrations of smoke solutions and the control.

Germination experiment

Seeds of each taxon were incubated in different concentrations of smoke solution for 24 h. A group of seeds from each taxon was also incubated in distilled water (pH 5.75) for 24 h to be used as a control for the aqueous smoke treatments. After the treatments were applied, seeds were sown in Petri dishes containing agar (0.7%) as substrate. Each treatment consisted of four replicates of 25 seeds. Petri dishes were placed in an incubator set at 20.0°C ($\pm 0.5^\circ\text{C}$) in darkness, i.e. favourable conditions for the germination of many Mediterranean species (Thanos *et al.*, 1991; Luna *et al.*, 2012). The seeds were monitored for germination on a daily basis during the first 10 d of the incubation, and then weekly until the end of the experiment.

The criterion of germination was the observation of visible radicle protrusion. Germinated seeds were counted and removed from the Petri dishes at every check. The experiment was finalized after 7 weeks of incubation. The viability of non-germinated seeds was determined by the cut test, and the seeds with an intact embryo were classified as viable.

Data analysis

For each taxon we included the germinated versus non-germinated seeds (including intact but not empty ones) in the analysis. The probability of germination stimulation by each smoke treatment was determined using analysis of deviance (GLM) with binomial error distribution (Moreira *et al.*, 2010). Mean germination time was calculated using the equation $\Sigma(nD)/\Sigma n$, where n is the number of seeds germinated on day D and D is the number of days from the beginning of the incubation period (Tompsett and Pritchard, 1998). Dunnett's test was used for comparison of mean germination time for smoke treatments and the control in each taxon. Data normality and homoscedasticity were tested using the Shapiro–Wilk and Bartlett's tests before each analysis, respectively. Since the prerequisites for parametric tests were obtained in all cases, germination time data were not transformed before analysis.

Results

Of the seven taxa examined, six showed significant increments in germination percentage in at least one smoke solution compared to the control (Table 2). Among the studied taxa, *Lavandula stoechas*, *Origanum onites* and *Teucrium lamiifolium* ssp. *stachyophyllum* had relatively higher germination percentages (>68%) in control conditions. On the other hand, *Stachys cretica* ssp. *smyrnaea* and *Teucrium divaricatum* ssp. *divaricatum* had very low germination percentages (<3%) in

Table 2. Mean (\pm SE) germination percentage in the control and after smoke treatments. The concentrations of different smoke treatments are shown as 1:1, 1:10 and 1:100. The significance of the pairwise comparison of each treatment with the corresponding control (GLM) is given (ns, not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$)

Taxa	Control	Smoke treatment		
		1:1	1:10	1:100
<i>L. stoechas</i>	68.6 \pm 2.3	88.9 \pm 5.2***	88.1 \pm 4.56**	94.6 \pm 3.2****
<i>O. onites</i>	83.1 \pm 4.7	92.0 \pm 2.3 ^{ns}	93.8 \pm 4.7*	97.0 \pm 1.8**
<i>P. bourgaei</i>	21.4 \pm 3.6	22.5 \pm 4.0 ^{ns}	40.9 \pm 3.5*	–
<i>S. thymbra</i>	47.0 \pm 4.4	86.5 \pm 5.6****	67.3 \pm 12.2*	77.0 \pm 3.6***
<i>S. cretica</i> ssp. <i>smyrnaea</i>	0.0 \pm 0.0	40.9 \pm 2.0****	11.0 \pm 3.0****	1.0 \pm 1.0 ^{ns}
<i>T. divaricatum</i> ssp. <i>divaricatum</i>	2.1 \pm 2.1	2.0 \pm 1.1 ^{ns}	1.0 \pm 1.0 ^{ns}	1.0 \pm 1.0 ^{ns}
<i>T. lamiifolium</i> ssp. <i>stachyophyllum</i>	75.5 \pm 3.4	77.5 \pm 5.6 ^{ns}	75.8 \pm 5.4 ^{ns}	97.7 \pm 1.3****

Table 3. Mean (\pm SE) germination time (in days) in the control and after smoke treatments. The concentrations of different smoke treatments are shown as 1:1, 1:10 and 1:100. The significance of the pairwise comparison of each treatment with the corresponding control (Dunnett's test) is given (ns, not significant; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). No statistical analysis was performed for *T. divaricatum* ssp. *divaricatum* and for *S. cretica* ssp. *smyrnaea* since very little and no germination was observed in the control, respectively.

Taxa	Control	Smoke treatment		
		1:1	1:10	1:100
<i>L. stoechas</i>	8.0 \pm 0.8	6.3 \pm 0.6 ^{ns}	5.3 \pm 0.3**	6.7 \pm 0.4 ^{ns}
<i>O. onites</i>	4.3 \pm 0.2	4.5 \pm 0.5 ^{ns}	4.9 \pm 0.3 ^{ns}	6.3 \pm 0.3**
<i>P. bourgaei</i>	39.7 \pm 3.4	37.6 \pm 1.5 ^{ns}	39.4 \pm 1.2 ^{ns}	–
<i>S. thymbra</i>	14.9 \pm 1.4	7.8 \pm 1.7*	7.7 \pm 1.9*	8.7 \pm 1.0*
<i>S. cretica</i> ssp. <i>smyrnaea</i>	–	2.9 \pm 0.2	3.4 \pm 0.6	–
<i>T. lamiifolium</i> ssp. <i>stachyophyllum</i>	14.8 \pm 0.6	16.8 \pm 1.2 ^{ns}	16.7 \pm 0.9 ^{ns}	7.6 \pm 0.4***

control conditions, suggesting the presence of a high degree of dormancy in these taxa (Table 2). Application of any smoke solution did not break the dormancy in *T. divaricatum* ssp. *divaricatum*, but *Stachys cretica* ssp. *smyrnaea* responded positively. In the taxa showing significant increments in germination percentages after aqueous smoke application, at least one single concentration of smoke solution did not stimulate the germination, except in *L. stoechas* and *Satureja thymbra* which responded positively to all aqueous smoke treatments. On the other hand, none of the smoke concentrations significantly decreased the germination percentages (Table 2).

L. stoechas, *S. thymbra* and *T. lamiifolium* ssp. *stachyophyllum* had faster germination in at least one smoke solution than in the control (Table 3). No significant difference in germination rate was found between the control and aqueous smoke treatments in the other taxa tested, except *O. onites* which had a significantly slower germination rate in smoke 1:100 treatment (Table 3). No result of mean germination time was obtained for *T. divaricatum* ssp. *divaricatum* for any treatment, due to very low germination percentages.

Discussion

Our study shows that aqueous smoke treatments improved germination of six out of the seven Mediterranean Lamiaceae taxa studied. Our results provide the first record of smoke-enhanced germination in four taxa endemic to the eastern Mediterranean Basin (*Origanum onites*, *Teucrium lamiifolium* ssp. *stachyophyllum*, *Phlomis bourgaei* and *Stachys cretica* ssp. *smyrnaea*). The records of germination stimulation by aqueous smoke in the latter two taxa are of additional importance since those taxa are regionally endemic to Turkey and the east Aegean islands, and

especially when it is considered that the high-level dormancy (zero germination in the control) in *S. cretica* ssp. *smyrnaea* was broken by smoke solutions.

Aqueous smoke application resulted in faster germination in three taxa, besides the stimulation of germination. One of the outcomes of earlier germination of seeds is earlier emergence of seedlings. This is likely to have ecological consequences at the plant community level since early emergence of seedlings has a competitive advantage over the species with late-emergent seedlings (De Luis *et al.*, 2008). In post-fire environments, similarly, early appearance of seedlings may give an advantage to a species to outcompete the others. The observed faster germination rate in aqueous smoke treatments in *L. stoechas*, *S. thymbra* and *T. lamiifolium* ssp. *stachyophyllum*, therefore, possibly contributes to the establishment success of these taxa after fire.

An increase in germination percentage and rate in aqueous smoke treatments has previously been observed in *L. stoechas* (Moreira *et al.*, 2010, 2012), similar to the results of our study. However, two studies (Crosti *et al.*, 2006; Çatav *et al.*, 2012) failed to show an increase in germination percentage in this species, possibly because of loss of dormancy due to longer seed storage time (Çatav *et al.*, 2012). Charred wood also stimulated germination in *L. stoechas* (Keeley and Baer-Keeley, 1999). Çatav *et al.* (2012) found an improvement in germination percentage in aqueous smoke treatment in a population of *S. thymbra* from around 100 km to the east of the seed collection area in the present study.

The importance of smoke in fire-induced germination in Mediterranean-type ecosystems of California (Keeley and Fotheringham, 1998), South Africa (Brown, 1993a) and Australia (Dixon *et al.*, 1995) has long been known. However, it had been overlooked in the Mediterranean Basin, based on the presence of few records of smoke- or charred-wood-stimulated

Table 4. List of the Lamiaceae species in which smoke-stimulated germination has been tested (including this study). The studies, conducted both with aqueous smoke solutions and with airborne smoke, are included in the table. GF is growth form and R is the germination response of the species to smoke

Species	GF	Region	R	References
Within Mediterranean-type ecosystems				
<i>Hemiandra pungens</i>	w	Australia	0	Dixon <i>et al.</i> , 1995
<i>Hemiandra pungens</i>	w	Australia	0	Roche <i>et al.</i> , 1997
<i>Lavandula latifolia</i>	w	MB	+	Moreira <i>et al.</i> , 2010
<i>Lavandula stoechas</i>	w	MB	0	Crosti <i>et al.</i> , 2006
<i>Lavandula stoechas</i>	w	MB	+	Moreira <i>et al.</i> , 2010
<i>Lavandula stoechas</i>	w	MB	+	Moreira <i>et al.</i> , 2012
<i>Lavandula stoechas</i>	w	MB	0	Çatav <i>et al.</i> , 2012
<i>Lavandula stoechas</i>	w	MB	+	The present study
<i>Origanum onites</i>	w	MB	+	The present study
<i>Phlomis bourgaei</i>	w	MB	+	The present study
<i>Prostanthera eurybioides</i>	w	Australia	-	Ainsley <i>et al.</i> , 2008
<i>Rosmarinus officinalis</i>	w	MB	+	Moreira <i>et al.</i> , 2010
<i>Rosmarinus officinalis</i>	w	MB	0	Reyes and Trabaud, 2009
<i>Salvia apiana</i>	w	Cal	+	Keeley and Fotheringham, 1998
<i>Salvia columbariae</i>	a	Cal	+	Keeley and Fotheringham, 1998
<i>Salvia leucophylla</i>	w	Cal	+	Keeley and Fotheringham, 1998
<i>Salvia mellifera</i>	w	Cal	+	Keeley and Fotheringham, 1998
<i>Satureja thymbra</i>	w	MB	+	Çatav <i>et al.</i> , 2012
<i>Satureja thymbra</i>	w	MB	+	The present study
<i>Sideritis angustifolia</i>	w	MB	0	Moreira <i>et al.</i> , 2010
<i>Stachys cretica</i> ssp. <i>smyrnaea</i>	p	MB	+	The present study
<i>Teucrium capitatum</i>	w	MB	0	Moreira <i>et al.</i> , 2010
<i>Teucrium divaricatum</i> ssp. <i>divaricatum</i>	w	MB	0	The present study
<i>Teucrium lamiifolium</i> ssp. <i>stachyophyllum</i>	p	MB	+	The present study
<i>Teucrium ronniiferi</i>	w	MB	0	Moreira <i>et al.</i> , 2010
<i>Thymus piperella</i>	w	MB	0	Moreira <i>et al.</i> , 2010
<i>Thymus vulgaris</i>	w	MB	+	Moreira <i>et al.</i> , 2010
Outside Mediterranean-type ecosystems				
<i>Ajuga australis</i>	p	Australia	+	Clarke <i>et al.</i> , 2000
<i>Monarda citriodora</i>	a	USA	0	Schwilk and Zavala, 2012
<i>Monarda citriodora</i>	p	USA	0	Chou <i>et al.</i> , 2012
<i>Monarda fistulosa</i>	p	USA	0	Jefferson <i>et al.</i> , 2008
<i>Nepeta rtanjensis</i>	p	Serbia	+	Todorović <i>et al.</i> , 2007
<i>Phlomis pungens</i>	p	Turkey	0	Tavşanoğlu <i>et al.</i> , in prep.
<i>Plectranthus parviflorus</i>	w	Australia	+	Tang <i>et al.</i> , 2003
<i>Prostanthera askania</i>	w	Australia	+	Tierney, 2006
<i>Pycnanthemum pilosum</i>	p	USA	+	Pennacchio <i>et al.</i> , 2005
<i>Pycnanthemum pilosum</i>	p	USA	+	Ervin <i>et al.</i> , 2010
<i>Pycnanthemum pilosum</i>	p	USA	0	Jefferson <i>et al.</i> , 2008
<i>Pycnanthemum virginianum</i>	p	USA	+	Ervin <i>et al.</i> , 2010
<i>Salvia azurea</i>	p	USA	-	Chou <i>et al.</i> , 2012
<i>Salvia coccinea</i>	p	USA	+	Schwilk and Zavala, 2012
<i>Salvia farinacea</i>	p	USA	+	Schwilk and Zavala, 2012
<i>Salvia iodantha</i>	w	Mexico	-	Zuloaga-Aguilar <i>et al.</i> , 2011
<i>Salvia lavanduloides</i>	w	Mexico	0	Zuloaga-Aguilar <i>et al.</i> , 2011
<i>Salvia penstemonoides</i>	p	USA	0	Schwilk and Zavala, 2012
<i>Salvia reflexa</i>	a	USA	-	Chou <i>et al.</i> , 2012
<i>Salvia syriaca</i>	p	Turkey	0	Tavşanoğlu <i>et al.</i> , in prep.
<i>Salvia thyrsoiflora</i>	w	Mexico	-	Zuloaga-Aguilar <i>et al.</i> , 2011
<i>Stachys annua</i> ssp. <i>annua</i>	p	Turkey	0	Tavşanoğlu <i>et al.</i> , in prep.
<i>Stachys byzantina</i>	p	Turkey	+	Tavşanoğlu <i>et al.</i> , in prep.
From unknown origin				
<i>Hemiandra</i> sp.	w	Australia	+	Cochrane <i>et al.</i> , 2002
<i>Melissa officinalis</i>	p	Iran?	0	Abdollahi <i>et al.</i> , 2010
<i>Microcorys eremophiloides</i>	w	Australia	+	Cochrane <i>et al.</i> , 2002
<i>Salvia stenophylla</i>	w	South Africa	+	Musarurwa <i>et al.</i> , 2010

a, Annual herb; p, perennial herb; w, woody; +, positive response; -, negative response; 0, neutral response; MB, Mediterranean Basin; Cal, California, USA; USA, United States of America.

germination in this region (Keeley, 1995; Keeley and Baer-Keeley, 1999; Buhk and Hensen, 2006), until more evidence was reported for Mediterranean Basin flora in recent years (Moreira *et al.*, 2010; Çatav *et al.*, 2012). Physiological seed dormancy (Baskin and Baskin, 2004; Karlsson and Milberg, 2008) and a water-permeable seed coat (Moreira *et al.*, 2010) are characteristics of the Lamiaceae family. Thus, the chemical compounds in smoke responsible for the stimulation of germination (such as karrikinolides) can pass the seed coat and break dormancy and/or improve germination in Lamiaceae species. Smoke-enhanced germination in a number of Lamiaceae species was previously demonstrated in fire-prone environments of the Mediterranean Basin (Moreira *et al.*, 2010) and California (Keeley and Fotheringham, 1998). Our results also support the hypothesis that germination in species of the Mediterranean Lamiaceae is improved by the existence of smoke chemicals. The Lamiaceae is one of the families in which a seeder mechanism is found in the Mediterranean Basin, as a result of phylogenetic clustering of fire-adapted traits (Verdú and Pausas, 2007). Fire-enhanced germination is one of these traits that allow Lamiaceae species to persist in frequently burned Mediterranean habitats. Indeed, 11 out of the 17 tested Lamiaceae species from Mediterranean ecosystems (including the present study) give a positive germination response to smoke (Table 4), while only five have no response and one has a negative response. On the other hand, studies performed with Lamiaceae species from outside the Mediterranean-type ecosystems showed a similar pattern ($\chi^2 = 2.1$, $df = 2$, $P > 0.05$) with 9 positive, 7 neutral and 4 negative responses (Table 4). This suggests that smoke-enhanced germination in Lamiaceae is not restricted to Mediterranean-type ecosystems; however, more studies are needed to clarify this conclusion.

Lower concentrations of smoke solutions improved germination in *T. lamiifolium* ssp. *stachyophyllum*, *O. onites* and *P. bourgaei*, while the concentrated solution (1:1) had no effect on the germination percentage in these taxa. Conversely, *S. cretica* ssp. *smyrnaea* positively responded to the concentrated smoke solution but not to the 1:100 smoke solution treatment. This suggests the presence of a species-specific germination response to aqueous smoke solutions with different concentrations. This supports the suggestion of Keeley *et al.* (2012) that reporting a lack of smoke-stimulated germination is inconclusive 'without conducting an experiment over a large concentration gradient'.

In conclusion, our results contribute to the current limited knowledge on the smoke-enhanced germination in Mediterranean Lamiaceae, and support the idea that smoke is an important germination cue for this family.

Acknowledgements

The seeds were collected in the field with the permission of the Ministry of Food, Agriculture and Livestock of Republic of Turkey (no. 1379, dated 9 March 2012). The experiments were conducted in the laboratories of the Department of Biology, Muğla Sıtkı Koçman University, Turkey. We thank Costas A. Thanos and three anonymous reviewers for their comments, which helped improve the manuscript.

Financial support

The fieldwork of this study was supported by Hacettepe University Scientific Researches Unit (project no. 012.D06.601.0013).

Conflicts of interest

None.

References

- Abdollahi, M., Aghavaisi, B., Asl, A.M. and Babolhvaeji, H. (2010) Comparative effects of plant-derived smoke and potassium nitrate on germination and post-germination parameters of four medicinal species. *Planta Medica* **76**, P386, DOI: 10.1055/s-0030-1264684.
- Adkins, S.W. and Peters, N.C.B. (2001) Smoke derived from burnt vegetation stimulates germination of arable weeds. *Seed Science Research* **11**, 213–222.
- Ainsley, P.J., Jones, M.K. and Erickson, T.E. (2008) Overcoming physiological dormancy in *Prostanthera eurybioides* (Lamiaceae), a nationally endangered Australian shrub species. *Australian Journal of Botany* **56**, 214–219.
- Baskin, J.M. and Baskin, C.C. (2004) A classification system for seed dormancy. *Seed Science Research* **14**, 1–16.
- Bond, W.J. and Keeley, J.E. (2005) Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems. *Trends in Ecology and Evolution* **20**, 387–394.
- Brown, J.K. and Smith, J.K. (2000) *Wildland fire in ecosystems: effects of fire on flora*. General Technical Report RMRS-GTR-42, Vol. 2. USDA Forest Service, Rocky Mountain Research Station, Ogden, Utah.
- Brown, N.A.C. (1993a) Promotion of germination of fynbos seeds by plant-derived smoke. *New Phytologist* **123**, 575–583.
- Brown, N.A.C. (1993b) Seed germination in the fynbos fire ephemeral, *Syncarpha vestita* (L.) B. Nord. is promoted by smoke, aqueous extracts of smoke and charred wood derived from burning the ericoid-leaved shrub, *Passerina vulgaris* Thoday. *International Journal of Wildland Fire* **3**, 203–206.
- Brown, N.A.C. and Van Staden, J. (1997) Smoke as a germination cue: a review. *Plant Growth Regulation* **22**, 115–124.
- Buhk, C. and Hensen, I. (2006) 'Fire seeders' during early post-fire succession and their quantitative importance in

- south-eastern Spain. *Journal of Arid Environments* **66**, 193–209.
- Çatav, Ş.S., Bekar, İ., Ateş, B.S., Ergan, G., Oymak, F., Ülker, E.D. and Tavşanoğlu, Ç. (2012) Germination response of five eastern Mediterranean woody species to smoke solutions derived from various plants. *Turkish Journal of Botany* **36**, 480–487.
- Chou, Y.F., Cox, R.D. and Wester, D.B. (2012) Smoke water and heat shock influence germination of shortgrass prairie species. *Rangeland Ecology and Management* **65**, 260–267.
- Clarke, P., Davison, E.A. and Fulloon, L. (2000) Germination and dormancy of grassy woodland and forest species: effects of smoke, heat, darkness and cold. *Australian Journal of Botany* **48**, 687–699.
- Cochrane, A., Kelly, A., Brown, K. and Cunneen, S. (2002) Relationships between seed germination requirements and ecophysiological characteristics aid the recovery of threatened native plant species in western Australia. *Ecological Management and Restoration* **3**, 47–60.
- Crosti, R., Ladd, P.G., Dixon, K.W. and Piotto, B. (2006) Post-fire germination: The effect of smoke on seeds of selected species from the central Mediterranean basin. *Forest Ecology and Management* **221**, 306–312.
- Davis, P.H. (Ed.) (1965–1985) *Flora of Turkey and the East Aegean Islands, Vols 1–9*. Edinburgh, Edinburgh University Press.
- Daws, M.I., Davies, J., Pritchard, H.W., Brown, N.A.C. and Van Staden, J. (2007) Butenolide from plant-derived smoke enhances germination and seedling growth of arable weed species. *Plant Growth Regulation* **51**, 73–82.
- De Luis, M., Verdú, M. and Raventós, J. (2008) Early to rise makes a plant healthy, wealthy, and wise. *Ecology* **89**, 3061–3071.
- Dixon, K.W., Roche, S. and Pate, J.S. (1995) The promotive effect of smoke derived from burnt native vegetation on seed germination of Western Australian plants. *Oecologia* **101**, 185–192.
- Ervin S.L., Daugherty M., Coutant N.E. and Coons J.M. (2010) Influence of smoke solutions on the germination of twenty prairie plant species. Available from <http://www.eiu.edu/biology/posters/2010-04.pdf> (accessed 7 May 2014).
- Gürdal, B. and Kültür, Ş. (2013) An ethnobotanical study of medicinal plants in Marmaris (Muğla, Turkey). *Journal of Ethnopharmacology* **146**, 113–126.
- Herranz, J.M., Ferrandis, P. and Martínez-Sánchez, J.J. (1998) Influence of heat on seed germination of seven Mediterranean Leguminosae species. *Plant Ecology* **136**, 95–103.
- Jäger, A.K., Light, M.E. and Van Staden, J. (1996) Effects of source of plant material and temperature on the production of smoke extracts that promote germination of light-sensitive lettuce seeds. *Environmental and Experimental Botany* **36**, 421–429.
- Jefferson, L.V., Pennacchio, M., Havens, K., Forsberg, B., Sollenberger, D. and Ault, J. (2008) Ex situ germination responses of Midwestern USA prairie species to plant-derived smoke. *American Midland Naturalist* **159**, 251–256.
- Karlsson, L.M. and Milberg, P. (2008) Variation within species and inter-species comparison of seed dormancy and germination of four annual *Lamium* species. *Flora – Morphology, Distribution, Functional Ecology of Plants* **203**, 409–420.
- Kavgacı, A., Čarni, A., Başaran, S., Başaran, M.A., Košir, P., Marinšek, A. and Šilc, U. (2010) Long-term post-fire succession of *Pinus brutia* forest in the east Mediterranean. *International Journal of Wildland Fire* **19**, 599–605.
- Kazanis, D. and Arianoutsou, M. (2004) Long-term post-fire vegetation dynamics in *Pinus halepensis* forests of Central Greece: A functional group approach. *Plant Ecology* **171**, 101–121.
- Keeley, J.E. (1995) Seed-germination patterns in fire-prone Mediterranean-climate regions. pp. 239–273 in Arroyo, M.T.K.; Zedler, P.H.; Fox, M.D. (Eds) *Ecology and biogeography of Mediterranean ecosystems in Chile, California and Australia*. New York, Springer-Verlag.
- Keeley, J.E. and Baer-Keeley, M. (1999) Role of charred wood, heat-shock, and light in germination of postfire phrygana species from the eastern Mediterranean basin. *Israel Journal of Plant Sciences* **47**, 11–16.
- Keeley, J.E. and Bond, W.J. (1997) Convergent seed germination in South African fynbos and Californian chaparral. *Plant Ecology* **133**, 153–167.
- Keeley, J.E. and Fotheringham, C.J. (1998) Smoke-induced seed germination in California chaparral. *Ecology* **79**, 2320–2336.
- Keeley, J.E. and Zedler, P.H. (1978) Reproduction of chaparral shrubs after fire: a comparison of sprouting and seeding strategies. *American Midland Naturalist* **99**, 142–161.
- Keeley, J.E., Bond, W.J., Bradstock, R.A., Pausas, J.G. and Rundel, P.W. (2012) *Fire in Mediterranean ecosystems: Ecology, evolution and management*. Cambridge, Cambridge University Press.
- Luna, B., Pérez, B., Torres, I. and Moreno, J.M. (2012) Effects of incubation temperature on seed germination of Mediterranean plants with different geographical distribution ranges. *Folia Geobotanica* **47**, 17–27.
- Moreira, B., Tormo, J., Estrelles, E. and Pausas, J.G. (2010) Disentangling the role of heat and smoke as germination cues in Mediterranean Basin flora. *Annals of Botany* **105**, 627–635.
- Moreira, B., Tavşanoğlu, Ç. and Pausas, J.G. (2012) Local versus regional intraspecific variability in regeneration traits. *Oecologia* **168**, 671–677.
- Musarurwa, H.T., Van Staden, J. and Makunga, N.P. (2010) In vitro seed germination and cultivation of the aromatic medicinal *Salvia stenophylla* (Burch. ex Benth.) provides an alternative source of α -bisabolol. *Plant Growth Regulation* **61**, 287–295.
- Naghbi, F., Mosaddegh, M., Motamed, S.M. and Ghorbani, A. (2005) Labiatae family in folk medicine in Iran: from ethnobotany to pharmacology. *Iranian Journal of Pharmaceutical Research* **2**, 63–79.
- Paula, S., Arianoutsou, M., Kazanis, D., Tavşanoğlu, Ç., Lloret, F., Buhk, C., Ojeda, F., Luna, B., Moreno, J.M., Rodrigo, A., Espelta, J.M., Palacio, S., Fernández-Santos, B., Fernandes, P.M. and Pausas, J.G. (2009) Fire-related traits for plant species of the Mediterranean Basin. *Ecology* **90**, 1420.
- Pausas, J.G. and Verdú, M. (2005) Plant persistence traits in fire-prone ecosystems of the Mediterranean basin: a phylogenetic approach. *Oikos* **109**, 196–202.

- Pennacchio, M., Jefferson, L.V. and Havens, K.** (2005) Smoke: promoting germination of a tallgrass prairie species. *Chicago Wilderness Journal* **3**, 14–19.
- Pierce, S.M., Esler, K. and Cowling, R.M.** (1995) Smoke-induced germination of succulents (Mesembryanthemaceae) from fire-prone and fire-free habitats in South Africa. *Oecologia* **102**, 520–522.
- Reyes, O. and Trabaud, L.** (2009) Germination behaviour of 14 Mediterranean species in relation to fire factors: smoke and heat. *Plant Ecology* **202**, 113–121.
- Roche, S., Koch, J.M. and Dixon, K.W.** (1997) Smoke enhanced seed germination for mine rehabilitation in the southwest of western Australia. *Restoration Ecology* **5**, 191–203.
- Schwilk, D.W. and Zavala, N.** (2012) Germination response of grassland species to plant-derived smoke. *Journal of Arid Environments* **79**, 111–115.
- Simpson, M.G.** (2010) *Plant systematics*. Amsterdam, Elsevier.
- Small, E.** (2006) *Culinary herbs*. Canada, NRC Research Press.
- Takhtajan, A.** (2009) *Flowering plants* (2nd edition). Berlin, Springer-Verlag.
- Tang, Y., Boulter, S.L. and Kitching, R.L.** (2003) Heat and smoke effects on the germination of seeds from soil seed banks across forest edges between subtropical rainforest and eucalypt forest at Lamington National Park, south-eastern Queensland, Australia. *Australian Journal of Botany* **51**, 227–237.
- Tavşanoğlu, Ç. and Gürkan, B.** (2014) Long-term post-fire dynamics of co-occurring woody species in *Pinus brutia* forests: the role of regeneration mode. *Plant Ecology* **215**, 355–365.
- Tavşanoğlu, Ç., Çatav, Ş.S. and Özüdoğru, B.** Fire in grasslands: Germination and early seedling growth in central Anatolian steppe (in prep.).
- Thanos, C.A., Georghiou, K., Douma, D.J. and Marangaki, C.J.** (1991) Photoinhibition of seed germination in Mediterranean maritime plants. *Annals of Botany* **68**, 469–475.
- Thanos, C.A., Georghiou, K., Kadis, C. and Pantazi, C.** (1992) Cistaceae: A plant family with hard seeds. *Israel Journal of Botany* **41**, 251–263.
- Tierney, D.A.** (2006) The effect of fire-related germination cues on the germination of a declining forest understorey species. *Australian Journal of Botany* **54**, 297–303.
- Todorović, S., Živković, S., Giba, Z., Grubišić, D. and Mišić, D.** (2007) Basic seed germination characteristics of the endemic species *Nepeta rtanjensis* (Lamiaceae). *Plant Species Biology* **22**, 205–210.
- Tompsett, P.B. and Pritchard, H.W.** (1998) The effect of chilling and moisture status on the germination, desiccation tolerance and longevity of *Aesculus hippocastanum* L. seed. *Annals of Botany* **82**, 249–261.
- Van Staden, J., Brown, N.A.C., Jäger, A.K. and Johnson, T.A.** (2000) Smoke as a germination cue. *Plant Species Biology* **15**, 167–178.
- Verdú, M. and Pausas, J.G.** (2007) Fire drives phylogenetic clustering in Mediterranean Basin woody plant communities. *Journal of Ecology* **95**, 1316–1323.
- Yeşilyurt, E.B. and Akaydin, G.** (2012) Endemic plants and their threat categories of Muğla province (Turkey). *Hacettepe Journal of Biology and Chemistry* **40**, 195–212.
- Zuloaga-Aguilar, S., Briones, O. and Orozco-Segovia, A.** (2011) Seed germination of montane forest species in response to ash, smoke and heat shock in Mexico. *Acta Oecologica* **37**, 256–262.