

Research Article

Cite this article: Coffman W, Barber T, Norsworthy JK, Kruger GR (2021) Effect of dicamba rate and application parameters on protoporphyrinogen oxidase inhibitor-resistant and -susceptible Palmer amaranth (*Amaranthus palmeri*) control. *Weed Technol.* **35**: 22–26. doi: [10.1017/wet.2020.59](https://doi.org/10.1017/wet.2020.59)

Received: 8 October 2019
Revised: 13 May 2020
Accepted: 24 May 2020
First published online: 9 June 2020

Associate Editor:

Prashant Jha, Iowa State University

Nomenclature:

Dicamba; Palmer amaranth, *Amaranthus palmeri* S. Wats.; cotton, *Gossypium hirsutum* L.; soybean, *Glycine max* (L.) Merr

Keywords:

Herbicide resistance; dicamba; PPO-resistance; nozzle type; carrier volume

Author for correspondence:

Wyatt Coffman, 1724 W. Altheimer Dr., Fayetteville, AR 72704.
Email: wcoffman@uark.edu

Effect of dicamba rate and application parameters on protoporphyrinogen oxidase inhibitor-resistant and -susceptible Palmer amaranth (*Amaranthus palmeri*) control

Wyatt Coffman¹ , Tom Barber², Jason K. Norsworthy³ and Greg R. Kruger⁴

¹Former Graduate Research Assistant, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA; ²Professor, Department of Crop, Soil, and Environmental Sciences, University of Arkansas Lonoke Agricultural Center, Lonoke, AR, USA; ³Professor and Elms Farming Chair of Weed Science, Department of Crop, Soil, and Environmental Sciences, University of Arkansas, Fayetteville, AR, USA and ⁴Associate Professor, Department of Agronomy and Horticulture, University of Nebraska-Lincoln, North Platte, NE, USA

Abstract

Throughout eastern Arkansas, Palmer amaranth resistant to protoporphyrinogen oxidase (PPO)-inhibiting herbicides (Group 14 herbicides) has become widespread. Most PPO-resistant Palmer amaranth biotypes possess a target-site mutation, but a metabolic resistance mechanism to fomesafen (Group 14) has also been identified. Once metabolic resistance manifests, plants may also be tolerant to other herbicides and sites of action. To evaluate whether varying spray parameters affected control of PPO-resistant Palmer amaranth in dicamba-tolerant crops, field trials were conducted in 2017 and 2018 at the Lon Mann Cotton Research Station near Marianna, AR, and on-farm in Marion, AR. The experiment included split plot factors of dicamba rate, nozzle type, and carrier volume, with a whole plot factor of population. Dicamba was applied at 560 or 1120 g ae ha⁻¹ through 110015 TTI or AirMix nozzles at 70 or 140 L ha⁻¹ to PPO-resistant or PPO-susceptible Palmer amaranth. Palmer amaranth control 14 d after treatment (DAT) was influenced by an interaction between population and carrier volume. PPO-resistant Palmer amaranth control 14 DAT was 81% regardless of carrier volume, compared with 90% and 95% control at 70 and 140 L ha⁻¹, respectively, of the PPO-susceptible population. An interaction between nozzle type and carrier volume influenced Palmer amaranth control 21 DAT, whereas AirMix nozzles at 140 L ha⁻¹ controlled Palmer amaranth at a greater level (94%) than any other nozzle and carrier volume combination (≤90%). An interaction between population and dicamba rate influenced the relative density of Palmer amaranth 21 DAT. PPO-resistant Palmer amaranth density was less affected by dicamba at either rate than PPO-susceptible Palmer amaranth, relative to the non-treated check. Results concur with those of other research that suggest PPO-resistant Palmer amaranth is harder to control with dicamba. Otherwise, increasing carrier volume affected overall Palmer amaranth control to a greater degree than any other factor.

Introduction

Protoporphyrinogen oxidase (PPO)-inhibiting herbicide-resistant Palmer amaranth was first confirmed in 2011 and is now widespread throughout the crop-producing region of Arkansas (Salas et al. 2016; Varanasi et al. 2018b). The resistant populations in this area mostly possess a target-site resistance to all PPO-inhibiting herbicides, as well as resistance to other common herbicides such as glyphosate and acetolactate synthase (ALS)-inhibiting chemistries (Heap 2019; Varanasi et al. 2018b). Some populations of PPO-resistant Palmer amaranth have been noted as being harder to control with other herbicides that are effective on PPO-susceptible Palmer amaranth (Houston et al. 2019; Schwartz-Lazaro et al. 2017; Steckel 2018).

In 2018, metabolic resistance of Palmer amaranth to fomesafen was confirmed in Arkansas (Varanasi et al. 2018a). Subsequently, metabolic resistance to the very-long-chain fatty acid inhibitor *S*-metolachlor was also identified in Arkansas (Brabham et al. 2019). While no mechanisms of resistance to dicamba have been identified in Arkansas Palmer amaranth, the discovery of metabolic resistance mechanisms to other herbicides in Arkansas suggests that metabolic resistance to more herbicide sites of action could be building (Yu and Powles 2014).

Dicamba-resistant cotton was released for commercial use in 2015 and dicamba-resistant soybean was released in 2016. With the release of these new varieties, certain label restrictions were required for the products approved for use in these cropping systems to limit the off-target movement of dicamba to sensitive species. These limitations include nozzle type and spray volume specifications, among others (Anonymous 2018a, 2018b).

© The Author(s), 2020. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Herbicide application is influenced by application pressure, orifice size, nozzle design, and characteristics of the spray solution. The droplet sizes a nozzle produces are commonly classified by the volume median diameter (VMD), or D_{v50} of spray droplets, which is the value of the median size of spray droplets produced (i.e., 50% of droplets are larger and 50% are smaller than this value). Increasing VMD can contribute to decreased particle drift when herbicides are applied, but in turn, it can decrease the efficacy of some herbicides (Meyer et al. 2016). Another way to classify droplets produced by a nozzle is by examining the relative span (RS) of the droplet spectrum. The RS is a unitless measurement that represents the total variation in droplet sizes produced by a nozzle, where a smaller number indicates less variation in droplet size. Herbicide droplet size can also be affected by the product being applied in the spray solution (Mueller and Womac 1997). When comparing the VMD of applications of glyphosate, glufosinate, and paraquat, Etheridge et al. (1999) determined that a smaller VMD was generated by glufosinate than the other two chemicals. Chemical mixtures can also play a role in altering the VMD of a spray solution. When glufosinate was applied alone with a Turbo TeeJet Induction (TTI) 11004 nozzle, a VMD of 617 μm was produced, but when glufosinate was mixed with glyphosate and dicamba and applied with the same nozzle type, a VMD of 877 μm was produced (Meyer et al. 2015).

Nozzles are designed to control spray angle, spray pattern, droplet size, and solution flow rate as precisely as possible. Nozzles are available that produce a variety of spray patterns, in a variety of orifice sizes (Anonymous 2014). Increased droplet size can be obtained by increasing the orifice size for any given nozzle (Nuyttens et al. 2007). In order to increase droplet size without altering orifice size or spray pressure, nozzles with an inlet above the orifice are produced. These are typically referred to as air induction or venturi-type nozzles and work by essentially impregnating spray droplets with air, making them larger and less likely to drift (Etheridge et al. 1999, 2001).

When water is sprayed through an AirMix 110015 nozzle at 276 kPa, a droplet size classification of medium (VMD 236–340 μm) is produced. At the same pressure with a TTI 110015 nozzle, a droplet size classification of ultra-coarse (VMD >665 μm) is produced. Daggupati (2007) found that AirMix 11003 nozzles covered 2.8, 4.6, and 6.9 percentage points more total ground area than TTI 11003 nozzles at 207, 276, and 344 kPa, respectively. Meyer et al. (2015) demonstrated that mixtures of dicamba and glyphosate do not vary from droplet size classifications obtained with water for two Venturi-type nozzles, one specifically being the TTI nozzle. Meyer et al. (2015) also found that increasing carrier volume from 94 to 187 L ha⁻¹ increased spray coverage of a dicamba + glyphosate solution by 7% when averaged over three nozzle types.

Although not as important for the control of horizontally structured broadleaf weeds, smaller droplets adhere better to upright grasses, and therefore provide better control (Etheridge et al. 2001; McKinlay et al. 1974). Droplet size also plays a vital role in control levels provided by contact herbicides. When glufosinate and paraquat were applied to broadleaf signalgrass [*Urochloa platyphylla* (Munro ex C. Wright) R.D. Webster] and common cocklebur (*Xanthium strumarium* L.) with air induction (AI) nozzles (coarser droplets) and flat-fan nozzles (finer droplets), decreased control was noted in treatments where AI nozzles were used (Etheridge et al. 2001). McKinlay et al. (1974) observed decreased paraquat efficacy on common sunflower (*Helianthus annuus* L.) as VMD increased. Meyer et al. (2015) also observed

a decrease in control of Palmer amaranth, hemp sesbania [*Sesbania herbacea* (Mill.) McVaugh], velvetleaf (*Abutilon theophrasti* Medik.), and barnyardgrass [*Echinochloa crus-galli* (L.) P. Beauv.] with glufosinate as droplet size increased. Conflicting conclusions exist on the effect of droplet size and synthetic auxin efficacy. 2,4-D efficacy has been shown to decrease with increases in VMD (McKinlay et al. 1972). These are similar findings to those reported by Ennis and Williamson (1963) and Way (1969), who observed that synthetic auxin efficacy increased as droplet size decreased. Meyer et al. (2015), however, noted no difference in efficacy of dicamba on Palmer amaranth, hemp sesbania, velvetleaf, and prickly sida (*Sida spinosa* L.) across VMD values ranging from 340 to 756 μm .

Another factor that can influence efficacy of a foliar-applied herbicide is the carrier volume, or amount of herbicide solution being applied per hectare (Knoche 1994). Creech et al. (2015) observed no difference in control of *Amaranthus* spp. when glyphosate was applied at 70, 94, 140, and 187 L ha⁻¹. However, in the same study, efficacy of 2,4-D on *Amaranthus* spp. and soybean increased with increases in carrier volume (Creech et al. 2015), which is similar to findings by Smith (1946). When dicamba was applied postemergence to actively growing weeds, Butts et al. (2018) observed a greater effect of droplet size on weed mortality with a carrier volume of 47 L ha⁻¹ than when dicamba was applied at a carrier volume of 187 L ha⁻¹. Because weed control can be affected by a variety of application factors and a metabolic resistance mechanism has been discovered in Arkansas Palmer amaranth, the objective of this research was to determine whether there were differences in control of Palmer amaranth between two populations when spray parameters were varied.

Materials and Methods

On-farm field experiments were conducted in Marion, AR, on a Dubbs silt loam (fine-silty, mixed, active, thermic Typic Hapludalfs) with a PPO-resistant population of Palmer amaranth and at the Lon Mann Cotton Research Station (LMCRS) near Marianna, AR, on a Zachary soil (fine-silty, mixed, active, thermic Typic Albaqualfs) with a PPO-susceptible population of Palmer amaranth in 2017 and 2018. The objective was to compare the efficacy of dicamba on two populations of Palmer amaranth when it is applied according to varying spray parameters. No crop was planted at either location in 2017. In 2018, cotton cultivar Deltapine® 1518B2XF (Bayer CropScience, St. Louis, MO) was planted at both locations at 9.8 seeds m⁻¹ of row with 96-cm row spacing to provide a crop canopy. Because cotton grows slowly in the early vegetative stages (personal observation) and no pre-emergence herbicide was used, cotton canopy had no impact on Palmer amaranth emergence or control. Plots for all experiments were 3.9 m wide by 9.1 m long, with only the center 1.95 m receiving herbicide applications, creating a weedy check between plots.

Experiments were designed as a simple randomized complete block with three factors for each location in each year. Factors evaluated were nozzle type, carrier volume, and dicamba rate, with a nontreated control included for the basis of comparison. The previously mentioned factors were considered fixed effects, while block and year were considered random effects. When population is taken into account as a fixed effect, the experiment becomes a split plot, four-factor factorial, with population being the whole plot factor and nozzle type, carrier volume, and dicamba rate being the split plot factors.

Table 1. Mean spray characteristics as influenced by dicamba rate, nozzle type, and carrier volume.^{a,b,c,d}

Dicamba rate	Nozzle	Carrier volume	Droplet spectra parameters							
			D _{v10}		D _{v50}		D _{v90}		Relative span	
g ae ha ⁻¹		L ha ⁻¹	µm	SE	µm	SE	µm	SE	-	SE
560	TTI	70	350	1	683	2	984	2	0.93	0.00
560	AirMix	70	156	1	342	2	565	2	1.20	0.01
560	TTI	140	383	2	720	1	1048	4	0.92	0.01
560	AirMix	140	170	1	362	1	579	1	1.13	0.00
1,120	TTI	70	357	2	692	1	994	4	0.92	0.00
1,120	AirMix	70	151	0	336	0	553	4	1.20	0.01
1,120	TTI	140	381	1	734	4	1076	10	0.95	0.01
1,120	AirMix	140	166	1	358	1	570	4	1.12	0.00

^aData are reported as means followed by the standard error of the mean.

^bAll treatments contained glyphosate at a rate of 870 g ae ha⁻¹

^cAbbreviations: D_{v10}, 10% of droplets are smaller than this value; D_{v50}, 50% of droplets are smaller than this value; D_{v90}, 90% of droplets are smaller than this value; SE, standard error; TTI, Turbo TeeJet Induction.

^dAll nozzles used were 110015 orifice size.

All herbicide treatments were applied to 15- to 20-cm tall Palmer amaranth using a Bowman Mudmaster (Bowman Manufacturing, Newport, AR) calibrated to deliver 140 L ha⁻¹ at 4.8 km h⁻¹ or 70 L ha⁻¹ at 9.6 km h⁻¹ with 276 kPa of pressure. Nozzle types evaluated were AirMix[®] 110015 (Greenleaf Technologies, Covington, LA) and TTI 110015 (TeeJet Technologies, Wheaton, IL), all at 48-cm nozzle spacing. Note that neither nozzle is approved for use on current dicamba labels (Anonymous 2018a, 2018b). The dicamba herbicide Engenia[®] (BASF Corporation, Florham Park, NJ) was applied at 560 g ae ha⁻¹ or 1,120 g ae ha⁻¹ in combination with glyphosate (Roundup PowerMAX[®] II herbicide, Bayer CropScience, St. Louis, MO) at 870 g ae ha⁻¹. At both sites, 80% to 90% of Palmer amaranth was GR (data not shown). Spray characteristics for each nozzle, herbicide, and carrier volume combination are displayed in Table 1. Plots were rated 14 and 21 days after application (DAA) for Palmer amaranth control on a scale of 0 to 100, with 0 being no Palmer amaranth injury and 100 being death of all Palmer amaranth. Densities of live Palmer amaranth m⁻² were also estimated at 21 DAA by counting the number of living Palmer amaranth in two 0.5-m⁻² quadrats placed randomly in each plot.

Droplet size spectra for each nozzle, carrier volume, and herbicide combination were analyzed in a low-speed wind tunnel at the University of Nebraska-Lincoln West Central Research and Extension Center in North Platte, NE. Laser diffraction was used to detect particle size distribution with a Sympatec Helos Vario KR particle size analyzer (Sympatec GmbH, Clausthal-Zellerfeld, Germany) equipped with an R7 lens. To analyze the width of the nozzle plume, a 121 linear actuator was used to move the nozzle across the laser. The laser was positioned 30 cm from the tip of the nozzle in a low-speed wind tunnel with speeds of 24 km h⁻¹ during testing. The same spray solutions were evaluated through the same nozzles in the wind tunnel that were evaluated in field experiments. Each treatment was replicated three times in accordance with American Society of Agricultural and Biological Engineers standard S572.1.

Statistical Analysis

Means were separated using ANOVA via the GLIMMIX procedure in SAS 9.4 (SAS Institute Inc., Cary, NC). A beta distribution was assumed for Palmer amaranth control and relative density of Palmer amaranth (Gbur et al. 2012). Palmer amaranth densities

Table 2. Significance of P-values for factor main effects and interactions for Palmer amaranth control and density averaged over site years.^{a,b}

Source	Control		
	14 DAT	21 DAT	Density 21 DAT
Population	0.1814	0.5210	<0.0001*
Nozzle type	0.0484*	0.1074	0.0963
Carrier volume	0.0063*	0.0097*	0.0463*
Dicamba rate	0.4704	0.0658	0.4155
Population × Nozzle type	0.4312	0.4918	0.4494
Population × Carrier volume	0.0014*	0.2257	0.3593
Population × Dicamba rate	0.7582	0.1126	0.0397*
Nozzle type × Carrier volume	0.0164*	0.0015*	0.1559
Nozzle type × Dicamba rate	0.3201	0.1764	0.5257
Carrier volume × Dicamba rate	0.6894	0.3024	0.5202
Population × Nozzle type × Carrier volume	0.0295*	0.3135	0.5936
Population × Nozzle type × Dicamba rate	0.7855	0.5186	0.7220
Population × Carrier volume × Dicamba rate	0.5732	0.1317	0.1375
Nozzle Type × Carrier volume × Dicamba rate	0.6853	0.7755	0.2041
Population × Nozzle type × Carrier volume × Dicamba rate	0.4567	0.4289	0.3226

^aAbbreviation: DAT, days after treatment.

^bAsterisks (*) indicate significant treatment effects.

were measured relative to the nontreated control to account for differences in natural weed density between experimental locations. Mean separation was based on Fisher's protected LSD ($P = 0.05$).

Results and Discussion

The effect of year was not significant for this experiment ($P = 0.4653$); therefore, data across years were analyzed together. Both Palmer amaranth control at 14 and 21 DAT and Palmer amaranth density 21 DAT were influenced by several two-way interactions and main effects (Table 2).

Palmer Amaranth Control

Palmer amaranth control 14 DAT was influenced by main effects of nozzle type and carrier volume; two-way interactions between

Table 3. Palmer amaranth control as influenced by significant interactions of population × nozzle type × carrier volume and nozzle type × carrier volume.^{a,b}

Factor	Control	
	14 DAT	21 DAT
	%	
Population × Nozzle type × Carrier volume		
Marion × AirMix × 70 L ha ⁻¹	83 ab	
Marion × AirMix × 140 L ha ⁻¹	83 b	
Marion × TTI × 70 L ha ⁻¹	82 b	
Marion × TTI × 140 L ha ⁻¹	80 b	
LMCRS × AirMix × 70 L ha ⁻¹	89 ab	
LMCRS × AirMix × 140 L ha ⁻¹	97 a	
LMCRS × TTI × 70 L ha ⁻¹	91 ab	
LMCRS × TTI × 140 L ha ⁻¹	92 ab	
Nozzle type × Carrier volume		
AirMix × 70 L ha ⁻¹		88 b
AirMix × 140 L ha ⁻¹		94 a
TTI × 70 L ha ⁻¹		90 b
TTI × 140 L ha ⁻¹		89 b

^aAbbreviations: DAT, days after treatment; LMCRS, Lon Mann Cotton Research Station near Marianna, AR.

^bMeans within a column followed by the same letter are not different according to Fisher's protected LSD at (P = 0.05). Means for nonsignificant interactions of Population × Carrier volume 21 DAT and Nozzle type × Carrier volume 14 DAT presented for informational purposes.

population and carrier volume, and nozzle type and carrier volume; and a three-way interaction between population, nozzle type, and carrier volume (Table 2). Means for the interaction between nozzle type and carrier volume 14 DAT were consistent with the later evaluation but are not displayed (Tables 2 and 3). Control of PPO-susceptible Palmer amaranth was 97% when dicamba was applied with AirMix nozzles at 140 L ha⁻¹, which was 14 percentage points higher than the same combination on PPO-resistant Palmer amaranth (Table 3). While the remaining three nozzle and carrier volume combinations do not control PPO-susceptible Palmer amaranth at different levels than any nozzle and carrier volume combination controls PPO-resistant Palmer amaranth, there is an overall trend of greater control of PPO-susceptible Palmer amaranth that can be observed (Table 3).

By 21 DAT, similar control was observed for the interaction of population × carrier volume, but control levels were not significantly different (Tables 2 and 3). The interaction between nozzle type and carrier volume was not significant at P = 0.05 14 DAT. There was a tendency (P = 0.0517), however, for Palmer amaranth control to be 5 to 6 percentage points higher with AirMix nozzles at 140 L ha⁻¹ than with other nozzle type and carrier volume combinations 14 DAT (Table 3). Greater control was likely observed with AirMix nozzles at 140 L ha⁻¹ because they produced smaller droplets (VMD = 360 μm) than TTI nozzles at the same carrier volume (VMD = 727 μm), and therefore provided greater coverage of the leaf surface (Table 1).

A main effect of carrier volume and an interaction between nozzle type and carrier volume were significant 21 DAT (Table 2). At this timing, applications made with AirMix nozzles at 140 L ha⁻¹ (VMD = 360 μm) controlled Palmer amaranth 94%, whereas applications made with TTI nozzles controlled Palmer amaranth 90% and 89% at 70 L ha⁻¹ (VMD = 688 μm) and 140 L ha⁻¹ (VMD = 727 μm), respectively (Tables 1 and 3). These results indicate that carrier volume was more important for Palmer amaranth control in this experiment when smaller droplets were being produced. Meyer et al. (2016) observed greater control of glyphosate-resistant Palmer amaranth with dicamba + glyphosate at 94 L ha⁻¹

Table 4. Palmer amaranth relative density 21 days after treatment as influenced by main effects of nozzle type and carrier volume, and the interaction of population × dicamba rate.^{a,b}

Factor	Density	
	% of nontreated	
Carrier volume		
70 L ha ⁻¹		12 A
140 L ha ⁻¹		9 B
Population × Dicamba rate		
Marion × 560 g ae ha ⁻¹		19 a
Marion × 1,120 g ae ha ⁻¹		14 b
LMCRS × 560 g ae ha ⁻¹		6 c
LMCRS × 1,120 g ae ha ⁻¹		7 c

^aDensities of Palmer amaranth in nontreated plots were as follows: Marion 2017, 31 plants m⁻²; LMCRS 2017, 40 plants m⁻²; Marion 2018, 51 plants m⁻²; LMCRS 2018, 36 plants m⁻².

^bAbbreviations: TTI, Turbo TeeJet Induction; LMCRS, Lon Mann Cotton Research Station near Marianna, AR.

^cMeans within a column followed by the same letter are not different according to Fisher's protected LSD at (P = 0.05).

(VMD = 385 μm) than at 187 L ha⁻¹ (VMD = 487 μm), but droplet size for the TTI nozzles was smaller than that observed here.

Palmer Amaranth Density

Relative densities of Palmer amaranth 21 DAT were influenced by main effects of population and carrier volume, and an interaction between population and dicamba rate (Table 2). Averaged over all other factors, densities of Palmer amaranth relative to the nontreated were 2 percentage points lower when dicamba was applied with AirMix nozzles (9%) than TTI nozzles (11%), suggesting the smaller droplets produced by AirMix nozzles (VMD = 336 to 362 μm), compared to TTI nozzles (VMD = 683 to 734 μm), probably increased dicamba absorption by the plants (Tables 1 and 4).

For the main effect of carrier volume, treatments applied at 140 L ha⁻¹ reduced Palmer amaranth densities to 9% relative to the nontreated, whereas treatments applied at 70 L ha⁻¹ reduced densities to 12% relative to the nontreated (Table 4). The influence of carrier volume suggests that applying dicamba at 140 L ha⁻¹ allows for greater coverage of the treated area than a carrier volume of 70 L ha⁻¹, again placing more dicamba on the leaf surface. The significant effects of nozzle type and carrier volume for relative Palmer amaranth density reflect the significant interaction between nozzle type and carrier volume for weed control 21 DAT, in that AirMix nozzles at 140 L ha⁻¹ provided greater control than all other nozzle type and carrier volume combinations due to greater coverage and smaller droplet sizes being produced (Tables 1, 3, and 4).

For the interaction between population and dicamba rate, the PPO-susceptible population at LMCRS was unaffected by dicamba rate. At this population, only 6% and 7% of treated Palmer amaranth, relative to the nontreated, survived dicamba application at 560 g ae ha⁻¹ and 1120 g ae ha⁻¹, respectively (Table 4). Dicamba at 560 g ae ha⁻¹ was likely so effective at LMCRS that no differences in density could be observed between the two rates. However, at Marion, with PPO-resistant Palmer amaranth, 19% of treated Palmer amaranth survived a dicamba application at 560 g ae ha⁻¹, and 14% of treated Palmer amaranth survived dicamba applied at 1,120 g ae ha⁻¹ (Table 4). Although the Palmer amaranth at Marion appeared to be controlled at comparable levels to those at LMCRS 21 DAT based on visible control ratings, relative density data indicate that Palmer amaranth at Marion was more difficult to kill.

Differences in weed densities between the two locations suggest that the Palmer amaranth population at Marion is more tolerant to dicamba than at LMCRS. Previous studies conducted at Marion also noted lower than expected control levels and density reduction with 560 g ae ha⁻¹ of dicamba (Houston et al. 2019). Schwartz-Lazaro et al. (2017) found that PPO-resistant Palmer amaranth populations from around the state of Arkansas were less sensitive to dicamba in the greenhouse than PPO-susceptible populations. These findings are not unlike other research that suggests that multiple postemergence applications of dicamba may be required to control PPO-resistant Palmer amaranth (Steckel 2018).

Practical Implications

In this experiment, carrier volume was the most important factor in Palmer amaranth control with dicamba. In general, treatments applied at a carrier volume of 140 L ha⁻¹ provided better control of Palmer amaranth than treatments applied at 70 L ha⁻¹, regardless of other factors. AirMix nozzles provided higher levels of Palmer amaranth mortality than did TTI nozzles, likely due to the smaller droplet size. However, current dicamba labels approved for post-emergence use state that dicamba must be applied through nozzles that produce extremely coarse or larger droplets for Engenia® and ultra-coarse droplets for Xtendimax™ (Anonymous 2018a, 2018b). By increasing carrier volume, applicators can mitigate reduced levels of weed control caused by using a nozzle producing coarser droplets.

Special attention should be paid to Palmer amaranth mortality with dicamba in fields where PPO-resistance is suspected. Metabolic resistance to S-metolachlor was recently confirmed in the Marion population and it is possible that this metabolic resistance could be the cause of reduced mortality of Palmer amaranth treated with dicamba at this location (Brabham et al. 2019). Because PPO-resistant populations have proven to be harder to control with dicamba, other weed control methods may need to be employed. Following best management practices to mitigate resistance is recommended to control PPO-resistant Palmer amaranth (Norsworthy et al. 2012).

Acknowledgments. No external funding was received to support this research. No conflicts of interest have been declared.

References

- Anonymous (2014) TeeJet® Technologies Product Catalog. TeeJet Publications Catalog 51A. 164 p
- Anonymous (2018a) Engenia® herbicide product label. BASF Publication No. 007969-00345.20180928e. Florham Park, NJ: BASF. 29 p
- Anonymous (2018b) Xtendimax™ application requirements—nozzles. <http://www.xtendimaxapplicationrequirements.com/Pages/nozzles.aspx>. Accessed: January 28, 2019
- Brabham C, Norsworthy JK, Houston MM, Varanasi VK, Barber T (2019) Confirmation of S-metolachlor resistance in Palmer amaranth (*Amaranthus palmeri*). *Weed Technol* 33:720–726
- Butts TR, Samples CA, Franca LX, Dodds DM, Reynolds DB, Adams JW, Zollinger RK, Howatt KA, Fritz BK, Hoffmann WC, Kruger GR (2018) Spray droplet size and carrier volume effect on dicamba and glufosinate efficacy. *Pest Manag Sci* 74:2020–2029
- Creech CF, Henry RS, Werle R, Sandell LD, Hewitt AJ, Kruger GR (2015) Performance of postemergence herbicides applied at different carrier volume rates. *Weed Technol* 29:611–624
- Daggupati NP (2007) Assessment of the varitarget nozzle for variable rate application of liquid crop protection products. MS thesis. Manhattan, KS: Kansas State University. 94 p
- Ennis WB, Williamson RE (1963) Influence of droplet size on effectiveness of low-volume herbicidal sprays. *Weeds* 11:67–72
- Etheridge RE, Hart WE, Hayes RM, Mueller TC (2001) Effect of venturi-type nozzles and application volume on postemergence herbicide efficacy. *Weed Technol* 15:75–80
- Etheridge RE, Womac AR, Mueller TC (1999) Characterization of the spray droplet spectra and patterns of four Venturi-type drift reduction nozzles. *Weed Technol* 13:765–770
- Gbur EE, Stroup WW, McCarter KS, Durham S, Young LJ, Christman M, West M, Kramer M (2012) Analysis of generalized linear mixed models in the agricultural and natural resources sciences. Madison, WI: American Society of Agronomy, Soil Science Society of America, Crop Science Society of America. 298 pp
- Heap I (2019) The International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org/>. Accessed: January 30, 2019
- Houston MM, Norsworthy JK, Barber T, Brabham C (2019) Field evaluation of preemergence and postemergence herbicides for control of protoporphyrinogen oxidase-resistant Palmer amaranth (*Amaranthus palmeri* S. Watson). *Weed Technol* 33:610–615
- Knoche M (1994) Effect of droplet size and carrier volume on performance of foliage-applied herbicides. *Crop Prot* 13:163–178
- McKinlay KS, Ashford R, Ford RJ (1974) Effects of drop size, spray volume, and dosage on paraquat toxicity. *Weed Sci* 22:31–34
- McKinlay KS, Brandt SA, Morse P, Ashford R (1972) Droplet size and phytotoxicity of herbicides. *Weed Sci* 20:450–452
- Meyer CJ, Norsworthy JK, Kruger GR, Barber LT (2015) Influence of droplet size on efficacy of the formulated products Engenia™, Roundup PowerMax®, and Liberty®. *Weed Technol* 29:641–652
- Meyer CJ, Norsworthy JK, Kruger GR, Barber LT (2016) Effect of nozzle selection and spray volume on droplet size and efficacy of Engenia tank-mix combinations. *Weed Technol* 30:377–390
- Mueller TC, Womac AR (1997) Effect of formulation and nozzle type on droplet size with isopropylamine and trimesium salts of glyphosate. *Weed Technol* 11:639–643
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Sci (SP1)* 60:31–62
- Nuytens D, Baetens K, De Schampheleire M, Sonck B (2007) Effect of nozzle type, size and pressure on spray droplet characteristics. *Biosyst Eng* 97:333–345
- Salas RA, Burgos NR, Tranel PJ, Singh S, Glasgow L, Scott RC, Nichols RL (2016) Resistance to PPO-inhibiting herbicides in Palmer amaranth from Arkansas. *Pest Manag Sci* 72:864–869
- Schwartz-Lazaro LM, Norsworthy JK, Scott RC, Barber LT (2017) Resistance of two Arkansas Palmer amaranth populations to multiple herbicide sites of action. *Crop Prot* 96:158–163
- Smith HH (1946) Quantitative aspects of aqueous spray applications of 2,4-D acid for herbicidal purposes. *Bot Gaz* 107:544–551
- Steckel LE (2018) Controlling multiple-resistant Palmer amaranth. <http://news.utcrops.com/2018/03/controlling-multiple-resistant-palmer-amaranth/>. Accessed: December 16, 2018
- Varanasi VK, Brabham C, Norsworthy JK (2018a) Confirmation and characterization of non-target site resistance to fomesafen in Palmer amaranth (*Amaranthus palmeri*). *Weed Sci* 66:702–709
- Varanasi VK, Brabham C, Norsworthy JK, Nie H, Young BG, Houston MH, Barber T, Scott RC (2018b) A statewide survey of PPO-inhibitor resistance and the prevalent target-site mechanisms in Palmer amaranth (*Amaranthus palmeri*) accessions from Arkansas. *Weed Sci* 66:149–158
- Way JM (1969) Some effects of droplet size and position on the response of lettuce to sub-lethal doses of MCPA. *Weed Res* 1:272–278
- Yu Q, Powles S (2014) Metabolism-based herbicide resistance and cross-resistance in crop weeds: a threat to herbicide sustainability and global crop production. *Plant Physiol* 166:1106–1118