

# THE PRO- $p$ -IWAHORI HECKE ALGEBRA OF A REDUCTIVE $p$ -ADIC GROUP III (SPHERICAL HECKE ALGEBRAS AND SUPERSINGULAR MODULES)

MARIE-FRANCE VIGNERAS

*UMR 7586, Institut de Mathématiques de Jussieu, 4 place Jussieu,  
Paris 75005, France* ([vigneras@math.jussieu.fr](mailto:vigneras@math.jussieu.fr))

(Received 26 May 2014; accepted 30 March 2015; first published online 3 June 2015)

*Abstract* Let  $R$  be a large field of characteristic  $p$ . We classify the supersingular simple modules of the pro- $p$ -Iwahori Hecke  $R$ -algebra  $\mathcal{H}$  of a general reductive  $p$ -adic group  $G$ . We show that the functor of pro- $p$ -Iwahori invariants behaves well when restricted to the representations compactly induced from an irreducible smooth  $R$ -representation  $\rho$  of a special parahoric subgroup  $K$  of  $G$ . We give an almost-isomorphism between the center of  $\mathcal{H}$  and the center of the spherical Hecke algebra  $\mathcal{H}(G, K, \rho)$ , and a Satake-type isomorphism for  $\mathcal{H}(G, K, \rho)$ . This generalizes results obtained by Ollivier for  $G$  split and  $K$  hyperspecial to  $G$  general and  $K$  special.

*Keywords:* group theory and generalizations; number theory

2010 *Mathematics subject classification:* Primary 20C08  
Secondary 22E50

## Contents

|          |   |            |
|----------|---|------------|
| <b>1</b> | <b>Introduction</b>   | <b>572</b> |
| <b>2</b> | <b>The characters of <math>\mathfrak{h}</math> and <math>\mathcal{H}^{\text{aff}}</math></b>        | <b>580</b> |
| <b>3</b> | <b>Distinguished representatives of <math>W_0 \backslash W</math></b>                               | <b>583</b> |
| <b>4</b> | <b><math>\mathfrak{h}</math>-eigenspace in <math>\eta \otimes_{\mathfrak{h}} \mathcal{H}</math></b> | <b>586</b> |
| <b>5</b> | <b>Centers</b>  | <b>592</b> |
| <b>6</b> | <b>Supersingular <math>\mathcal{H}</math>-modules</b>   | <b>594</b> |
| <b>7</b> | <b>Pro-<math>p</math>-Iwahori invariants and compact induction</b>                                  | <b>601</b> |
|          | <b>References</b>   | <b>607</b> |

### 1. Introduction

Let  $p$  be a prime number, let  $F$  be a finite extension of  $\mathbb{Q}_p$  or  $\mathbb{F}_p((T))$ , and let  $G$  be the group of rational points of a connected reductive  $F$ -group.

#### 1.1.

The smooth representations of  $G$  over an algebraically closed field  $C$  of characteristic  $p$  have been the subject of many investigations in recent years, in the modulo  $p$  Langlands program. The pro- $p$ -Iwahori invariant functor  $V \mapsto V^{I(1)}$  relates the representations of  $G$  to the modules of the pro- $p$ -Iwahori Hecke  $C$ -algebra  $\mathcal{H} = \mathcal{H}_C(G, I(1))$  studied in [13–15]. The  $I(1)$ -invariant functor and the theory of  $\mathcal{H}$ -modules play an increasingly important role in the representation theory of  $G$  modulo  $p$ . They are the key to the proof of the change of weight in the recent classification of irreducible smooth  $C$ -representations of  $G$  in terms of supersingular ones (a forthcoming work by Abe *et al.* [1]). The supersingular smooth irreducible  $C$ -representations  $\pi$  of  $G$  and their  $I(1)$ -invariant remain mysterious, but the supersingular simple  $\mathcal{H}$ -modules are classified in this paper, and the supersingularity of  $\pi^{I(1)}$  and of  $\pi$  are related. A variant of the modulo  $p$  Langlands program seems to exist for  $\mathcal{H}$ -modules. Grosse-Kloenne [5] constructed a functor from finite-dimensional  $\mathcal{H}_C(GL(n, \mathbb{Q}_p), I(1))$ -modules to finite-dimensional smooth  $C$ -representations of  $\text{Gal}_{\mathbb{Q}_p}$ , inducing a bijection between the simple supersingular  $\mathcal{H}_C(GL(n, F), I(1))$ -modules of dimension  $n$  and the irreducible smooth  $C$ -representations of  $\text{Gal}_F$  (the absolute Galois group of  $F$ ) of dimension  $n$  as in [9, 14].

In this paper, we prove that the  $I(1)$ -invariant functor behaves well when restricted to compactly induced representations  $\text{c-Ind}_K^G \rho$ , where  $\rho$  is an irreducible smooth  $C$ -representation of a special parahoric subgroup  $K$  of  $G$ . The vector space  $\rho^{I(1)}$  has dimension 1, and the pro- $p$ -Iwahori Hecke  $C$ -algebra  $\mathfrak{h} = H_C(K, I(1))$  of  $K$  acts on  $\rho^{I(1)}$  by a character  $\eta$ . The  $\mathcal{H}$ -module  $(\text{c-Ind}_K^G \rho)^{I(1)}$  is isomorphic to  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$ , and the spherical algebra  $\text{End}_{CG}(\text{c-Ind}_K^G \rho)$  is isomorphic to the algebra  $\text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H})$ . This paper is devoted to the study of the modules  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  and of the spherical Hecke algebras  $\text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H})$ . In the last section, we transfer our results from  $\mathcal{H}$  to the group  $G$  using the  $I(1)$ -invariant functor.

Let  $\rho$  be an irreducible smooth  $C$ -representation of  $K$ , and let  $\eta, \eta_1$  be two arbitrary characters of  $\mathfrak{h}$ . We obtain the following:

(i) *Isomorphisms*

$$(\text{c-Ind}_K^G \rho)^{I(1)} \simeq \rho^{I(1)} \otimes_{\mathfrak{h}} \mathcal{H}, \quad \text{End}_{CG}(\text{c-Ind}_K^G \rho) \simeq \text{End}_{\mathcal{H}}(\rho^{I(1)} \otimes_{\mathfrak{h}} \mathcal{H}).$$

(ii) *A Satake-type isomorphism for the spherical Hecke algebra  $\mathcal{H}(\mathfrak{h}, \eta) = \text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H})$ .*

(iii) *A basis of the space of intertwiners  $\text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H})$ .*

(iv) *An almost-isomorphism from the center of  $\mathcal{H}$  to the center of  $\mathcal{H}(\mathfrak{h}, \eta)$  (an isomorphism between finite index affine subalgebras).*

(v) *The classification of the supersingular simple  $\mathcal{H}$ -modules.*

When  $G$  is split and  $K$  hyperspecial, Ollivier proved (i), (ii), (iv) and (v). We follow her method. The alcove walk bases of  $\mathcal{H}$  and the product formula [12, 15] allow us to simplify her method and to extend it to  $G$  general and  $K$  special. Analogs of 2, 3 were proved for  $G$  in [6, 7] and 5 for  $G$  remains a wide-open question.

In the rest of this introduction, we consider the content of 2, 3, 4, 5.

After [13, 14], a generalization of  $\mathcal{H}_C(G, I(1))$  was introduced in [12] when  $G$  is split, and in [15] for  $G$  general, in order to study it. This is an algebra  $\mathcal{H}_R(q_s, c_{\bar{s}})$  over a commutative ring  $R$  with two sets of parameters  $(q_s), (c_{\bar{s}})$ . The properties of this algebra are often proved by reduction to  $(q_s) = (1)$  (this changes the parameters  $(c_{\bar{s}})$ ), and transferred to  $\mathcal{H}_R(0, c_{\bar{s}})$  by specialization to  $(q_s) = (0)$ . The algebra  $\mathcal{H}_R(q_s, c_{\bar{s}})$  contains a natural finite-dimensional subalgebra  $\mathfrak{h}_R(q_s, c_{\bar{s}})$ .

In 1.2 and 1.3, we recall the basic properties of  $\mathcal{H}_R(q_s, c_{\bar{s}})$  used in this work and the dictionary between  $\mathfrak{h}_R(q_s, c_{\bar{s}}), \mathcal{H}_R(q_s, c_{\bar{s}})$  and  $\mathcal{H}_R(K, I(1)), \mathcal{H}_R(G, I(1))$  [15, 16]. Theorems 1.2, 1.3, 1.4, and 1.5 are proved for  $\mathfrak{h}_R(0, c_{\bar{s}}), \mathcal{H}_R(0, c_{\bar{s}})$ , and are given in 1.4. They apply to the algebras  $\mathcal{H}_R(K, I(1)), \mathcal{H}_R(G, I(1))$  when  $R$  has characteristic  $p$ .

1.2.

Let  $\mathcal{W} = (\Sigma, \Delta, \Omega, \Lambda, \nu, W, Z_k, W(1))$  be data consisting of the following:

- (i) a reduced root system  $\Sigma$  of basis  $\Delta$  associated with the finite Weyl Coxeter system  $(W_0, S)$  of an affine Weyl Coxeter system  $(W^{\text{aff}}, S^{\text{aff}})$  acting on a real vector space  $V$  of dual of basis  $\Delta$ , with a  $W_0$ -invariant scalar product;
- (ii) three commutative groups,  $\Omega$  and  $\Lambda$  finitely generated, and  $Z_k$  finite;
- (iii) a group  $W = W^{\text{aff}} \rtimes \Omega = \Lambda \rtimes W_0$  which is a semi-direct product of subgroups in two different ways,  $\Omega$  acting on  $(W^{\text{aff}}, S^{\text{aff}})$  and  $W_0$  on  $\Lambda$ . The length  $\ell$  and the Bruhat order  $\leq$  of  $(W^{\text{aff}}, S^{\text{aff}})$  extend trivially to  $W = W^{\text{aff}} \rtimes \Omega$ ;
- (iv) a  $W_0$ -equivariant homomorphism  $\nu : \Lambda \rightarrow V$  such that the action of  $W^{\text{aff}}$  on  $V$  and the action of  $\Lambda$  on  $V$  by translation  $v \mapsto v + \nu(\lambda)$  for  $\lambda \in \Lambda, v \in V$ , extend to an action of  $W$  by affine automorphisms permuting the set of affine hyperplanes  $\mathfrak{H} = \{\text{Ker}(\alpha + n), | \alpha + n \in \Sigma^{\text{aff}} = \Sigma + \mathbb{Z}\}$ ;
- (v) a system of the representatives of  $W_0$  in  $\Lambda$ :

$$\Lambda^+ := \{\mu \in \Lambda \mid \nu(\mu) \in \overline{\mathfrak{D}}^+\},$$

where  $\overline{\mathfrak{D}}^+ = \{x \in V \mid 0 \leq \alpha(x), \alpha \in \Delta\}$  is the dominant closed Weyl chamber;

- (vi) an extension  $1 \rightarrow Z_k \rightarrow W(1) \rightarrow W \rightarrow 1$ .

*Notation.* The inverse image in  $W(1)$  of a subset  $X$  of  $W$  is denoted by  $X(1)$ , and  $\tilde{w}$  denotes an element of  $W(1)$  of image  $w \in W$ . For  $c \in R[Z_k]$ , the conjugate of  $c$  by  $\tilde{w}$  depends only on  $w$ , and is denoted  $w \bullet c := \tilde{w}c\tilde{w}^{-1}$ . The dominant Weyl chamber  $\mathfrak{D}^+ = \{x \in V \mid 0 < \alpha(x), \alpha \in \Delta\}$  is open. The dominant alcove  $\mathfrak{C}^+$  is the connected component  $\mathfrak{D}^+ \cap (V - \bigcup_{H \in \mathfrak{H}} H)$  of vertex  $0 \in V$ . The set  $\Sigma^{\text{aff}, +}$  of positive affine roots is the set of  $\gamma \in \Sigma^{\text{aff}}$  positive on  $\mathfrak{C}^+$ . The action of  $W$  on  $V$  defines by functoriality an action of  $W$  on  $\Sigma^{\text{aff}}$ .

We will often suppose that  $\Lambda$  contains a subgroup  $\Lambda_T$  satisfying the following.

(T1)  $\Lambda = \bigsqcup_{y \in Y} \Lambda_T y$  for a finite set  $Y$ .

(T2)  $\Lambda_T$  is  $W_0$ -stable.

(T3) There exists a central subgroup  $\tilde{\Lambda}_T$  of  $\Lambda(1)$  normalized by  $W_0(1)$  such that the quotient map  $\Lambda(1) \rightarrow \Lambda$  induces a group isomorphism  $\tilde{\Lambda}_T \xrightarrow{\sim} \Lambda_T$  sending  $\tilde{w}\tilde{\mu}\tilde{w}^{-1}$  to  $w\mu w^{-1}$  if  $\tilde{w} \in W_0(1)$  lifts  $w \in W_0$  and  $\tilde{\mu} \in \tilde{\Lambda}_T$  lifts  $\mu \in \Lambda_T$ .

Let  $(q_{\tilde{s}}, c_{\tilde{s}})_{\tilde{s} \in S^{\text{aff}}(1)}$  be a set of elements in  $R \times R[Z_k]$  satisfying  $q_{\tilde{s}'} = q_{\tilde{s}}$ ,  $c_{\tilde{s}'} = w \bullet c_{\tilde{s}}$  if  $\tilde{s}' = \tilde{w}\tilde{s}\tilde{w}^{-1} \in S^{\text{aff}}(1)$ ,  $\tilde{w} \in W(1)$ , and  $q_{t\tilde{s}} = q_{\tilde{s}}$ ,  $c_{t\tilde{s}} = tc_{\tilde{s}}$  if  $t \in Z_k$ . As  $q_{\tilde{s}}$  depends only on the image  $s \in S^{\text{aff}}$  of  $\tilde{s}$ , we denote also  $q_{\tilde{s}} = q_s$ .

There is a unique  $R$ -algebra  $\mathcal{H} = \mathcal{H}_R(\mathcal{W}, q_s, c_{\tilde{s}})$ , free of basis  $(T_{\tilde{w}})_{\tilde{w} \in W(1)}$ , with product satisfying

(i) the braid relations:

$$T_{\tilde{w}}T_{\tilde{w}'} = T_{\tilde{w}\tilde{w}'}, \quad \text{if } \tilde{w}, \tilde{w}' \in W(1), \ell(w) + \ell(w') = \ell(ww'), \tag{1}$$

allowing one to identify  $R[\Omega(1)]$  to a subalgebra of  $\mathcal{H}$ ;

(ii) the quadratic relations:

$$T_{\tilde{s}}T_{\tilde{s}}^* = q_s\tilde{s}^2, \quad \text{if } \tilde{s} \in S^{\text{aff}}(1), T_{\tilde{s}}^* = T_{\tilde{s}} - c_{\tilde{s}}. \tag{2}$$

This is called the Iwahori–Matsumoto presentation of  $\mathcal{H}_R(\mathcal{W}, q_s, c_{\tilde{s}})$ .

The  $R$ -submodule of basis  $(T_{\tilde{w}})_{\tilde{w} \in W_0(1)}$  is a finite subalgebra  $\mathfrak{h} = \mathfrak{h}_R(\mathcal{W}, q_s, c_{\tilde{s}})$ .

The  $R$ -submodule of basis  $(T_{\tilde{w}})_{\tilde{w} \in W^{\text{aff}}(1)}$  is a subalgebra  $\mathcal{H}^{\text{aff}}$ . The  $R$ -algebra  $\mathcal{H}^{\text{aff}}$  is an algebra like  $\mathcal{H}$  with  $\Omega$  trivial, and  $\mathcal{H}$  is isomorphic to the twisted tensor product

$$x \otimes y \mapsto xy : \mathcal{H}^{\text{aff}} \otimes'_{R[Z_k]} R[\Omega(1)] \rightarrow \mathcal{H} \tag{3}$$

of its subalgebras  $R[\Omega(1)]$  and  $\mathcal{H}^{\text{aff}}$ . The algebra  $\mathcal{H}$  admits an involutive  $R$ -automorphism  $\iota$ , equal to the identity on  $R[\Omega(1)]$  and such that [15, Proposition 4.23]

$$\iota(T_{\tilde{s}}) := -T_{\tilde{s}}^* \quad \text{for } s \in S^{\text{aff}}. \tag{4}$$

All the orientations that we consider are spherical [15]. For the orientation  $o$  associated to an (open) Weyl chamber  $\mathfrak{D}_o$ , the  $o$ -positive side of the affine hyperplane  $\text{Ker}(\alpha + n)$  is the set of  $x \in V$  where  $\alpha(x) + n > 0$ , if  $\alpha \in \Sigma$  takes positive values on  $\mathfrak{D}_o$ . The dominant orientation  $o$ , denoted by  $o^+$ , is associated to the dominant Weyl chamber  $\mathfrak{D}^+$ , and the anti-dominant orientation, denoted by  $o^-$ , to the anti-dominant Weyl chamber  $-\mathfrak{D}^+ = \mathfrak{D}^-$ . The orientation associated to the Weyl chamber  $w^{-1}(\mathfrak{D}_o)$ ,  $w \in W_0$ , is denoted by  $o \bullet w$ . For  $w \in W$  of projection  $w_0 \in W_0$ , the orientation  $o \bullet w_0$  is also denoted by  $o \bullet w$ . We have  $o \bullet \lambda = o$  for  $\lambda \in \Lambda$ . We set

$$S_o^{\text{aff}} := \{s \in S^{\text{aff}} \mid \mathfrak{C}^+ \text{ is in the } o\text{-positive side of } H_s\}, \quad S_o := S \cap S_o^{\text{aff}}, \tag{5}$$

where  $H_s$  is the affine hyperplane of  $V$  fixed by  $s$  and  $\mathfrak{C}^+$  the dominant alcove (Notation). There exists a unique set of bases  $(E_o(\tilde{w}))_{\tilde{w} \in W(1)}$  of  $\mathcal{H}$ , parameterized by

the orientations  $o$ , satisfying [15, §5.3]

$$E_o(\tilde{s}) := T_{\tilde{s}} \text{ if } s \in S^{\text{aff}} - S_o^{\text{aff}}, \quad E_o(\tilde{s}) := T_{\tilde{s}}^* \text{ if } s \in S_o^{\text{aff}}, \tag{6}$$

and the product formula, for  $\tilde{w}, \tilde{w}' \in W(1)$ ,

$$E_o(\tilde{w})E_{o \bullet w}(\tilde{w}') = E_o(\tilde{w}\tilde{w}') \quad \text{if } \ell(w) + \ell(w') = \ell(ww'). \tag{7}$$

In particular, for  $\tilde{\lambda}, \tilde{\lambda}' \in \Lambda(1)$ ,

$$E_o(\tilde{\lambda})E_o(\tilde{\lambda}') = E_o(\tilde{\lambda}\tilde{\lambda}') \quad \text{if } \nu(\lambda), \nu(\lambda') \text{ belong to a same closed Weyl chamber.} \tag{8}$$

We have  $E_o(\lambda) = T_{\lambda}$  when  $\nu(\lambda) \in \overline{D}_o$ .

The basis  $(E_o(\tilde{w}))_{\tilde{w} \in W(1)}$  is called an alcove walk basis; the alcove walk bases generalize the integral Bernstein bases defined in [11, 14].

The  $R$ -submodule of basis  $(E_o(\tilde{\lambda}))_{\tilde{\lambda} \in \Lambda(1)}$  is a subalgebra  $\mathcal{A}_o$  of  $\mathcal{H}$  containing the subalgebra  $\mathcal{A}_o^+$  of basis  $(E_o(\tilde{\lambda}))_{\tilde{\lambda} \in \Lambda^+(1)}$ , isomorphic to  $R[\Lambda^+(1)]$ .

If  $q_s = 0$  for all  $s \in S^{\text{aff}}$ , then for  $\tilde{w}, \tilde{w}' \in W(1)$  such that  $\ell(w) + \ell(w') > \ell(ww')$  we have  $E_o(\tilde{w})E_{o \bullet w}(\tilde{w}') = 0$ ; in particular,  $E_o(\tilde{\lambda})E_o(\tilde{\lambda}') = 0$  if  $\tilde{\lambda}, \tilde{\lambda}' \in \Lambda(1)$ , and  $\nu(\lambda), \nu(\lambda')$  do not belong to the same closed Weyl chamber.

**1.3.**

Let  $F$  be a local field of finite residue field  $k$  with  $q$  elements and of characteristic  $p$ , and  $p_F$  a generator of the maximal ideal of the ring of integers  $\mathcal{O}_F$  of  $F$ . Let  $G, T, Z$ , and  $N$  be respectively the  $F$ -rational points of a connected reductive  $F$ -group, a maximal  $F$ -split subtorus, its centralizer, and its normalizer. Let  $\mathfrak{C}^+$  be an open alcove of the semi-simple apartment of  $G$  defined by  $T$ , let  $x_0$  be a special vertex of the closed alcove  $\overline{\mathfrak{C}}^+$ , and let  $I, I(1), K$ , be respectively the Iwahori subgroup of  $G$  fixing  $\mathfrak{C}^+$ , its pro- $p$ -Sylow subgroup, and the parahoric subgroup of  $G$  fixing  $x_0$ .

We associate to  $G, T, Z, N, I, I(1), K$  the data

$$(\mathcal{W} = (\Sigma, \Delta, \Omega, \Lambda, \nu, W, Z_k, W(1)); (q_s, c_{\tilde{s}})),$$

and a group  $\Lambda_T$ , satisfying the properties given in §1.2 with  $R = \mathbb{Z}$ , as follows.

The apartment defined by  $T$  identifies with a Euclidean real vector space  $V$ . The set  $S^{\text{aff}}$  of orthogonal reflections with respect to the walls of  $\mathfrak{C}^+$  generates an affine Coxeter system  $(W^{\text{aff}}, S^{\text{aff}})$ , given by a based reduced root system  $(\Sigma, \Delta)$ . The action of  $N$  on the apartment transfers to an action on  $V$ . The subgroup  $Z$  acts by translations  $(z, x) \mapsto x + \nu_Z(z)$ ,  $(z, x) \in Z \times V$ , for an homomorphism  $\nu_Z : Z \rightarrow V$  satisfying  $\alpha \circ \nu_Z(t) = -\alpha(t)$  for  $t \in T$  and  $\alpha$  in the root system  $\Phi$  of  $T$  in  $G$ . There is a surjective map  $\alpha \mapsto e_{\alpha} : \Phi \rightarrow \Sigma$ , where  $e_{\alpha}$  is a positive integer for all  $\alpha \in \Phi$ .

Let  $T_0 := T \cap K$  (the maximal compact subgroup of  $T$ ),  $Z_0 := K \cap Z$  (the parahoric subgroup of  $Z$ ), and let  $Z_0(1)$  be the pro- $p$ -Sylow subgroup of  $Z_0$ . Then

$$\begin{aligned} \Lambda_T &:= T/T_0, & \Lambda &:= Z/Z_0, & \Lambda(1) &:= Z/Z_0(1), & Z_k &:= Z_0/Z_0(1), \\ W_0 &:= N/Z, & W &:= N/Z_0, & W(1) &:= N/Z_0(1). \end{aligned}$$

The homomorphism  $\nu_Z$  and the action of  $N$  on  $V$  are trivial on  $Z_0$ . They induce an homomorphism  $\nu : \Lambda \rightarrow V$  and an action of  $W$  on  $N$ . The monoid  $\Lambda^+$  represents the

orbits of  $W_0$  in  $\Lambda$  [7, 6.3] and the double cosets  $K \backslash G / K$ . The groups  $W, W(1)$  represent the double cosets  $I \backslash G / I, I(1) \backslash G / I(1)$ . The group  $\Omega$  is the  $W$ -stabilizer of the alcove  $\mathfrak{C}^+$ . We denote by  $\tilde{w}$  an element of  $W(1)$  of image  $w$  in  $W$ , and we call  $\tilde{w}$  a lift of  $w$ .

For  $s \in S^{\text{aff}}$ , let  $K_s$  be the parahoric subgroup of  $G$  fixing the face of  $\mathfrak{C}^+$  fixed by  $s$ . The quotient of  $K_s$  by its pro- $p$ -radical is the group  $G_{s,k}$  of rational points of a  $k$ -reductive connected group of rank 1. The image of  $I(1)$  in  $G_{s,k}$  is the group  $U_{s,k}$  of rational points of the unipotent radical of a  $k$ -Borel subgroup  $Z_k U_{s,k}$  of opposite group  $Z_k \overline{U}_{s,k}$ . It is known that  $s$  admits a lift  $n_s \in N \cap K_s$  of image in  $G_{s,k}$  belonging to the group  $\langle U_{s,k}, \overline{U}_{s,k} \rangle$  generated by  $U_{s,k} \cup \overline{U}_{s,k}$ . The image of  $n_s$  in  $W(1)$  is called an admissible lift of  $s$ . We set  $Z_{k,s} = Z_k \cap \langle U_{s,k}, \overline{U}_{s,k} \rangle$ .

For  $s \in S^{\text{aff}}$ ,  $\tilde{s}$  an admissible lift of  $s$ , and  $t \in Z_k$ , let

$$q_s = [In_s I : I] \text{ is a power of } q, \quad c_s := (q_s - 1) |Z_{k,s}|^{-1} \sum_{z \in Z_{k,s}} z,$$

and  $c_{t\tilde{s}} = \sum_{z \in Z_{k,s}} c_{\tilde{s}}(z)tz$ , for positive integers  $c_{\tilde{s}}(z) = c_{\tilde{s}}(z^{-1})$  of sum  $q_s - 1$ , constant on each coset modulo  $\{xs(x)^{-1} \mid x \in Z_k\}$ , and  $c_{\tilde{s}} \equiv c_s \pmod p$  as in [15, Theorem 2.2].

The cocharacter group  $X_*(T)$  of  $T$  is isomorphic to  $\Lambda_T$  and embeds in  $\Lambda(1)$  by the map  $\mu \mapsto \mu(p_F)^{-1} : X_*(T) \rightarrow Z$  followed by the quotient maps of  $Z$  onto  $\Lambda$  and  $\Lambda(1)$ . Remembering the sign  $-$  in the definition of  $\nu$ ,

$$\mu \in \Lambda_T^+ \Leftrightarrow \alpha(\mu(p_F)) \in O_F \quad \text{for all } \alpha \in \Delta.$$

We identify  $\mu$  with its image in  $\Lambda_T$ , and  $\tilde{\mu}$  denotes its image in  $\Lambda(1)$ .

For a commutative ring  $R$ , the pro- $p$ -Iwahori Hecke  $R$ -algebra  $\mathcal{H}_R(G, I(1))$  is isomorphic to the algebra  $\mathcal{H}_R(q_s, c_{\tilde{s}})$  associated to this data.

The pro- $p$ -Iwahori Hecke  $R$ -algebra  $\mathcal{H}_R(K, I(1))$  of  $K$  is a subalgebra of  $\mathcal{H}_R(G, I(1))$  isomorphic to the finite subalgebra  $\mathfrak{h}(q_s, c_{\tilde{s}})$  of  $\mathcal{H}$ .

The Iwahori Hecke  $R$ -algebra  $\mathcal{H}_R(G, I)$  is an algebra  $\mathcal{H}$  associated to the same data except that  $Z_k = \{1\}, W(1) = W, c_s = q_s - 1$ .

The group  $G$  is split  $\Leftrightarrow T = Z \Rightarrow \Lambda_T = \Lambda$ . The group  $G$  is quasi-split  $\Leftrightarrow Z$  is the  $F$ -points of an  $F$ -torus  $\Rightarrow \Lambda(1)$  is commutative. The group  $G$  is semi-simple  $\Leftrightarrow \text{Ker } \nu$  is finite  $\Rightarrow \Omega$  is finite and  $\nu$  is injective on  $\Lambda_T$ .

The quotient of  $K$  by its pro- $p$ -radical  $K(1)$  is the group  $G_k$  of  $k$ -rational points of a connected reductive  $k$ -group. The images in  $G_k$  of  $T_0, Z_0, I$ , and  $I(1)$  are the groups  $T_k, Z_k, B_k$ , and  $U_k$  of  $k$ -rational points of a maximal  $k$ -split torus, its centralizer (a  $k$ -torus), a Borel  $k$ -subgroup containing the maximal  $k$ -split torus, and its unipotent radical.

The finite Hecke algebras  $\mathcal{H}_R(K, I(1))$  and  $\mathcal{H}_R(G_k, U_k)$  are isomorphic.

The condition  $q_s = 0$  for all  $s \in S^{\text{aff}}$  means that the characteristic of  $R$  is  $p$ . Then,

$$c_{t\tilde{s}} = -|Z_{k,s}|^{-1} \sum_{z \in Z_{k,s}} tz,$$

and the irreducible smooth  $R$ -representations  $\rho$  of  $K$  are trivial on  $K(1)$ ; they identify with the irreducible  $R$ -representations of  $G_k$ , in bijection with the characters of  $\mathcal{H}_R(G_k, U_k)$  by the  $U_k$ -invariant functor  $\rho \mapsto \rho^{U_k}$  for  $R$  as in 1.4.

1.4.

For the remainder of this article, unless otherwise specified, we are in the setting of § 1.2 with  $q_s = 0$  for all  $s \in S^{\text{aff}}$ , and  $R$  is a field containing a root of unity of order the exponent of  $Z_k$ .

*Notation.* We denote by  $\hat{Z}_k$  the group of  $R$ -characters of  $Z_k$ . For a character  $\chi \in \hat{Z}_k$ , a character  $\eta$  of  $\mathfrak{h}$ , and a character  $\Xi$  of  $\mathcal{H}^{\text{aff}}$ , we set

$$S_\chi^{\text{aff}} := \{s \in S^{\text{aff}} \mid \chi(c_{\tilde{s}}) \neq 0\}, \quad S_\chi := S_\chi^{\text{aff}} \cap S, \tag{9}$$

$$S_\eta := \{s \in S \mid \eta(T_{\tilde{s}}) \neq 0\}, \quad S_\Xi^{\text{aff}} := \{s \in S^{\text{aff}} \mid \Xi(T_{\tilde{s}}) \neq 0\}. \tag{10}$$

These sets are independent of the choice of the lift  $\tilde{s}$  of  $s$ . For  $(\tilde{w}, \chi) \in W(1) \times \hat{Z}_k$  we denote by  $\chi^w \in \hat{Z}_k$  the character  $\chi^w(t) = \chi(\tilde{w}t\tilde{w}^{-1})$  for  $t \in Z_k$ . The subgroup generated by a subset  $X$  of a group is denoted by  $\langle X \rangle$ . For  $\lambda \in \Lambda$  we set

$$\Delta_\lambda := \{\alpha \in \Delta \mid \alpha \circ \nu(\lambda) = 0\}, \quad S_\lambda := \{s_\alpha \mid \alpha \in \Delta_\lambda\}. \tag{11}$$

We recall from § 1.2 the  $R$ -algebra  $\mathfrak{h}$  associated to the finite Coxeter system  $(W_0, S)$  and the extension  $1 \rightarrow Z_k \rightarrow W_0(1) \rightarrow W_0 \rightarrow 1$ , of basis  $(T_{\tilde{w}})_{\tilde{w} \in W_0(1)}$  satisfying the braid relations and the quadratic relations  $T_{\tilde{s}}(T_{\tilde{s}} - c_{\tilde{s}}) = 0$  for  $\tilde{s} \in S(1)$ .

**Theorem 1.1** (The characters of  $\mathfrak{h}$ ). (a) *The characters  $\eta$  of  $\mathfrak{h}$  are in bijection with the pairs  $(\chi, J)$ , where  $\chi \in \hat{Z}_k$  and  $J \subset S_\chi$ ,  $\chi = \eta|_{Z_k}$ , and  $J = S_\eta$ .*

(b) *For any  $\eta$ , there exists an orientation  $o$  such that the equivalent properties  $S_\eta = S_\chi \cap S_o \Leftrightarrow \eta(E_o(\tilde{s})) = 0$ , for all  $s \in S$ , hold true. We set  $\chi_o := \eta$ .*

(c) *For two characters  $\eta_1, \eta$  of  $\mathfrak{h}$ , there exists an orientation  $o$  such that  $\eta_1 = (\chi_1)_o$ ,  $\eta = \chi_o$  if and only if*

$$S_\eta \cap S_{\chi_1} = S_{\eta_1} \cap S_\chi.$$

For a reduced decomposition of  $\tilde{w} = \tilde{s}_1 \dots \tilde{s}_r$  of  $W(1)$ , the element  $c_{\tilde{w}} = c_{\tilde{s}_1} \dots c_{\tilde{s}_r}$  of  $R[Z_k]$  does not depend on the choice of the reduced decomposition [15, Propositions 4.13(ii) and 4.22].

**Theorem 1.2** (A basis of the intertwiners). *Let  $\eta_1, \eta$  be two characters of  $\mathfrak{h}$  of restrictions  $\chi_1, \chi$  to  $Z_k$ .*

(a)  *$\eta_1$  is contained in  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  (is a submodule) if and only if*

$$\chi_1 = \chi^\lambda, \quad S_{\eta_1} \cap S_\lambda = S_\eta \cap S_\lambda, \quad \text{for some } \lambda \in \Lambda^+.$$

(b) *For  $\lambda \in \Lambda^+$  satisfying (a), there exists a non-zero  $\mathcal{H}$ -intertwiner*

$$\Phi_{\tilde{\chi}} : 1 \otimes 1 \mapsto 1 \otimes \mathcal{E}_{\tilde{\chi}} : \eta_1 \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \eta \otimes_{\mathfrak{h}} \mathcal{H}, \quad \mathcal{E}_{\tilde{\chi}} := \sum_{w_0 \in Y_\lambda} \chi_1(c_{\tilde{w}_0})^{-1} \otimes T_{\tilde{\chi}\tilde{w}_0},$$

where  $Y_\lambda = \{w_0 \in \langle S_{\chi_1} - S_{\eta_1} \rangle \mid \chi_1^{w_0} = \chi_1, \ell(\lambda w_0) = \ell(\lambda) - \ell(w_0)\}$ , and  $\tilde{w}_0$  is a lift of  $w_0$ ; note that  $\chi_1(c_{\tilde{w}_0})^{-1} \otimes T_{\tilde{\chi}\tilde{w}_0}$  does not depend on the choice of the lift.  $(\Phi_{\tilde{\chi}})$ , for  $\lambda \in \Lambda^+$  satisfying (a), is a basis of  $\text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H})$ .

(c) If  $o$  satisfies (d) and  $\lambda \in \Lambda^+$  satisfies (a), there exists a non-zero  $\mathcal{H}$ -intertwiner

$$\Phi_{o, \tilde{\lambda}} : 1 \otimes 1 \mapsto 1 \otimes E_o(\tilde{\lambda}) : (\chi_1)_o \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \chi_o \otimes_{\mathfrak{h}} \mathcal{H}.$$

$(\Phi_{o, \tilde{\lambda}})$ , for  $\lambda \in \Lambda^+$  satisfying (a), is a basis of  $\text{Hom}_{\mathcal{H}}((\chi_1)_o \otimes_{\mathfrak{h}} \mathcal{H}, \chi_o \otimes_{\mathfrak{h}} \mathcal{H})$ .

We note that  $\chi_1(c_{\tilde{w}_0})^{-1} \otimes T_{\tilde{\lambda}\tilde{w}_0} \in \eta \otimes_{\mathfrak{h}} \mathcal{H}$  does not depend on the choice of the lift  $\tilde{w}_0$  of  $w_0 \in Y_{\lambda}$ . We set

$$\Lambda_{\chi} := \{\lambda \in \Lambda \mid \chi^{\lambda} = \chi\}, \quad \text{resp. } \Lambda_{\chi}^+ := \Lambda^+ \cap \Lambda_{\chi}. \tag{12}$$

The idempotent  $e_{\chi} := |Z_k|^{-1} \sum_{t \in Z_k} \chi(t)^{-1} t$  of  $R[Z_k]$  is central in  $R[\Lambda_{\chi}(1)]$ , and the  $R$ -linear map

$$\chi \otimes_{R[Z_k]} R[\Lambda_{\chi}(1)] \rightarrow e_{\chi} R[\Lambda_{\chi}(1)] \quad 1 \otimes \tilde{\lambda} \mapsto e_{\chi} \tilde{\lambda} \quad (\lambda \in \Lambda_{\chi}) \tag{13}$$

is an isomorphism. Any  $R$ -algebra  $A$  with a basis  $(a_{\tilde{\lambda}})_{\lambda \in \Lambda_{\chi}^+}$  satisfying

$$a_{\tilde{\lambda}} a_{\tilde{\lambda}'} = \chi(t) a_{\tilde{\lambda}''} \quad \text{for } \lambda, \lambda', \lambda'' \in \Lambda_{\chi}^+, t \in Z_k, \tilde{\lambda} \tilde{\lambda}' = t \tilde{\lambda}'', \tag{14}$$

is canonically isomorphic to the algebra  $e_{\chi} R[\Lambda_{\chi}^+(1)]$  with its natural basis  $(e_{\chi} \tilde{\lambda})_{\lambda \in \Lambda_{\chi}^+}$ .

For an orientation  $o$ , the  $R$ -submodule  $\mathcal{A}_{o, \chi}^+$  of basis  $(E_o(\tilde{\lambda}))_{\tilde{\lambda} \in \Lambda_{\chi}^+(1)}$  is a subalgebra of  $\mathcal{H}$ . The algebra  $\chi \otimes_{R[Z_k]} \mathcal{A}_{o, \chi}^+$  of basis  $(1 \otimes E_o(\tilde{\lambda}))_{\lambda \in \Lambda_{\chi}^+}$  is an  $R$ -algebra with a basis satisfying (14).

A spherical Hecke algebra is the algebra of  $\mathcal{H}$ -intertwiners of a right  $\mathcal{H}$ -module  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  induced from a character  $\eta$  of  $\mathfrak{h}$ , by analogy with the reductive  $p$ -adic groups

$$\mathcal{H}(\mathfrak{h}, \eta) := \text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H}).$$

Theorem 1.2 with  $\eta_1 = \eta$  becomes the following.

**Theorem 1.3** (A Satake-type isomorphism for the spherical algebra). (a) A basis of the spherical Hecke algebra  $\mathcal{H}(\mathfrak{h}, \eta)$  is  $(\Phi_{\tilde{\lambda}})_{\lambda \in \Lambda_{\chi}^+}$ , where

$$\Phi_{\tilde{\lambda}} : 1 \otimes 1 \mapsto 1 \otimes \mathcal{E}_{\tilde{\lambda}} : \eta \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \eta \otimes_{\mathfrak{h}} \mathcal{H}, \quad \mathcal{E}_{\tilde{\lambda}} := \sum_{w_0 \in Y_{\lambda}} \chi(c_{w_0}) \otimes T_{\tilde{\lambda}w_0},$$

$$Y_{\lambda} = \{w_0 \in \langle S_{\chi} - S_{\eta} \rangle \mid \chi^{w_0} = \chi, \ell(\lambda w_0) = \ell(\lambda) - \ell(w_0)\}.$$

(b) Let  $o$  be an orientation such that  $\eta = \chi_o$ . For  $\lambda \in \Lambda_{\chi}^+$ , there exists an injective  $\mathfrak{h}$ -intertwiner

$$\Phi_{o, \tilde{\lambda}} : 1 \otimes 1 \mapsto 1 \otimes E_o(\tilde{\lambda}) : \eta \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \eta \otimes_{\mathfrak{h}} \mathcal{H}.$$

$(\Phi_{o, \tilde{\lambda}})_{\lambda \in \Lambda_{\chi}^+}$  is a basis of the spherical Hecke algebra  $\mathcal{H}(\mathfrak{h}, \eta)$  satisfying (14), inducing an isomorphism

$$\mathcal{H}(\mathfrak{h}, \eta) \simeq e_{\chi} R[\Lambda_{\chi}^+(1)].$$

We suppose now that  $\Lambda_T$  exists. The center  $\mathcal{Z}$  of  $\mathcal{H}$  is the algebra  $\mathcal{A}_o^{W(1)}$  of  $W(1)$ -invariants of  $\mathcal{A}_o$ , and is a free  $R$ -module of basis

$$E(\tilde{C}) = \sum_{\tilde{\lambda} \in \tilde{C}} E_o(\tilde{\lambda}) \tag{15}$$

$(E(\tilde{C}))$  is independent of the choice of  $o$ ) for all finite conjugacy classes  $\tilde{C}$  of  $W(1)$ . We denote by  $\tilde{C}(\mu)$  the  $W(1)$ -conjugacy class of  $\tilde{\mu}$  for  $\mu \in \Lambda_T^+$ . The  $R$ -subspace of



basis  $(E(\tilde{C}(\mu)))_{\mu \in \Lambda_T^+}$  is a central subalgebra  $\mathcal{Z}_T$  of  $\mathcal{H}$  which has better properties than  $\mathcal{Z}$ .

A central element  $x \in \mathcal{Z}$  induces naturally a  $\mathcal{H}$ -intertwiner of  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$ :

$$\Phi_x : 1 \otimes h \mapsto 1 \otimes xh = 1 \otimes hx \quad \text{for } h \in \mathcal{H}. \tag{16}$$

It is straightforward to check that  $\Phi_x$  belongs to the center  $\mathcal{Z}(\eta, \mathfrak{h})$  of  $\mathcal{H}(\eta, \mathfrak{h})$ . The  $R$ -subspace of basis  $(\Phi_{E(\tilde{C}(\mu))})_{\mu \in \Lambda_T^+}$  is a central subalgebra  $\mathcal{Z}(\eta, \mathcal{H})_T$  of the spherical algebra  $\mathcal{H}(\eta, \mathfrak{h})$ .

**Theorem 1.4** (Almost-isomorphism between the centers of  $\mathcal{H}$  and  $\mathcal{H}(\eta, \mathfrak{h})$ ). *We suppose that  $\Lambda_T$  exists. Let  $\eta$  be a character of  $\mathfrak{h}$ .*

- (a)  $\mathcal{Z}_T$  is a finitely generated central  $R$ -subalgebra of  $\mathcal{H}$ , and  $\mathcal{H}$  is a finitely generated  $\mathcal{Z}_T$ -module. This is also true for  $(\mathcal{Z}(\eta, \mathcal{H})_T, \mathcal{H}(\eta, \mathfrak{h}))$  instead of  $(\mathcal{Z}_T, \mathcal{H})$ .
- (b)  $\Phi_{E(\tilde{C}(\mu))} = \Phi_{o, \tilde{\mu}}$  for  $\mu \in \Lambda_T^+$  and any orientation  $o$  such that  $\eta = \chi_o$ .  
The linear map  $\tilde{\mu} \mapsto \Phi_{E(\tilde{C}(\mu))} : R[\tilde{\Lambda}_T^+] \rightarrow \mathcal{Z}(\eta, \mathcal{H})_T$  is an algebra isomorphism.
- (c) The map  $x \mapsto \Phi_x : \mathcal{Z} \rightarrow \mathcal{Z}(\eta, \mathcal{H})$  restricts to an isomorphism  $\mathcal{Z}_T \rightarrow \mathcal{Z}(\eta, \mathcal{H})_T$ .

We prove (a) over any commutative ring  $R$ .

We transfer these results to the group  $G$ . The spherical Hecke algebra  $\mathcal{H}_R(G, K, \rho) = \text{End}_{RG} \text{c-Ind}_K^G \rho$  of an irreducible smooth representation  $\rho$  of  $K$  with  $\mathcal{H}_R(K, I(1))$  acting by  $\eta$  on  $\rho^{I(1)}$  is isomorphic to  $\mathcal{H}(\eta, \mathfrak{h})$  by the pro- $p$ -Iwahori invariant functor. We denote by  $\mathcal{Z}_R(G, K, \rho)_T$  the algebra corresponding to  $\mathcal{Z}(\eta, \mathcal{H})_T$ . We denote by  $\mathcal{H}_R(Z^+, Z_0, \chi)$  the  $R$ -algebra of elements in the Hecke algebra  $\mathcal{H}_R(Z^+, Z_0, \chi)$  with support contained in the monoid  $Z^+$  of  $z \in Z$  with  $v_Z(z)$  dominant.

From Theorem 1.3 we obtain an algebra isomorphism

$$S_o : \mathcal{H}_R(G, K, \rho) \rightarrow \mathcal{H}_R(Z^+, Z_0, \chi) \tag{17}$$

for each orientation  $o$  such that  $\eta = \chi_o$ . This isomorphism restricts to an isomorphism, independent of the choice of  $o$ ,

$$S_T : \mathcal{Z}_R(G, K, \rho)_T \rightarrow \mathcal{H}_R(T^+, T_0, \chi). \tag{18}$$

Let  $\pi$  be a smooth  $R$ -representation of  $G$  such that  $\text{Hom}_R(\rho, \pi)$  contains a  $\mathcal{Z}_R(G, K, \rho)_T$ -eigenvector  $A$  of eigenvalue  $\xi$ , seen as an homomorphism  $\tilde{\Lambda}_T^+ \rightarrow R$  (Theorem 1.4). From Theorem 1.4, for  $v \in \rho^{I(1)}$  non-zero and  $\mu \in \Lambda_T^+$ ,

$$\xi(\tilde{\mu})A(v) = A(v)E_o(\tilde{\mu}) = A(v)E(\tilde{C}(\mu)).$$

**Theorem 1.5** (Supersingularity in  $G$  and in  $\mathcal{H}$ ). *The eigenvalue  $\xi$  of the  $\mathcal{Z}_R(G, K, \rho)_T$ -eigenvector  $A \in \text{Hom}_R(\rho, \pi)$  is supersingular if and only if the submodule  $A(v)\mathcal{H}$  of  $\pi^{I(1)}$  is supersingular.*

We recall that an homomorphism  $\tilde{\Lambda}_T^+ \rightarrow R$  is called supersingular if it vanishes on the non-invertible elements, and that a simple right  $\mathcal{H}$ -module  $M$  is called supersingular if  $ME(\tilde{C}) = 0$  for all finite conjugacy classes  $\tilde{C}$  in  $W(1)$  with positive length [13, Definition 1]. This is equivalent to  $ME(\tilde{C}(\mu)) = 0$  for all non-invertible  $\mu \in \tilde{\Lambda}_T^+$ .

In a forthcoming article, we will study the parabolic induction for  $\mathcal{H}$ -modules; we hope to prove that the isomorphism  $\mathcal{S}_o$  (17) is the Satake isomorphism of [7] for a good choice of  $o$  such that  $\eta = \chi_o$  (this was proved by Ollivier [10, Theorem 5.5]), when  $G$  is split with a simply connected derived group, and  $K$  is hyperspecial; as  $Z = T$ , we have  $\mathcal{S}_o = \mathcal{S}_T$ , and that an irreducible smooth admissible representation  $\pi$  is supersingular if and only if  $\pi^{I(1)}$  contains a supersingular module (this was proved by Ollivier for  $G = GL(n, F)$  and  $PGL(n, F)$  [11, Theorem 5.26]).

Finally, we classify the supersingular simple finite-dimensional  $\mathcal{H}$ -modules (proved by Ollivier when  $G$  is split, and  $K$  is hyperspecial [11, Corollary 5.15]).

For a character  $\Xi$  of  $\mathcal{H}^{\text{aff}}$ , the  $R$ -subalgebra  $\mathcal{H}_\Xi$  of  $\mathcal{H}$  generated by  $\mathcal{H}^{\text{aff}}$  and the  $\Omega(1)$ -fixator of  $\Xi$ ,

$$\Omega(1)_\Xi := \{u \in \Omega(1) \mid \Xi(uhu^{-1}) = \Xi(h) \text{ for } h \in \mathcal{H}^{\text{aff}}\},$$

is identified by (3) with the twisted tensor product  $\mathcal{H}^{\text{aff}} \otimes_{R[Z_k]} R[\Omega(1)_\Xi] \rightarrow \mathcal{H}_\Xi$ . For a simple finite-dimensional  $R$ -representation  $\sigma$  of  $\Omega(1)_\Xi$  equal to  $\Xi$  on  $Z_k$ , let

$$M(\Xi, \sigma) := (\Xi \otimes \sigma) \otimes_{\mathcal{H}_\Xi} \mathcal{H} \tag{19}$$

be the right  $\mathcal{H}$ -module induced from the right  $\mathcal{H}_\Xi$ -module  $\Xi \otimes \sigma$ . The induced module  $M(\Xi, \sigma)$  is finite dimensional. Two pairs  $(\Xi_1, \sigma_1), (\Xi_2, \sigma_2)$  are called conjugate by an element  $u \in \Omega(1)$  if

$$\Xi_1(uhu^{-1}) = \Xi_2(h), \sigma_1(uvu^{-1}) = \sigma_2(v) \quad \text{for } (h, v) \in \mathcal{H}^{\text{aff}} \times u^{-1}\Omega_\Xi(1)u.$$

The affine Coxeter system  $(W^{\text{aff}}, S^{\text{aff}})$  is the direct product of the irreducible affine Coxeter systems  $(W_i^{\text{aff}}, S_i^{\text{aff}})_{1 \leq i \leq r}$  associated to the irreducible components  $(\Sigma_i, \Delta_i)_{1 \leq i \leq r}$  of the based reduced root system  $(\Sigma, \Delta)$ . The  $R$ -submodule of basis  $(T_{\tilde{w}})_{\tilde{w}_i \in W_i^{\text{aff}}(1)}$  is a subalgebra  $\mathcal{H}_i^{\text{aff}}$  of  $\mathcal{H}^{\text{aff}}$ . The algebras  $\mathcal{H}_i^{\text{aff}}$  are called the irreducible components of  $\mathcal{H}^{\text{aff}}$ .

**Theorem 1.6** (Supersingular simple modules). (a) *The characters  $\Xi$  of  $\mathcal{H}^{\text{aff}}$  are in bijection with the pairs  $(\chi, J)$ , where  $\chi \in \hat{Z}_k$  and  $J \subset S_\chi^{\text{aff}}$ ,  $\chi = \Xi|_{Z_k}$ , and  $J = S_\Xi^{\text{aff}}$  (10). When  $S_\Xi^{\text{aff}} = S^{\text{aff}}$ ,  $\Xi$  is called a sign character, and the character  $\Xi \circ \iota$  (4) is called a trivial character.*

- (b) *A character  $\Xi$  of  $\mathcal{H}^{\text{aff}}$  is supersingular if and only if it is not a sign or trivial character on each irreducible component of  $\mathcal{H}^{\text{aff}}$ .*
- (c) *A finite-dimensional right  $\mathcal{H}$ -module is supersingular if and only if it is isomorphic to  $M(\Xi, \sigma)$ , where  $\Xi$  is a supersingular character of  $\mathcal{H}^{\text{aff}}$  and  $\sigma$  is a simple finite-dimensional  $R$ -representation  $\sigma$  of  $\Omega(1)_\Xi$  equal to  $\Xi$  on  $Z_k$ .*
- (d)  *$M(\Xi_1, \sigma_1) \simeq M(\Xi_2, \sigma_2)$  if and only if  $(\Xi_1, \sigma_1), (\Xi_2, \sigma_2)$  are  $\Omega(1)$ -conjugate.*

## 2. The characters of $\mathfrak{h}$ and $\mathcal{H}^{\text{aff}}$

**Proposition 2.1.** *A simple  $\mathfrak{h}$ -module has dimension 1.*

**Proof.** The finite-dimensional  $R$ -algebra  $\mathfrak{h}$  is generated by  $Z_k$  and  $T_{\tilde{s}}$  for all  $s \in S$ . By the hypothesis on  $R$  (§ 1.4), a right simple  $\mathfrak{h}$ -module is finite dimensional and contains an eigenvector  $v$  of  $Z_k$ . Following the argument of [4, Theorem 6.10], we choose  $w$  in the finite group  $W_0$  of maximal length such that  $vT_{\tilde{w}} \neq 0$ , and we show that  $RvT_{\tilde{w}}$  is  $\mathfrak{h}$ -stable.

$RvT_{\tilde{w}}$  is stable by  $T_t$ , because  $T_{\tilde{w}}T_t = (w \bullet t)T_{\tilde{w}}$  for  $t \in Z_k$ .

$RvT_{\tilde{w}}$  is stable by  $T_{\tilde{s}}$ , because

- if  $\ell(ws) = \ell(w) + 1$ ,  $vT_{\tilde{w}}T_{\tilde{s}} = vT_{w\tilde{s}}$  and by the hypothesis on  $w$ ,  $vT_{w\tilde{s}} = 0$ ;
- if  $\ell(ws) = \ell(w) - 1$ ,  $T_{\tilde{w}}T_{\tilde{s}} = T_{\tilde{w}\tilde{s}^{-1}}T_{\tilde{s}}^2 = T_{\tilde{w}\tilde{s}^{-1}}c_{\tilde{s}}T_{\tilde{s}} = T_{w\tilde{s}^{-1}}T_{\tilde{s}}c_{\tilde{s}} = (w \bullet c_{\tilde{s}})T_{\tilde{w}}$ . We used that  $T_{\tilde{s}}$  and  $c_{\tilde{s}}$  commute. □

**Proposition 2.2.** *The characters  $\eta$  of  $\mathfrak{h}$  are in bijection with the pairs  $(\chi, J)$ , where  $\chi \in \hat{Z}_k$  and  $J \subset S_\chi$  (9), by the recipe*

$$\eta|_{Z_k} = \chi, \quad S_\eta = \{s \in S \mid \eta(T_{\tilde{s}}) \neq 0\} = J.$$

We have  $\eta(T_{\tilde{s}}) = \chi(c_{\tilde{s}})$  if  $s \in J$ .

The characters  $\Xi$  of  $\mathcal{H}^{\text{aff}}$  are in bijection with the pairs  $(\chi, J)$ , where  $\chi \in \hat{Z}_k$  and  $J \subset S_\chi^{\text{aff}}$ , by the recipe

$$\Xi|_{Z_k} = \chi, \quad S_\Xi^{\text{aff}} = \{s \in S^{\text{aff}} \mid \Xi(T_{\tilde{s}}) \neq 0\} = J.$$

We have  $\Xi(T_{\tilde{s}}) = \chi(c_{\tilde{s}})$  if  $s \in J$ .

The set  $J$  is independent of the choice of the lift  $\tilde{s}$  of  $s$ . We call  $(\chi, J)$  the parameters of the character. The restriction to  $\mathfrak{h}$  of the character  $\Xi$  of  $\mathcal{H}^{\text{aff}}$  with parameters  $(\chi, S_\Xi^{\text{aff}})$  is the character of parameters  $(\chi, S_\Xi^{\text{aff}} \cap S)$ .

**Proof.** The proposition follows from the Iwahori–Matsumoto presentation in both cases. If  $\eta|_{Z_k} = \chi$ , we have

$$\eta(T_{\tilde{s}})(\eta(T_{\tilde{s}}) - \chi(c_{\tilde{s}})) = 0$$

for  $s \in S$ . We can replace  $\eta, S$  by  $\Xi, S^{\text{aff}}$ . □

The involutive automorphism  $\iota$  of  $\mathcal{H}$  (4) has the property for  $s \in S$  that

$$\eta(T_{\tilde{s}}) = 0 \Leftrightarrow \eta \circ \iota(T_{\tilde{s}}) = \eta(c_{\tilde{s}}).$$

The same holds for  $(\Xi, S^{\text{aff}})$  instead of  $(\eta, S)$ .

**Lemma 2.3.** *Let  $\eta$  be a character with parameters  $(\chi, S_\eta)$  of  $\mathfrak{h}$ . Then  $\eta \circ \iota$  is a character of  $\mathfrak{h}$  with parameters  $(\chi, S_\chi - S_\eta)$ . We can replace  $\eta, S, \mathfrak{h}$  by  $\Xi, S^{\text{aff}}, \mathcal{H}^{\text{aff}}$ .*

Let  $o$  be an orientation. We recall the notation (5), (6), (9), (10).

**Lemma 2.4.** *Let  $\eta$  be a character of  $\mathfrak{h}$  with parameters  $(\chi, S_\eta)$ . Then  $S_\eta = S_\chi \cap S_o \Leftrightarrow \eta(E_o(\tilde{s})) = 0$  for all  $s \in S$ . When this holds true, we denote  $\eta = \chi_o$ .*

We can replace  $(\eta, \mathfrak{h}, S, \chi_o)$  by  $(\Xi, \mathcal{H}^{\text{aff}}, S^{\text{aff}}, \chi_o^{\text{aff}})$ .

**Proof.** We compare the values of  $E_o(\tilde{s})$  and  $\eta(T_{\tilde{s}})$  for  $s \in S$ :

$$\begin{aligned} E_o(\tilde{s}) &= T_{\tilde{s}} \Leftrightarrow s \in S - S_o, \\ &= T_{\tilde{s}} - c_{\tilde{s}} \Leftrightarrow s \in S_o, \\ \eta(T_{\tilde{s}}) &= 0 \Leftrightarrow s \in S - S_\eta, \\ &= \chi(c_{\tilde{s}}) \neq 0 \text{ if } s \in S_\eta. \end{aligned}$$

We see that

- if  $s \in S - S_\chi$ , then  $\eta(E_o(\tilde{s})) = \eta(T_{\tilde{s}}) = \chi(c_{\tilde{s}}) = 0$ ;
- if  $s \in S_\chi - S_\eta$ , then  $\eta(E_o(\tilde{s})) = \eta(T_{\tilde{s}}) = 0 \Leftrightarrow s \notin (S_\chi - S_\eta) \cap S_o$ ;
- if  $s \in S_\eta$ , then  $\eta(E_o(\tilde{s})) = 0 \Leftrightarrow s \in S_\eta \cap S_o$ .

Hence we obtain the lemma for  $\eta$ . The proof is the same for  $\Xi$ . □

**Example 2.5.** For the dominant orientation  $o^+$ ,  $S_{o^+}^{\text{aff}} = S$ , and the parameters of  $\chi_{o^+}$  and of  $\chi_{o^+}^{\text{aff}}$  are  $(\chi, S_\chi)$ .

For the anti-dominant orientation  $o^-$ ,  $S_{o^-}^{\text{aff}} = S^{\text{aff}} - S$ , and the parameters of  $\chi_{o^-}$  are  $(\chi, \emptyset)$ , while those of  $\chi_{o^-}^{\text{aff}}$  are  $(\chi, S_\chi^{\text{aff}} - S_\chi)$ .

**Lemma 2.6.** (i) Any subset of  $S$  is equal to  $S_o$  for some orientation  $o$ .

A character  $\eta$  of  $\mathfrak{h}$  of restriction  $\chi$  to  $Z_k$  is equal to  $\chi_o$  for some orientation  $o$ , and

$$\eta = \chi_o \Leftrightarrow S_o \cap S_\chi = S_\eta.$$

(ii) Two  $R$ -characters  $\eta_1, \eta$  of  $\mathfrak{h}$  of parameters  $(\chi_1, S_{\eta_1}), (\chi, S_\eta)$  are equal to  $(\chi_1)_o, \chi_o$  for some orientation  $o$  if and only if

$$S_{\eta_1} \cap S_\chi = S_\eta \cap S_{\chi_1}.$$

In this case,  $\eta_1 = (\chi_1)_o$  and  $\eta = \chi_o \Leftrightarrow S_o \cap (S_{\chi_1} \cup S_\chi) = S_{\eta_1} \cup S_\eta$ .

**Proof.** (i) Let  $w_o \in W_0$ . For  $\alpha \in \Delta$ , the root in  $\{\alpha, -\alpha\}$  positive on  $w_o^{-1}(\mathfrak{D}^+)$  is equal to  $\alpha_o = \alpha$  if  $w_o(\alpha) > 0$  and  $\alpha_o = -\alpha$  if  $w_o(\alpha) < 0$ ; hence

$$s_\alpha \in S_o \Leftrightarrow w_o(\alpha) > 0.$$

For a subset  $X$  of  $S$ , we have  $X = S_o$  for the orientation  $o = o^+ \bullet w_o$  of Weyl chamber  $\mathfrak{D}_o = w_o^{-1}(\mathfrak{D}^+)$ , where  $w_o$  is the longest element of the group  $\langle S - X \rangle$  ( $w = 1$  if  $S = X$ ).

(ii)  $S_o \cap S_{\chi_1} = S_{\eta_1}$  and  $S_o \cap S_\chi = S_\eta$  imply that  $S_o \cap S_{\chi_1} \cap S_\chi = S_{\eta_1} \cap S_\chi = S_\eta \cap S_{\chi_1}$ . If  $S_{\eta_1} \cap S_\chi = S_\eta \cap S_{\chi_1}$ , then  $S_o \cap (S_{\chi_1} \cup S_\chi) = S_{\eta_1} \cup S_\eta$  implies that  $S_o \cap S_{\chi_1} = S_{\eta_1}$  and  $S_o \cap S_\chi = S_\eta$ . □

**Definition 2.7.** A character of  $\mathfrak{h}$  not vanishing on  $T_{\tilde{s}}$  for all  $s \in S$  is called a twisted sign character, and its image by the involution  $\iota$  is called a twisted trivial character.

We make the same definition for  $\mathcal{H}^{\text{aff}}, S^{\text{aff}}$  replacing  $\mathfrak{h}, S$ .

The twisted sign characters  $\eta$  are never 0 on  $T_{\tilde{w}}$  for  $w \in W_0$ . The algebra  $\mathfrak{h}$  admits no twisted sign or trivial characters when  $c_{\tilde{s}} = 0$  for some  $s \in S$ . They are equal to  $\chi_{o^+}$ , where  $\chi \in \hat{Z}_k$  satisfies  $S_\chi = S$ .

The twisted trivial characters  $\eta$  vanish on  $T_{\tilde{w}}$  for all  $w \in W_0$ . They are equal to  $\chi_{o^-}$ , where  $\chi \in \hat{Z}_k$  satisfies  $S_\chi = S$ .

The same remarks can be made for  $\mathcal{H}^{\text{aff}}, (W^{\text{aff}}, S^{\text{aff}})$  replacing  $\mathfrak{h}, (W_0, S)$ .

### 3. Distinguished representatives of $W_0 \backslash W$

We recall a well-known lemma for the affine Coxeter system  $(W^{\text{aff}}, S^{\text{aff}})$  extended to the group  $W = W^{\text{aff}} \rtimes \Omega$ .

For  $s \in S^{\text{aff}}$ , we denote by  $A_s$  the unique positive affine root such that  $s(A_s)$  is negative. We have  $s(A_s) = -A_s$  [8, 1.3.11]. When  $s \in S$  we write  $A_s = \alpha_s$ .

**Lemma 3.1.** (1) For  $(s, w) \in S^{\text{aff}} \times W$ , we have

$$\ell(ws) = 1 + \ell(w) \Leftrightarrow w(\alpha_s) > 0.$$

(2) For  $v \leq w$  in  $W$  and  $s \in S^{\text{aff}}$ , we have

- (a) either  $sv \leq w$  or  $sv \leq sw$ ;
- (b) either  $v \leq sw$  or  $sv \leq sw$ .

**Proof.** We recall that  $W = W^{\text{aff}} \rtimes \Omega$ . Let  $(s, u, w) \in S^{\text{aff}} \times \Omega \times W^{\text{aff}}$ .

- (1) We have  $\ell(uws) = \ell(ws)$ ,  $\ell(uw) = \ell(w)$ , and  $\ell(ws) = \ell(w) + 1 \Leftrightarrow w(\alpha_s) > 0$  [8, 1.13.13]. By definition (§ 1.2) an affine root is positive if and only if it is positive on the dominant alcove  $\mathfrak{C}^+$ . As the group  $\Omega$  normalizes  $\mathfrak{C}^+$ , it normalizes the set of positive affine roots, in particular  $w(\alpha_s) > 0 \Leftrightarrow (uw)(\alpha_s) > 0$ .
- (2) Let  $(v, u') \in W^{\text{aff}} \times \Omega$ . By definition of the Bruhat–Chevalley partial order [14, Ap. 2],  $vu' \leq wu$  is equivalent to  $u' = u, v \leq w$ . In  $W^{\text{aff}}$  [8, 1.3.19],
  - (a) either  $sv \leq w$  or  $sv \leq sw$ ;
  - (b) either  $v \leq sw$  or  $sv \leq sw$ .

We multiply (a) and (b) by  $u$  on the right without changing  $\leq$ . □

**Remark 3.2.** As  $\ell(w) = \ell(w^{-1})$  and  $v \leq w \Leftrightarrow v^{-1} \leq w^{-1}$ , in Lemma 3.1(1) we also have  $\ell(sw) = 1 + \ell(w) \Leftrightarrow w^{-1}(\alpha_s) > 0$ , and in Lemma 3.1(2), (a) and (b) can be replaced by

- (c) either  $vs \leq w$  or  $vs \leq ws$ ;
- (d) either  $v \leq ws$  or  $vs \leq ws$ .

We introduce now a distinguished set  $\mathcal{D}$  of representatives of  $W_0 \backslash W$ .

**Proposition 3.3.** The three sets

$$\mathcal{D}_1 = \{d \in W \mid d^{-1}(\alpha) > 0 \text{ for all } \alpha \in \Sigma^+\},$$

$$\begin{aligned} \mathcal{D}_2 &= \{\lambda w_0 \mid (\lambda, w_0) \in \Lambda^+ \times W_0, \ell(\lambda w_0) = \ell(\lambda) - \ell(w_0)\}, \\ \mathcal{D}_3 &= \{d \in W \mid \ell(w_0 d) = \ell(w_0) + \ell(d) \text{ for all } w_0 \in W_0\}, \end{aligned}$$

are equal, and will be denoted by  $\mathcal{D}$ . The cosets  $W_0 d$ , for  $d \in \mathcal{D}$ , are disjoint of union  $W$ .

**Proof.** The set  $\mathcal{D}_1$  is also equal to

$$\{d \in W \mid \ell(sd) = \ell(d) + 1 \text{ for all } s \in S\}, \tag{20}$$

because one can restrict to  $\alpha \in \Delta$  in the definition of  $\mathcal{D}_1$  and, for  $s \in S$ ,  $d^{-1}(\alpha_s) > 0 \Leftrightarrow \ell(sd) = \ell(d) + 1$  (Remark 3.2). Let  $w \in W$  not in  $\mathcal{D}_1$ . There exists  $s \in S$  with  $\ell(sw) = \ell(w) - 1$ . Then  $w_1 = sw$  satisfies  $\ell(w) = 1 + \ell(w_1)$ . We reiterate, and after finitely many steps we obtain  $(w_0, d) \in W_0 \times \mathcal{D}_1$  such that  $w = w_0 d$ ,  $\ell(w) = \ell(w_0) + \ell(d)$ . The pair  $(w_0, d)$  is unique. Indeed, for  $d, d'$  in  $\mathcal{D}_1$  with  $d'd^{-1} \in W_0$ , for all  $\alpha \in \Delta$  we have  $d'd^{-1}(\alpha) = \gamma \in \Sigma$ , and  $d^{-1}(\alpha) = d'^{-1}(\gamma)$  is positive as  $d \in \mathcal{D}_1$ ; hence  $\gamma > 0$  as  $d' \in \mathcal{D}_1$ . This implies  $d = d'$ . We deduce that  $\mathcal{D}_1$  is a set of representatives of  $W_0 \backslash W$ , that  $d \in \mathcal{D}_1$  is the unique element of minimal length in  $W_0 d$ , and that  $\mathcal{D}_1 \subset \mathcal{D}_3$ . This implies that  $\mathcal{D}_1 = \mathcal{D}_3$ .

We now compare the sets  $\mathcal{D}_1$  and  $\mathcal{D}_2$ . For  $(\lambda, w_0) \in \Lambda \times W_0$ , we deduce from Lemma 3.1 (see [15, Corollary 5.11]) that

$$\ell(\lambda w_0) = \ell(\lambda) - \ell(w_0) \Leftrightarrow \alpha \circ \nu(\lambda) > 0 \quad \text{for all } \alpha \in \Sigma^+ \cap w_0(\Sigma^-). \tag{21}$$

On the other hand, for all  $\alpha \in \Sigma^+$ ,  $(\lambda w_0)^{-1}(\alpha) = w_0^{-1}(\alpha) + \alpha \circ \nu(\lambda)$  is positive if and only if

$$w_0^{-1}(\alpha) > 0, \alpha \circ \nu(\lambda) \geq 0 \quad \text{or} \quad w_0^{-1}(\alpha) < 0, \alpha \circ \nu(\lambda) > 0 \tag{22}$$

[15, (36)]. Comparing (21) and (22), we deduce that  $\mathcal{D}_1 = \mathcal{D}_2$ . □

**Remark 3.4.** (i) The distinguished set  $\mathcal{D}^{\text{aff}}$  of representatives of  $W_0 \backslash W^{\text{aff}}$  given by Proposition 3.3 applied to  $W^{\text{aff}}$  is equal to  $\mathcal{D}^{\text{aff}} = \mathcal{D} \cap W^{\text{aff}}$ , and  $\mathcal{D} = \mathcal{D}^{\text{aff}} \Omega$ .

(ii) The distinguished set  $\mathcal{D}$  of representatives of  $W_0 \backslash W^{\text{aff}}$  can be inductively constructed: it is the set of  $\lambda w_0 \in \mathcal{D}$  for  $\lambda \in \Lambda^+$  and  $w_0 \in W_0$ , such that  $w_0 = 1$  or  $w_0$  has a reduced decomposition  $w_0 = s_1 \dots s_r$  ( $s_i \in S$ ), such that

$$\ell(\lambda s_1 \dots s_{i+1}) = \ell(\lambda s_1 \dots s_i) - 1 \quad \text{for } 1 \leq i \leq r.$$

Note that  $\lambda s \in \mathcal{D} \Leftrightarrow \alpha_s \circ \nu(\lambda) > 0$  when  $s \in S$ .

We denote by  $w_1$  the unique element of maximal length in the finite Weyl group  $W_0$ .

**Lemma 3.5.** Let  $\lambda, \mu \in \Lambda^+$ . The double  $W_0$ -coset  $W_0 \lambda W_0$  has a unique element  $w_\lambda$  of maximal length,

$$w_\lambda = w_1 \lambda, \quad \ell(w_\lambda) = \ell(w_1) + \ell(\lambda) \quad \text{and} \quad \lambda \leq \mu \Leftrightarrow w_\lambda \leq w_\mu.$$

The set  $W_0 \lambda W_0 \cap \mathcal{D}$  is equal to  $\mathcal{D}(\lambda) = \{\lambda w_0 \mid w_0 \in W_0, \ell(\lambda w_0) = \ell(\lambda) - \ell(w_0)\}$ .

**Proof.** The coset  $W_0d$  of  $d \in \mathcal{D}$  contains a unique element of maximal length, equal to  $w_1d$ ,  $\ell(w_1d) = \ell(w_1) + \ell(d)$ . For  $\lambda \in \Lambda^+$ , the set  $\mathcal{D} \cap W_0\lambda W_0$  contains a unique element of maximal length, equal to  $\lambda$  (Remark 3.4(ii)). Hence  $W_0\lambda W_0$  contains a unique element  $w_\lambda$  of maximal length, equal to  $w_1\lambda$  and  $\ell(w_\lambda) = \ell(w_1) + \ell(\lambda)$ . As  $w_\mu = w_1\mu$ ,  $\ell(w_\mu) = \ell(w_1) + \ell(\mu)$ , the equivalence  $\lambda \leq \mu \Leftrightarrow w_1\lambda \leq w_1\mu$  is clear. We have  $\mathcal{D}(\lambda) = \lambda W_0 \cap \mathcal{D}$  (Proposition 3.3), and  $\mu \in W_0\lambda W_0 \Leftrightarrow \mu = w\lambda w^{-1}$  for some  $w \in W_0 \Leftrightarrow \mu = \lambda$ , as  $\Lambda^+$  represents the orbits of  $W_0$  in  $\Lambda$  [7, 6.3].  $\square$

**Lemma 3.6.** Let  $(\lambda, w_0) \in \Lambda^+ \times W_0$ ,  $d = \lambda w_0 \in \mathcal{D}$ , and let  $\mu \in \Lambda^+$ .

- (1) For  $s \in S^{\text{aff}}$ ,  $ds \notin \mathcal{D} \Leftrightarrow dsd^{-1} \in S \Rightarrow \ell(ds) = \ell(d) + 1$ .
- (2) For  $s \in S$  and  $ds \in \mathcal{D}$ , we have  $\ell(ds) = \ell(d) + 1 \Leftrightarrow \ell(w_0s) = \ell(w_0) - 1$ .
- (3) For  $(w, d') \in W_0 \times \mathcal{D}$ , we have  $d \leq wd' \Rightarrow d \leq d'$ .
- (4) For  $s \in S$  such that  $ds \in \mathcal{D}$ , we have  $d \leq \mu \Rightarrow ds \leq \mu$ .
- (5) We have  $d \leq w_\mu \Leftrightarrow d \leq \mu \Leftrightarrow \lambda \leq \mu$ .

**Proof.** (1) Let  $s \in S^{\text{aff}}$ . By (20) and Remark 3.2,

$$ds \notin \mathcal{D} \Leftrightarrow (ds)^{-1}(\alpha) < 0 \text{ for some } \alpha \in \Delta.$$

As  $d^{-1}(\beta) > 0$  for all  $\beta \in \Sigma^+$ , and  $dsd^{-1} \in W^{\text{aff}}$ , we have

$$s((d^{-1}(\alpha))) < 0 \Leftrightarrow d^{-1}(\alpha) = A_s \Leftrightarrow \alpha = d(A_s) \Leftrightarrow s_\alpha = dsd^{-1}.$$

We have  $\ell(ds) = \ell(d) + 1$  by Lemma 3.1(1).

(2) Let  $s \in S$  with  $ds \in \mathcal{D}$ . Then

$$\ell(ds) = \ell(d) + 1 \Leftrightarrow \ell(\lambda) - \ell(w_0s) = \ell(\lambda) - \ell(w_0) + 1 \Leftrightarrow \ell(w_0s) = \ell(w_0) - 1.$$

(3)  $d \leq wd'$  and  $s \in S$  imply that  $d \leq swd'$  or  $sd \leq swd'$  by Lemma 3.1(2); as  $d < sd$ , we obtain

$$d \leq wd' \Rightarrow d \leq swd'.$$

If  $w \neq 1$ , we choose  $s$  such that  $sw < w$ . Repeating the procedure, we obtain  $d \leq d'$  by induction on the length of  $w \in W_0$ .

(4) As  $d \leq \mu$ ,  $ds \leq \mu$  or  $ds \leq \mu s$  by Lemma 3.1(2). When  $\mu s < \mu$ , we obtain  $ds \leq \mu$ . Suppose that  $\mu s > \mu$  and  $ds \leq \mu s$ . By Lemma 3.1(1),

$$\begin{aligned} \ell(\mu s) = \ell(mu) + 1 &\Leftrightarrow \mu(\alpha_s) = \alpha_s - \alpha_s \circ v(\mu) > 0 \Leftrightarrow \alpha_s \circ v(\mu) \leq 0, \\ &\Leftrightarrow \alpha_s \circ v(\mu) = 0 \Leftrightarrow v(\mu) \text{ fixed by } s \Leftrightarrow \mu s = s\mu u, u \in \Lambda \cap \Omega. \end{aligned}$$

We deduce that  $ds \leq s\mu u$ . By (3),  $ds \leq \mu u$ , because  $ds, \mu u \in \mathcal{D}$ . As  $\Lambda$  is commutative,  $ds \leq u\mu$ . For  $w \in W$ , there is a unique element  $u_w \in \Omega$  such that  $w \in u_w W^{\text{aff}}$ . By the definition of the Bruhat–Chevalley order,  $d \leq \mu$ ,  $ds \leq u\mu$  imply that  $u_d = u_\mu = uu_\mu$ . We deduce that  $u = 1$ ,  $ds \leq \mu$ .

(5) The implications  $d \leq w_\mu \Leftrightarrow d \leq \mu \Leftrightarrow \lambda \leq \mu$  are obvious, because  $d \leq \lambda$ ,  $\mu \leq w_\mu$ . The implication  $d \leq w_\mu \Rightarrow d \leq \mu$  follows from (3), because  $w_\mu = w_1\mu$  (Lemma 3.5) and  $\mu \in \mathcal{D}$ . The implication  $d \leq \mu \Rightarrow \lambda \leq \mu$  follows from (4) reiterated finitely many times for  $s \in S$  such that  $\ell(ds) = \ell(d) + 1$  if  $d \neq \lambda$  (Remark 3.4(ii)).  $\square$

**Remark 3.7.** Results similar to Proposition 3.3 and Lemma 3.6 are already in [9, Proposition 2.5, Lemma 2.6, Proposition 2.7], [10, Lemma 2.4], [11, Proposition 1.3], when  $W$  is the Iwahori Weyl group of a split reductive  $p$ -adic group  $G$ .

**Lemma 3.8.** *In Lemma 3.6, for  $s \in S$  and  $\Delta_\lambda$  as in (11),*

$$ds \notin \mathcal{D} \Leftrightarrow dsd^{-1} = w_0s w_0^{-1} \in S_\lambda \Leftrightarrow w_0(\alpha_s) \in \Delta_\lambda \Leftrightarrow w_0(\alpha_s) \in \Sigma^+, w_0(\alpha_s) \circ \nu(\lambda) = 0.$$

*This implies that  $\ell(w_0s) = \ell(w_0) + 1$  and  $\ell(ds) = \ell(d) + 1 = \ell(\lambda) - \ell(w_0s) + 2$ .*

**Proof.** By Lemma 3.6(1),  $ds \notin \mathcal{D} \Leftrightarrow d(\alpha_s) = \lambda w_0(\alpha_s) = w_0(\alpha_s) - w_0(\alpha_s) \circ \nu(\lambda) \in \Delta \Leftrightarrow w_0(\alpha_s) \in \Delta, w_0(\alpha_s) \circ \nu(\lambda) = 0 \Leftrightarrow w_0(\alpha_s) \in \Delta_\lambda$ . In the proof of Lemma 3.6(1), we saw that  $dsd^{-1} = s_{w_0(\alpha_s)} = w_0s w_0^{-1}$ . Note that  $ds \notin \mathcal{D}$  implies that  $\ell(ds) = \ell(d) + 1 = \ell(\lambda) - \ell(w_0) + 1 \neq \ell(\lambda) - \ell(w_0s)$ . Hence  $\ell(w_0s) = \ell(w_0) + 1, \ell(ds) = \ell(\lambda w_0s) = \ell(\lambda) - \ell(w_0s) + 2$ .

By (22),  $ds \in \mathcal{D} \Leftrightarrow \alpha \circ \nu(\lambda) > 0$  for all  $\alpha \in \Sigma^+ \cap w_0s(\Sigma^-)$ . We have

$$\begin{aligned} \Sigma^+ \cap w_0s(\Sigma^-) &= (\Sigma^+ \cap w_0(\Sigma^-)) - \{w_0(-\alpha_s)\} \text{ if } w_0(\alpha_s) \in \Sigma^-, \\ &= (\Sigma^+ \cap w_0(\Sigma^-)) \cup \{w_0(\alpha_s)\} \text{ if } w_0(\alpha_s) \in \Sigma^+, \end{aligned}$$

because, for  $\gamma \in \Sigma^+$ , we have  $sw_0^{-1}(\gamma) < 0$  if and only if  $\gamma \in \{w_0(\alpha_s)\} \cup (w_0(\Sigma^-) - \{w_0(-\alpha_s)\})$ , as recalled at the beginning of this section. As  $d \in \mathcal{D}$ , we have  $\alpha \circ \nu(\lambda) > 0$  for all  $\alpha \in \Sigma^+ \cap w_0(\Sigma^-)$ . We deduce that  $ds \notin \mathcal{D} \Leftrightarrow w_0(\alpha_s) \in \Sigma^+, w_0(\alpha_s) \circ \nu(\lambda) = 0$ .  $\square$

#### 4. $\mathfrak{h}$ -eigenspace in $\eta \otimes_{\mathfrak{h}} \mathcal{H}$

**Proposition 4.1.** *For any choice of lift  $\tilde{d}$  of  $d \in \mathcal{D}$  in  $\mathcal{D}(1)$ , the left  $\mathfrak{h}$ -module  $\mathcal{H}$  is free of basis  $(T_{\tilde{d}})_{d \in \mathcal{D}}$ , and the right  $\mathfrak{h}$ -module  $\mathcal{H}$  is free of basis  $(T_{\tilde{d}^{-1}})_{d \in \mathcal{D}}$ .*

**Proof.** To the set  $\mathcal{D}$  of distinguished representatives of the right  $W_0$ -cosets in  $W$  is associated a disjoint union  $W(1) = \bigsqcup_{d \in \mathcal{D}} W_0(1)\tilde{d}$ . Hence  $\mathcal{H}$  admits the  $R$ -bases

$$(T_{w\tilde{d}})_{w \in W_0(1), d \in \mathcal{D}} \text{ and } (T_{\tilde{d}^{-1}w})_{w \in W_0(1), d \in \mathcal{D}}.$$

A basis of  $\mathfrak{h}$  is  $(T_w)_{w \in W_0(1)}$ . By the braid relations,  $T_{w\tilde{d}} = T_w T_{\tilde{d}}$  and  $T_{\tilde{d}^{-1}w} = T_{\tilde{d}^{-1}} T_w$ , because  $\ell(wd) = \ell(w) + \ell(d)$ .  $\square$

**Remark 4.2.** An element of  $\mathcal{H}$  can be written as a sum  $\sum_{d \in \mathcal{D}} h_{\tilde{d}} T_{\tilde{d}}$ , where  $h_{\tilde{d}} \in \mathfrak{h}$ , and, for  $t \in Z_k$ ,

$$h_{\tilde{d}} T_{\tilde{d}} = h_{t\tilde{d}} T_{t\tilde{d}} = h_{t\tilde{d}} h_t T_{\tilde{d}}, \quad h_{\tilde{d}} = h_{t\tilde{d}} h_t.$$

The monoid  $\Lambda^+$  represents the orbits of  $W_0$  in  $\Lambda$ , and the double  $(W_0, W_0)$ -cosets of  $W$ , because  $W = \Lambda \rtimes W_0$ . The  $(\mathfrak{h}, \mathfrak{h})$ -module  $\mathcal{H}$  is the direct sum

$$\mathcal{H} = \bigoplus_{\lambda \in \Lambda^+} \mathfrak{h}(\lambda) \tag{23}$$

of the  $(\mathfrak{h}, \mathfrak{h})$ -submodules  $\mathfrak{h}(\lambda)$  of  $R$ -basis  $(T_w)_{w \in W_0(1)\tilde{\lambda}W_0(1)}$ . We set  $\mathcal{D}(\lambda) := W_0\lambda W_0 \cap \mathcal{D}$ .



**Corollary 4.3.** *Let  $\lambda \in \Lambda^+$ . The left  $\mathfrak{h}$ -module  $\mathfrak{h}(\lambda)$  is free of basis  $(T_{\bar{d}})_{d \in \mathcal{D}(\lambda)}$ , and the right  $\mathfrak{h}$ -module  $\mathfrak{h}(\lambda)$  is free of basis  $(T_{\bar{d}^{-1}})_{d \in \mathcal{D}(\lambda^{-1})}$ .*

Let  $\eta$  be a character of  $\mathfrak{h}$  of parameters  $(\chi, S_\eta)$ . Let  $\lambda \in \Lambda^+$ . By Corollary 4.3, an  $R$ -basis of  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  is

$$(1 \otimes T_{\bar{d}})_{d \in \mathcal{D}(\lambda)}. \tag{24}$$

When the algebra  $\mathcal{H}$  arises from a split reductive  $p$ -adic group  $G$ , Ollivier proved that the right  $\mathfrak{h}$ -module  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  has multiplicity 1 (private communication by email March 2014). This property is general, and the characters of  $\mathfrak{h}$  contained in  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  admit the following description.

**Proposition 4.4.** *Let  $\eta_1$  be a character of  $\mathfrak{h}$  of parameters  $(\chi_1, S_{\eta_1})$ . The  $\eta_1$ -eigenspace of  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  is not 0 if and only if  $(\eta_1, \eta, \lambda)$  satisfies*

$$\chi_1 = \chi^\lambda, \quad S_{\eta_1} \cap S_\lambda = S_\eta \cap S_\lambda.$$

When  $(\eta_1, \eta, \lambda)$  satisfies these conditions, the  $\eta_1$ -eigenspace of  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  has dimension 1 and is generated by  $1 \otimes \mathcal{E}_\lambda$  (defined in Theorem 1.2).

**Proof.** Let  $\mathcal{E} \in \eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$ . We write (24)  $\mathcal{E} = \sum_{d \in \mathcal{D}(\lambda)} a_{\bar{d}} \otimes T_{\bar{d}}$ , where  $a_{\bar{d}} \in R$ , and, for  $t \in Z_k$ ,

$$a_{\bar{d}} \otimes T_{\bar{d}} = a_{t\bar{d}} \otimes T_{t\bar{d}} = \chi(t)a_{t\bar{d}} \otimes T_{\bar{d}}, \quad a_{\bar{d}} = \chi(t)a_{t\bar{d}}.$$

For  $(w, t) \in W \times Z_k$  and a lift  $\tilde{w}$  of  $w$  in  $W(1)$ , using the notation of §§ 1.2 and 1.4,

$$(1 \otimes T_{\tilde{w}})T_t = 1 \otimes (w \bullet t)T_{\tilde{w}} = \chi^w(t) \otimes T_{\tilde{w}}. \tag{25}$$

Using Proposition 2.2 and (25),  $\mathcal{E}$  is an  $\mathfrak{h}$ -eigenvector of  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  with eigenvalue  $\eta_1$  if and only if  $\mathcal{E}$  satisfies

$$\mathcal{E} = \sum_{d \in \mathcal{D}(\lambda), \chi^d = \chi_1} a_{\bar{d}} \otimes T_{\bar{d}} \neq 0, \tag{26}$$

$$\mathcal{E}T_{\bar{s}} = 0 \quad \text{for } s \in S - S_{\eta_1}, \quad \mathcal{E}T_{\bar{s}} = \chi_1(c_{\bar{s}})\mathcal{E} \quad \text{for } s \in S_{\eta_1}. \tag{27}$$

The space  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  does not contain a  $\mathfrak{h}$ -eigenvector with eigenvalue  $\eta_1$  when the set  $X = \{d \in \mathcal{D}(\lambda), \chi^d = \chi_1\}$  is empty, and the proposition is obviously true. When  $\nu(\lambda) = 0$ , we have  $\mathcal{D}(\lambda) = \{\lambda\}$  by Lemma 3.5, and the proposition is true, because it is clearly true when  $X = \{\lambda\}$ .

We suppose that  $\nu(\lambda) \neq 0$ . For  $s \in S$ , the set  $X$  is the disjoint union of the subsets

$$X_1(s) = \{d \in X \mid \ell(ds) = \ell(d) + 1, ds \in \mathcal{D}\},$$

$$X_2(s) = \{d \in X \mid ds \notin \mathcal{D}\},$$

$$X_3(s) = \{d \in X \mid \ell(ds) = \ell(d) - 1\}.$$

In  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$ , we have

$$(1 \otimes T_{\bar{d}})T_{\bar{s}} = 1 \otimes T_{\bar{d}}T_{\bar{s}} = \begin{cases} 1 \otimes T_{\bar{d}\bar{s}} & (d \in X_1(s)) \\ \eta(T_{\bar{d}\bar{s}\bar{d}^{-1}}) \otimes T_{\bar{d}} & (d \in X_2(s)) \\ \chi_1(c_{\bar{s}}) \otimes T_{\bar{d}} & (d \in X_3(s)). \end{cases}$$

Indeed, if  $\ell(ds) = \ell(d) + 1$ , the braid relations imply that  $T_{\tilde{d}}T_{\tilde{s}} = T_{\tilde{d}\tilde{s}}$ . If  $ds \notin \mathcal{D}$ , by Lemma 3.6,  $dsd^{-1} \in S$ ,  $T_{\tilde{d}\tilde{s}} = T_{\tilde{d}\tilde{s}\tilde{d}^{-1}\tilde{d}} = T_{\tilde{d}\tilde{s}\tilde{d}^{-1}}T_{\tilde{d}}$ . If  $\ell(ds) = \ell(d) - 1$ , the braid and quadratic relations imply that  $T_{\tilde{d}}T_{\tilde{s}} = T_{\tilde{d}\tilde{s}^{-1}}T_{\tilde{s}}^2 = T_{\tilde{d}\tilde{s}^{-1}}c_{\tilde{s}}T_{\tilde{s}} = \tilde{d}c_{\tilde{s}}\tilde{d}^{-1}T_{\tilde{d}\tilde{s}^{-1}}T_{\tilde{s}} = \tilde{d}c_{\tilde{s}}\tilde{d}^{-1}T_{\tilde{d}}$ .

Multiplying (26) by  $T_{\tilde{s}}$  on the right,

$$\mathcal{E}T_{\tilde{s}} = \sum_{d \in X_1(s)} a_{\tilde{d}} \otimes T_{\tilde{d}\tilde{s}} + \sum_{d \in X_2(s)} \eta(T_{\tilde{d}\tilde{s}\tilde{d}^{-1}})a_{\tilde{d}} \otimes T_{\tilde{d}} + \sum_{d \in X_3(s)} \chi_1(c_{\tilde{s}})a_{\tilde{d}} \otimes T_{\tilde{d}}.$$

As  $X_1(s)s = X_3(s)$ , the expansion of  $\mathcal{E}T_{\tilde{s}}$  in the basis (24) of  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$  is

$$\mathcal{E}T_{\tilde{s}} = \sum_{d \in X_2(s)} \eta(T_{\tilde{d}\tilde{s}\tilde{d}^{-1}})a_{\tilde{d}} \otimes T_{\tilde{d}} + \sum_{d \in X_3(s)} (a_{\tilde{d}(\tilde{s})^{-1}} + \chi_1(c_{\tilde{s}})a_{\tilde{d}}) \otimes T_{\tilde{d}}. \tag{28}$$

Relations (27) are equivalent to the following.

For  $d \in X_2(s)$ ,

$$\eta(T_{\tilde{d}\tilde{s}\tilde{d}^{-1}})a_{\tilde{d}} = 0 \quad \text{if } s \in S - S_{\eta_1}, \quad \eta(T_{\tilde{d}\tilde{s}\tilde{d}^{-1}})a_{\tilde{d}} = \chi_1(c_{\tilde{s}})a_{\tilde{d}} \quad \text{if } s \in S_{\eta_1}. \tag{29}$$

For  $d \in X_1(s)$ ,

$$0 = \chi_1(c_{\tilde{s}})a_{\tilde{d}} \quad \text{if } s \in S_{\eta_1}.$$

For  $d \in X_3(s)$ ,

$$a_{\tilde{d}(\tilde{s})^{-1}} = \chi_1(c_{\tilde{s}})a_{\tilde{d}} \quad \text{if } s \in S - S_{\eta_1}, \quad a_{\tilde{d}(\tilde{s})^{-1}} = 0 \quad \text{if } s \in S_{\eta_1}.$$

The relations for  $d \in X_3(s) = X_1(s)s^{-1}$  are equivalent to the following.

For  $d \in X_1(s)$ ,

$$a_{\tilde{d}} = \chi_1(c_{\tilde{s}})a_{\tilde{d}\tilde{s}} \quad \text{if } s \in S - S_{\eta_1}, \quad a_{\tilde{d}} = 0 \quad \text{if } s \in S_{\eta_1}.$$

The relations associated to  $\bigcup_{s \in S} (X_1(s) \cup X_3(s))$  are equivalent to

$$a_{\tilde{d}} = 0 \quad \text{if } d \in \bigcup_{s \in S_{\eta_1}} X_1(s). \tag{30}$$

$$a_{\tilde{d}} = \chi_1(c_{\tilde{s}})a_{\tilde{d}\tilde{s}} \quad \text{if } d \in \bigcup_{s \in S - S_{\eta_1}} X_1(s). \tag{31}$$

As  $\nu(\lambda) \neq 0$ , we have  $X = \bigcup_{s \in S} (X_1(s) \cup X_3(s))$ , because  $d = \lambda w_0 \in \mathcal{D}(\lambda)$ ,  $w_0 \in W_0$  (Lemma 3.5), satisfies  $ds \notin \mathcal{D}(\lambda)$  for all  $s \in S$  if and only if  $w_0(\alpha_s) \in \Sigma^+$ ,  $w_0(\alpha_s) \circ \nu(\lambda) = 0$  for all  $s \in S$  (Lemma 3.8), and this is equivalent to  $\nu(\lambda) = 0$ .

For  $d = \lambda w_0 \in X$  and  $\tilde{d} = \tilde{\lambda}\tilde{w}_0$ , the relations (30), (31) are equivalent to

$$a_{\tilde{d}} = \chi_1(c_{\tilde{w}_0})^{-1}a_{\tilde{\lambda}} \quad \text{if } w_0 \text{ in } \langle S_{\chi_1} - S_{\eta_1} \rangle, \quad a_{\tilde{d}} = 0 \text{ otherwise.} \tag{32}$$

With the notation  $\mathcal{E}_{\tilde{\lambda}}, Y_{\tilde{\lambda}}$  introduced in Theorem 1.2, (32) implies that  $\mathcal{E} = a_{\tilde{\lambda}} \otimes \mathcal{E}_{\tilde{\lambda}}$ . If  $\eta_1$  is contained in  $\eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$ , then  $a_{\tilde{\lambda}} \neq 0$ , the multiplicity of  $\eta_1$  is 1, and  $\chi_1 = \chi^{\tilde{\lambda}}$ .

To end the proof of the proposition, we show that the conditions associated to  $\bigcup_{s \in S} X_2(s)$  on  $\mathcal{E} = 1 \otimes \mathcal{E}_{\tilde{\lambda}}$  are

$$S_{\lambda} - S_{\eta_1} = S_{\lambda} - S_{\eta}. \tag{33}$$

Relation (29) for  $d \in X_2(s)$  is always true if  $a_{\tilde{d}} = 0$ . For  $\mathcal{E} = 1 \otimes \mathcal{E}_{\tilde{\lambda}}$ , we have  $a_{\tilde{d}} \neq 0 \Leftrightarrow d \in \lambda Y_{\tilde{\lambda}}$ . By Lemma 3.8,  $d \in \lambda Y_{\tilde{\lambda}} \cap X_2(s) \Leftrightarrow d = \lambda w_0$ , where

$$w_0 \in \langle S_{\chi_1} - S_{\eta_1} \rangle, \quad \ell(\lambda w_0) = \ell(\lambda) - \ell(w_0), \quad \chi_1^{w_0} = \chi_1, \quad dsd^{-1} = w_0 s w_0^{-1} \in S_{\tilde{\lambda}}.$$

For  $s_d = dsd^{-1} \in S_{\tilde{\lambda}}$  and  $\tilde{s}_d = \tilde{d}\tilde{s}_d\tilde{d}^{-1}$ , we have  $\chi(c_{\tilde{s}_d}) = \chi(\tilde{d}c_{\tilde{s}_d}\tilde{d}^{-1}) = \chi^d(c_{\tilde{s}}) = \chi_1(c_{\tilde{s}})$ . The conditions associated to  $\bigcup_{s \in S} X_2(s)$  are as follows: for all  $d \in \lambda Y_{\tilde{\lambda}} \cap X_2(s)$ ,

$$s_d \in S - S_{\eta} \quad \text{if } s \in S - S_{\eta_1} \quad \text{and} \quad s_d \in S_{\eta} \quad \text{if } s \in S_{\eta_1}; \tag{34}$$

that is,  $s \in S_{\eta_1} \Leftrightarrow s_d \in S_{\eta}$  when  $s \in S, d \in \lambda Y_{\tilde{\lambda}} \cap X_2(s)$ . They are equivalent to (33); that is,  $s \in S_{\eta_1} \Leftrightarrow s \in S_{\eta}$  when  $s \in S_{\tilde{\lambda}}$ , because, for  $d \in \lambda Y_{\tilde{\lambda}} \cap X_2(s)$ , we have  $s_d \in S_{\tilde{\lambda}}$ , and  $\langle s, S_{\chi_1} - S_{\eta_1} \rangle = \langle s_d, S_{\chi_1} - S_{\eta_1} \rangle$ ; hence  $s_d \in S_{\eta_1} \Leftrightarrow s \in S_{\eta_1}$ .  $\square$

Let  $\eta, \eta_1$  be two characters of  $\mathfrak{h}$  of parameters  $(\chi, S_{\eta}), (\chi_1, S_{\eta_1})$ , and let  $o, o_1$  be an orientation such that  $\eta = \chi_o, \eta_1 = (\chi_1)_{o_1}$ .

By the decomposition (23), the  $\mathfrak{h}$ -module  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  is a direct sum of  $\mathfrak{h}$ -submodules:

$$\eta \otimes_{\mathfrak{h}} \mathcal{H} = \bigoplus_{\lambda \in \Lambda^+} \eta \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda). \tag{35}$$

**Proposition 4.5.** *The character  $\eta_1$  of  $\mathfrak{h}$  is contained in  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  if and only if there exists  $\lambda$  such that  $(\eta, \eta_1, \lambda)$  satisfies*

$$\lambda \in \Lambda^+, \quad \chi_1 = \chi^{\lambda}, \quad S_{\eta_1} \cap S_{\tilde{\lambda}} = S_{\eta} \cap S_{\tilde{\lambda}}.$$

The  $\eta_1$ -eigenspace of  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  admits the  $R$ -basis  $(1 \otimes \mathcal{E}_{\tilde{\lambda}})$  for all  $\lambda$  such that  $(\eta, \eta_1, \lambda)$  satisfies these conditions.

For  $(\eta, \eta_1, \lambda)$  as in Proposition 4.5, we denote by  $\Phi_{\tilde{\lambda}}$  the  $\mathcal{H}$ -intertwiner

$$\Phi_{\tilde{\lambda}} : 1 \otimes 1 \mapsto 1 \otimes \mathcal{E}_{\tilde{\lambda}} : \eta_1 \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \eta \otimes_{\mathfrak{h}} \mathcal{H}.$$

**Corollary 4.6.** *An  $R$ -basis of  $\text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H})$  is  $(\Phi_{\tilde{\lambda}})$  for all  $\lambda$  such that  $(\eta, \eta_1, \lambda)$  satisfies the conditions of Proposition 4.5.*

Taking  $\eta = \eta_1$ , and recalling the  $\Lambda^+$ -fixator  $\Lambda_{\tilde{\lambda}}^+$  of  $\chi$  (12), we obtain the following.

**Corollary 4.7.**  *$(\Phi_{\tilde{\lambda}})_{\lambda \in \Lambda_{\tilde{\lambda}}^+}$  is a basis of the spherical Hecke algebra  $\mathcal{H}(\eta, \mathfrak{h})$ .*

To obtain a basis of the spherical Hecke algebra satisfying (14), for an orientation  $o$  we construct  $\mathfrak{h}$ -eigenvectors of the form

$$1 \otimes E_o(\tilde{\lambda}) \in \chi_o \otimes_{\mathfrak{h}} \mathcal{H}$$

with  $\tilde{\lambda} \in \Lambda^+(1)$ , where, as in § 1.2,  $(E_o(\tilde{w}))_{\tilde{w} \in W(1)}$  is the alcove walk basis of  $\mathcal{H}$  associated to  $o$  [15, § 5.3 Corollary 5.26], and the character  $\chi_o$  of  $\mathfrak{h}$  is as in Lemma 2.4.

**Lemma 4.8.** *Let  $\lambda \in \Lambda$ . We have, in  $\chi_o \otimes_{\mathfrak{h}} \mathcal{H}$ ,*

$$1 \otimes E_o(\tilde{\lambda}) - 1 \otimes T_{\tilde{\lambda}} \in \sum_d R \otimes T_{\tilde{d}},$$

where  $d$  runs over the elements of  $\mathcal{D}$  satisfying  $d < \lambda$  and  $\chi^d = \chi^{\lambda}$ . If  $\lambda \in \Lambda^+$ , then  $1 \otimes E_o(\tilde{\lambda}) \neq 0$  is a  $Z_k$ -eigenvector of eigenvalue  $\chi^{\lambda}$ .

**Proof.** For  $t \in Z_k$ , we have [15, Example 5.30]  $E_o(\tilde{\lambda})T_t = T_{\lambda(t)}E_o(\tilde{\lambda})$ ,  $T_{\tilde{\lambda}}T_t = T_{\lambda(t)}T_{\tilde{\lambda}}$ ; hence  $1 \otimes E_o(\tilde{\lambda})T_t = \chi^\lambda(t) \otimes E_o(\tilde{\lambda})$ ,  $(1 \otimes T_{\tilde{\lambda}})T_t = \chi^\lambda(t) \otimes T_{\tilde{\lambda}}$ . With the disjoint decomposition  $W(1) = \bigcup_{d \in \mathcal{D}} W_0(1)d$  and the triangular decomposition of  $E_o(\tilde{\lambda})$  in the basis  $(T_{\tilde{w}})_{\tilde{w} \in W(1)}$  of  $\mathcal{H}$  [15, Corollary 5.26], if  $1 \otimes E_o(\tilde{\lambda}) \neq 0$  is a  $Z_k$ -eigenvector of eigenvalue  $\chi^\lambda$ , we have

$$1 \otimes E_o(\tilde{\lambda}) - 1 \otimes T_{\tilde{\lambda}} \in \sum_{d \in \mathcal{D}, \chi^d = \chi^\lambda} \sum_{\tilde{w} \in W_0(1), wd < \lambda} R \otimes T_{\tilde{w}d}.$$

As  $\ell(wd) = \ell(w) + \ell(d)$ , by the braid relations,  $1 \otimes T_{\tilde{w}d} = 1 \otimes T_{\tilde{w}}T_d = \eta(T_{\tilde{w}}) \otimes T_d$ ,

$$\sum_{\tilde{w} \in W_0(1), wd < \lambda} R(1 \otimes T_{\tilde{w}d}) = R(1 \otimes T_{\tilde{\lambda}}).$$

As  $d < wd$  for  $w \in W_0$ , we deduce that

$$1 \otimes E_o(\tilde{\lambda}) - 1 \otimes T_{\tilde{\lambda}} \in \sum_{d \in \mathcal{D}, \chi^d = \chi^\lambda, d < \lambda} R \otimes T_d.$$

For  $\lambda \in \Lambda^+$ ,  $1 \otimes E_o(\tilde{\lambda})$  is not 0, because  $\Lambda^+ \subset \mathcal{D}$ , and  $(1 \otimes T_d)_{d \in \mathcal{D}}$  is a basis of  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$  (Proposition 4.1). □

**Lemma 4.9.** *Let  $\lambda \in \Lambda$ . Then  $1 \otimes E_o(\tilde{\lambda}) \in \chi_o \otimes_{\mathfrak{h}} \mathcal{H}$  is a  $\mathfrak{h}$ -eigenvector of eigenvalue  $(\chi_1)_{o_1}$  if and only if  $1 \otimes E_o(\tilde{\lambda}) \neq 0$  and*

$$\chi_1 = \chi^\lambda, \quad 1 \otimes E_o(\tilde{\lambda})E_{o_1}(\tilde{s}) = 0 \quad \text{for all } s \in S.$$

**Proof.** By Lemma 4.8(ii),  $1 \otimes E_o(\tilde{\lambda})$  is a  $\mathfrak{h}$ -eigenvector with eigenvalue  $\eta_1$  if and only if  $1 \otimes E_o(\tilde{\lambda}) \neq 0$ , and  $\chi_1 = \chi^\lambda$ ,  $(1 \otimes E_o(\tilde{\lambda}))E_{o_1}(\tilde{s}) = 0$  for all  $s \in S$  (Lemma 2.4). We have  $(1 \otimes E_o(\tilde{\lambda}))E_{o_1}(\tilde{s}) = 1 \otimes E_o(\tilde{\lambda})E_{o_1}(\tilde{s})$ . □

**Lemma 4.10.** *Let  $\lambda \in \Lambda^+$ . Then  $1 \otimes E_o(\tilde{\lambda})$  is a  $\mathfrak{h}$ -eigenvector of eigenvalue  $(\chi^\lambda)_o$  if and only if  $\eta(E_o(\tilde{s})) = 0$  for all  $s \in S$  such that  $\ell(\lambda s) = 1 + \ell(\lambda)$ .*

**Proof.** Let  $s \in S$ .

If  $\ell(\lambda s) = \ell(\lambda) - 1$ , then  $E_o(\tilde{\lambda})E_o(\tilde{s}) = 0$  by the product formula.

If  $\ell(\lambda s) = \ell(\lambda) + 1$ , then  $E_o(\tilde{\lambda})E_o(\tilde{s}) = E_o(\lambda\tilde{s}) = E_o(\tilde{s}\tilde{s}^{-1}\lambda\tilde{s}) = E_o(\tilde{s})E_{o_{\bullet s}}(\tilde{s}^{-1}\lambda\tilde{s})$ .

The latter equality follows from the fact that the length is constant on a  $W_0$ -orbit in  $\Lambda$ . It implies that  $1 \otimes E_o(\tilde{s})E_{o_{\bullet s}}(\tilde{s}^{-1}\lambda\tilde{s}) = \eta(E_o(\tilde{s})) \otimes E_o(\tilde{\lambda})$ . Apply Lemmas 4.8 and 4.9. □

**Proposition 4.11.** *Let  $\lambda \in \Lambda^+$ . Then,*

$1 \otimes E_o(\tilde{\lambda})$  is a  $\mathfrak{h}$ -eigenvector in  $\chi_o \otimes_{\mathfrak{h}} \mathcal{H}$  of eigenvalue  $(\chi^\lambda)_o$ , and  $\mathcal{E}_{\tilde{\lambda}}$  is the component of  $1 \otimes E_o(\tilde{\lambda})$  in  $\chi_o \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda)$ .

**Proof.** Use Lemmas 2.4 and 4.10 for the first assertion. The non-zero components of  $1 \otimes E_o(\tilde{\lambda})$  in the direct decomposition (35) are  $\mathfrak{h}$ -eigenvectors of eigenvalue  $(\chi^\lambda)_o$ . Apply Proposition 4.4 and Lemma 4.8 for the second assertion. □

**Corollary 4.12.** *If  $o = o_1$  (Lemma 2.6), an  $R$ -basis of  $\text{Hom}_{\mathcal{H}}((\chi_1)_o \otimes_{\mathfrak{h}} \mathcal{H}, \chi_o \otimes_{\mathfrak{h}} \mathcal{H})$  is  $(1 \otimes E_o(\tilde{\lambda}))$  for all  $\lambda$  such that  $(\chi_o, (\chi_1)_o, \lambda)$  satisfies the conditions of Proposition 4.5.*

**Proposition 4.13.** *For each  $\lambda \in \Lambda_{\chi}^+$ , we have an injective  $\mathcal{H}$ -intertwiner*

$$\Phi_{o, \tilde{\lambda}} : 1 \otimes 1 \mapsto 1 \otimes E_o(\tilde{\lambda}) : \chi_o \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \chi_o \otimes_{\mathfrak{h}} \mathcal{H}.$$

$(\Phi_{o, \tilde{\lambda}})_{\lambda \in \Lambda_{\chi}^+}$  is an  $R$ -basis satisfying (14) of the spherical Hecke algebra  $\mathcal{H}(\chi_o, \mathfrak{h})$ .

**Proof.** By Corollary 4.12 and the product formula (8),  $(\Phi_{o, \tilde{\lambda}})_{\lambda \in \Lambda_{\chi}^+}$  is an  $R$ -basis of  $\mathcal{H}(\chi_o, \mathfrak{h})$  satisfying (14).

If  $\Phi_{o, \tilde{\lambda}}$  is not injective,  $\text{Ker } \Phi_{o, \tilde{\lambda}}$  contains a simple character  $\eta_1$  of  $\mathfrak{h}$ , and  $\Phi_{o, \tilde{\lambda}} \circ \Phi_1 = 0$  for some non-zero  $\Phi_1 \in \text{End}_{\mathfrak{h}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H})$ .

Expanding  $\Phi_1(1 \otimes 1) = \sum_{\mu \in \Lambda^+} a_{\tilde{\mu}} \otimes E_o(\tilde{\mu})$ ,  $a_{\tilde{\mu}} \in R$ , in the basis  $(1 \otimes E_o(\tilde{\mu}))_{\mu \in \Lambda^+}$  of  $\eta \otimes_{\mathfrak{h}} \mathcal{H}$ , and using the product formula  $E_o(\tilde{\lambda})E_o(\tilde{\mu}) = E_o(\tilde{\lambda}\tilde{\mu})$ , the decomposition of  $(\Phi_{o, \tilde{\lambda}} \circ \Phi_1)(1 \otimes 1)$  in this basis is

$$\sum_{\mu \in \Lambda^+} \Phi_{o, \tilde{\lambda}}(a_{\tilde{\mu}} \otimes E_o(\tilde{\mu})) = \sum_{\mu \in \Lambda^+} a_{\tilde{\mu}} \otimes E_o(\tilde{\lambda})E_o(\tilde{\mu}) = \sum_{\mu \in \Lambda^+} a_{\tilde{\mu}} \otimes E_o(\tilde{\lambda}\tilde{\mu}).$$

We have  $\Phi_1 \neq 0 \Leftrightarrow \Phi_1(1 \otimes 1) \neq 0 \Leftrightarrow a_{\tilde{\mu}} \neq 0$  for some  $\mu \in \Lambda^+ \Leftrightarrow (\Phi_{o, \tilde{\lambda}} \circ \Phi_1)(1 \otimes 1) \neq 0 \Leftrightarrow \Phi_{o, \tilde{\lambda}} \circ \Phi_1 \neq 0$ . □

**Corollary 4.14.**  $1 \otimes E_o(\tilde{\lambda}) = 0$  in  $\chi_o \otimes_{\mathfrak{h}} \mathcal{H}$  if  $\lambda \in \Lambda - \Lambda^+$ .

**Proof.** Let  $\lambda \in \Lambda - \Lambda^+$ . We choose  $\mu \in \Lambda_{\chi}^+$  not 0. Then  $\Phi_{o, \tilde{\mu}}$  of  $\text{End}_{\mathfrak{h}} \eta \otimes_{\mathfrak{h}} \mathcal{H}$  is injective (Proposition 4.13) and  $\Phi_{o, \tilde{\mu}}(1 \otimes E_o(\tilde{\lambda})) = 1 \otimes E_o(\tilde{\mu})E_o(\tilde{\lambda})$ . As  $\mu, \lambda$  belong to different closed Weyl chambers,  $E_o(\tilde{\mu})E_o(\tilde{\lambda}) = 0$ ; hence  $1 \otimes E_o(\tilde{\lambda}) = 0$ . □

More generally, if  $(\chi_o, (\chi_1)_o, \lambda)$  satisfies the conditions of Proposition 4.5, we have the non-zero  $\mathcal{H}$ -intertwiner

$$\Phi_{o, \tilde{\lambda}} : 1 \otimes 1 \mapsto 1 \otimes E_o(\tilde{\lambda}) : (\chi_1)_o \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow \chi_o \otimes_{\mathfrak{h}} \mathcal{H}.$$

An  $R$ -basis of  $\text{Hom}_{\mathcal{H}}((\chi_1)_o \otimes_{\mathfrak{h}} \mathcal{H}, \chi_o \otimes_{\mathfrak{h}} \mathcal{H})$  is  $(\Phi_{o, \tilde{\lambda}})$  for all  $\lambda$  such that  $(\chi_o, (\chi_1)_o, \lambda)$  satisfies the conditions of Proposition 4.5.

We fix  $x_1 \in \Lambda$  such that  $\chi_1 = \chi^{x_1}$ . For  $\lambda \in \Lambda$ ,  $\chi_1 = \chi^{\lambda x_1} \Leftrightarrow \lambda \in \Lambda_{\chi}$ . We embed  $\text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H})$  into the algebra  $e_{\chi} R[\Lambda_{\chi}]$  (§1.4) by the  $R$ -linear map

$$S_{\eta_1, \eta, \tilde{x}_1} : \text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H}) \rightarrow e_{\chi} R[\Lambda_{\chi}], \tag{36}$$

$$\Phi_{o, \tilde{\lambda}\tilde{x}_1} \mapsto e_{\chi} \tilde{\lambda} \quad (\lambda \in \Lambda_{\chi} \cap \Lambda^+ x_1^{-1}), \tag{37}$$

where  $\tilde{\lambda}, \tilde{x}_1 \in \Lambda(1)$  lift  $\lambda, x_1$ . If  $\eta = \eta_1$  and  $\tilde{x}_1 = 1$ , the map  $S_{\eta, \eta, 1} = S_{\eta, \eta}$  embeds the spherical Hecke algebra  $\mathcal{H}(\eta, \mathfrak{h}) = \text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H})$  into the algebra  $e_{\chi} R[\Lambda_{\chi}]$

$$S_{\eta, \eta} : \mathcal{H}(\eta, \mathfrak{h}) \rightarrow e_{\chi} R[\Lambda_{\chi}]. \tag{38}$$

**Lemma 4.15.** *The composition*

$$(A, B) \mapsto B \circ A : \text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H}) \times \text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H}) \rightarrow \text{Hom}_{\mathcal{H}}(\eta_1 \otimes_{\mathfrak{h}} \mathcal{H}, \eta \otimes_{\mathfrak{h}} \mathcal{H}),$$

corresponds to the product  $S_{\eta_1, \eta, \tilde{x}_1}(A \circ B) = S_{\eta, \eta}(B)S_{\eta_1, \eta, \tilde{x}_1}(A)$  in  $e_{\chi}R[\Lambda_{\chi}]$ .

**Proof.** For  $\lambda \in \Lambda_{\chi}^+$  and  $\lambda_1 \in \Lambda^+$ ,  $\chi^{\lambda_1} = \chi_1$ ,  $S_{\eta_1} \cap S_{\lambda_1} = S_{\eta} \cap S_{\lambda_1}$ , we have

$$\Phi_{o, \tilde{\lambda}} \circ \Phi_{o, \tilde{\lambda}_1}(1 \otimes 1) = \Phi_{o, \tilde{\lambda}}(1 \otimes E_o(\tilde{\lambda}_1)) = 1 \otimes E_o(\tilde{\lambda})E_o(\tilde{\lambda}_1) = 1 \otimes E_o(\tilde{\lambda}\tilde{\lambda}_1),$$

by the product formula (8). Hence  $\Phi_{o, \tilde{\lambda}} \circ \Phi_{o, \tilde{\lambda}_1} = \Phi_{o, \tilde{\lambda}\tilde{\lambda}_1}$  and  $S_{\eta_1, \eta, \tilde{x}_1}(\Phi_{o, \tilde{\lambda}} \circ \Phi_{o, \tilde{\lambda}_1}) = e_{\chi} \tilde{\lambda}\tilde{\lambda}_1(\tilde{x}_1)^{-1}$ . As  $e_{\chi}$  is a central idempotent of  $R[\Lambda_{\chi}]$ , we have  $e_{\chi} \tilde{\lambda}\tilde{\lambda}_1(\tilde{x}_1)^{-1} = e_{\chi} \tilde{\lambda}e_{\chi} \tilde{\lambda}_1(\tilde{x}_1)^{-1} = S_{\eta, \eta}(\Phi_{o, \tilde{\lambda}})S_{\eta_1, \eta, \tilde{x}_1}(\Phi_{o, \tilde{\lambda}_1})$ . □

### 5. Centers

We make the same hypotheses as in § 1.2, and we suppose that  $\Lambda_T$  exists.

As  $\tilde{\Lambda}_T$  is central in  $\Lambda(1)$ , the action of  $W(1)$  on  $\tilde{\Lambda}_T$  factorizes through an action of  $W_0$ , and the  $R$ -module  $\mathcal{A}_o(\Lambda_T)$  of basis  $(E_o(\tilde{\mu}))_{\mu \in \Lambda_T}$  is a  $W_0$ -stable subalgebra of the center  $\mathcal{Z}_o$  of  $\mathcal{A}_o$ , for any orientation  $o$ . The quotient map  $\Lambda_T(1) \rightarrow \Lambda_T$  of splitting  $\mu \mapsto \tilde{\mu}$  is  $W_0$ -equivariant. For  $\mu \in \Lambda_T$  of  $W_0$ -conjugacy class  $C(\mu)$ , and  $\tilde{C}(\mu)$  the  $W_0$ -conjugacy class of  $\tilde{\mu}$ , the set  $\nu(C(\mu))$  contains a single element in the dominant closed Weyl chamber, and

$$\ell(\mu) = 0 \Leftrightarrow \nu(\mu) = 0 \Leftrightarrow \mu \in \Lambda_T^{W_0} \Leftrightarrow \tilde{C}(\mu) = \tilde{\mu}. \tag{39}$$

By axiom (T1) (1.2), a  $W(1)$ -conjugacy class  $\tilde{C}$  is finite if and only if  $\tilde{C} \subset \Lambda(1)$ .

In the following theorem,  $R$  is any commutative ring.

**Theorem 5.1.** *The center  $\mathcal{Z}$  of  $\mathcal{H}_R(q_s, c_{\tilde{s}})$  is the algebra  $\mathcal{A}_o^{W(1)}$  of  $W(1)$ -invariants of  $\mathcal{A}_o$ , equal to the algebra  $\mathcal{Z}_o^{W_0}$  of the  $W_0$ -invariants of the center  $\mathcal{Z}_o$  of  $\mathcal{A}_o$ . The center  $\mathcal{Z}$  is a free  $R$ -module of basis (independent of the choice of the orientation  $o$ )*

$$E(\tilde{C}) = \sum_{\tilde{\lambda} \in \tilde{C}} E_o(\tilde{\lambda}) \quad \text{for } \tilde{C} \text{ running through the finite conjugacy classes of } W(1).$$

The involution  $\iota$  of  $\mathcal{H}$  satisfies, for any finite conjugacy class  $\tilde{C}$  of  $W(1)$ ,

$$\iota(E(\tilde{C})) = (-1)^{\ell(C)} E(\tilde{C}). \tag{40}$$

The algebra  $\mathcal{Z}_T = \mathcal{A}_o(\Lambda_T)^{W_0}$  of  $W_0$ -invariants of  $\mathcal{A}_o(\Lambda_T)$  is a central subalgebra of  $\mathcal{H}$ , and a free  $R$ -module of basis  $(E(\tilde{C}(\mu)))_{\mu \in \Lambda_T^+}$ .

The  $\mathcal{Z}_T$ -modules  $\mathcal{Z}$  and  $\mathcal{H}_R(q_s, c_{\tilde{s}})$  are finitely generated.

When the ring  $R$  is noetherian, the  $R$ -algebras  $\mathcal{Z}_T, \mathcal{Z}$ , and  $\mathcal{H}_R(q_s, c_{\tilde{s}})$  are finitely generated.

**Proof.** The steps of the proof are as follows.

- (1) The center  $\mathcal{Z}_o$  of  $\mathcal{A}_o$  is a free  $R$ -module of basis  $E_o(\tilde{c}) = \sum_{\tilde{\lambda} \in \tilde{c}} E_o(\tilde{\lambda})$  for all conjugacy classes  $\tilde{c}$  of  $\Lambda(1)$ .
- (2)  $\sum_{\tilde{\lambda} \in \tilde{c}} E_o(\tilde{\lambda})$  does not depend on the orientation  $o$ , and the center  $\mathcal{Z}$  is equal to  $\mathcal{A}_{o^-}^{W(1)}$  for the anti-dominant orientation  $o^-$ .
- (3) (a) The  $\mathcal{A}_o(\Lambda_T)^{W_0}$ -module  $\mathcal{A}_o(\Lambda_T)$  is finitely generated, and if  $R$  is noetherian the algebra  $\mathcal{A}_o(\Lambda_T)^{W_0}$  is finitely generated.  
 (b) The left  $\mathcal{A}_o(\Lambda_T)$ -module  $\mathcal{A}_o$  is finitely generated.  
 (c) The left  $\mathcal{A}_o$ -module  $\mathcal{H}_R(q_s, c_{\tilde{s}})$  is finitely generated.

The theorem is proved for the pro- $p$ -Iwahori Hecke algebra  $\mathcal{H}_R(G, I(1))$ , where the assertions on  $\mathcal{Z}_T$  are not formulated but are implicit in the proof. Properties (1), (2), (3)(a), (b) and (40) admit exactly the same proofs as in [16, Propositions 2.3, 2.7, Lemma 2.15 and Proposition 3.3]. The same is true for the property (3)(c) [16, Lemma 2.17], once we have strengthened the finiteness property [14, 1.6.3], [16, Lemma 2.16]. This is done in Lemma 5.3 below. As in [16, added in proof], this is a variant of the finiteness of the set of minimal elements in a subset  $L$  of  $\mathbb{Z}^n$  ( $n > 0$ ) [12, Lemma 4.2.18]. □

Let  $L$  be a group isomorphic to  $\mathbb{Z}^n$ . For  $a = (a_i), b = (b_i) \in \mathbb{Z}^n$ , we write  $b \leq a$  if  $|a_i| = |b_i| + |a_i - b_i|$  for all  $i$ . We write  $b < a$  if  $a \neq b, b \leq a$ ; we say that  $a \in L$  is minimal if  $b \in L, b \leq a$  implies that  $b = a$ .

**Lemma 5.2.** (1) Let  $a \in L$ . There exists  $b \in L$  minimal such that  $b \leq a$ .

(2) The set  $L_{min}$  of minimal elements in  $L$  is finite.

**Proof.** We have  $|a_i| = |b_i| + |a_i - b_i| \Leftrightarrow b_i = 0$  or  $a_i b_i > 0, |b_i| \leq |a_i|$ .

(1) If  $a$  is not minimal in  $L$ , we choose  $b < a$  and we reiterate. The processes stops after finitely many steps, because  $b < a$  implies that  $|b_i| \leq |a_i|$  for  $1 \leq i \leq n$ , and  $|b_i| \in \mathbb{N}$ .

(2) Suppose that  $L_{min}$  is infinite. If the set  $\{a_i \mid a \in L_{min}\}$  is finite,  $a_i$  is constant for  $a$  in an infinite subset of  $L_{min}$ . If the set  $\{a_i \mid a \in L_{min}\}$  is infinite,  $L_{min}$  contains a sequence  $(a(m))_{m \in \mathbb{N}}$  such that  $(a(m)_i)_{m \in \mathbb{N}}$  is strictly increasing positive or strictly decreasing negative. Hence  $L_{min}$  contains a sequence  $(a(m))_{m \in \mathbb{N}}$  such that, for all  $1 \leq i \leq n$ ,  $(a(m)_i)_{m \in \mathbb{N}}$  is either constant, or strictly increasing positive or strictly decreasing negative. For all  $i$  in the non-empty set where  $(a(m)_i)_{m \in \mathbb{N}}$  is not constant, we have  $a(m)_i a(m+1)_i > 0, |a(m)_i| < |a(m+1)_i|$  for all  $m \in \mathbb{N}$ . Hence  $a(m) < a(m+1)$  for all  $m \in \mathbb{N}$ . This contradicts the minimality of the  $a(m)$ . □

By axiom (T1),  $W = \bigsqcup_{(y, w_0) \in Y \times W_0} \Lambda_T y w_0$ . For  $(y, w_0) \in Y \times W_0$ , let

$$L(y, w_0) = \{\vec{\ell}(w) = (\ell_\gamma(w))_{\gamma \in \Sigma^+} \mid w \in \Lambda_T y w_0\},$$

where  $\ell(w) = \sum_{\gamma \in \Sigma^+} |\ell_\gamma(w)|$  and  $\ell_\gamma(w)$  as in [15, Propositions 5.7 and 5.9]. By Lemma 5.2, the set  $L(y, w_0)_{min}$  is finite. Let  $X_*(y, w_0)$  be a finite subset of  $\Lambda_T$  such that

$$L(y, w_0)_{min} = \{\vec{\ell}(w) \mid w \in X_*(y, w_0) y w_0\}.$$

Let  $X$  be the finite subset  $\bigcup_{(y,w_0) \in Y \times W_0} X_*(y, w_0)y$  of  $\Lambda$ . We have

$$\ell(w) = \ell(ww'^{-1}) + \ell(w') \quad \text{for } w, w' \in \Lambda w_0, \quad \vec{\ell}(w') \leq \vec{\ell}(w),$$

[16, Proof of Lemma 2.16(18)]. This implies the following.

**Lemma 5.3.** *For any  $(\lambda, w_0) \in \Lambda \times W_0$  there exists  $x \in X$  such that*

$$\lambda x^{-1} \in \Lambda_T, \quad \ell(\lambda w_0) = \ell(\lambda x^{-1}) + \ell(x w_0).$$

For a central element  $x$  of  $\mathcal{H}$ , the  $\mathcal{H}$ -intertwiner

$$\Phi_x : 1 \otimes h \mapsto 1 \otimes xh = 1 \otimes hx \quad \text{for } h \in \mathcal{H}. \tag{41}$$

is central in  $\mathcal{H}(\chi_o, \mathfrak{h})$  by Proposition 4.13 and

$$\begin{aligned} \Phi_x \circ \Phi_{o,\tilde{\lambda}}(1 \otimes 1) &= \Phi_x(1 \otimes E_{o,\tilde{\lambda}}) = 1 \otimes x E_{o,\tilde{\lambda}} \\ &= 1 \otimes E_{o,\tilde{\lambda}} x = \Phi_{o,\tilde{\lambda}}(1 \otimes x) = \Phi_{o,\tilde{\lambda}} \circ \Phi_x(1 \otimes 1). \end{aligned}$$

We denote by  $\mathcal{Z}(\chi_o, \mathfrak{h})$  the center of  $\mathcal{H}(\chi_o, \mathfrak{h})$ . The homomorphism

$$x \mapsto \Phi_x : \mathcal{Z} \rightarrow \mathcal{Z}(\chi_o, \mathfrak{h}) \tag{42}$$

may be not injective or not surjective.

**Proposition 5.4.** (1) *For  $\mu \in \Lambda_T^+$ , we have  $1 \otimes E(\tilde{C}(\mu)) = 1 \otimes E_o(\tilde{\mu})$  and  $\Phi_{E(\tilde{C}(\mu))} = \Phi_{o,\tilde{\mu}}$ .*

(2)  *$(\Phi_{o,\tilde{\mu}})_{\mu \in \Lambda_T^+}$  is a basis, independent of  $o$ , satisfying (14) of a central subalgebra  $\mathcal{Z}_T(\eta, \mathfrak{h})$  of the spherical algebra  $\mathcal{H}(\eta, \mathfrak{h})$ , and  $\mathcal{H}(\eta, \mathfrak{h})$  is a finitely generated  $\mathcal{Z}_T(\eta, \mathfrak{h})$ -module.*

**Proof.** (1) From Corollary 4.14,

$$1 \otimes E(\tilde{C}(\mu)) = \sum_{\tilde{\lambda} \in \tilde{C}(\mu) \cap \Lambda^+(1)} 1 \otimes E_o(\tilde{\lambda}) \quad \text{in } \chi_o \otimes \mathcal{H}.$$

For  $\mu \in \Lambda_T^+$  we have  $\tilde{C}(\mu) \cap \Lambda^+(1) = \{\tilde{\mu}\}$ . Hence  $1 \otimes E(\tilde{C}(\mu)) = 1 \otimes E_o(\tilde{\mu})$  and  $\Phi_{E(\tilde{C}(\mu))} = \Phi_{o,\tilde{\mu}}$ .

(2) The canonical isomorphism  $\mathcal{H}(\eta, \mathfrak{h}) \rightarrow e_\chi R[\Lambda_\chi^+]$  associated to the basis  $(\Phi_{o,\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  (Proposition 4.13) sends  $\mathcal{Z}_T(\eta, \mathfrak{h})$  to  $e_\chi R[\Lambda_T^+]$ , and  $e_\chi R[\Lambda_\chi^+]$  is a finitely generated  $e_\chi R[\Lambda_T^+]$ -module. □

### 6. Supersingular $\mathcal{H}$ -modules

We make the same hypotheses as in § 1.2 and we suppose that  $\Lambda_T$  exists. We construct different filtrations of  $\mathcal{H}$  which are all equivalent when the ring  $R$  is noetherian.

**Lemma 6.1.** *The  $R$ -module  $\mathcal{F}_{o,n}$  of basis  $\{E_o(\tilde{w}) \mid \tilde{w} \in W(1), \ell(w) \geq n\}$  for  $n \in \mathbb{N}$  is a right ideal of  $\mathcal{H}$ , for any orientation  $o$ .*



**Proof.** We have  $\mathcal{F}_{o,n}\mathcal{H} \subset \mathcal{F}_{o,n}$ , because, for  $\tilde{w} \in W(1)$ , a basis of  $\mathcal{H}$  is  $(E_{o\bullet w}(\tilde{w}'))_{\tilde{w}' \in W(1)}$ , and  $E_o(w)E_{o\bullet w}(\tilde{w}') = E_o(\tilde{w}\tilde{w}')$  if  $\ell(w) + \ell(w') = \ell(w\tilde{w}')$  and 0 otherwise.  $\square$

The length is constant on the projection  $C$  in  $W$  of a finite  $W(1)$ -conjugacy class  $\tilde{C}$ , and is denoted by  $\ell(\tilde{C}) = \ell(C)$ .

**Lemma 6.2.** *The  $R$ -module  $\mathcal{Z}_{\ell>0}$  of basis  $E(\tilde{C})$  for the finite  $W(1)$ -conjugacy classes  $\tilde{C}$  of positive length is an ideal of the center  $\mathcal{Z}$  of  $\mathcal{H}$ , stable by the involutive  $R$ -automorphism  $\iota$  (4).*

**Proof.** Let  $\tilde{C}_1, \tilde{C}_2$  be two finite  $W(1)$ -conjugacy classes. They are contained in  $\Lambda(1)$ . By the product formula,

$$E(\tilde{C}_1)E(\tilde{C}_2) = \sum_{\tilde{C}} a_{\tilde{C}} E(\tilde{C}), \tag{43}$$

where  $\tilde{C}$  runs over finite conjugacy classes with  $\ell(C) = \ell(C_1) + \ell(C_2)$ . The stability by  $\iota$  follows from (40).  $\square$

It is more convenient to replace the center  $\mathcal{Z}$  of  $\mathcal{H}$  by the central subalgebra  $\mathcal{Z}_T$  of basis  $(E(\tilde{C}(\mu)))_{\mu \in X^+(T)}$  which admits better properties.

**Lemma 6.3.** *We have*

$$\mathcal{Z}_T = \mathcal{R}_T \oplus \mathcal{Z}_{T,\ell>0},$$

where  $\mathcal{R}_T$  is the algebra of basis  $(T_{\tilde{\mu}})_{\mu \in \Lambda_T^{w_0}}$ , isomorphic to  $R[\Lambda_T^{w_0}]$ , and  $\mathcal{Z}_{T,\ell>0}$  is the ideal of  $\mathcal{Z}_T$  of basis  $(E(\tilde{C}(\mu)))_{\mu \in \Lambda_T^+, \ell(\mu)>0}$ .

The algebras  $\mathcal{R}_T$  and  $\mathcal{Z}_{T,\ell>0}$  are stable by the involutive automorphism  $\iota$ .

**Proof.** The proof is straightforward.  $\square$

The  $R$ -module  $\mathcal{F}_{T,o,n}$  of basis  $(E_o(\tilde{\mu}))_{\mu \in \Lambda_T, \ell(\mu) \geq n}$  is contained in  $\mathcal{F}_{o,n}$  and contains  $(\mathcal{Z}_{T,\ell>0})^n$ .

**Proposition 6.4.** *When  $R$  is noetherian, the filtrations of  $\mathcal{H}$*

$$((\mathcal{Z}_{T,\ell>0})^n \mathcal{H})_{n \in \mathbb{N}}, \quad ((\mathcal{Z}_{\ell>0})^n \mathcal{H})_{n \in \mathbb{N}}, \quad (\mathcal{F}_{T,o,n})_{n \in \mathbb{N}} \mathcal{H}, \quad (\mathcal{F}_{o,n})_{n \in \mathbb{N}},$$

are equivalent.

We have  $(\mathcal{Z}_{T,\ell>0})^n \mathcal{H} \subset (\mathcal{Z}_{\ell>0})^n \mathcal{H} \subset \mathcal{F}_{o,n}$ . The last inclusion uses the product formula, the equality  $\tilde{E}(C) = \tilde{E}_o(C)$ , and that  $(E_o(w))_{w \in W(1)}$  is a basis of  $\mathcal{H}$ . The noetherianity of  $R$  is used only for the proof (Lemma 6.7) of the property (which implies the proposition):

$$\text{for } n \in \mathbb{N} \text{ there exists } n' \in \mathbb{N} \text{ such that } \mathcal{F}_{o,n'} \subset (\mathcal{Z}_{T,\ell>0})^n \mathcal{H}.$$

This property follows from the next three lemmas.

**Lemma 6.5.**  *$E(\tilde{C}(\mu))^n E_o(\tilde{\mu}) = E_o(\tilde{\mu}^{n+1})$  for  $\mu \in \Lambda_T$  and  $n > 0$ .*

**Proof.** By the product formula,  $E(C(\tilde{\mu}))E_o(\tilde{\mu}) = E_o(\tilde{\mu}^2)$ , because  $\tilde{\mu}$  is the only element of  $\tilde{C}(\mu)$  sent by  $\nu$  in the same closed Weyl chamber as  $\nu(\mu)$ . By induction on  $n$ ,

$$\begin{aligned} E(\tilde{C}(\mu))^{n+1} E_o(\tilde{\mu}) &= E(\tilde{C}(\mu))E(\tilde{C}(\mu))^n E_o(\tilde{\mu}) = E(\tilde{C}(\mu))E_o(\tilde{\mu}^{n+1}) \\ &= E(\tilde{C}(\mu))E_o(\tilde{\mu})E_o(\tilde{\mu}^n) = E_o(\tilde{\mu}^2)E_o(\tilde{\mu}^n) = E_o(\tilde{\mu}^{n+2}). \end{aligned} \quad \square$$

**Lemma 6.6.** *There exists a positive integer  $a$  such that, for any positive integer  $n$ ,*

$$E_o(\mu) \in \mathcal{Z}_{T, \ell > 0}^n \mathcal{A}_o$$

if  $\mu \in \Lambda_T$  satisfies  $\ell(\mu) \geq na$ .

**Proof.** Let  $\overline{\mathcal{D}}$  be a closed Weyl chamber. We choose  $\mu_1, \dots, \mu_r$  in  $\Lambda_T - \Lambda_T^{W_0}$  such that  $\nu(\mu_1), \dots, \nu(\mu_r)$  generate the monoid  $\nu(\Lambda_T) \cap \overline{\mathcal{D}}$ . We show that

$$E_o(\mu) \in \mathcal{Z}_{T, \ell > 0}^n \mathcal{A}_o,$$

if  $\mu \in \Lambda_T, \nu(\mu) \in \overline{\mathcal{D}}$  and  $\ell(\mu) > n(\ell(\mu_1) + \dots + \ell(\mu_r))$ . Clearly, this implies the lemma.

Let  $\mu = \mu_1^{n_1} \dots \mu_r^{n_r} u$  with  $u \in (\Lambda_T)^{W_0}, n_1, \dots, n_r$  in  $\mathbb{N}$ . We have  $\ell(\mu_i) \neq 0$  for  $1 \leq i \leq r$  and  $\ell(\mu) = n_1 \ell(\mu_1) + \dots + n_r \ell(\mu_r)$ . Changing the numerotation, we suppose that  $n_1 > n$ , and obtain

$$E_o(\mu) = E_o(\mu_1)^{n_1} h, \quad h = E_o(\mu_2)^{n_2} \dots E_o(\mu_r)^{n_r} T_u \in \mathcal{A}_o.$$

By Lemma 6.5,  $E_o(\mu_1)^{n_1} = E(\tilde{C}(\mu_1))^{n_1-1} E_o(\mu_1)$ . Hence  $E_o(\mu) \in E(\tilde{C}(\mu_1))^n \mathcal{A}_o \subset \mathcal{Z}_{T, \ell > 0}^n \mathcal{A}_o$ .  $\square$

**Lemma 6.7.** *When  $R$  is noetherian, for every positive integer  $n > 0$  there exists a positive integer  $n' > 0$  such that  $\mathcal{F}_{o, n'} \subset (\mathcal{Z}_{T, \ell > 0})^n \mathcal{H}$ .*

**Proof.** By Lemma 5.3, we can choose a finite subset  $X \subset \Lambda$  such that, for  $(\lambda, w_0) \in \Lambda \times W_0$ , we have  $\ell(\lambda w_0) = \ell(\lambda x^{-1}) + \ell(x w_0)$  for some  $x \in X$  with  $\mu = \lambda x^{-1} \in \Lambda_T$ . By the product formula,  $E_o(\lambda w_0) = E_o(\mu)E_o(x w_0)$ . If

$$\ell(\lambda w_0) \geq n' = na + \max\{\ell(xw) \mid (x, w) \in X \times W_0\},$$

we have  $\ell(\mu) \geq na$ . Taking  $a$  as in Lemma 6.6,  $E_o(\mu) \in (\mathcal{Z}_{T, \ell > 0})^n \mathcal{A}_o$ ; hence  $E_o(\lambda w_0) \in (\mathcal{Z}_{T, \ell > 0})^n \mathcal{H}$ . As  $(\lambda, w_0)$  was arbitrary, we get the lemma.  $\square$

We define  $\mathcal{F}_{o, n}^{\text{aff}}$  as  $\mathcal{F}_{o, n}$ , with  $W(1)$  replaced by  $W^{\text{aff}}(1)$ . The isomorphism (3) restricts to an isomorphism

$$\mathcal{F}_{o, n}^{\text{aff}} \otimes_{R[Z_k]} R[\Omega(1)] \simeq \mathcal{F}_{o, n}. \tag{44}$$

The based root system  $(\Phi, \Delta)$  is the finite disjoint union of irreducible based root systems  $(\Phi_i, \Delta_i)$  for  $1 \leq i \leq r$ , the Coxeter affine Weyl group  $(W^{\text{aff}}, S^{\text{aff}})$  is the product of the irreducible Coxeter affine Weyl groups  $(W_i^{\text{aff}}, S_i^{\text{aff}})$ , and  $W^{\text{aff}}(1)$  is an extension

$$1 \rightarrow Z_k \rightarrow W^{\text{aff}}(1) \rightarrow \prod_i W_i^{\text{aff}} \rightarrow 1.$$

The algebras  $\mathcal{H}_i^{\text{aff}}$  defined by  $(\Phi_i, \Delta_i)$  identify with the subalgebras of basis  $(T_w)_{w \in W_i^{\text{aff}}(1)}$  of  $\mathcal{H}^{\text{aff}}$ , called the irreducible components of  $\mathcal{H}^{\text{aff}}$ .

**Lemma 6.8.** *The filtrations of  $\mathcal{H}^{\text{aff}}$*

$$(\mathcal{F}_{o,n}^{\text{aff}})_{n \in \mathbb{N}}, \quad \left( \sum_i \mathcal{F}_{i,o,n}^{\text{aff}} \mathcal{H}^{\text{aff}} \right)_{n \in \mathbb{N}}$$

are equivalent.

**Proof.** The length of  $w_i \in W_i^{\text{aff}}$  seen as an element of  $(W_i^{\text{aff}}, S_i^{\text{aff}})$  or of  $(W^{\text{aff}}, S^{\text{aff}})$  is the same; hence

$$\mathcal{F}_{i,o,n}^{\text{aff}} \subset \mathcal{F}_{o,n}^{\text{aff}}.$$

For  $w \in W^{\text{aff}}$  of components  $w_i \in W_i^{\text{aff}}$ , we have  $\ell(w) = \sum_i \ell(w_i)$  and  $E_o(w) = \prod_i E_o(w_i)$  by the product formula, and the factors  $E_o(w_i)$  commute. If  $\ell(w) \geq nr$ , at least one component  $w_i$  satisfies  $\ell(w_i) \geq n$ ; hence

$$\mathcal{F}_{o,n}^{\text{aff}} \subset \sum_i \mathcal{F}_{i,o,n}^{\text{aff}} \mathcal{H}^{\text{aff}}. \quad \square$$

**Proposition 6.9.** *Let  $M$  be a right  $\mathcal{H}$ -module, and let  $o$  be an orientation. The following properties are equivalent.*

- (1) *There exists a positive integer  $n$  such that  $M\mathcal{F}_{o,n} = 0$ .*
- (2) *There exists a positive integer  $n$  such that  $M(\mathcal{Z}_{\ell>0})^n = 0$ .*
- (3) *There exists a positive integer  $n$  such that  $M(\mathcal{Z}_{T,\ell>0})^n = 0$ .*
- (4) *There exists a positive integer  $n$  such that  $M\mathcal{F}_{T,o,n} = 0$ .*
- (5) *There exists a positive integer  $n$  such that  $M\mathcal{F}_{o,n}^{\text{aff}} = 0$ .*
- (6) *There exists a positive integer  $n$  such that  $M\mathcal{F}_{i,o,n}^{\text{aff}} = 0$  for  $1 \leq i \leq r$ .*

**Proof.** The isomorphism (44) shows that  $M\mathcal{F}_{o,n} = 0 \Leftrightarrow M\mathcal{F}_{o,n}^{\text{aff}} = 0$ , because the action of  $\Omega(1)$  is invertible. Applying Proposition 6.4 and Lemma 6.8, the properties are equivalent. □

**Definition 6.10.** A right  $\mathcal{H}$ -module  $M$  is called supersingular if it is not 0 and satisfies the properties of Proposition 6.9.

For future reference, we present the properties of the supersingular right  $\mathcal{H}$ -modules  $M$  deduced easily from Proposition 6.9 and Lemma 6.3, as a proposition. For a right  $\mathcal{H}$ -module  $M$ , we have the right  $\mathcal{H}$ -module  $\iota(M)$ , equal to  $M$  with  $h \in \mathcal{H}$  acting by  $\iota(h)$ .

**Proposition 6.11.** (1) *The category of supersingular right  $\mathcal{H}$ -modules is stable by subquotients, by extensions, and by finite sums.*

- (2) *A right  $\mathcal{H}$ -module is supersingular if and only if it is supersingular as a right  $\mathcal{H}^{\text{aff}}$ -module.*
- (3) *A right  $\mathcal{H}$ -module generated by a supersingular right  $\mathcal{H}^{\text{aff}}$ -submodule is supersingular.*

- (4) A right  $\mathcal{H}^{\text{aff}}$ -module is supersingular if and only if it is supersingular as a right  $\mathcal{H}_i^{\text{aff}}$ -module for all the irreducible components  $\mathcal{H}_i^{\text{aff}}$  of  $\mathcal{H}^{\text{aff}}$ .
- (5) A right  $\mathcal{H}$ -module  $M$  is supersingular if and only if  $\iota(M)$  is supersingular.
- (6) A simple right  $\mathcal{H}$ -module  $M$  is supersingular if and only if  $M\mathcal{Z}_{\ell>0} = 0 \Leftrightarrow M\mathcal{Z}_{T,\ell>0} = 0$ .

The properties in (vi) are also equivalent to  $M\mathcal{F}_{T,o,1} = 0$ . See Remark 6.16.

The classification of the supersingular simple  $\mathcal{H}$ -modules reduces to the classification of the supersingular characters of  $\mathcal{H}^{\text{aff}}$ . For the algebra  $\mathcal{H}(G, I(1))$ , this was a conjecture for  $G = GL(n, F)$  [13] proved in [11, Proposition 5.10] for  $G$  split.

**Proposition 6.12.** *A supersingular right  $\mathcal{H}$ -module  $M$  contains a character of  $\mathcal{H}^{\text{aff}}$ .*

**Proof.** A non-zero element of  $M$  generates a right  $\mathfrak{h}$ -module containing a character of  $\mathfrak{h}$  (Proposition 2.1). We choose a  $\mathfrak{h}$ -eigenvector  $v \in M$  of eigenvalue  $\eta$ . Let  $(\chi, S_\eta)$  be the parameters of  $\eta$  (Proposition 2.2). As  $M$  is supersingular, there exists a positive integer  $n$  such that  $M\mathcal{F}_{o,n} = 0$ . We choose  $d \in \mathcal{D}$  of maximal length satisfying  $vE_o(\tilde{d}) \neq 0$  (Proposition 3.3). We show that  $vE_o(\tilde{d})$  is a  $\mathcal{H}^{\text{aff}}$ -eigenvector. Let  $(t, s) \in Z_k \times S^{\text{aff}}$ .

We have  $vE_o(\tilde{d})T_t = vT_{dt}d^{-1}E_o(\tilde{d}) = \chi(dt d^{-1})vE_o(\tilde{d}) = \chi^d(t)vE_o(\tilde{d})$ .

For the computation of  $vE_o(\tilde{d})T_{\tilde{s}}$ , we distinguish three cases.

(1)  $\ell(ds) = \ell(d) - 1$ . Then  $E_o(\tilde{d}) = T_t E_o(\tilde{d}\tilde{s})E_o(\tilde{s})$ , where  $t \in Z_k, t\tilde{d}\tilde{s}^2 = \tilde{d}$ .

If  $E_o(\tilde{s}) = T_{\tilde{s}} - c_{\tilde{s}}$ , we have  $E_o(\tilde{s})T_{\tilde{s}} = (T_{\tilde{s}} - c_{\tilde{s}})T_{\tilde{s}} = 0$ .

If  $E_o(\tilde{s}) = T_{\tilde{s}}$ , we have  $E_o(\tilde{s})T_{\tilde{s}} = T_{\tilde{s}}^2 = c_{\tilde{s}}T_{\tilde{s}} = c_{\tilde{s}}E_o(\tilde{s})$ ; as  $E_o(\tilde{d}\tilde{s})c_{\tilde{s}} = (ds \bullet c_{\tilde{s}})E_o(\tilde{d}\tilde{s}) = d \bullet c_{\tilde{s}}E_o(\tilde{d}\tilde{s})$ , we deduce that  $vE_o(\tilde{d})T_{\tilde{s}} = 0$  or  $\chi(d \bullet c_{\tilde{s}})vE_o(\tilde{d}) = \chi^d(c_{\tilde{s}})vE_o(\tilde{d})$ .

(2)  $\ell(ds) = \ell(d) + 1$  and  $ds \in Z_k\mathcal{D}$ . Either  $E_o(\tilde{d})T_{\tilde{s}} = E_o(\tilde{d})E_o(\tilde{s}) = E_o(\tilde{d}\tilde{s})$  or  $E_o(\tilde{d})T_{\tilde{s}} = E_o(\tilde{d})(E_o(\tilde{s}) + c_{\tilde{s}}) = E_o(\tilde{d}\tilde{s}) + (d \bullet c_{\tilde{s}})E_o(\tilde{d})$ . By the maximality of  $\ell(d)$ ,  $vE_o(\tilde{d}\tilde{s}) = 0$  and  $vE_o(\tilde{d})T_{\tilde{s}} = 0$  or  $\chi(d \bullet c_{\tilde{s}})vE_o(\tilde{d}) = \chi^d(c_{\tilde{s}})vE_o(\tilde{d})$ .

(3)  $\ell(ds) = \ell(d) + 1$  and  $ds \notin Z_k\mathcal{D}$ . Let  $s_d \in S$  such that  $\tilde{d}\tilde{s} = \tilde{s}_d\tilde{d}$  (Lemma 3.8). Either  $E_o(\tilde{d})T_{\tilde{s}} = E_o(\tilde{d})E_o(\tilde{s}) = E_o(\tilde{d}\tilde{s}) = E_o(\tilde{s}_d\tilde{d}) = E_o(\tilde{s}_d)E_o(\tilde{d})$  or  $E_o(\tilde{d})T_{\tilde{s}} = E_o(\tilde{d})(E_o(\tilde{s}) + c_{\tilde{s}}) = E_o(\tilde{d}\tilde{s}) + E_o(\tilde{d})c_{\tilde{s}} = (E_o(\tilde{s}_d) + d \bullet c_{\tilde{s}})E_o(\tilde{d})$ . Hence  $vE_o(\tilde{d})T_{\tilde{s}} = \eta(E_o(\tilde{s}_d))vE_o(\tilde{d})$  or  $\eta(E_o(\tilde{s}_d) + d \bullet c_{\tilde{s}})vE_o(\tilde{d}) = (\eta(E_o(\tilde{s}_d)) + \chi(d \bullet c_{\tilde{s}}))vE_o(\tilde{d}) = (\eta(E_o(\tilde{s}_d)) + \chi^d(c_{\tilde{s}}))vE_o(\tilde{d})$ . □

The compatibility of supersingularity for  $\mathcal{H}$  and  $\mathcal{H}^{\text{aff}}$  (Proposition 6.9) and Proposition 6.12 imply the following.

**Corollary 6.13.** (1) *A simple supersingular right  $\mathcal{H}^{\text{aff}}$ -module has dimension 1.*

(2) *A simple right  $\mathcal{H}$ -module is supersingular*

*if and only if it contains a supersingular character of  $\mathcal{H}^{\text{aff}}$ ;*

*if and only if any simple right  $\mathcal{H}^{\text{aff}}$ -submodule is a supersingular character of  $\mathcal{H}^{\text{aff}}$ .*

The classification of the supersingular characters of  $\mathcal{H}^{\text{aff}}$ , given in Theorem 6.15 after technical Lemma 6.14, follows from the classification of the characters of  $\mathcal{H}^{\text{aff}}$  (Proposition 2.2). The classification was done for  $\mathcal{H}(G, I(1))$  in [13] for  $G = GL(n, F)$  and in [11, Lemma 5.11 and Theorem 5.13] for  $G$  split.

Let  $\Xi$  be a character of  $\mathcal{H}^{\text{aff}}$ ,  $\chi$  a character of  $Z_k$ , and  $o$  an orientation such that  $\Xi|_{\mathfrak{h}} = \chi_o$  (Lemma 2.4). Let  $w_o \in W_0$  such that the Weyl chamber of  $o$  is  $w_o^{-1}(\mathfrak{D}^+)$ . For a subset  $J$  of  $S$ , let  $w_J$  be the longest element of the subgroup of  $W_0$  generated by  $J$ .

**Lemma 6.14.** (1)  $\Xi(E(\tilde{C}(\mu))) = \Xi(E_o(\tilde{\mu}))$  for  $\mu \in \Lambda_T^+$ .

(2) If  $S^{\text{aff}} - S = \{s_0\}$  and  $\lambda \in \Lambda^+$  has positive length, we have

- (i)  $\ell(s_0\lambda) = -1 + \ell(\lambda)$ ;
- (ii)  $E_o(\tilde{s}_0) = T_{\tilde{s}_0} \Leftrightarrow w_o(\alpha_0) \in \Sigma^+$ , where  $\alpha_0$  is the highest root of  $\Sigma^+$ ;
- (iii)  $E_o(\tilde{\lambda}) = T_{\tilde{s}_0} E_{o \bullet s_0}(\tilde{s}_0^{-1}\tilde{\lambda})$  if  $w_o(\alpha_0) \in \Sigma^+$ ;
- (iv)  $w_J(\alpha_0) \in \Sigma^+ \Leftrightarrow J \neq S$ .

**Proof.** (1) The character  $\xi$  factorizes through the canonical homomorphism

$$h \mapsto 1 \otimes h : \mathcal{H}^{\text{aff}} \rightarrow \xi|_{\mathfrak{h}} \otimes_{\mathfrak{h}} \mathcal{H}^{\text{aff}},$$

and  $1 \otimes E(\tilde{C}(\mu)) = 1 \otimes E_o(\tilde{\mu})$  in  $\chi_o \otimes_{\mathfrak{h}} \mathcal{H}^{\text{aff}}$  by Proposition 5.4.

(2) The hypothesis means that the root system  $\Sigma$  is irreducible. The highest positive root  $\alpha_0 \in \Sigma^+$  has the following well-known properties:  $-\alpha_0 + 1$  is a simple affine root and  $s_0 = s_{-\alpha_0+1}$ ,  $0 < -\alpha_0(x) + 1 < 1$  for  $x \in \mathfrak{C}^+$ .

(i)  $\ell(s_0\lambda) = -1 + \ell(\lambda) \Leftrightarrow \mathfrak{C}^+$  and  $\mathfrak{C}^+ + \nu(\lambda)$  are not on the same side of  $\text{Ker}(-\alpha_0 + 1)$  [15, Example 5.4]. This is equivalent to  $-\alpha_0(x + \nu(\lambda)) + 1 = -\alpha_0(x) + 1 - \alpha_0 \circ \nu(\lambda)$  is negative for  $x \in \mathfrak{C}^+ \Leftrightarrow \alpha_0 \circ \nu(\lambda) \geq 1$ , which is true, because  $\alpha_0 \circ \nu(\lambda) \in \mathbb{N}_{>0}$  as  $\lambda \in \Lambda^+$  has positive length [15, Corollary 5.11].

(ii) By (6),  $E_o(\tilde{s}_0) = T_{\tilde{s}_0} \Leftrightarrow \mathfrak{C}^+$  is on the  $o$ -negative side of  $\text{Ker}(-\alpha_0 + 1)$ . By [15, Definition 5.16], this means that  $-\alpha_0$  is  $o$ -negative, because  $-\alpha_0 + 1$  is positive on  $\mathfrak{C}^+$ . The root  $-\alpha_0$  is  $o$ -negative if and only if  $\alpha_0$  is positive on the Weyl chamber  $w_o^{-1}(\mathfrak{D}^+)$  of  $o$ . This is true if and only if  $w_o(\alpha_0) \in \Sigma^+$ .

(iii) For any orientation  $o$ ,  $E_o(\tilde{\lambda}) = E_o(\tilde{s}_0)E_{o \bullet s_0}(\tilde{s}_0^{-1}\tilde{\lambda})$  by the product formula and  $\ell(\lambda) = 1 + \ell(s_0\lambda)$  (i). Apply (ii).

(iv) Let  $S = J \cup J'$ . We have  $\alpha_0 = (\sum_{s \in J} n_s \alpha_s) + (\sum_{s \in J'} n_s \alpha_s)$  with  $n_s \in \mathbb{N}_{>0}$ , and  $w_J(\alpha_0) = -(\sum_{s \in J} n_s \alpha_s) + (\sum_{s \in J'} n_s w_J(\alpha_s))$ . If  $J' = \emptyset$ , then  $w_J(\alpha_0) \notin \Sigma^+$ . If  $J' \neq \emptyset$ , for any  $s \in J'$ , the root  $w_J(\alpha_s)$  is positive and does not belong to the group generated by  $J$ . The decomposition of  $w_J(\alpha_0)$  on the basis  $(\alpha_s)_{s \in S}$  has a positive coefficient; i.e.,  $w_J(\alpha_0) \in \Sigma^+$ . □

**Theorem 6.15.** A character of  $\mathcal{H}^{\text{aff}}$  is supersingular if and only if its restriction to each irreducible component of  $\mathcal{H}^{\text{aff}}$  is not a twisted sign or trivial character.

**Proof.** The involutive automorphism  $\iota$  of  $\mathcal{H}^{\text{aff}}$  respects supersingularity and exchanges a twisted sign character and a twisted trivial character (Definition 2.7). For  $s \in S^{\text{aff}}$  and a character  $\xi$  of  $\mathcal{H}^{\text{aff}}$ ,  $\xi$  vanishes on  $T_s$  or  $\iota(T_s)$  (Proposition 2.2). Let  $\mu \in \Lambda_T^+$  of positive length. We have  $\xi(E(\tilde{C}(\mu))) = \xi(E_o(\tilde{\mu}))$  for any orientation  $o$  (Lemma 6.14) and  $E_o(\tilde{\mu}) = T_\mu$  when  $o$  is dominant [15, Example 5.30].

(i) A twisted sign character is not 0 on  $T_w$  for all  $w \in W(1)$  of positive length; hence it is not 0 on  $E(\tilde{C}(\mu))$ , and it is not supersingular. Applying  $\iota$ , a twisted trivial character is not supersingular.

(ii) It remains to prove that, when  $\mathcal{H}^{\text{aff}}$  is irreducible, i.e.,  $S^{\text{aff}} - S = \{s_0\}$ , a character  $\xi$  of  $\mathcal{H}^{\text{aff}}$  different from a twisted sign or trivial character is supersingular.

Applying  $\iota$ , it suffices to prove it when  $\xi(T_{s_0}^\vee) = 0$ . The set  $J = S - \{s \in S \mid \xi(T_s^\vee) \neq 0\}$  is different from  $S$ , because  $\xi$  is not a twisted sign character. Let  $o$  be the orientation of Weyl chamber  $w_J^{-1}(\mathfrak{D}^+)$ . By Lemma 2.6, the restriction of  $\xi$  to  $\mathfrak{h}$  is of the form  $\chi_o$ , because  $S_o = \{s \in S \mid \xi(T_s^\vee) \neq 0\}$  (5). Applying Lemma 6.14, we obtain, for any  $\mu \in \Lambda_T^+$  of positive length,

$$E_o(s_0) = T_{\tilde{s}_0}, \quad E_o(\tilde{\mu}) = T_{\tilde{s}_0} E_{o \bullet s_0}((\tilde{s}_0)^{-1} \tilde{\mu}), \quad \xi(E(C(\tilde{\mu}))) = \xi(E_o(\tilde{\mu})) = 0.$$

Hence  $\xi$  is supersingular. □

**Remark 6.16.** We can complete Proposition 6.11(6): a simple  $\mathcal{H}$ -module  $M$  is supersingular if and only if  $M\mathcal{F}_{T,o,1} = 0$ . This follows from Corollary 6.13 and part (ii) in the proof of Theorem 6.15.

Clifford’s theory studies classically the induction of representations from normal subgroups. We give a “Clifford’s theory style” proposition to describe the simple finite-dimensional  $\mathcal{H}$ -modules containing a character of  $\mathcal{H}^{\text{aff}}$ , as in [13, Proposition 3], [11, Lemma 5.12] for the algebra  $\mathcal{H}(G, I(1))$  when  $G$  is split.

Let  $R$  be a field, and let  $A$  be an  $R$ -algebra of the form  $A = JB$ , where  $J$  is an ideal of  $A$  and  $B$  a subalgebra of  $A$  equal to the  $R$ -algebra  $R[G]$  of a group  $G$ .

Let  $\Xi : J \rightarrow R$  be a character of  $J$  with a  $G$ -fixator  $G_\Xi = \{g \in G \mid \Xi^g = \Xi\}$  of  $\Xi$  of finite index in  $G$ , where  $\Xi^g$  is the character  $j \mapsto \Xi^g(j) = \Xi(gjg^{-1})$  of  $J$ .

Let  $V$  be a finite-dimensional right  $R[G_\Xi]$ -module, where the group  $J \cap G$  acts by  $\Xi|_{J \cap G}$ . For  $g \in G$ , we denote by  $V^g$  the right  $R[g^{-1}G_\Xi g]$ -module  $V$ , where  $g^{-1}hg$  acts by  $h$  for  $h \in G_\Xi$ .

We extend  $V$  to a right  $A_\Xi = JR[G_\Xi]$ -module, where  $J$  acts by  $\Xi$ , denoted by  $\Xi \otimes V$ . We induce  $\Xi \otimes V$  to a right  $A$ -module

$$I(\Xi, V) = (\Xi \otimes V) \otimes_{A_\Xi} A.$$

**Proposition 6.17.** *Let  $R, A, J, G, \Xi, V$  be as above. We suppose  $V$  to be simple. We have the following.*

- (i)  $I(\Xi, V)$  is finite dimensional and is a simple right  $A$ -module.
- (ii) A finite-dimensional simple right  $A$ -module containing  $\Xi$  as a  $J$ -module is isomorphic to  $I(\Xi, V)$  for some  $V$ .
- (iii)  $I(\Xi_1, V_1) \simeq I(\Xi_2, V_2)$  if and only if  $(\Xi_2, V_2) = (\Xi_1^g, V_1^g)$ , for some element  $g \in G$ .

**Proof [11, Lemma 5.12].**  $\Xi \otimes V$  is finite dimensional and is a simple  $A_\Xi$ -module, because its restriction to the subalgebra  $R[G_\Xi]$  satisfies these properties. The left  $A_\Xi$ -module  $A = \bigoplus_{g \in G_\Xi \backslash G} A_\Xi g$  is free of finite rank. The restriction of  $I(\Xi \otimes V)$  to  $J$  is isomorphic

to a direct sum  $\bigoplus^{\dim_R V} \bigoplus_{g \in G_{\Xi} \backslash G} \Xi^g$ , and  $I(\Xi, V) = \bigoplus_{g \in G_{\Xi} \backslash G} (\Xi^g \otimes V^g)$  is equal to the direct sum of all the conjugates of  $\Xi \otimes V$  by  $G$ . The dimension of  $I(\Xi \otimes V)$  is finite, equal to  $[G : G_{\Xi}] \dim_R V$ . The restriction to  $J$  of a non-zero  $A$ -submodule of  $I(\Xi \otimes V)$  contains a submodule isomorphic to  $\bigoplus_{g \in G_{\Xi} \backslash G} \Xi^g$ ; hence its  $\Xi$ -isotypic component is not 0. The  $\Xi$ -isotypic component of  $I(\Xi \otimes V)$  is the simple  $A_{\Xi}$ -module  $\Xi \otimes V$ . Therefore  $I(\Xi \otimes V)$  is a simple  $A$ -module.

Let  $M$  be a finite-dimensional simple right  $A$ -module with a non-zero  $\Xi$ -isotypic component as a  $J$ -module. The  $\Xi$ -isotypic component is an  $A_{\Xi}$ -module of the form  $\Xi \otimes V'$  for some finite-dimensional right  $R[G_{\Xi}]$ -module  $V'$ . The non-zero  $R[G_{\Xi}]$ -module  $V'$  contains a simple submodule  $V$ . The module  $\Xi \otimes V$  is isomorphic to an  $A_{\Xi}$ -submodule of  $M$ , and  $I(\Xi \otimes V)$  is isomorphic to an  $A$ -submodule of  $M$ . As  $M$  is simple,  $M = I(\Xi, V)$ .

The restriction of  $I(\Xi \otimes V)$  to  $J$  shows that  $I(\Xi \otimes V)$  determines the  $G$ -orbit of  $\Xi$ , the  $\Xi$ -isotypic part of  $I(\Xi \otimes V)$  determines  $V$ , and the  $\Xi^g$ -isotypic part of  $I(\Xi \otimes V)$  is  $\Xi^g \otimes V^g$  for  $g \in G$ . This implies that  $I(\Xi_1, V_1) \simeq I(\Xi_2, V_2)$  if and only if  $(\Xi_2, V_2) = (\Xi_1^g, V_1^g)$ , for some  $g \in G$ . □

We can apply Proposition 6.17 to the  $R$ -algebra  $A = \mathcal{H}$ , its ideal  $J = \mathcal{H}^{\text{aff}}$ , the group  $G = \Omega(1)$ , an arbitrary character  $\Xi$  of  $\mathcal{H}^{\text{aff}}$ , and a finite-dimensional simple right  $R[\Omega(1)]$ -module  $V$  such that  $Z_k$  acts on  $V$  by the character  $\Xi|_{Z_k}$ . As a subgroup of  $\Omega$  of finite index acts trivially on  $V$ , the fixator  $\Omega(1)_{\Xi}$  of  $\Xi$  has a finite index in  $\Omega(1)$ .

Corollary 6.13, Theorem 6.15, and Proposition 6.17 imply the following.

**Theorem 6.18.** *The supersingular simple finite-dimensional right  $\mathcal{H}$ -modules are isomorphic to the  $\mathcal{H}$ -modules  $I(\Xi, V)$ , where*

- (i)  $\Xi$  is a character of  $\mathcal{H}^{\text{aff}}$  different from a twisted sign or trivial character on each irreducible component of  $\mathcal{H}^{\text{aff}}$ ,
- (ii)  $V$  is a simple finite-dimensional right  $R[\Omega(1)_{\Xi}]$ -module, where  $Z_k$  acts by  $\Xi|_{Z_k}$ .

Two  $\mathcal{H}$ -modules  $I(\Xi_1, V_1), I(\Xi_2, V_2)$  are isomorphic if and only if the pairs  $(\Xi_1, V_1), (\Xi_2, V_2)$  are  $\Omega(1)$ -conjugate.

### 7. Pro- $p$ -Iwahori invariants and compact induction

We use the notation of 1.3, and  $R$  is as in 1.4. The algebras  $\mathcal{H}$  and  $\mathfrak{h}$  denote the pro- $p$ -Iwahori Hecke algebra  $\mathcal{H}_R(G, I(1))$  and  $\mathcal{H}_R(K, I(1))$ .

Let  $\rho$  be an irreducible smooth  $R$ -representation of  $K$ , let  $v \in \rho^{I(1)}$  not 0, let  $\eta$  be the character of  $\mathfrak{h}$  on  $\rho^{I(1)}$ , let  $\chi$  be the restriction of  $\eta$  to  $Z_k$ , and let  $o$  be an orientation such that  $\eta = \chi_o$  (Lemma 2.4).

We show that the pro- $p$ -Iwahori invariant functor behaves well on compact induced representations of  $G$ , generalizing the results of Ollivier [10, Corollary 3.14] proved when  $G$  is split.

By Cabanes [3, Theorem 2], the  $I(1)$ -invariant functor  $\rho \mapsto \rho^{I(1)}$  gives an equivalence

- from the category of finite-dimensional  $R$ -representations  $\rho$  of  $K$  trivial on  $K(1)$ , such that  $\rho$  and its dual  $\rho^*$  are generated by  $I(1)$ ;
- to the category of finite-dimensional right  $\mathfrak{h}$ -modules.

**Remark 7.1.** For  $n \in N \cap K$  of image  $w \in W_o(1)$ , the action on  $\rho^{I(1)}$  of the basis element  $T_w \in \mathfrak{h}$  is

$$vT_w = \sum_{\gamma \in I(1) \backslash I(1)nI(1)} \gamma^{-1}v = \eta(T_w)v.$$

The action of  $Z_k$  on  $\rho^{I(1)}$  arising from the action of  $Z_0 \subset I$  normalizing  $I(1)$  and the action of  $Z_k$  embedded in the Hecke algebra  $\mathfrak{h}$  on  $\rho^{I(1)}$  are inverse from each other.

Let

$$\mathbf{c}\text{-Ind}_K^G \rho$$

be the compactly induced representation of  $G$  by right translations on the space of compactly supported functions  $f : G \rightarrow V(\rho)$  satisfying  $f(k_1g) = \rho(k_1)f(g)$  for  $k_1 \in K$  and  $g \in G$ . Let

$$[1, v]_K \in (\mathbf{c}\text{-Ind}_K^G \rho)^{I(1)}$$

be the function of support  $K$  and value  $v$  at 1. The representation  $\mathbf{c}\text{-Ind}_K^G \rho$  is generated by  $[1, v]_K$ , and  $\dim_R \rho^{I(1)} = 1$ .

**Proposition 7.2.** *The  $\mathcal{H}$ -equivariant linear map*

$$\rho^{I(1)} \otimes_{\mathfrak{h}} \mathcal{H} \rightarrow (\mathbf{c}\text{-Ind}_K^G \rho)^{I(1)} \quad 1 \otimes 1 \mapsto [1, v]_K$$

*is an isomorphism.*

We explain the strategy of the proof, which reduces the proposition to the next lemma.

The disjoint union of  $W$  into  $W_0$ -cosets corresponds to a disjoint union of  $G$  into  $(K, I)$ -cosets. A  $(K, I)$ -coset is equal to a  $(K, I(1))$ -coset. We have

$$G = \bigcup_{d \in \mathcal{D}} KdI = \bigcup_{d \in \mathcal{D}} K\tilde{d}I(1), \tag{45}$$

where, for  $d$  in the distinguished set  $\mathcal{D}$  of representatives of  $W_0 \backslash W$  (Proposition 3.3),  $KdI = K\tilde{d}I(1)$  denotes the double coset  $Kn_{\tilde{d}}I = Kn_{\tilde{d}}I(1)$ ,  $\tilde{d} \in \mathcal{D}(1)$  lifting  $d$ , and  $n_{\tilde{d}} \in N$  lifting  $\tilde{d}$ , with  $n_1 = 1$ . The space  $(\mathbf{c}\text{-Ind}_K^G \rho)^{I(1)}$  is the direct sum

$$(\mathbf{c}\text{-Ind}_K^G \rho)^{I(1)} = \bigoplus_{d \in \mathcal{D}} (\mathbf{c}\text{-Ind}_K^{KdI} \rho)^{I(1)} \tag{46}$$

of the subspaces of functions in  $(\mathbf{c}\text{-Ind}_K^G \rho)^{I(1)}$  with support contained in  $KdI$ , for  $d \in \mathcal{D}$ . The pro- $p$ -Iwahori Hecke algebra is the direct sum

$$\mathcal{H} = \bigoplus_{d \in \mathcal{D}} \mathfrak{h}T_{\tilde{d}} \tag{47}$$

of the left  $\mathfrak{h}$ -modules  $\mathfrak{h}T_{\tilde{d}}$  of functions in  $\mathcal{H}$  with support contained in  $KdI$ , for  $d \in \mathcal{D}$ . We denote by  $\eta$  the character of  $\mathfrak{h}$  on  $\rho^{I(1)}$ , and by  $f_{\tilde{d}}$  the function in  $(\mathbf{c}\text{-Ind}_K^G \rho)^{I(1)}$  of support  $KdI$  with  $f(n_{\tilde{d}}) = v$ , for  $d \in \mathcal{D}$ . We have  $f_1 = [1, v]_K$ . The proposition follows from the following lemma.



**Lemma 7.3.** (i) For  $d \in \mathcal{D}$ , we have  $K(1)(K \cap n_{\bar{d}}I(1)n_{\bar{d}}^{-1}) = I(1)$ .

(ii) A basis of  $(\mathfrak{c}\text{-Ind}_K^G \rho)^{I(1)}$  is  $(f_{\bar{d}})_{d \in \mathcal{D}}$ .

(iii)  $f_{\bar{d}} = f_1 T_{\bar{d}}$  for  $d \in \mathcal{D}$ .

(iv)  $f_1$  is a  $\mathfrak{h}$ -eigenvector in  $(\mathfrak{c}\text{-Ind}_K^G \rho)^{I(1)}$  of eigenvalue  $\eta$ .

**Proof.** (1) We denote by  $I'$  the subgroup of  $I(1)$  generated by  $U \cap I = U \cap K$  and  $U^- \cap I$ . We have  $I(1) = Z_0(1)I'$  and  $Z_0(1) = K(1) \cap Z_0$ . The lemma follows from the inclusion

$$U \cap I \subset n_{\bar{d}}I'n_{\bar{d}}^{-1},$$

because  $K(1)(K \cap n_{\bar{d}}I(1)n_{\bar{d}}^{-1}) = K(1)(K \cap n_{\bar{d}}I'n_{\bar{d}}^{-1})$  is a pro- $p$ -subgroup of  $K$  and  $I(1) = K(1)(U \cap I)$  is a pro- $p$ -Sylow subgroup of  $K$ . The group  $U \cap I$  is the product of the groups  $U_{\alpha,0} = U_{\alpha} \cap K$  for all  $\alpha$  in the set  $\Phi_{red}^+$  of positive reduced roots of associated root subgroup  $U_{\alpha}$ . By Proposition 3.3 and § 1.3,  $d^{-1}(e_{\alpha}\alpha)$  is positive on  $\mathfrak{C}^+$ . As  $e_{\alpha}$  is a positive integer,  $d^{-1}(\alpha)$  is positive on  $\mathfrak{C}^+$ . By [15, §§ 3.3 and 3.5],  $n_{\bar{d}}^{-1}U_{\alpha,0}n_{\bar{d}} = U_{d^{-1}(\alpha)}$ . As  $d^{-1}(\alpha)$  is positive on  $\mathfrak{C}^+$ ,  $U_{d^{-1}(\alpha)} \subset I'$ . Hence  $U_{\alpha,0} \subset n_{\bar{d}}I'n_{\bar{d}}^{-1}$ .

(2) By (46) and  $f_{n_{\bar{d}}} \in (\mathfrak{c}\text{-Ind}_K^{KdI(1)} \rho)^{I(1)}$ , it suffices to prove that the dimension of  $(\mathfrak{c}\text{-Ind}_K^{Kn_{\bar{d}}I(1)} \rho)^{I(1)}$  is 1. The value at  $n_{\bar{d}}$  gives a linear map

$$(\mathfrak{c}\text{-Ind}_K^{KdI(1)} \rho)^{I(1)} \rightarrow \rho^{K \cap n_{\bar{d}}I(1)n_{\bar{d}}^{-1}},$$

because  $kf(n_{\bar{d}}) = f(kn_{\bar{d}}) = f(n_{\bar{d}}n_{\bar{d}}^{-1}kn_{\bar{d}})$  for  $k \in K$ . The map is clearly injective, and  $\rho^{K \cap n_{\bar{d}}I(1)n_{\bar{d}}^{-1}} = \rho^{I(1)}$ , because  $\rho$  is trivial on  $K(1)$  and (1). As  $\dim_R \rho^{I(1)} = 1$ , we have  $\dim_R (\mathfrak{c}\text{-Ind}_K^{Kn_{\bar{d}}I(1)} \rho)^{I(1)} = 1$ .

(3) We show that the support of the function  $f_1 T_{n_{\bar{d}}}$  is contained in  $KdI(1)$  and that the value at  $n_{\bar{d}}$  of  $f_1 T_{n_{\bar{d}}}$  is  $v$ .

For  $g \in G$ , we have

$$f_1 T_g = \sum_{\gamma \in I(1) \backslash I(1)gI(1)} \gamma^{-1} f_1,$$

and  $\gamma^{-1} f_1(x) = f_1(x\gamma^{-1})$  for  $x \in G$ . The support of  $f_1$  is  $K$ , and the support of  $f_1 T_g$  is contained in  $KgI(1)$ .

In particular, the support of the function  $f_1 T_{n_{\bar{d}}}$  is contained in  $KdI(1)$ . We have

$$(f_1 T_{n_{\bar{d}}})(n_{\bar{d}}) = \sum_{\gamma \in I(1) \backslash I(1)n_{\bar{d}}I(1)} f_1(n_{\bar{d}}\gamma^{-1}) = \sum_{u \in (K \cap n_{\bar{d}}I(1)n_{\bar{d}}^{-1}) / (I(1) \cap n_{\bar{d}}I(1)n_{\bar{d}}^{-1})} f_1(u).$$

By (1), this is equal to  $f_1(1) = v$ .

(4) For  $k \in K$ , the support of the function  $f_1 T_k$  is contained in  $K$ , and

$$(f_1 T_k)(1) = \sum_{\gamma \in I(1) \backslash I(1)kI(1)} f_1(\gamma^{-1}) = \sum_{\gamma \in I(1) \backslash I(1)kI(1)} \gamma^{-1} f_1(1) = \eta(T_k)v.$$

Therefore  $f_1 T_k = \eta(T_k) f_1$  for  $k \in K$ . □

**Remark 7.4.** For  $\lambda \in \Lambda$ , the isomorphism of Proposition 7.2 restricts to a right  $\mathfrak{h}$ -module isomorphism

$$\rho^{I(1)} \otimes_{\mathfrak{h}} \mathfrak{h}(\lambda) \rightarrow (\text{c-Ind}_K^{K\lambda K} \rho)^{I(1)}.$$

**Proposition 7.5.** Let  $\rho_1, \rho$  be two irreducible smooth  $R$ -representations of  $K$ . The  $I(1)$ -invariant map

$$\text{Hom}_{RG}(\text{c-Ind}_K^G \rho_1, \text{c-Ind}_K^G \rho) \rightarrow \text{Hom}_{\mathcal{H}}((\text{c-Ind}_K^G \rho_1)^{I(1)}, (\text{c-Ind}_K^G \rho)^{I(1)})$$

is an isomorphism.

To explain the strategy of the proof, we recall the adjunction isomorphisms

$$\begin{aligned} \text{Hom}_{RG}(\text{c-Ind}_K^G \rho_1, \pi) &\simeq \text{Hom}_{RK}(\rho_1, \pi) = \text{Hom}_{RK}(\rho_1, \pi^{K(1)}), \\ \text{Hom}_{\mathcal{H}}(\rho_1^{I(1)} \otimes_{\mathfrak{h}} \mathcal{H}, \pi^{I(1)}) &\simeq \text{Hom}_{\mathfrak{h}}(\rho_1^{I(1)}, \pi^{I(1)}), \end{aligned}$$

for any smooth  $R$ -representation  $\pi$  of  $G$ . The  $I(1)$ -invariant map

$$\text{Hom}_{RG}(\text{c-Ind}_K^G \rho_1, \pi) \rightarrow \text{Hom}_{\mathcal{H}}((\text{c-Ind}_K^G \rho_1)^{I(1)}, \pi^{I(1)})$$

is an isomorphism if and only if the  $I(1)$ -invariant map

$$\text{Hom}_K(\rho_1, \pi^{K(1)}) \rightarrow \text{Hom}_{\mathfrak{h}}(\rho_1^{I(1)}, \pi^{I(1)}) \tag{48}$$

is an isomorphism, by Proposition 7.2. The map (48) is injective, because  $\rho^{I(1)}$  generates  $\rho$ , but it is not surjective in general. The proposition says that the map (48) is surjective if  $\pi = \text{c-Ind}_K^G \rho$ .

The dominant monoid  $\Lambda^+$  represents the cosets  $K \backslash G / K$  (see 1.3). The anti-dominant monoid  $\Lambda^-$  has the same property and is more convenient now. The representation of  $K$  on  $\text{c-Ind}_K^G \rho$  is a direct sum

$$\text{c-Ind}_K^G \rho = \bigoplus_{\lambda \in \Lambda^-} \text{c-Ind}_K^{K\lambda K} \rho,$$

where  $\text{c-Ind}_K^{K\lambda K} \rho$  is the space of functions in  $\text{c-Ind}_K^G \rho$  with support in  $K\lambda K$ . We will prove that, for all  $\lambda \in \Lambda^-$ , the  $I(1)$ -invariant map

$$\text{Hom}_K(\rho_1, (\text{c-Ind}_K^{K\lambda K} \rho)^{K(1)}) \rightarrow \text{Hom}_{\mathfrak{h}}(\rho_1^{I(1)}, (\text{c-Ind}_K^{K\lambda K} \rho)^{I(1)}) \tag{49}$$

is an isomorphism. A representation of  $K$  trivial on  $K(1)$  generated by its  $I(1)$ -invariant vectors identifies with a representation of the finite reductive group  $G_k$  generated by its  $U_k$ -invariant vectors (using the notation of § 1.3). We describe  $(\text{c-Ind}_K^{K\lambda K} \rho)^{K(1)}$  as a representation of  $G_k$ . Let  $z \in Z^-$  lifting  $\lambda$ . We have  $KzK = K\lambda K$  and by [7, Proposition 6.13] the group

$$K_\lambda = K(1)(K \cap z^{-1}Kz)$$

is the inverse image in  $K$  of a parabolic subgroup  $P_k = M_k N_k$  of  $G_k$  containing  $B_k$ , of unipotent radical  $N_k$  equal to the image in  $G_k$  of  $\langle \bigcup_{\alpha \in \Phi^+, \alpha \circ v(z) < 0} U_{\alpha, 0} \rangle$  as  $v(z)$  is anti-dominant and  $\langle \alpha, z \rangle = \langle \alpha, -v(z) \rangle$  in the notation of [7, 6.11]; it is a parahoric

subgroup of  $G$  of pro- $p$ -radical  $K_\lambda(1) = K(1)(K \cap z^{-1}K(1)z)$ . Let  $\rho_z$  be the representation of  $K \cap z^{-1}Kz$  on the space  $V(\rho)$  of  $\rho$  such that  $\rho_z(k) = \rho(zkz^{-1})$ . The map  $f \mapsto \phi : k \mapsto f(zk) : \text{Ind}_K^{KzK} \rho \rightarrow \text{Ind}_{K \cap z^{-1}Kz}^K \rho_z$  is a  $K$ -equivariant isomorphism. It restricts to a  $K$ -equivariant isomorphism

$$(\text{Ind}_K^{KzK} \rho)^{K(1)} \rightarrow (\text{Ind}_{K \cap z^{-1}Kz}^K \rho_z)^{K(1)} = \text{Ind}_{K_\lambda}^K (\rho_z^{K(1) \cap z^{-1}Kz}),$$

where the natural representation of  $K \cap z^{-1}Kz$  on  $\rho_z^{K(1) \cap z^{-1}Kz}$  is extended to a representation of  $K_\lambda$  trivial on  $K(1)$ . The representation  $\rho_z^{K(1) \cap z^{-1}Kz}$  of  $K_\lambda$  identifies to the representation  $\rho_z^{N_k}$  of  $P_k$  on the space  $V(\rho^{N_k})$  of  $\rho^{N_k}$  such that  $\rho_z(m) = \rho(zmz^{-1})$  for  $m$  in the group  $M_0 = \langle Z_0, \bigcup_{\alpha \in \Phi, \alpha \circ v(z)=0} U_{\alpha,0} \rangle$ . The representation  $\text{Ind}_{K_\lambda}^K (\rho_z^{K(1) \cap z^{-1}Kz})$  identifies to  $\text{Ind}_{P_k}^{G_k} (\rho_z^{N_k})$ . The representation  $\rho_z^{N_k}$  of  $P_k$  is irreducible [2]. The  $U_k$ -invariant functor

$$\text{Hom}_{G_k}(\rho_1, \text{Ind}_{P_k}^{G_k} (\rho_z^{N_k})) \rightarrow \text{Hom}_{\mathfrak{h}}(\rho_1^{U_k}, (\text{Ind}_{P_k}^{G_k} (\rho_z^{N_k}))^{U_k}) \tag{50}$$

is an isomorphism, by Cabanes’s equivalence recalled at the beginning of this section, because  $\text{Ind}_{P_k}^{G_k} (\rho_z^{N_k})$  and its contragredient are generated by their  $U_k$ -invariant vectors. This is a general property proved in the next lemma.

**Lemma 7.6.** *Let  $\tau$  be an irreducible  $R$ -representation of  $P_k$  trivial on  $N_k$ . The representation  $\text{Ind}_{P_k}^{G_k} \tau$  of  $G_k$  and its contragredient are isomorphic to a subrepresentation and to a quotient of  $\text{Ind}_{U_k}^{G_k} 1$ . In particular, they are generated by their  $U_k$ -invariant vectors. Their socle and their heads are multiplicity free.*

**Proof.** A representation of  $G_k$  is generated by its  $U_k$ -invariant vectors if and only if it is a quotient of a direct sum of representations isomorphic to  $\text{Ind}_{U_k}^{G_k} 1$ .

The representation  $\text{Ind}_{P_k}^{G_k} \tau$  is a quotient of  $\text{Ind}_{U_k}^{G_k} 1$ , because it is generated by a  $U_k$ -invariant vector (a function in  $\text{Ind}_{P_k}^{G_k} \tau$  of support  $P_k$  with non-zero value in  $\tau^{U_k \cap M_k}$ ).

The inflation of  $\tau$  to  $P_k$  is contained in  $\text{Ind}_{U_k}^{P_k} 1$ . By transitivity of the induction,  $\text{Ind}_{P_k}^{G_k} \tau$  is contained in  $\text{Ind}_{U_k}^{G_k} 1$ .

The contragredient representation  $(\text{Ind}_{P_k}^{G_k} \tau)^*$  is a subrepresentation and a quotient of  $\text{Ind}_{U_k}^{G_k} 1$ , because  $\text{Ind}_{U_k}^{G_k} 1$  is isomorphic to its contragredient, the contragredient permutes the irreducible  $R$ -representations of  $M_k$ , and it commutes with the parabolic induction.

The socle of a subrepresentation of  $\text{Ind}_{U_k}^{G_k} 1$  is contained in the socle of  $\text{Ind}_{U_k}^{G_k} 1$ . The socle of  $\text{Ind}_{U_k}^{G_k} 1$  is multiplicity free, because  $\dim \rho_{U_k} = 1$ , and by adjunction  $\text{Hom}_{G_k}(\rho, \text{Ind}_{U_k}^{G_k} 1) \simeq \text{Hom}_{U_k}(\rho_{U_k}, 1)$  for any irreducible  $R$ -representation  $\rho$  of  $G_k$  of  $U_k$ -coinvariants  $\rho_{U_k}$ .

The contragredient of the socle is the head of the contragredient. □

With (51) and the  $I(1)$ -invariant functor (Proposition 7.5 for  $\rho_1 = \rho$ ), we transfer our results on the spherical algebra  $\mathcal{H}(\eta, \mathfrak{h})$  to the spherical algebra  $\mathcal{H}_R(G, K, \rho)$ , which is the convolution algebra of compactly supported functions

$$\phi : G \rightarrow \text{End}_R(V(\rho)) \text{ satisfying } \phi(k_1 g k_2) = \rho(k_1) \phi(g) \rho(k_2) \text{ for } k_1, k_2 \in K, g \in G.$$

It is isomorphic to the algebra  $\text{End}_{RG} \text{c-Ind}_K^G \rho$  by the map sending  $\phi$  to the  $RG$ -intertwiner  $E_\phi$  of  $\text{c-Ind}_K^G \rho$  defined by

$$E_\phi(f_1)(g) = \phi(g)(v) \quad (g \in G). \tag{51}$$

The spherical Hecke  $R$ -algebra  $\mathcal{H}_R(G, K, \rho)$  admits a natural basis [7, 7.3]  $(\mathcal{F}_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$ , where

$$\mathcal{F}_{\tilde{\lambda}} \text{ has support } K\lambda K \quad \text{and} \quad \mathcal{F}_{\tilde{\lambda}}(\tilde{\lambda})(v) = v. \tag{52}$$

The basis  $(\mathcal{F}_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  does not satisfy (14) in general. The basis (52) for the spherical Hecke algebra  $\mathcal{H}_R(Z, Z_0, \chi)$  is denoted by  $(\tau_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi}$ ,

$$\tau_{\tilde{\lambda}} \text{ has support } Z_0\lambda \quad \text{and} \quad \tau_{\tilde{\lambda}}(\tilde{\lambda})(v) = v.$$

The basis (52) for the central spherical Hecke subalgebra  $\mathcal{H}_R(T, T_0, \rho^{I(1)})$  is  $(\tau_{\tilde{\mu}})_{\mu \in \Lambda_T}$ , and the  $\mathcal{H}_R(T, T_0, \rho^{I(1)})$ -module  $\mathcal{H}_R(Z, Z_0, \rho^{I(1)})$  is finitely generated. We denote by  $\mathcal{H}_R(T^+, T_0, \rho^{I(1)}) \subset \mathcal{H}_R(Z^+, Z_0, \rho^{I(1)})$  the subalgebras of bases  $(\tau_{\tilde{\mu}})_{\mu \in \Lambda_T^+}$  and  $(\tau_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$ . The basis  $(\tau_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  satisfies (14).

**Theorem 7.7.** *The  $R$ -algebras*

$$\mathcal{H}_R(G, K, \rho) \simeq \text{End}_{RG} \text{c-Ind}_K^G \rho \simeq \text{End}_{\mathcal{H}}(\eta \otimes_{\mathfrak{h}} \mathcal{H}) = \mathcal{H}(\eta, \mathfrak{h})$$

are isomorphic via (51) and the  $I(1)$ -invariant functor (Proposition 7.5).

The basis  $(\mathcal{F}_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  of  $\mathcal{H}_R(G, K, \rho)$  (52) corresponds to the basis  $(\mathcal{E}_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  of  $\mathcal{H}(\eta, \mathfrak{h})$  (Proposition 4.4).

The basis  $(\phi_{o, \tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  of  $\mathcal{H}_R(G, K, \rho)$  corresponding to the basis  $(\Phi_{o, \tilde{\lambda}})_{\lambda \in \Lambda_\chi^+}$  of  $\mathcal{H}(\eta, \mathfrak{h})$  (Proposition 4.13) satisfies (14).

For  $\mu \in \Lambda_T^+$ ,  $\phi_{\tilde{\mu}} = \phi_{o, \tilde{\mu}}$  does not depend on the choice of  $o$ .

$(\phi_{\tilde{\mu}})_{\mu \in \Lambda_T^+}$  is a basis of a central subalgebra  $\mathcal{Z}_R(G, K, \rho)_T$  of  $\mathcal{H}_R(G, K, \rho)$ , and  $\mathcal{H}_R(G, K, \rho)$  is a finitely generated  $\mathcal{Z}_R(G, K, \rho)_T$ -module (Proposition 5.4).

**Remark 7.8.** The  $RG$ -endomorphism of  $\text{c-Ind}_K^G \rho$  corresponding to  $\phi_{\tilde{\mu}}$  sends  $[1, v]_K$  to  $[1, v]_K E_o(\tilde{\mu})$  for any orientation  $o$  such that  $\eta = \chi_o$  (Propositions 7.2 and 4.13).

We denote by  $\mathcal{A}_{o, T}^+$  the  $R$ -algebra of basis  $(1 \otimes E_o(\tilde{\mu}))_{\mu \in \Lambda_T^+}$ .

**Corollary 7.9.** *We have an  $R$ -algebra isomorphism*

$$(\phi_{o, \tilde{\lambda}})_{\lambda \in \Lambda_\chi^+} \mapsto (\tau_{\tilde{\lambda}})_{\lambda \in \Lambda_\chi^+} : \mathcal{H}_R(G, K, \rho) \xrightarrow{S_o} \mathcal{H}_R(Z^+, Z_0, \chi)$$

restricting to an isomorphism  $\mathcal{Z}_R(G, K, \rho)_T \xrightarrow{S_T} \mathcal{H}_R(T^+, T_0, \chi)$  independent of  $o$ . We have the  $R$ -algebra isomorphisms

$$\begin{aligned} \mathcal{Z}_T &\rightarrow \mathcal{Z}_R(G, K, \rho)_T \xrightarrow{S_T} \mathcal{H}_R(T^+, T_0, \chi) \rightarrow \mathcal{A}_{o, T}^+ \rightarrow R[\tilde{\Lambda}_T^+] \rightarrow R[\Lambda_T^+] \\ (E(\tilde{C}(\mu)))_{\mu \in \Lambda_T^+} &\rightarrow (\phi_{\tilde{\mu}})_{\mu \in \Lambda_T^+} \rightarrow (\tau_{\tilde{\mu}})_{\mu \in \Lambda_T^+} \rightarrow (E_o(\tilde{\mu}))_{\mu \in \Lambda_T^+} \rightarrow (\tilde{\mu})_{\mu \in \Lambda_T^+} \rightarrow (\mu)_{\mu \in \Lambda_T^+}. \end{aligned}$$

When the group  $G$  is split,  $(Z^+, Z_0) = (T^+, T_0)$  and  $\mathcal{Z}_R(G, K, \rho)_T = \mathcal{H}_R(G, K, \rho)$ .

Theorem 1.5 in §1 follows from Corollary 7.9 and the next proposition. The  $R$ -characters  $\xi$  of  $\Lambda_T^+$  identify with the characters of the  $R$ -algebras isomorphic to  $R[\tilde{\Lambda}_T^+]$  in Corollary 7.9. We write

$$\xi(\tau_{\tilde{\mu}}) = \xi(E(\tilde{C}(\mu))) = \xi(\phi_{\tilde{\mu}}) = \xi(E_o(\tilde{\mu})) = \xi(\tilde{\mu}) = \xi(\mu)$$

for  $\mu \in \Lambda_T^+$ . Let  $\pi$  be a smooth  $R$ -representation of  $G$ . We suppose that  $\pi|_K$  contains  $\rho$ .

**Proposition 7.10.** *Let  $A \in \text{Hom}_{RK}(\rho, \pi)$  be non-zero, and let  $\mu \in \Lambda_T^+$ . We have*

$$(A\phi_{\tilde{\mu}})(v) = A(v)E_o(\tilde{\mu}) = A(v)E(\tilde{C}(\mu)).$$

*In particular, if  $A$  is a  $\mathcal{Z}_R(G, K, \rho)_T$ -eigenvector in  $\text{Hom}_{RK}(\rho, \pi)$  of eigenvalue  $\xi$ ,*

$$\xi(\tilde{\mu})A(v) = A(v)E_o(\tilde{\mu}) = A(v)E(\tilde{C}(\mu)).$$

**Proof.** By the adjunction isomorphism,  $A$  and  $A\phi_{\tilde{\mu}}$  correspond to the  $RG$ -intertwiners  $\text{c-Ind}_K^G \rho \rightarrow \pi$  sending  $[1, v]_K$  to  $A(v)$  and to  $A(v)E_o(\tilde{\mu})$  (Remark 7.8). We deduce that  $(A\phi_{\tilde{\mu}})(v) = A(v)E_o(\tilde{\mu})$ .

The  $\mathcal{H}$ -isomorphism  $(\text{c-Ind}_K^G \rho)^{I(1)} \rightarrow \chi_o \otimes_{\mathfrak{o}} \mathcal{H}$  of Proposition 7.2 sends  $[1, v]_K E(\tilde{C}(\mu))$  to  $1 \otimes E(\tilde{C}(\mu))$ . By Proposition 5.4,  $1 \otimes E(\tilde{C}(\mu)) = 1 \otimes E_o(\tilde{\mu})$ . Hence  $[1, v]_K E(\tilde{C}(\mu)) = [1, v]_K E_o(\tilde{\mu})$ . Applying the  $\mathcal{H}$ -intertwiner  $(\text{c-Ind}_K^G \rho)^{I(1)} \rightarrow \pi^{I(1)}$  corresponding to  $A$  sending  $[1, v]_K$  to  $A(v)$ , we deduce that  $A(v)E_o(\tilde{\mu}) = A(v)E(\tilde{C}(\mu))$ .

If  $A$  is a  $\mathcal{Z}_R(G, K, \rho)_T$ -eigenvector in  $\text{Hom}_{RK}(\rho, \pi)$  of eigenvalue  $\xi$  (Corollary 7.9), we have  $A\phi_{\tilde{\mu}} = \xi(\phi_{\tilde{\mu}})A$  for  $\mu \in \Lambda_T^+$  (Theorem 7.7). □

For  $J \subset \Delta$ , we denote by  $\mu_J$  an element of  $\Lambda_T^+$  such that  $\alpha \circ v(\mu_J) > 0$  for all  $\alpha \in \Delta - J$ .

**Remark 7.11.** Let  $\xi$  be an  $R$ -character of  $\Lambda_T^+$ . The character  $\xi$  is called supersingular if it satisfies the following three equivalent properties.

- (1)  $\xi(\mu) = 0$  for all  $\mu \in \Lambda_T^+$  non-invertible in  $\Lambda_T^+$ .
- (2)  $\xi(\mu_J) = 0$  for any  $J \neq \Delta$ .
- (3) For some  $n \geq 1$ ,  $\xi(\mu) = 0$  for all  $\mu \in \Lambda_T^+$  with  $\ell(\mu) > n$ .

In Proposition 7.10, the eigenvalue  $\xi$  of  $A$  is supersingular if and only if the module  $A(v)\mathcal{H}$  is supersingular (Definition 6.10).

**Acknowledgements.** I am grateful to N. Abe, G. Henniart, and R. Ollivier for helpful discussions. I thank the organizer of the conferences in Mumbai (D. Prasad) and in Berlin (E. Grosse-Kloenne) for their interest and for giving me the opportunity to present this work, and the Institut de Mathématiques de Jussieu for its support. I thank the anonymous referee for having read my article so carefully and for his/her comments.

**References**

1. N. ABE, G. HENNIART, F. HERZIG AND M.-F. VIGNÉRAS, A classification of admissible irreducible modulo  $p$  representations of reductive  $p$ -adic groups, preprint, 2014.
2. M. CABANES, Irreducible modules and Levi supplements, *J. Algebra* **90** (1984), 84–97.

3. M. CABANES, A criterion of complete reducibility and some applications, in *Représentations linéaires des groupes finis*, (ed. M. CABANES), Astérisque, pp. 181–182. (1990) 93–112.
4. M. CABANES AND M. ENGUEHARD, *Representation theory of finite reductive groups* (Cambridge University Press, 2004).
5. E. GROSSE-KLÖNNE, From pro- $p$ -Iwahori Hecke modules to  $(\phi, \Gamma)$ -modules, preprint, 2013.
6. G. HENNIART AND M.-F. VIGNERAS, Comparison of compact induction with parabolic induction, *Pac. J. Math.* **260**(2) (2012), 457–495.
7. G. HENNIART AND M.-F. VIGNERAS, A Satake isomorphism for representations modulo  $p$  of reductive groups over local fields, *J. Reine Angew. Math.* **701** (2015), 33–75.
8. S. KUMAR, Kac–Moody groups, their flag varieties and representation theory, *Progr. Math.* **204** (2002).
9. R. OLLIVIER, Parabolic Induction and Hecke modules in characteristic  $p$  for  $p$ -adic  $GL_n$ , *ANT* **4**(6) (2010), 701–742.
10. R. OLLIVIER, An Inverse Satake isomorphism in characteristic  $p$ , *Selecta Math.*, preprint, 2012.
11. R. OLLIVIER, Compatibility between Satake and Bernstein isomorphisms in characteristic  $p$ , preprint, 2013.
12. N. A. SCHMIDT, *Generische pro- $p$ -algebren* (Dilpomarbeit, Berlin, 2009).
13. M.-F. VIGNÉRAS, On a numerical Langlands correspondence modulo  $p$  with the pro- $p$ -Iwahori Hecke ring, *Math. Ann.* **331**(3) 1432–1807. Erratum **333**(3) (2005), 699–701.
14. M.-F. VIGNÉRAS, Algèbres de Hecke affines génériques, *J. Represent. Theory* **10** (2006), 1–20.
15. M.-F. VIGNÉRAS, The pro- $p$ -Iwahori Hecke algebra of a reductive  $p$ -adic group I, preprint, 2013.
16. M.-F. VIGNÉRAS, The pro- $p$ -Iwahori Hecke algebra of a reductive  $p$ -adic group II, *Muenster J. Math.* **7** (2014), 363–379.