

Gestational Weight Gain and Maternal and Neonatal Outcomes in Term Twin Pregnancies in Obese Women

Jeffrey A. Gavard and Raul Artal

Department of Obstetrics, Gynecology, and Women's Health, Saint Louis University School of Medicine, St. Louis, MO, USA

Limited data is available that estimates the effect of gestational weight gain on maternal and neonatal outcomes in term twin pregnancies in obese women. A historical cohort study of 831 obese (BMI ≥ 30.0 kg/m²) women in Missouri delivering 1,662 liveborn, term (≥ 37 weeks gestation) twin infants in 1998–2005 was conducted. Three gestational weight gain categories were examined: <25 pounds, 25–42 pounds, and >42 pounds. Adjusted odds ratios were calculated with multiple logistic regression, using the 2009 Institute of Medicine provisional guideline of 25–42 pounds as the reference group. Significant increasing trends with gestational weight gain were found for preeclampsia ($p < .05$), larger twin birth weight ($p < .01$), smaller twin birth weight ($p < .001$), and infants weighing >2,500 grams ($p < .001$). Significant increasing trends for preeclampsia and for cesarean delivery were found in concordant twin pairs (smaller twin >80% of birth weight of larger twin). Women who gained >42 pounds had a borderline significantly higher odds of preeclampsia than women who gained 25–42 pounds (adjusted OR 1.72; 95% CI 1.00–2.99, $p = .052$). No significant differences were found for 1-min Apgar score <4, 5-min Apgar score <7, or infant mortality ≤ 1 year. Our study suggests that increasing gestational weight gain is associated with larger infants but increased risk of preeclampsia and cesarean delivery in term twin pregnancies in obese women. Limiting gestational weight gain could reduce the risk of preeclampsia and cesarean delivery. Prospective studies of other study populations and maternal/infant outcomes are needed to evaluate the efficacy of the Institute of Medicine guideline.

■ **Keywords:** gestational weight gain, twins, obese, maternal outcomes, neonatal outcomes, epidemiology

Twins account for 3.3% of all live births in the United States (Martin et al., 2012) and are associated with high infant morbidity and mortality (Kogan et al., 2000; Mathews & MacDorman, 2006). Maternal obesity has been found to be significantly related to an increased risk of preeclampsia and other adverse outcomes (Sebire et al., 2001; Weiss et al., 2004). The Institute of Medicine (IOM) in 2009 issued a provisional guideline for gestational weight gain of 25–42 pounds for term twin pregnancies in obese women (BMI ≥ 30.0 kg/m²) that reflected the interquartile range (25th–75th percentile) of cumulative weight gain among obese women who delivered twins at 37–42 weeks gestation (Institute of Medicine and National Research Council, 2009). Limited data is available, however, documenting the efficacy of this provisional guideline on maternal/neonatal outcomes. A study of 29 obese women found no difference in adverse outcomes between women whose gestational weight gain was below the IOM guideline and women whose gesta-

tional weight gain met or exceeded the IOM guideline (Fox et al., 2010). Another study found no difference in outcomes between women whose gestational weight gain was below, equal to, or exceeded IOM guidelines; normal, overweight, and obese women, however, were included in this study (Fox et al., 2011). The purpose of this study was to estimate the effect of gestational weight gain on maternal/neonatal outcomes in term (≥ 37 weeks gestation) twin pregnancies in obese women using the IOM provisional guideline of 25–42 pounds as the reference group.

RECEIVED 31 October 2013; ACCEPTED 3 December 2013. First published online 15 January 2014.

ADDRESS FOR CORRESPONDENCE: Jeffrey A. Gavard, Department of Obstetrics, Gynecology, and Women's Health, Saint Louis University School of Medicine, 6420 Clayton Road, Suite 290, St. Louis, MO 63117, USA. E-mail: gavardja@slu.edu

Materials and Methods

A population-based historical cohort study was conducted using data from the Missouri maternally linked birth and fetal death certificate registry. The methods and algorithms used within the Missouri vital records system have been previously described, are considered reliable, and have been used to validate national data sets of the United States that use matching and linking procedures (Herman et al., 1997; Martin et al., 2003). The study population included all obese pregnant women (BMI ≥ 30.0 kg/m²) residing in Missouri who delivered liveborn, term (clinical estimate of gestational age ≥ 37 weeks) twin infants without congenital abnormalities in 1998–2005. Pre-pregnancy BMI was calculated from self-reported weight and height recorded on the birth certificate. Gestational weight gain was abstracted from the medical chart or provided by the physician. Three categories of gestational weight gain were examined: <25 pounds (includes women who lost weight), 25–42 pounds, and >42 pounds. The 25–42 pound group served as the reference category for all comparisons, as it is the current IOM provisional guideline for weight gain for term twin pregnancies in obese women. The <25-pound group represented women gaining less than the IOM provisional guideline or losing weight during pregnancy, while the >42-pound group represented women whose weight gain exceeded the IOM provisional guideline.

Maternal and neonatal outcomes of this study included preeclampsia, cesarean delivery, instrumental delivery (forceps or vacuum), infant birth weight, infants weighing $\leq 2,500$ grams, 1-min Apgar score <4, 5-min Apgar score <7, and infant mortality ≤ 1 year. Preeclampsia was defined on the birth certificate as pregnancy-induced hypertension occurring >20 weeks gestation that was characterized by either blood pressure >140/90 mm Hg or an increase in systolic blood pressure ≥ 30 mm Hg or an increase in diastolic blood pressure ≥ 15 mm Hg over baseline on two measurements taken six hours apart that was accompanied by generalized and overt proteinuria or edema. Potential confounders that could impact the risk of the above maternal/neonatal outcomes that were adjusted for in the study included maternal age (≤ 25 years, 26–35 years, >35 years), race (non-Hispanic white, black, Hispanic, other), education (<12 years, 12 years, >12 years), pre-pregnancy body mass index (Class I Obese [≥ 30.0 –<35.0 kg/m²], Class II Obese [≥ 35.0 –<40.0 kg/m²], Class III Obese [≥ 40.0 kg/m²]), socio-economic status (using enrollment in Medicaid; the Women, Infants, and Children public health program; or food stamp programs as a proxy, as income was not reported on the birth certificate), smoking, parity (0, 1+ prior live births), diabetes, chronic hypertension, and gestational age at delivery. Diabetes was defined on the birth certificate as insulin-dependent diabetes or other diabetes. The latter could be either pre-existing or gestational diabetes. Women were combined into a single diabetes group for purposes of analysis.

Continuous variables were expressed as medians due to lack of normality of the distributions. Categorical variables were expressed as numbers and percentages. Differences in demographic characteristics, medical/obstetrical history, and maternal/neonatal outcomes by gestational weight gain categories were assessed using chi-square test for categorical variables, including the linear-by-linear test for trend. The non-parametric Kruskal-Wallis test was used for all continuous variables. The multivariate predictability of gestational weight gain categories for maternal/neonatal outcomes was examined using multiple logistic regression. Multivariate findings reflect adjustment for the above potential confounders in the model. Maternal/neonatal outcomes also were assessed by gestational weight gain categories according to concordant and discordant twin pairs. Discordance was defined as birth weight $\geq 20\%$ according to the formula: (birth weight of larger twin – birth weight of smaller twin)/birth weight of larger twin $\times 100$. The 20% level for birth weight discordance was chosen because it is commonly used in twin studies and has been shown to be approximately the level above which perinatal morbidity significantly increases (Amaru et al., 2004; Breathnach et al., 2011). A *p* value of <.05 was used for statistical significance. All analyses were performed using SPSS version 19.0 for Windows.

This research did not involve interaction with subjects and the existing publicly available data contained no identifying private information. The Saint Louis University Institutional Review Board deemed that this study did not require formal review according to federal regulation 45 CFR 46.102.

Results

The study population of 1,662 twins was born to 405 (48.7%) Class I Obese women, 223 (26.8%) Class II Obese women, and 203 (24.4%) Class III Obese women (Table 1). Over 70% of the women were non-Hispanic white. Approximately 60% of the women lived in poverty and 27.8% of the mothers were nulliparous. Only 38.9% of the women followed the Institute of Medicine's provisional guideline of weight gain during pregnancy of 25–42 pounds. The percentage of women gaining <25 pounds during pregnancy was 30.8%, while 30.3% of the women gained >42 pounds. Eleven of the 256 (4.3%) women in the <25-pound group lost weight during pregnancy, with a range of weight loss of 4–16 pounds and a median weight loss of 8 pounds. Preeclampsia, cesarean delivery, and instrumental delivery occurred in 10.5%, 68.0%, and 6.9% of the women, respectively. Women who gained >42 pounds during pregnancy were more likely to be non-white than women who gained <25 pounds or 25–42 pounds. Women who gained <25 pounds were less likely to be Class I Obese and nulliparous than women who gained 25–42 pounds or >42 pounds.

TABLE 1
Maternal Characteristics for 831 Obese Women Who Had Twin Pregnancies in Missouri, 1998–2005

Characteristic	Maternal weight gain during pregnancy (lb)						p value
	<25 ^a (n = 256)		25–42 ^b (n = 323)		>42 ^c (n = 252)		
	n	%	n	%	n	%	
Maternal age (years)							
≤25	79	30.9	111	34.4	98	38.9	
26–35	145	56.6	183	56.7	139	55.2	.08
>35	32	12.5	29	9.0	15	6.0	
Maternal race							
White (non-Hispanic)	196	76.9	242	74.9	154	61.4	
Black	47	18.4	72	22.3	86	34.3	<.001
Hispanic	6	2.4	5	1.5	8	3.2	
Other	6	2.4	4	1.2	3	1.2	
Maternal education (years)							
<12	36	14.2	38	11.8	47	18.7	
12	86	33.9	127	39.4	88	35.1	.14
>12	132	52.0	157	48.8	116	46.2	
Pre-pregnancy body mass index (kg/m ²)							
Class I Obese (≥30.0–<35.0)	97	37.9	168	52.0	140	55.6	
Class II Obese (≥35.0–<40.0)	74	28.9	92	28.5	57	22.6	<.001
Class III Obese (≥40.0)	85	33.2	63	19.5	55	21.8	
Socio-economic status ^d	154	60.4	179	55.4	162	64.3	.10
Smoked during pregnancy	43	16.8	39	12.1	45	17.9	.12
Adequacy of prenatal care ^e							
Inadequate	22	8.8	19	6.1	14	5.7	
Intermediate	2	0.8	5	1.6	3	1.2	.66
Adequate	18	7.2	24	7.7	24	9.8	
Adequate plus	207	83.1	265	84.7	204	83.3	
Nulliparous	53	20.7	93	28.8	85	33.7	<.01
Anemia	10	3.9	3	0.9	8	3.2	.06
Cardiac disease	1	0.4	0	0.0	0	0.0	.33
Renal disease	0	0.0	0	0.0	1	0.4	.32
Diabetes (any type)	17	6.6	25	7.7	19	7.5	.87
Chronic hypertension	13	5.1	12	3.7	7	2.8	.40
Preeclampsia	21	8.2	30	9.3	36	14.3	<.05
Cesarean delivery	162	63.3	224	69.3	179	71.0	.06
Instrumental delivery (forceps or vacuum)	18	7.0	15	4.6	24	9.5	.27

Note: ^a Maternal race was unknown for one woman who gained <25 pounds during pregnancy; maternal education was unknown for two women who gained <25 pounds during pregnancy; socio-economic status was unknown for one woman who gained <25 pounds during pregnancy; adequacy of prenatal care was unknown for seven women who gained <25 pounds during pregnancy.

^b Maternal education was unknown for one woman who gained 25–42 pounds during pregnancy; adequacy of prenatal care was unknown for 10 women who gained 25–42 pounds during pregnancy.

^c Maternal race was unknown for one woman who gained >42 pounds during pregnancy; maternal education was unknown for one woman who gained >42 pounds during pregnancy; adequacy of prenatal care was unknown for seven women who gained >42 pounds during pregnancy.

^d Enrollment in Medicaid; Women, Infants, and Children's Program; or food stamps.

^e The Kotelchuck Index of Adequacy of Prenatal Care defines the levels as inadequate (<50% of expected prenatal care visits), intermediate (≥50%–<80% of expected prenatal care visits), adequate (≥80%–<110% of expected prenatal care visits), and adequate plus (≥110% of expected prenatal care visits).

A significant increasing trend with gestational weight gain was found for preeclampsia ($p < .05$). An increasing trend with gestational weight gain for cesarean delivery was of borderline significance ($p = .06$).

Significant increasing trends with gestational weight gain were found for infant birth weight ($p < .001$), larger twin birth weight ($p < .01$), smaller twin birth weight ($p < .001$), and average twin birth weight ($p < .001$, Table 2). The weight difference in twin pairs, however, was relatively constant across gestational weight gain categories ($p = .77$). A significant inverse trend with increasing gestational weight gain was found for infants weighing ≤2,500 grams ($p < .001$). No significant differences were found for 1-min Apgar score <4, 5-min Apgar score <7, or infant mortality within 1 year.

The incidence and multivariate findings for preeclampsia and infants weighing ≤2,500 grams by gestational weight gain category are given in Table 3. Women who gained >42 pounds had a borderline significantly higher odds of preeclampsia than women who gained 25–42 pounds (adjusted OR 1.72; 95% CI 1.00–2.99, $p = .052$). Women who gained <25 pounds had significantly higher odds of an infant weighing ≤2,500 grams than women who gained 25–42 pounds (adjusted OR 1.74; 95% CI 1.28–2.36, $p < .001$). Analyses by obesity class showed that women who gained >42 pounds always had an elevated, although non-significant, odds of preeclampsia than women who gained 25–42 pounds (data not shown). Class I Obese women who gained <25 pounds had significantly higher odds of an infant weighing ≤2,500 grams than women who gained

TABLE 2
Outcomes for 1,662 Twins Born to 831 Obese Women in Missouri, 1998–2005

Characteristic	Maternal weight gain during pregnancy (lb)						p value
	<25 ^a (n = 512)		25–42 ^b (n = 646)		>42 ^c (n = 504)		
	n	%	n	%	n	%	
Median gestational age at delivery (weeks)	37		38		38		<.01
Median infant birth weight (grams)	2,718		2,835		2,835		<.001
Infant birth weight ≤2,500 grams	144	28.1	122	18.9	87	17.3	<.001
Median larger twin birth weight (grams)	2,863		2,977		2,977		<.01
Median smaller twin birth weight (grams)	2,580		2,665		2,715		<.001
Median difference in twin weight (grams)	225.5		227.0		221.0		.77
Median average twin weight (grams)	2,712		2,835		2,851		<.001
1-min Apgar score <4	7	1.4	13	2.0	9	1.8	.61
5-min Apgar score <7	3	0.6	10	1.6	3	0.6	.98
Infant mortality ≤1 year	2	0.4	1	0.2	4	0.8	.33

Note: ^a One-min and 5-min Apgar scores were unknown for one infant whose mother gained <25 pounds during pregnancy.
^b One-min and 5-min Apgar scores were unknown for three infants whose mothers gained 25–42 pounds during pregnancy.
^c One-min and 5-min Apgar scores were unknown for two infants whose mothers gained >42 pounds during pregnancy.

TABLE 3
Incidence and Multivariate Risk of Preeclampsia and Infant Birth Weight ≤2,500 Grams by Pregnancy Weight Gain for 1,662 Twins Born to 831 Obese Women in Missouri, 1998–2005

Maternal/neonatal Outcome	Weight gain during pregnancy (lb)	Incidence		Trend	Multivariate ^a		
		n	%	p value	Odds ratio	95% CI	p value
Preeclampsia	<25 ^b (n = 256)	21	8.2	<.05	0.88	0.47–1.66	.70
	25–42 (n = 323)	30	9.3		1.00	Reference	Reference
	>42 (n = 252)	36	14.3		1.72	1.00–2.99	.05
Infant birth weight ≤2,500 grams	<25 ^b (n = 512)	144	28.1	<.001	1.74	1.28–2.36	<.001
	25–42 (n = 646)	122	18.9		1.00	Reference	Reference
	>42 (n = 504)	87	17.3		0.75	0.54–1.06	.10

Note: ^a Adjusted for maternal age, race, education, pre-pregnancy body mass index, socio-economic status, smoking, parity, anemia, diabetes, chronic hypertension, adequacy of prenatal care, sex of twin, twin discordance, and gestational age at delivery.
^b Includes women who lost weight during pregnancy.
 Note: CI = confidence interval.

25–42 pounds (adjusted OR 2.04; 95% CI 1.28–3.26, *p* < .01). Class III Obese women who gained >42 pounds had significantly lower odds of an infant weighing ≤2,500 grams than women who gained 25–42 pounds (adjusted OR 0.39; 95% CI 0.17–0.91, *p* < .05).

Significant increasing trends with gestational weight gain were found for preeclampsia (*p* < .05) and cesarean delivery (*p* < .05) in concordant twin pairs (Table 4). Significant inverse trends with increasing gestational weight gain were found for infants weighing ≤2,500 grams in both concordant and discordant twin pairs (*p* < .05). Only 73/831 (8.8%) of the twin pairs were discordant, likely rendering the risk pattern of preeclampsia in discordant twin pairs non-significant due to small sample size.

Discussion

This study focuses on the association of gestational weight gain with maternal/neonatal outcomes in term (≥37 weeks gestation) twin pregnancies in obese women using the revised 2009 IOM provisional guideline of 25–42 pounds. Significant increasing trends across the three gestational weight gain categories of <25 pounds, 25–42 pounds, and >42 pounds were found for infant birth weight, larger

twin birth weight, smaller twin birth weight, and average twin birth weight. A significant inverse trend with increasing gestational weight gain was found for infants weighing ≤2,500 grams. These findings are consistent with those of other studies involving twin pregnancies in obese women (Fox et al., 2011; Gonzalez-Quintero et al., 2012). Another study noted that larger twin birth weight and smaller twin birth weight increased in obese women whose gestational weight gain met or exceeded the IOM guideline compared to women whose gestational weight gain was below the IOM guideline; the sample size of only 29 obese women, however, precluded the finding of significant differences (Fox et al., 2010). Only 38.9% of the women in our study followed the Institute of Medicine’s provisional guideline of weight gain during pregnancy of 25–42 pounds, which is similar to that found in a national sampling of women in 28 states and New York City participating in the population-based surveillance system (PRAMS; Chu & D’Angelo, 2009).

Increased gestational weight gain is a significant risk factor for preeclampsia in obese women (Kabiru & Raynor, 2004). Preeclampsia also occurs with much higher frequency in twin pregnancies than in singleton gestations. In one multicenter trial examining the efficacy of low-dose aspirin for the prevention of preeclampsia, the rates of

TABLE 4
Outcomes by Twin Discordance^a for 1,662 Twins Born to 831 Obese Women in Missouri, 1998–2005

Characteristic	Maternal weight gain during pregnancy (lb)						p value
	<25 (n = 256)		25–42 (n = 323)		>42 (n = 252)		
	n	%	n	%	n	%	
Twin concordance (birth weight >80%) ^a							
Concordant twin pairs	230	89.8	293	90.7	235	93.3	.18
Discordant twin pairs	26	10.2	30	9.3	17	6.7	
Preeclampsia							
Concordant twin pairs	18	7.8	26	8.9	32	13.6	<.05
Discordant twin pairs	3	11.5	4	13.3	4	23.5	.31
Cesarean delivery							
Concordant twin pairs	142	61.7	200	68.3	166	70.6	<.05
Discordant twin pairs	20	76.9	24	80.0	13	76.5	.99
Instrumental delivery							
Concordant twin pairs	16	7.0	14	4.8	23	9.8	.23
Discordant twin pairs	2	7.7	1	3.3	1	5.9	.73
Birth weight ≤2,500 grams: Either twin							
Concordant twin pairs	83	36.1	68	23.2	60	25.5	<.05
Discordant twin pairs	22	84.6	26	86.7	9	52.9	<.05

Note: ^a Twin discordance defined $\geq 20\%$ in birth weight according to the formula: $(\text{Birth weight of larger twin} - \text{Birth weight of smaller twin}) / \text{Birth weight of larger twin} \times 100$.

preeclampsia were found to be 12.7% in twin pregnancies and 4.9% in singleton gestations (Sibai et al., 2000). A population-based study of birth certificates in Washington state found that twin pregnancies had a 3.5 increased risk of preeclampsia compared to singleton pregnancies (Coonrod et al., 1995). The combination of twin gestation and maternal obesity could amplify the risk for preeclampsia. Efforts to identify and minimize risk factors for preeclampsia in twin pregnancies in obese women should be pursued. Such factors may include limiting gestational weight gain in such pregnancies.

Previous studies of twin pregnancies in obese women have demonstrated that increasing gestational weight gain is associated with larger infants without accompanying increases in complications. The rate of gestational hypertension and gestational diabetes did not significantly increase with increasing gestational weight gain in one study (Fox et al., 2010) and the rate of gestational hypertension, preeclampsia, and gestational diabetes did not significantly increase with increasing gestational weight gain in a second study (Fox et al., 2011). The former study again included only 29 obese women. The latter study included 170 women, of whom 117 had a normal pre-pregnancy BMI and the remaining 53 women had an overweight or obese pre-pregnancy BMI. Analyses were performed across all three pre-pregnancy BMI classes simultaneously or only in women with a normal pre-pregnancy BMI. The exact number of obese women in the latter study was unknown and a separate analysis relating gestational weight gain with maternal and neonatal outcomes in obese women was not performed.

A strength of our study was that we had a relatively large sample size of 831 obese mothers and 1,662 twins from which to calculate trends in maternal and neonatal outcomes across gestational weight gain categories. A sig-

nificant increasing trend with gestational weight gain was found for preeclampsia. Such a finding has not been reported in previous studies. Women who gained >42 pounds had a borderline significantly higher odds of preeclampsia than women who gained 25–42 pounds. Elevated, although non-significant, adjusted odds of preeclampsia were found in all obesity classes for women who gained >42 pounds compared to women who gained 25–42 pounds. It is likely that the small number of preeclampsia cases rendered the study underpowered to detect significant differences between gestational weight gain categories within obesity classes. Prospective studies with larger sample sizes are needed to determine if increasing trends with gestational weight gain for preeclampsia exist by obesity class, to corroborate the increased risk of preeclampsia for women gaining above the IOM provisional guideline of 42 pounds in all obesity classes, and to further examine the association between gestational weight gain and preeclampsia and cesarean delivery by concordant and discordant twin pairs.

The risk of preeclampsia, cesarean delivery, and infants weighing $\leq 2,500$ grams was always much higher for discordant twin pairs than for concordant twin pairs at a particular level of gestational weight gain. This corroborates earlier findings for discordant twin pairs on these and other endpoints (Amaru et al., 2004; Breathnach et al., 2011). The risk patterns for preeclampsia and cesarean delivery in concordant twin pairs indicate that increasing gestational weight gain is significantly associated with increased risk for these outcomes in obese women when twins are similar in size at delivery. The risk pattern for preeclampsia in discordant twin pairs had an increasing trend with gestational weight gain, but was non-significant due to few preeclampsia cases. In contrast, the risk of cesarean delivery was relatively constant across gestational weight gain categories in discordant twin pairs. The amount of gestational

weight gain apparently is not significantly associated with the risk of cesarean delivery in obese women when twins are sufficiently dissimilar in size at delivery.

Additional strengths of our study include a historical cohort design that permitted calculation of incidence rates of maternal/neonatal outcomes and the availability of and adjustment for many potential confounding variables that could impact maternal/neonatal outcomes. Limitations of our study include those that are inherent with the use of birth certificate records, such as incomplete or inaccurate reporting of data; self-reported information, such as height and weight; and potential under-reporting of medical conditions, such as chronic hypertension, heart disease, and renal disease. Preeclampsia has been found to match very well between birth certificates with a check-box format (such as used in Missouri), and hospital discharge data, with 85–97% of the former being present in the latter (Frost et al., 1984). This corroboration, and that repeated blood pressure measurements with accompanying proteinuria or edema are necessary for a diagnosis of preeclampsia, lessen the potential of an inflated rate of preeclampsia due to artificial elevation of blood pressure that is recorded when a standard size sphygmomanometer cuff is used in obese women (Fonseca-Reyes et al., 2003). Self-reported maternal weight, as appears on Missouri birth certificates and on which pre-pregnancy BMI classes were based, also has been found to be highly correlated with and similar to clinically recorded weight (Lederman & Paxton, 1998). Self-reported BMI has been found to match very well with clinically measured BMI in obese (Brunner Huber, 2007) and combined overweight and obese samples (Nawaz et al., 2001). Gestational age was determined by clinical estimate, which has been found to be a more accurate measure of gestational age at delivery than length of pregnancy calculated using the last menstrual period (Callaghan & Dietz, 2010). Our study only examined the endpoints of preeclampsia, cesarean delivery, instrumental delivery, indices of birth weight, Apgar scores, and infant mortality. The impact of gestational weight gain on other adverse short- and long-term maternal and infant outcomes such as gestational diabetes, gestational hypertension, neonatal intensive care unit admission, childhood obesity, childhood cognitive performance, and post-partum weight retention remains to be determined. Our study population consisted of liveborn, term (≥ 37 weeks gestation) twin infants free of congenital abnormalities, and our findings may not be generalizable to other obese pregnant populations. Only term infants were included in our study because the IOM provisional guideline of 25–42 pounds was based on twin pregnancies in obese women who delivered at 37–42 weeks gestation. Other studies involving twin pregnancies in obese women have used a rate of weight gain of 0.68 pounds/week that is based on the minimal IOM provisional guideline of 25 pounds divided by the minimal duration of 37 weeks for a term pregnancy, which enables other maternal/neonatal outcomes, such as pre-term birth,

to be evaluated (Fox et al., 2010; Gonzalez-Quintero et al., 2012). Diet, physical activity, and chorionicity were also potential confounders that could not be assessed.

Our findings are consistent with other studies that report larger infants with increased gestational weight gain in twin pregnancies in obese women. Our study, however, found that this occurred with an accompanying increased risk of preeclampsia and cesarean delivery, particularly in concordant twin pairs. Limiting gestational weight gain should be considered in view of the significant complications of preeclampsia and cesarean delivery. Our findings need corroboration from prospective studies of twin pregnancies in obese women that estimate the effect of gestational weight gain on preeclampsia and cesarean delivery for composite samples of obese women and by obesity class. Prospective studies of other study populations and other short- and long-term maternal and infant outcomes also are needed to evaluate the efficacy of the IOM guideline.

Acknowledgments

The original source of data used in this study was from the Missouri Department of Health and Senior Services, Section of Epidemiology for Public Health Practice. The analyses, interpretations, and conclusions of this study are those of the authors and not the Missouri Department of Health and Senior Services, Section of Epidemiology for Public Health Practice.

References

- Amaru, R. C., Bush, M. C., Berkowitz, R. L., Lapinski, R. H., & Gaddipari, S. (2004). Is discordant growth in twins an independent risk factor for adverse neonatal outcome? *Obstetrics and Gynecology*, *103*, 71–76.
- Breathnach, F. M., McAuliffe, F. M., Geary, M., Daly, S., Higgins, J. R., Dornan, J., . . . Malone, F. D. (2011). Definition of intertwin birth weight discordance. *Obstetrics and Gynecology*, *118*, 94–103.
- Brunner Huber, L. R. (2007). Validity of self-reported height and weight in women of reproductive age. *Maternal and Child Health Journal*, *11*, 137–144.
- Callaghan, W. M., & Dietz, P. M. (2010). Differences in birth weight for gestational age distributions according to the measures used to assign gestational age. *American Journal of Epidemiology*, *171*, 826–836.
- Chu, S. Y., & D'Angelo, D. V. (2009). Gestational weight gain among US women who deliver twins, 2001–2006. *American Journal of Obstetrics and Gynecology*, *200*, 390.e1–390.e6.
- Coonrod, D. V., Hickok, D. E., Zhu, K., Easterling, T. R., & Daling, J. R. (1995). Risk factors for preeclampsia in twin pregnancies: A population-based cohort study. *Obstetrics and Gynecology*, *85*, 645–650.
- Fonseca-Reyes, S., de Alba-Garcia, J. G., Parra-Carrillo, J. Z., & Paczka-Zapata, J. A. (2003). Effect of standard cuff on blood pressure readings in patients with obese arms. How

- frequent are arms of a 'large circumference'? *Blood Pressure Monitoring*, 8, 101–106.
- Fox, N. S., Rebarber, A., Roman, A. S., Klauser, C. K., Peress, D., & Saltzman, D. H. (2010). Weight gain in twin pregnancies and adverse outcomes: Examining the 2009 Institute of Medicine guidelines. *Obstetrics and Gynecology*, 116, 100–106.
- Fox, N. S., Saltzman, D. H., Kurtz, H., & Rebarber, A. (2011). Excessive weight gain in term twin pregnancies: Examining the 2009 Institute of Medicine definitions. *Obstetrics and Gynecology*, 118, 1000–1004.
- Frost, F., Starzyk, P., George, S., & McLaughlin, J. F. (1984). Birth complication reporting: The effect of birth certificate design. *American Journal of Public Health*, 74, 505–506.
- Gonzalez-Quintero, V. H., Kathiresan, A. S. Q., Tudela, F. J., Rhea, D., Desch, C., & Istwan, N. (2012). The Association of gestational weight gain per Institute of Medicine guidelines and prepregnancy body mass index on outcomes of twin pregnancies. *American Journal of Perinatology*, 29, 435–440.
- Herman, A. A., McCarthy, B. J., Bakewell, J. M., Ward, R. H., Mueller, B. A., Maconochie, N. E., . . . Skjaerven, R. (1997). Data linkage methods used in maternally-linked birth and infant death surveillance data sets from the United States (Georgia, Missouri, Utah, and Washington State), Israel, Norway, Scotland, and Western Australia. *Paediatric and Perinatal Epidemiology*, 11, 5–22.
- Institute of Medicine and National Research Council. (2009). *Weight gain during pregnancy: Reexamining the guidelines*. Washington, DC: National Academies Press.
- Kabiru, W., & Raynor, B. D. (2004). Obstetric outcomes associated with increase in BMI category during pregnancy. *American Journal of Obstetrics and Gynecology*, 191, 928–932.
- Kogan, M. D., Alexander, G. R., Kotelchuck, M., MacDorman, M. F., Buekens, P., Martin, J. A., & Papiernik, E. (2000). Trends in twin birth outcomes and prenatal care utilization in the United States, 1981–1997. *Journal of the American Medical Association*, 284, 335–341.
- Lederman, S. A., & Paxton, A. (1998). Maternal reporting of prepregnancy weight and birth outcome: Consistency and completeness compared to clinical record. *Maternal and Child Health Journal*, 2, 123–126.
- Martin, J., Curtin, S., Saulnier, M., & Mousavi, J. (2003). *Development of the matched multiple birth file: 1995–1998* (Matched Multiple Birth File CD-ROM series 21, no 13a). Hyattsville, MD: National Center for Health Statistics.
- Martin, J. A., Hamilton, B. E., Ventura, S. J., Osterman, M. J. K., Wilson, E. C., & Mathews, T. J. (2012). Births: Final data for 2010. *National Vital Statistics Reports*, 61, 1–72.
- Mathews, T. J., & MacDorman, M. F. (2006). Infant mortality statistics from the 2003 period linked birth/infant death data set. *National Vital Statistics Reports*, 54, 1–29.
- Nawaz, H., Chan, W., Abdulrahman, M., Larson, D., & Katz, D. L. (2001). Self-reported weight and height: Implications for obesity research. *American Journal of Preventive Medicine*, 20, 294–298.
- Sebire, N. J., Jolly, M., Harris, J. P., Wadsworth, J., Joffe, M., Beard, R. W., . . . Robinson, S. (2001). Maternal obesity and pregnancy outcome: A study of 287,213 pregnancies in London. *International Journal of Obesity and Related Metabolic Disorders*, 25, 1175–1182.
- Sibai, B. M., Hauth, J., Caritis, S., Lindheimer, M. D., MacPherson, C., Klebanoff, M., . . . McNellis, D. (2000). Hypertensive disorders in twin versus singleton gestations. National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *American Journal of Obstetrics and Gynecology*, 182, 938–942.
- Weiss, J. L., Malone, F. D., Emig, D., Bali, R. H., Nyberg, D. A., Comstock, C. H., . . . D'Alton, M. E. (2004). Obesity, obstetric complications, and cesarean delivery rate: A population-based screening study. *American Journal of Obstetrics and Gynecology*, 190, 1091–1097.