

## ***Cetraria steppae* Savicz is conspecific with *Cetraria aculeata* (Schreb.) Fr. according to morphology, secondary chemistry and ecology**

**Olga NADYEINA, Tetiana LUTSAK, Oleg BLUM, Volodymyr GRAKHOV and Christoph SCHEIDEGGER**

**Abstract:** Eurasian *Cetraria steppae* and the more widely distributed *C. aculeata* are two lichen species traditionally distinguished by Eastern European and Spanish lichenologists on the basis of their morphological and ecological characteristics. Other specialists, however, consider them puzzling. This paper aims to evaluate the taxonomic status of these members of the *C. aculeata* group and thereby to clarify their conservation status in Ukraine. Morphological, chemical and ecological features of specimens originating from populations in different regions of Ukraine were tested and compared with the main characteristics commonly used for the species delimitation. Neither morphological nor chemical traits were found to correlate with ecological characteristics on a small geographical scale. Variation in the norstictic acid content detected in 256 individuals from 13 populations in Ukraine showed no correlation with the morphological characteristics that are currently used for species delimitation. These morphological features appear to vary continuously and did not support subdivision among the specimens studied. We hypothesize that *C. steppae* and *C. aculeata* are conspecific, and provide a formal synonymy. Specimens with norstictic acid are regarded as a different chemotype. Possible evolutionary and adaptive roles of norstictic acid in *C. aculeata* s. lat. are discussed. Based on current and historical data, we consider *C. aculeata* s. lat. as vulnerable in Ukraine, according to the IUCN criteria for regional Red List assessment.

**Key words:** IUCN, lichens, norstictic acid, populations, Red Data Book, Red List, Ukraine

*Accepted for publication 4 July 2013*

---

### **Introduction**

Members of the *Cetraria aculeata* group are fruticose, brownish black lichens and they include *C. aculeata* (Schreb.) Fr., *C.*

*australiensis* W. A. Weber ex Kärnefelt, *C. crespoae* (Barreno & Vázquez) Kärnefelt, *C. muricata* (Ach.) Eckfeldt, *C. odontella* (Ach.) Ach., ‘*C. panamericana*’ and *C. steppae* (Savicz) Kärnefelt (Kärnefelt *et al.* 1992; Thell *et al.* 2002; Printzen *et al.* 2013). Of these, *C. aculeata* and *C. steppae* have been considered as two closely related species (Savicz 1924; Kärnefelt 1986; Kärnefelt *et al.* 1993). *Cetraria aculeata* s. lat. was the subject of special taxonomic and phylogenetic studies (Kärnefelt 1986; Kärnefelt *et al.* 1992, 1993; Thell *et al.* 2000, 2002; Printzen *et al.* 2013). The phylogeography of its symbionts has been investigated (Fernández-Mendoza *et al.* 2011; Domaschke *et al.* 2012; Printzen *et al.* 2013; Fernández-Mendoza & Printzen 2013), as well as the anatomy and ecophysiology of extreme morphotypes (Pérez-Ortega *et al.* 2012), and the bacterial communities within geographically distinct populations (Printzen

---

O. Nadyeina: M. G. Kholodny Institute of Botany, Lichenology and Bryology Department, Kyiv, Ukraine; and Biodiversity and Conservation Biology, Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland. Email: nadyeina@gmail.com

T. Lutsak: Taras Shevchenko National University of Kyiv, Department of Botany, Kyiv, Ukraine; and Senckenberg Research Institute and Natural History Museum, Frankfurt on Main, Germany.

O. Blum: M. M. Gryshko National Botanical Garden of the Academy of Science of Ukraine, Bioindication and Chemosystematics Laboratory, Kyiv, Ukraine.

V. Grakhov: M. M. Gryshko National Botanical Garden of the Academy of Sciences of Ukraine, HPLC Center, Kyiv, Ukraine.

C. Scheidegger: Biodiversity and Conservation Biology, Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland.

TABLE 1. Features used for the delimitation of *C. aculeata* and *C. steppae* according to Savicz's (1924) and Kärnefelt's (1986) descriptions.

Character	<i>Cetraria aculeata</i> (Schreb.) Fr.	<i>Cetraria steppae</i> (Savicz) Kärnefelt	Reference
Thallus vagrancy	Attached tufts	Different mode of life from <i>C. aculeata</i>	Savicz 1924
	Branches forming $\pm$ shrubby tufts, which arise directly from the substratum; also tends to develop spherical thalli, composed of more branches than thalli of <i>C. steppae</i>	Branches forming $\pm$ spherical tufts composed of a few branches due to the vagrant mode of life	Kärnefelt 1986
Thallus specific features	Glossy, solid, inflexible, fragile	Matt, smooth, soft and flexible, not fragile	Savicz, 1924
	Glossy to matt; black, brown to light brown; varying fragility		Kärnefelt 1986
Thallus size		Not indicated	Savicz 1924
	2–4(–10) cm high	1–3 cm high	Kärnefelt 1986
Branch size	Long and thin: 0.3–1.0 mm wide	Short and thick: 0.4–2.0 mm wide	Savicz 1924
	Main branches c. 1(–4) mm wide; terminal branches c. 0.3–1.0 mm wide	Main branches c. 0.5–2.0(–4.0) mm wide; terminal branches c. 0.1–5.0 mm wide	Kärnefelt 1986
Isidioid projections	Tiny, rarely present	Absent	Savicz 1924
		Absent or present	Kärnefelt 1986
Chemistry	K–	K–, C–, I–	Savicz 1924
	K–; norstictic acid absent (TLC)	K+; norstictic acid present (TLC)	Kärnefelt 1986

*et al.* 2012). However, the taxonomic status of *C. steppae* as a distinct species has been questioned (Fernández-Mendoza *et al.* 2011; Lutsak *et al.* 2012) as its morphology varies greatly and its distribution range overlaps with that of *C. aculeata* (Poelt 1969; Randle & Saag 2006).

Distinguishing *Cetraria steppae* and *C. aculeata* as distinct species was traditionally based on visual morphological aspects (Savicz 1924; Oxner 1937; Poelt 1969), which include a vagrant mode of life, the surface characteristics of the thalli and morphological parameters of the main and terminal branches (Table 1).

Mereschkowsky was the first to pay attention to vagrancy and the small number of branches among specimens from Crimea (Ukraine). His description is, however, very brief: “Thallus libertus, opacus, laciniis paullulum minus attenuatis” (Mereschkowsky 1921). The author of *C. steppae* at the species level is Savicz, who also gave a concise description (Table 1). He emphasized that both taxa are distinct species without intermediate forms, which implied that *C. steppae* occurs in the southern steppe biotopes and *C. aculeata* in the north (Savicz 1924). This statement guided lichenologists to distinguish the two species, when

the vagrant mode of life or the thickness of branches were not well developed. In the 1980s, Kärnefelt performed a wide-scale survey of the brown *Cetraria* species (Kärnefelt 1986) and revised the traits for morphological distinction of the two species. He introduced a new feature, the presence of norstictic acid in *C. steppae*, which was not considered in the original description of *C. steppae* (Savicz 1924; Table 1) and became crucial for the delimitation of the two species. Moreover, Kärnefelt extended the concept of *C. steppae* and also included specimens with isidioid projections and black, glossy thalli (Table 1). After being described by Savicz (1924), the species was reported in other parts of southern Ukraine (Oxner 1937, 1993; Blum 1996; Blum *et al.* 2009) and Russia (Zhurbenko 1996; Moutchnik & Zavarzin 2005; Golubkova *et al.* 2008; Urbanavichus 2010), Kazakhstan (Wagner & Spribille 2005), Turkey (John 1999; John & Breuss 2004; Yazıcı *et al.* 2010), Iran (Sohrabi & Alstrup 2007; Seaward *et al.* 2008), and Spain (Llimona & Hladun 2001).

In our analysis of 247 herbarium specimens of *C. aculeata* s. lat., we were not able to classify all specimens as belonging to one or other of the two taxa because of contradictions or uncertainties in diagnostic traits (Table 1). The features traditionally used to distinguish them seemed to be quite subjective, such as the degree of vagrancy or the thickness of the branches. The vagrant life form and spherical tuft-like thalli are often difficult to designate, both on herbarium specimens and in the field. Specimens kept in herbaria are usually carefully separated, and could look 'artificially vagrant' (Fig. 1A & C). In the field, the recognizable vagrant thalli grow on bare soil or sand. However, a few centimetres away, individuals may often be attached, which changes their degree of vagrancy (Fig. 1B & D). Delimitation of individuals is also contentious because thalli in vagrant tufts or in mostly attached mats are often densely entangled (Fahselt 2008).

The above-mentioned difficulties motivated us to check the morphological variability within and between populations sampled from different regions in Ukraine. The present study aimed to test characters tradition-

ally used for the delimitation of *C. steppae* and *C. aculeata*, and to elucidate their taxonomic value. We also wanted to study the impact of ecological and geographical factors on the morphological and chemical characteristics within this group. We hypothesized that specimens belonging to *C. steppae* contain norstictic acid in the medulla (Kärnefelt 1986), have wider thallus branches, smaller thalli, a vagrant lifestyle, and a distribution limited to the south of Ukraine (Savicz 1924). The aim was also to test the correlation between the presence of norstictic acid and the content of carbonates in the substrata of arid areas (Hauck *et al.* 2010). Finally, we assess and discuss the conservation status of *C. aculeata* s. lat. according to the IUCN Red List criteria for the next edition of the Red Data Book of Ukraine.

## Materials and Methods

### Materials and sampling design

A total of 503 specimens of *C. aculeata* s. lat. from Ukraine were studied. First, we examined 247 specimens from different herbaria (KW, KHER, LE), according to their morphological, chemical and ecological characteristics (Fig. 1E). We then collected 256 specimens from 13 local populations, representing different natural zones of Ukraine during 2010–2011. Their characteristics are summarized in Table 2 and locations shown in Fig. 1E. All specimens were classified *a priori* as *Cetraria aculeata* s. lat. A tuft c. 5–10 cm<sup>2</sup> was considered to be one specimen ('thallus'). To analyze the chemical variability among neighbouring thalli (which are probably vegetative clones), we sampled thalli at 0.5 m intervals along transects 10–25 m in length and 0.5 m in width within each population. As a result, 5–41 specimens per population were collected according to their availability, amounting to 256 thalli in total. To study the morphological variability within the local populations, we selected a subset of 122 specimens, sampled at distances of >1 m along the same transects, resulting in 5–18 specimens per population (Table 2). Because the species reproduces only asexually in Ukraine (no specimens with fruit bodies have been found so far in this area), we selected 1 m as the minimum distance to reduce the clones in our dataset because the range of dispersal through thallus fragments is considered to be less than 1 m (Heinken 1999).

### Morphological study

A subset of 122 air-dried specimens, collected from the 13 local populations mentioned above, was selected for morphometric measurements (Tables 2 & 3). For each specimen, we measured the size of the whole thallus,




FIG. 1. Morphological diversity and distribution of *Cetraria aculeata* s. lat. A, Crimea, Dzhankoj region, norstictic acid minor; B, Luhansk region, norstictic acid minor; C, type specimen (LE 220), Kherson region, norstictic acid major; D, Donetsk region, norstictic acid major; E, distribution in Ukraine, based on the analysis of 247 specimens (KW, KHER, LE) tested by TLC, numbers correspond to the sampling sites of the populations given in Table 2. Arrows indicate specimens selected for HPLC analysis (listed in Table 4); red circles, norstictic acid present; blue triangles, norstictic acid absent. Scales: A–D = 1 cm.

TABLE 2. Populations of *Cetraria aculeata* s. lat. examined with characteristics of their sites (natural zones of Ukraine: F – polissia (forest), FS – forest-steppe, S – steppe, SSA – steppe semi-arid, SA – steppe arid, CM – Crimean Mountains; Habitats: O – open places, clearings among pine forest plantation, G – grasslands and heaths, Y – yayla or mountain top, deforested by grazing).

Site No.	Population	Altitude, m a.s.l.	Natural zone of Ukraine	Habitat	Substratum	Number of specimens selected for morphometric study (n = 122)	Total number of specimens analyzed (including TLC, n = 256)
1	Shatsk*	170	F	O	Sand	11	41
2	Ratno	145	F	O	Sand	5	12
3	Platonivka	90	FS	O	Sand	9	16
4	Trehizbenka**	107	S	G	Sand	10	25
5	Stanychno-Luhansk**	55	S	O	Sand	10	20
6	Provallia**	170	S	O	Soil	10	21
7	Kamjany Mohyly†	160	S	O	Granite	14	22
8	Anratsit	140	S	O	Soil	6	15
9	Burkuty	15	SA	G	Sand	15	5
10	Kazantyp‡	10	SSA	G	Limestone	11	23
11	Bondarenkovo	55	SSA	G	Soil	7	14
12	Arabat Split¶	0	SA	G	Sand	6	11
13	Chatyr Dag	1340	CM	Y	Limestone	18	21

\* Shatsk Lakes National Nature Park; \*\* Branch of Luhansk Steppe Nature Reserve; † Branch of Ukrainian Nature Steppe Reserve; ‡ Kazantyp Nature Reserve; ¶ Arabat Split Botanical Reserve.

the width of the main branches at the basal bifurcations, and the width of terminal branches below the last bifurcation (Table 3). Five measurements of each trait per specimen were performed. The morphometric measurements are presented as (min–) mean  $\pm$  Standard Deviation (–max). The presence or absence of isidioid projections was recorded for each specimen (Table 3).

### Chemical analyses

Each of the 247 herbarium specimens was tested by spot reactions with 10% aqueous solution of potassium hydroxide (K) and dissolved crystals of paraphenylenediamine in ethanol (Pd). The lichen substances were also analyzed using the microcrystal test method (Orange *et al.* 2001) for a subset of 17 selected specimens (Table 4).

Each of the 503 specimens studied were also analyzed by thin-layer chromatography (TLC), according to standard protocols (White & James 1985; Orange *et al.* 2001) using *Sorbifil* UV plates (Russia) and solvent A.

High performance liquid chromatography (HPLC) was applied to a subset of 17 specimens, which were selected according to the following criteria: 1) specimens originating from different parts of Ukraine (the Carpathians, north-west, south-east and south Ukraine, including Crimea); 2) individuals originating from localities where, according to the TLC analysis, specimens

with and without norstictic acid were present. Localities for specimens analyzed by HPLC are indicated in Fig. 1E with arrows and listed in Table 4. HPLC analyses were carried out on the basis of methods described in Yoshimura *et al.* (1994), Gupta *et al.* (2007) and Manojlović *et al.* (2010). The content of the lichen substances was quantified according to a 10-point scale using peak heights at 224 nm and 1000 mAU detector signal range for each specimen. A peak height between 500–1000 mAU was considered as “major” (5–10 points), 300–500 mAU as “medium” (3–5 points), 30–300 mAU as “minor” (1–3 points) and less than 30 mAU as “traces” (1 point).

### Statistical analyses

Basic statistics for the morphometric parameters of the *C. aculeata* s. lat. specimens collected from all 13 populations, an analysis of variance (ANOVA) and the principal component analysis (PCA) were run in JMP 9 (SAS Institute Inc., Cary, North Carolina). The coefficient of variation (CV) was calculated for the entire dataset and each population (Table 3). Values of CV were classified in five categories: minor variation (0–10%), little variation (11–20%), average variation (21–40%), high variation (41–60%) and very high variation (61–100%). To

TABLE 3. *Morphometric (n = 122) and chemical (n = 256) characteristics of the populations of C. aculeata s. lat. studied.*

Site No.	Population	Width of the main branch (mm)		Width of the terminal branch (mm)		Thallus size (mm)		Thalli with isidioid projections, %	Thalli with norstictic acid, %
		Mean $\pm$ SD	CV, %	Mean $\pm$ SD	CV, %	Mean $\pm$ SD	CV, %		
1	Shatsk	1.29 $\pm$ 0.16	12.45	0.47 $\pm$ 0.08	16.00	3.27 $\pm$ 1.26	38.42	36	0
2	Ratno	1.51 $\pm$ 0.28	18.19	0.40 $\pm$ 0.10	24.85	6.43 $\pm$ 2.00	31.05	60	0
3	Platonivka	1.44 $\pm$ 0.36	24.63	0.48 $\pm$ 0.06	12.92	3.57 $\pm$ 1.06	29.64	100	100
4	Trehizbenka	1.15 $\pm$ 0.23	20.37	0.46 $\pm$ 0.06	13.41	3.20 $\pm$ 0.74	23.23	30	100
5	Stanychno-Luhansk	1.20 $\pm$ 0.35	29.40	0.50 $\pm$ 0.10	20.23	3.21 $\pm$ 0.38	11.96	20	90
6	Provallia	1.57 $\pm$ 0.17	10.81	0.50 $\pm$ 0.03	6.82	2.26 $\pm$ 0.60	26.50	10	80
7	Kamjany Mohyly	1.62 $\pm$ 0.34	20.84	0.50 $\pm$ 0.03	6.08	4.33 $\pm$ 1.19	27.43	43	64
8	Anratsit	1.56 $\pm$ 0.16	10.36	0.50 $\pm$ 0.05	10.00	2.92 $\pm$ 0.26	8.85	100	17
9	Burkuty	1.25 $\pm$ 0.26	21.13	0.44 $\pm$ 0.08	18.13	4.47 $\pm$ 1.21	27.00	40	100
10	Kazantyp	1.32 $\pm$ 0.23	17.76	0.53 $\pm$ 0.06	11.93	4.19 $\pm$ 1.38	32.92	64	73
11	Bondarenkovo	1.41 $\pm$ 0.37	26.34	0.50 $\pm$ 0.02	4.89	3.40 $\pm$ 0.92	27.04	29	100
12	Arabat Split	1.75 $\pm$ 0.25	14.42	0.55 $\pm$ 0.06	10.64	3.05 $\pm$ 0.45	14.77	0	100
13	Chatyr Dag	1.28 $\pm$ 0.30	23.56	0.47 $\pm$ 0.06	12.15	4.45 $\pm$ 0.90	20.18	100	22
Total in the dataset		1.39 $\pm$ 0.32	22.74	0.49 $\pm$ 0.07	13.81	3.73 $\pm$ 1.31	34.99	52	62

TABLE 4. HPLC study of selected specimens of *C. aculeata* s. lat. from Ukraine

Herbarium No.	Locality, collector, year	TLC data*	HPLC data**
LE220 (Type)	Kherson reg., Chaplynsky distr. <i>Oxner</i> , 1923	NST, L	NST (8), L (2), PL (1)
KW5770	Kherson reg., Hola Pristan distr. <i>Makarevych</i> , 1950	NST, L, PL	NST (4), L (4), PL (2)
KW65049	Kherson reg., Bilozerka distr. <i>Nadyeina</i> , 2010	NST, L	NST (3), L (5), PL (1)
KW5824	Crimea, Dzhankoi reg. <i>Osmanova</i> , 1932	NST, L, PL	NST (9), L (6), PL (2)
KW66928	Luhansk reg., Sverdlovsk distr. <i>Vasylyuk</i> , 2010	L, PL	NST (2), L (3), PL (1)
KW66930	Luhansk reg., Sverdlovsk distr. <i>Vasylyuk</i> , 2010	NST, L	NST (2), L (8)
KW64472	Luhansk reg., Sverdlovsk distr. <i>Nadyeina</i> , 2005	L	NST (1), L (6)
KW64477	Luhansk reg., Sverdlovsk distr. <i>Nadyeina</i> , 2005	NST, L	NST (2–3), L (3), PL (1)
KW5855	Donetsk reg., Volodarsk distr. <i>Oxner &amp; Kopachevska</i> , 1954	L, PL	NST (1), L (5), PL (1)
KW5856	Donetsk reg., Volodarsk distr. <i>Oxner &amp; Kopachevska</i> , 1954	NST, L	NST (1), L (4), PL (1)
KW64475	Donetsk reg., Shakhtarsk distr. <i>Nadyeina</i> , 2006	L, PL	NST (2), L (6)
KW64476	Donetsk reg., Shakhtarsk distr. <i>Nadyeina</i> , 2006	NST	NST (9)
KW33190	Kyrovograd reg., Holovanivka distr. <i>Kondratyuk</i> , 1983	NST, L	NST (10)
KW27645	Mykolaiv reg., Bratsk distr. <i>Makarevych</i> , 1950	NST, L	NST (6), L (6), PL (1)
KW33618	Dnipropetrovsk reg. and distr. <i>Hajova</i> , 1960	NST, L, PL	NST (2), L (5), PL (1)
KW33628	Volyn reg., Kamin-Kashirsk distr. <i>Bradis</i> , 1940	L, PL	NST (1), L (2), PL (1)
KW62763	Ivano-Frankivsk reg., Verkhovina distr. <i>Blum</i> , 1978	L, PL	NST (1), L (2), PL (1)

\*NST – norstictic acid, L – lichesterinic acid, PL – protolichesterinic acid; \*\* content of substances indicated in brackets are: 1 – traces, 2–3 – minor, 4–5 – medium, 6–10 – major.

examine the effects of environmental, geographical, chemical and morphological variables on the quantitative morphometric thallus parameters (such as width of the main and terminal branches, and size of the thalli), we performed an analysis of variance (ANOVA). Nominal variables included in ANOVA were: occurrence in one of the 13 populations and five natural zones of Ukraine, in three specific habitats, at four substrata and three classes of altitude above sea level (Table 2), the presence of norstictic acid detected by TLC and presence of isidioid projections (Table 3). Only values with high support are shown in Table 5 and discussed. The variables included in the PCA were: width of the main branch, width of the terminal branch and size of the thalli. The resulting PCA biplots were selected according to the maximum variance of projection along each component (Fig. 3).

To analyze the pairwise correlation between the presence and absence of norstictic acid and the geographical distance between *C. aculeata* specimens, we performed a Mantel test with 99 permutations, as implemented in Genalex (Peakall & Smouse 2006; Fig. 4). In total, 256 specimens of the *C. aculeata* s. lat. analyzed by TLC (Table 2) were included in this study.

The distribution map for the *C. aculeata* s. lat. specimens studied was drawn in ArcGis.10 (<http://www.esri.com/software/arcgis>). The natural zonation of Ukraine

follows Prydatko (1998; <http://biomodel.info/training-package/ukraine-nature-agricultural-zoning>), and is shown in Fig. 1E.

## Results

### Morphology

We found a strong correlation between the size of thalli and the presence of isidioid projections (Fig. 2, Table 5). Thalli without isidioid projections were smaller than thalli with this trait. However, we found no clear regional differentiation in the frequency of isidioid projections. The percentage of specimens with isidioid projections varied greatly among populations (0–100% of specimens per population, Table 3). While every specimen investigated in the populations of Antratsit and Chatyr Dag had isidioid projections, none of the specimens from Arabat Split Botanical Reserve had this characteristic (Table 3).




FIG. 2. Relationships among the 122 specimens of *Cetraria aculeata* group between thallus size and presence or absence of isidioid projections (A,  $t = -3.751$ ,  $P = 0.0003$ ,  $df = 120$ ), presence or absence of norstictic acid (B,  $t = 1.996$ ,  $P = 0.048$ ,  $df = 74$ ) and substratum groups (C,  $t = 2.222$ ,  $P = 0.029$ ,  $df = 76$ ), and between width of terminal branches and presence or absence of norstictic acid (D,  $t = -2.146$ ,  $P = 0.033$ ,  $df = 120$ ).

Our systematically sampled dataset revealed rather small differences in morphometric traits, that are essential for species identification, such as the width of the main and terminal branches or the size of the thalli (Table 3). The CV of these characteristics showed little or only average variation in the entire dataset, as well as within each population. How-

ever, an ANOVA applied to the morphometric variables of *C. aculeata* s. lat. in relation to regional effects revealed significant differences between populations (Table 5). Continuous variation within the three morphometric characteristics analyzed was also confirmed by PCA (Fig. 3). The first two components (axes) of PCA explained 77.8%



TABLE 5. Results of an analysis of variance testing the effect of environmental, geographical and morphological variables on the morphometric parameters of *Cetraria aculeata* s. lat. individuals representing different parts of Ukraine (only significant values are shown, n = 122).

Morphometric parameters	Source of variation	Sum of Squares	df	F	P
Width of main branch	Population	1.22	4	4.00	0.0046
	Substratum	0.47	1	6.22	0.0142
Width of terminal branch	Population	0.06	4	3.54	0.0094
Size of the thalli	Population	41.31	4	10.21	<0.0001
	Isidia	6.16	1	6.10	0.0152

of variance (the two axes explained 43.4% and 34.4% of the variance).

### Chemistry

The analysis of the specimens by TLC showed that the presence of norstictic acid among the populations varied considerably (Table 3). Specimens distributed in the Forest zone of Ukraine and in the Carpathians always lacked norstictic acid (Table 3, Fig. 1E). Additionally, we detected licheterinic and/or protolicheterinic acids in the medulla of some of the specimens, as have others in the literature (Kärnefelt 1986; Vainshtein *et al.* 1990). In contrast to the northern populations of *C. aculeata* s. lat., some of the specimens from the southern populations in the Steppe zone and the Crimean Mountains contained norstictic acid (17–100% of specimens per population; Table 3).

HPLC revealed a very broad range of norstictic acid content in each specimen (Table 4). The specimens collected from the same localities contained very different quantities, and some had only traces which could not be detected by TLC. Surprisingly, traces of norstictic acid were detected by HPLC even in specimens from northern Ukraine and the Carpathians, where no norstictic acid was detected by TLC (Table 4). With HPLC we also detected traces of fumarprotocetraric and connorstictic acids in several specimens, which have never been reported before for *C. aculeata* s. lat. Rangiformic acid was mentioned once for *C. aculeata* from New Zealand (Galloway 1985), and occasionally

minor amounts of nephrosterinic and isonephrosterinic acids have been detected by HPLC (Thell & Kärnefelt 2011).

The analyzed specimens of *C. aculeata* s. lat. showed no reactions with spot and microcrystal tests. As a control, we tested spot reactions and conducted microcrystal tests with K and Pd to detect norstictic acid in other lichen species: *Aspicilia cinerea* (L.) Körb., *Cladonia symphyocarpia* (Flörke) Fr. and *Pleurosticta acetabulum* (Neck.) Elix & Lumbsch. These species developed red spots with K or Pd and displayed typical reddish crystals in the microcrystal tests. Therefore, we conclude that, in the case of *Cetraria aculeata* s. lat., spot reactions and the microcrystal test cannot be used to detect norstictic acid.

### Relationship between morphology, chemistry, geographical and environmental factors

We found a significant correlation between thallus size and the presence of norstictic acid, as detected by TLC (Fig. 2). Samples containing norstictic acid were slightly smaller than samples where norstictic acid could not be detected by TLC. At the same time, specimens lacking norstictic acid had narrower terminal branches than specimens with norstictic acid (Fig. 2). Specimens growing on bare soil and sand had smaller thalli than specimens collected on granite or limestone outcrops (Fig. 2).

All three morphometric thallus variables differed between the populations of *C. aculeata* s. lat. The width of the main thallus branch also depended on the substratum,




FIG. 3. PCA results based on the morphometric characteristics of the *C. aculeata* s. lat. in Ukraine,  $n = 122$ . A, specimens differing chemically are shown; B, specimens with and without isidioid projections are marked; C, specimens preferring different substrata are distinguished; D, specimens belonging to different populations are indicated.

and the presence of isidioid projections was correlated with thallus size (Table 5). No correlations were found between morphometric features and the five natural zones of Ukraine, the three specific habitats, or the three classes of altitude above sea level (Table 5).

A continuous scatter plot from a PCA of the morphometric features of the *C. aculeata* s. lat. specimens (Fig. 3) had no structure when compared to the presence of isidioid projections, norstictic acid content, substrata preference and sampling in different populations.

We found a weak positive correlation between the presence of norstictic acid and a large geographical distance up to 800 km.

The overall linear fit of the Mantel test for geographical distance and chemical identity was  $7.015^{-7}$  ( $P < 0.001$ , Fig. 4B), but only  $5.385^{-4}$  ( $P = 0.6$ , Fig. 4A) for short distances up to 50 m.

## Discussion

### Taxonomic status

Our analyses of the morphological, chemical, ecological and geographical characteristics of *C. aculeata* s. lat. specimens, originating from 13 populations (Tables 2 & 3), showed a continuous variation in morphometric features, as well as in the presence of isidioid projections and norstictic acid. Differences




FIG. 4. Linear fit of Mantel test of geographical distance and chemical identity (absence or presence of norstictic acid) among pairs of specimens,  $n = 256$ . A, small distances up to 50 m, parameters estimated with a linear fit of  $5.385^{-4}$ ,  $P < 0.6$ ; B, large distances up to 800 km, parameters estimated with a linear fit of  $7.015^{-7}$ ,  $P < 0.001$ .

between populations in the morphometric traits examined (Tables 3 & 5) explained local infraspecific variability and did not support subdivisions among the specimens of *Cetraria aculeata* s. lat. we studied (Fig. 3D).

Seven out of 13 populations were homogeneous in norstictic acid content when analyzed by TLC, while the other six populations contained chemically distinct individuals (Table 3). We found a general trend for specimens lacking norstictic acid to have larger thalli and narrower terminal branches (Fig. 2B & D). Therefore the presence of norstictic acid depends on the size of the thallus, which reflects the ontogenetic stage, and thus should not be considered for taxonomic purposes. An analogous trend has been found for perlatolic and usnic acids in another terricolous lichen species, *Cladonia stellaris* (Opiz) Pouzar & Vězda (Fahsel 1984). We hypothesize that norstictic acid biosynthesis is active during the early stages of thallus development, appearing shortly after thallus fragmentation, and declines or becomes deactivated later, once the thalli have reached a certain size. We may also speculate that norstictic acid plays an important role in the complexation with heavy metals occurring in the substratum, which would otherwise be toxic for the small thalli. The ability of lichen thalli to incorporate

heavy metal ions into oxalates is well-known, and seems to be a way of coping with the effects of toxic elements (Jones & Wilson 1985; Purvis 1996). Norstictic acid is also known to form complexes with other toxic metals, including copper (Purvis *et al.* 1987).

We found no correlation between the presence of norstictic acid and any of the environmental factors analyzed (Table 5), even though we expected specimens with norstictic acid (revealed by TLC) to prefer geographically different regions (according to Kärnefelt 1986) or to occur mainly on carbonaceous substrata (according to Hauck *et al.* 2010). The distribution pattern of norstictic acid in *C. aculeata* s. lat. did not correlate with the natural zones of Ukraine, habitats, substrata or different altitudinal levels. However, specimens with norstictic acid (TLC) were limited to southern regions in Ukraine (the forest-steppe, steppe zones, and the Crimean Mountains), where they co-occur with specimens lacking norstictic acid. In contrast, specimens without norstictic acid (TLC) grow throughout all of Ukraine (Table 3, Fig. 1E). The sporadic presence of specimens with norstictic acid in southern, but not northern, areas supports the hypothesis that norstictic acid plays an adaptive role in lichens growing in an arid climate (Hauck *et al.* 2010). A similar trend has been identified

in the terricolous lichen *Flavocetraria nivalis* (L.) Kärnefelt & A. Thell, which has higher concentrations of usnic acid in colder environments (Bjerke *et al.* 2004), and for the epiphytic *Pseudevernia furfuracea* (L.) Zopf, which contains olivetoric or physodic acids, depending on climate (Martellos 2003). These support Hale's theory (Hale 1956) that the synthesis of lichen secondary compounds is governed by climatic gradients. Nevertheless, several other studies have found no evidence for a climatic dependency of lichens secondary chemistry (Halvorsen & Bendiksen 1982; Fahselt 1984).

Mantel tests revealed a significant spatial autocorrelation between the norstictic acid content (as detected by TLC) and the geographical distance between the individuals sampled (Fig. 4B). This reflects regional differences in norstictic acid content (as measured by HPLC) with only traces of norstictic acid (not detectable by TLC) in northern Ukraine and partly higher concentrations in the southern regions. However, we found no autocorrelation at local scales (Fig. 4A). This indicates that local habitat characteristics do not strongly influence the presence or absence of norstictic acid in *C. aculeata* s. lat. Pérez-Ortega *et al.* (2012) studied extreme cases of phenotypic plasticity in *C. aculeata* s. lat., including both distinctly vagrant specimens with thick thalli and 'typical' *C. aculeata* from the same habitats in Spain. These thick vagrant specimens did not contain norstictic acid and thus, according to Kärnefelt (1986), were classified as *C. aculeata* s. lat. (Pérez-Ortega *et al.* 2012). The study of Pérez-Ortega *et al.* (2012) revealed anatomical and physiological differences between the thick vagrant and 'typical' individuals of *C. aculeata* s. lat. from the same habitats. The authors found, however, shared haplotypes between those morphs and thus emphasized that the vagrant morphs are adaptations to aridity. We consider this as further evidence for the morpho/chemical variability within populations of *C. aculeata* s. lat. being due to ecological (microclimate) and not taxonomic reasons.

The recent worldwide multigene studies of *Cetraria aculeata* s. lat. populations (Fernández-

Mendoza *et al.* 2011; Fernández-Mendoza & Printzen 2013; Printzen *et al.* 2013)<sup>1</sup> revealed pleistocene diversification within the *C. aculeata* group into several genetic clusters. The only small population from Ukraine (Crimean Mountains) of five individuals with norstictic acid was included in the last two studies, and it appeared to be related to the Mediterranean phylogenetic clade. However, more extensive sampling of *C. aculeata* s. lat. from Ukraine, based on a subset of the specimens used in our study, revealed shared haplotypes with other genetic clusters, but no relationship between the genetic structures of populations with norstictic acid production (Lutsak *et al.* 2012).

Further evidence for the incomplete speciation within *C. aculeata* s. lat. comes from the gene flow between different phylogenetic clusters within *C. aculeata* (Fernández-Mendoza & Printzen 2013), and shared haplotypes among different operational taxonomic units, such as *C. steppae*, *C. crespoae* and *C. aculeata* s. str. represented from geographically distinct areas (Printzen *et al.* 2013). The speciation seems to be very slow for taxa with vegetative reproduction only, where natural selection is determined by random mutations but not recombinations. *Cetraria aculeata* s. lat. includes several genetically distinct clades related to the phylogeographical history of the species (Fernández-Mendoza & Printzen 2013; Printzen *et al.* 2013), but these do not correspond to secondary chemistry or morphological characteristics. Higher levels of norstictic acid production in *C. aculeata* s. lat. in southern Ukraine may be connected to the aridity of glacial permafrost (Hewitt 1999). Acid production seems to be selectively activated depending on the microclimate and the developmental stage of thalli.

Possible reasons for the chemical variation in *C. aculeata* s. lat. populations may be the large-scale geographical and climatic dependence of gene expression, photobiont switching and microbial inhabitants (Muggia *et al.*

<sup>1</sup>All representatives of *C. aculeata* group were pooled together and treated as "*C. aculeata* s. lat." in Printzen *et al.* (2013), but not in other studies.

2009; Bates *et al.* 2011; Fernández-Mendoza *et al.* 2011; Hodkinson *et al.* 2012). In a recent study, Printzen *et al.* (2012) demonstrated that the composition of alphaproteobacterial communities in geographically distant populations of *C. aculeata* is affected by environmental factors, but no relationship between these communities and secondary metabolite production has so far been found.

Our results support a decision to merge *C. steppae* Savicz with *C. aculeata* (Schreb.) Fr. and to treat both names as synonyms. The features currently used for the delimitation at species level are not appropriate because the differences in the morphological features are too small, the presence and quantity of norstictic acid varies greatly, and the substrata and habitats occupied by lichens are very diverse. *Cetraria steppae* Savicz may be regarded as a chemotype only, reflecting an infraspecific variation, even though norstictic acid has been detected in the type material of *C. steppae* by both TLC and HPLC (Table 4). It is commonly accepted that variation in the concentrations of particular secondary lichen substances does not provide sufficient grounds for taxonomic differentiation (Hawksworth 1976; Elix & Stocker-Wörgötter 2008). As a consequence, we recommend formally synonymizing *C. steppae* Savicz with *C. aculeata* (Schreb.) Fr.:

### ***Cetraria aculeata* (Schreb.) Fr.**

*Nov. Sched. Critic. Lich.* 4: 32 (1826).—*Lichen aculeatus* Schreb., *Spicil. Flor. Lipsiens.* 125 (1771).

*Cetraria steppae* (Savicz) Kärnefelt, *Bryologist* 96: 400 (1993).—*Coelocaulon steppae* (Savicz) Barreno & Vázquez, *Lazaroa* 3: 329 (1982).—*Cornicularia steppae* Savicz, *Notul. Syst. Inst. Cryptog. Horti bot. Petropol.* 3: 187 (1924); type: Ukraine, Askania-Nova, Kherson reg., 1924, Oxner (LE, lectotype!).

*Coelocaulon stepposa* (Mereschk.) S. Y. Kondr., *Flora Lyshajnykiv Ukraini* 2(2): 296 (1993).—*Cetraria tenuissima* f. *stepposa* Mereschk., *Annals and Magaz. Nat. History* 9(8): 249 (1921); type: Russia meridio-orientalis, Tauria, *Mereschkowsky* (KAZ).

### **Conservation status: Ukrainian case**

We suggest including *Cetraria aculeata* s. lat. in the next edition of the Red Data Book of Ukraine, which will replace the entry of *C.*

*steppae* (Blum 1996; Blum *et al.* 2009). Including *C. aculeata* s. lat. in the Red Data Book will highlight the importance of conserving sand dunes in the Forest natural zone of Ukraine, as well as rock outcrops in the Crimean and Carpathian Mountains, where the species is even rarer than in the steppes. *Cetraria aculeata* already has an approved national conservation status in neighbouring areas (Lőkös & Tóth 1996; Moutchnik & Zavarzin 2005; Liška *et al.* 2008). Moreover, the other member of *C. aculeata* s. lat., which is difficult to designate in the field, is *C. muricata* (Printzen *et al.* 2013), a species that is also of conservation concern in neighbouring countries (Golubkov & Kobzar 2005; Liška *et al.* 2008).

To assess the national threat category of *C. aculeata* s. lat., we took into consideration that the area of occupancy by this taxon in Ukraine has been constant during recent decades. Just two localities have disappeared due to habitat destruction in the Odessa and Kharkiv regions during the last 100 years (according to herbaria data from KW and KHER), which corresponds to 2.4% of the total number of localities known for the taxon. However, the species inhabit vulnerable biotopes and recovery after disturbance is expected to be very slow because of vagrancy and the species' spatially limited vegetative reproduction (Heinken 1999).

We assessed the total national area of occupancy of *C. aculeata* s. lat. as nearly 0.136 km<sup>2</sup> by adding: 1) the area of occupancy of 13 local populations (each covered very small patches of up to 1000 m<sup>2</sup>) equal to 0.013 km<sup>2</sup>, and 2) the area of occupancy of the other 123 localities known in Ukraine, as documented in herbaria (KW, KHER), which is equal to at least 0.123 km<sup>2</sup>. Thus, the total national area of occupancy of *C. aculeata* s. lat. is considerably less than the 20 km<sup>2</sup> recommended for the 'vulnerable' threat category (VU) according to criterion D2 of IUCN (IUCN Standards and Petitions Subcommittee 2011). Finally, the threat category of *C. aculeata* s. lat. in Ukraine is assessed as vulnerable (VU). This corresponds with the status of *C. steppae* in the current edition of the Red Data Book of Ukraine

(Blum *et al.* 2009) and Russia (Golubkova *et al.* 2008).

We thank Prof. O.Ye. Khodosovtsev (Kherson) for valuable advice and for sending herbarium materials (KHER), as well as Prof. S.Ya. Kondratyuk (Kyiv), Prof. C. Printzen (Frankfurt on Main) and three anonymous referees for their useful comments. We are grateful to Prof. M. P. Andreev (Saint-Petersburg) for sending the type specimen (LE220); Prof. V.P. Heluta and O.V. Vasylyuk (Kyiv) for sampling the Ratno and Antratsit populations; Dr. F. Fernández-Mendoza (Frankfurt on Main) for help with the sampling design for population study; Dr. L. Lőkös (Budapest), Dr. A. Senkandeshler (Izmir), Dr. A. Tsurikov (Hornsea), and Dr. A. Yatsina (Minsk) for providing several references; and Dr. S. Dingwall (Birmensdorf) for revising the English of the manuscript. We thank the Rufford SGF for partial funding of this study, as well as the Swiss State Secretariat for Education and Research (SER).

#### REFERENCES

- Bates, S. T., Cropsey, G. W. G., Caporaso, G. J., Knight, R. & Fierer, N. (2011) Bacterial communities associated with the lichen symbiosis. *Applied and Environmental Microbiology* **77**: 1309–1314.
- Bjerke, J. W., Joly, D., Nilsen, L. & Brossard, T. (2004) Spatial trends in usnic acid concentrations of the lichen *Flavocetraria nivalis* along local climatic gradients in the Arctic (Kongsfjorden, Svalbard). *Polar Biology* **27**: 409–417.
- Blum, O. B. (1996) Lichen species have to be protected in Ukrainian flora. In *Red Book of Ukraine: Plants* (Y. R. Sheliag-Sosonko, ed.): 503–530. Kyiv: Ukrayinska Entsiklopedia [In Ukrainian].
- Blum, O. B., Kondratyuk, S. Y. & Khodosovtsev, O. Y. (2009) Lichens. In *Red Book of Ukraine: Plants* (Y. P. Didukh, ed.): 728–780. Kyiv: Globalconsulting [In Ukrainian].
- Domaschke, S., Fernández-Mendoza, F., García, M. A., Martín, M. P. & Printzen, C. (2012) Low genetic diversity in Antarctic populations of the lichen-forming ascomycete *Cetraria aculeata* and its photobiont. *Polar Research* **31**: 1–13.
- Elix, J. A. & Stocker-Wörgötter, E. (2008) Biochemistry and secondary metabolites. In *Lichen Biology. 2nd Ed* (T. H. Nash III, ed.): 104–133. Cambridge: Cambridge University Press.
- Fahselt, D. (1984) Interthalline variability in levels of lichen products within stands of *Cladonia stellaris*. *Bryologist* **87**: 50–56.
- Fahselt, D. (2008) Individuals, populations and population ecology. In *Lichen Biology. 2nd Ed* (T. H. Nash III, ed.): 252–273. Cambridge: Cambridge University Press.
- Fernández-Mendoza, F. & Printzen, C. (2013) Pleistocene expansion of the bipolar lichen *Cetraria aculeata* into the Southern hemisphere. *Molecular Ecology* **22**: 1961–1983.
- Fernández-Mendoza, F., Domaschke, S., García, M. A., Jordan, P., Martín, M. P. & Printzen, C. (2011) Population structure of mycobionts and photobionts of the widespread lichen *Cetraria aculeata*. *Molecular Ecology* **20**: 1208–1232.
- Galloway, D. J. (1985) *Flora of New Zealand. Lichens*. Wellington, New Zealand: Hasselberg.
- Golubkov, V. V. & Kobzar, N. N. (2005) Lichens. In *Red Data Book of the Republic of Belarus: Plants* (L. I. Khoruzhik, ed.): 355–383. Minsk: Belarusskaia Entsiklopedia imeni Petrusia Brovka [In Russian].
- Golubkova, N. S., Dombrovskaia, A. V., Istomina, N. B., Makarova, I. I., Makryi, T. V., Muchnik, E. Ye., Pchelkin, A. V., Sedelnikova, N. V., Skirina, I. F., Tolpisheva, T. U. *et al.* (2008) Lichens. In *Red Book of the Russian Federation: Plants and Fungi* (Y. P. Trutnev, ed.): 703–752. Moskva: Tovari-shestvo nauchnyh izdanij KMK [In Russian].
- Gupta, V. K., Darokar, M. P., Shanker, K., Negi, A., Shrivastava, S. K., Gupta, M. M. & Khanuja, S. P. S. (2007) Rapid and sensitive HPLC method for the determination of polyphenols in various lichen species of Himalayan origin. *Journal of Liquid Chromatography and Related Technologies* **30**: 97–111.
- Hale, M. E. (1956) Chemical strains of the lichen *Parmelia furfuracea*. *American Journal of Botany* **43**: 456–459.
- Halvorsen, R. & Bendiksen, E. (1982) The chemical variation of *Pseudevernia furfuracea* in Norway. *Nordic Journal of Botany* **2**: 371–380.
- Hauck, M., Jürgens, S.-R. & Leuschner, C. (2010) Norstictic acid: correlations between its physico-chemical characteristics and ecological preferences of lichens producing this depsidone. *Environmental and Experimental Botany* **68**: 309–313.
- Hawksworth, D. L. (1976) Lichen chemotaxonomy. In *Lichenology: Progress and Problems* (D. H. Brown, D. L. Hawksworth & R. H. Bailey, eds): 139–184. London: Academic Press.
- Heinken, T. (1999) Dispersal patterns of terricolous lichens by thallus fragments. *Lichenologist* **31**: 603–612.
- Hewitt, G. M. (1999) Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society* **68**: 87–112.
- Hodkinson, B. P., Gottel, N. R., Schadt, C. W. & Lutzoni, F. (2012) Photoautotrophic symbiont and geography are major factors affecting highly structured and diverse bacterial communities in the lichen microbiome. *Environmental Microbiology* **14**: 147–161.
- IUCN Standards and Petitions Subcommittee (2011) *Guidelines for Using the IUCN Red List Categories and Criteria. Version 9.0*. Prepared by the Standards and Petitions Subcommittee. Online access: <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- John, V. (1999) Lichenes Anatolici Exsiccati. Fasc. 1–3 (no. 1–75). *Arnoldia* **16**: 1–44.
- John, V. & Breuss, O. (2004) Flechten der Östlichen Schwarzmeer-Region in der Türkei. *Herzogia* **17**: 137–155.

- Jones, D. & Wilson, M. J. (1985) Chemical activity of lichens on mineral surfaces – a review. *International Biodeterioration* **21**: 99–104.
- Kärnefelt, I. (1986) The genera *Bryocaulon*, *Coelocaulon* and *Cornicularia* and formerly associated taxa. *Opera Botanica* **86**: 1–90.
- Kärnefelt, I., Mattsson, J.-E. & Thell, A. (1992) Evolution and phylogeny of cetrarioid lichens. *Plant Systematics and Evolution* **183**: 113–160.
- Kärnefelt, I., Mattsson, J.-E. & Thell, A. (1993) The lichen genera *Arctocetraria*, *Cetraria*, and *Cetrariella* (*Parmeliaceae*) and their presumed evolutionary affinities. *Bryologist* **96**: 394–404.
- Liška, J., Palice, Z. & Slavíková, Š. (2008) Checklist and Red List of lichens of the Czech Republic. *Preslia* **80**: 151–182.
- Llimona, X. & Hladun, N. L. (2001) Checklist of the lichens and lichenicolous fungi of the Iberian Peninsula and Balearic Islands. *Bocconea* **14**: 1–581.
- Lökös, L. & Tóth, E. (1996) Red list of lichens of Hungary (a proposal). In *Proceedings of the “Research, Conservation, Management” Conference* (E. Tóth & R. Horváth, eds): 337–343. Aggtelek, Hungary: Aggtelek National Park Directorate.
- Lutsak, T., Fernández-Mendoza, F., Nadyeina, O., Pérez-Ortega, S. & Printzen, C. (2012) *Cetraria steppae*: a lichen species of central Asian steppes, or part of a wider Mediterranean clade of *C. aculeata*? In *Abstracts of the 21st International Symposium “Biodiversity and Evolutionary Biology” of the German Botanical Society (DBG), 16–19 September, 2012, Mainz, Germany*, p. 126.
- Manojlović, N. T., Vasiljević, P., Jusković, M., Najman, S., Janković, S. & Milenković-Andjelković, A. (2010) HPLC analysis and cytotoxic potential of extracts from the lichen, *Thamnolia vermicularis* var. *subuliformis*. *Journal of Medicinal Plants Research* **4**: 817–823.
- Martellos, S. (2003) The distribution of the two chemical varieties of the lichen *Pseudevernia furfuracea* in Italy. *Plant Biosystems* **137**: 29–34.
- Mereschkowsky, C. (1921) Diagnoses of some lichens. *Annals and Magazine of Natural History* **9**: 249.
- Moutchnik, E. & Zavarzin, A. (2005) Lichens in the Red Data Books in Russia. *Folia Cryptogamica Estonica* **41**: 59–80.
- Muggia, L., Schmitt, I. & Grube, M. (2009) Lichens as treasure chests of natural products. *Sim News* **59**: 85–97.
- Orange, A., James, P. W. & White, F. J. (2001) *Microchemical Methods for the Identification of Lichens*. London: British Lichen Society.
- Oxner, A. M. (1937) *Handbook of the Lichens of Ukrainian Republic*. Kyiv: Vydavnytstvo AN URSR [In Ukrainian].
- Oxner, A. M. (ed.) (1993) *Flora of the Lichens of Ukraine. Vol. 2*. Kyiv: Naukova Dumka [In Ukrainian].
- Peakall, R. & Smouse, P. E. (2006) GenAlEx 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* **6**: 288–295.
- Pérez-Ortega, S., Fernández-Mendoza, F., Raggio, J., Vivas, M., Ascaso, C., Sancho, L. G., Printzen, C. & de los Ríos, A. (2012) Phenotypic plasticity in lichens: insights on the biology of extreme modifications in *Cetraria aculeata* (lichenized Ascomycota). *Annals of Botany* **109**: 1133–1148.
- Poelt, J. (1969) *Bestimmungsschlüssel Europäischer Flechten*. Lehre: Verlag J. Cramer.
- Printzen, C., Fernández-Mendoza, F., Muggia, L., Berg, G. & Grube, M. (2012) Alphaproteobacterial communities in geographically distant populations of the lichen *Cetraria aculeata*. *FEMS Microbiology Ecology* **82**: 316–325.
- Printzen, C., Domaschke, S., Fernández-Mendoza, F. & Pérez-Ortega, S. (2013) Biogeography and ecology of *Cetraria aculeata*, a widely distributed lichen with a bipolar distribution. *MycKeys* **6**: 33–53.
- Prydatko, V. (ed.) (1998) *Nature-Agricultural Zoning of Ukraine*. Kyiv: GUGKK.
- Purvis, O. W. (1996) Interactions of lichens with metals. *Science Progress* **79**: 283–309.
- Purvis, O. W., Elix, J. A., Broomhead, J. A. & Jones, G. C. (1987) The occurrence of copper-norstictic acid in lichens from cupriferous substrata. *Lichenologist* **19**: 193–203.
- Randlane, T. & Saag, A. (2006) Cetrarioid lichens in Europe – an identification key for the species. In *Central European Lichens – Diversity and Threat* (A. Lackovičová, A. Guttova, E. Lisicka & P. Lizoň, eds): 75–84. Ithaca: Mycotaxon Ltd.
- Savicz, V. P. (1924) De lichene terrestri novo *Cornicularia steppae* mihi nec non lichene *Cornicularia tenuissima*. *Notulae Systematicae ex Instituto Cryptogamico Horti Botanici Petropol.* **3**: 185–188 [In Russian].
- Seaward, M. R. D., Sipman, H. J. M. & Sohrabi, M. (2008) A revised checklist of lichenized, lichenicolous and allied fungi for Iran. *Sauteria* **15**: 459–520.
- Sohrabi, M. & Alstrup, V. (2007) Additions to the lichen mycota of Iran from East Azerbaijan Province. *Mycotaxon* **100**: 145–148.
- Thell, A. & Kärnefelt, I. (2011) *Cetraria*. *Nordic Lichen Flora* **4**: 37–43.
- Thell, A., Stenroos, S. & Myllys, L. (2000) A DNA-study of the *Cetraria aculeata* and *C. islandica* groups. *Folia Cryptogamica Estonica* **32**: 95–106.
- Thell, A., Stenroos, S., Feuerer, T., Kärnefelt, I., Myllys, L. & Hyvönen, J. (2002) Phylogeny of cetrarioid lichens (*Parmeliaceae*) inferred from ITS and  $\beta$ -tubulin sequences, morphology, anatomy and secondary chemistry. *Mycological Progress* **1**: 335–354.
- Urbanavichus, G. P. (2010) *A Checklist of the Lichen Flora of Russia*. St. Petersburg: Nauka.
- Vainshtein, E. A., Ravinskaia, A. P. & Shapiro, I. A. (1990) *Handbook on Lichen Chemotaxonomy: Methodological Guide*. Leningrad: Nauka [In Russian].
- Wagner, V. & Spribille, T. (2005) *Preliminary Checklist of Lichens of Kazakhstan*. Online access: <http://www.geobotanik.uni-goettingen.de/spribille/>
- White, F. J. & James, P. W. (1985) A new guide to microchemical techniques for the identification of lichen substances. *Bulletin of the British Lichen Society* **57**: 1–41.

- Yazıcı, K., Aptroot, A., Aslan, A., Etayo, J., Spier, L. & Karagöz Y. (2010) Lichenized and lichenicolous fungi from nine different areas in Turkey. *Mycotaxon* **111**: 113–116.
- Yoshimura, I., Kinoshita, Y., Yamamoto, Y., Hunecks, S. & Yamadan, Y. (1994) Analysis of secondary metabolites from lichen by high performance liquid chromatography with a photodiode array detector. *Phytochemical Analysis* **5**: 197–205.
- Zhurbenko, M. P. (1996) *Cetraria* Ach. emend. Kärnef. In *Handbook of the Lichens of Russia. Vol. 6. Alectoriaceae, Parmeliaceae, Stereocaulaceae* (N. S. Golubkova, ed.): 41–44. St-Petersburg: Nauka [In Russian].