

DE RHAM COHOMOLOGY OF LOCAL COHOMOLOGY MODULES: THE GRADED CASE

TONY J. PUTHENPURAKAL

Abstract. Let K be a field of characteristic zero, and let $R = K[X_1, \dots, X_n]$. Let $A_n(K) = K\langle X_1, \dots, X_n, \partial_1, \dots, \partial_n \rangle$ be the n th Weyl algebra over K . We consider the case when R and $A_n(K)$ are graded by giving $\deg X_i = \omega_i$ and $\deg \partial_i = -\omega_i$ for $i = 1, \dots, n$ (here ω_i are positive integers). Set $\omega = \sum_{k=1}^n \omega_k$. Let I be a graded ideal in R . By a result due to Lyubeznik the local cohomology modules $H_I^i(R)$ are holonomic $(A_n(K))$ -modules for each $i \geq 0$. In this article we prove that the de Rham cohomology modules $H^*(\partial; H_I^*(R))$ are concentrated in degree $-\omega$; that is, $H^*(\partial; H_I^*(R))_j = 0$ for $j \neq -\omega$. As an application when $A = R/(f)$ is an isolated singularity, we relate $H^{n-1}(\partial; H_{(f)}^1(R))$ to $H^{n-1}(\partial(f); A)$, the $(n-1)$ th Koszul cohomology of A with respect to $\partial_1(f), \dots, \partial_n(f)$.

Let K be a field of characteristic zero, and let $R = K[X_1, \dots, X_n]$. We consider R graded with $\deg X_i = \omega_i$ for $i = 1, \dots, n$; here ω_i are positive integers. Set $\mathfrak{m} = (X_1, \dots, X_n)$. Let I be a graded ideal in R . The local cohomology modules $H_I^*(R)$ are clearly graded R -modules. Let $A_n(K) = K\langle X_1, \dots, X_n, \partial_1, \dots, \partial_n \rangle$ be the n th Weyl algebra over K . By a result due to Lyubeznik (see [3, Section 2.2.d]), the local cohomology modules $H_I^i(R)$ are holonomic $(A_n(K))$ -modules for each $i \geq 0$. We can consider $A_n(K)$ graded by giving $\deg \partial_i = -\omega_i$ for $i = 1, \dots, n$.

Let N be a graded left $(A_n(K))$ -module. Now $\partial = \partial_1, \dots, \partial_n$ are pairwise commuting K -linear maps, so we can consider the de Rham complex $K(\partial; N)$. Notice that the de Rham cohomology modules $H^*(\partial; N)$ are in general only graded K -vector spaces. They are finite-dimensional if N is holonomic (see [1, Chapter 1, Theorem 6.1]). In particular, $H^*(\partial; H_I^*(R))$ are finite-dimensional graded K -vector spaces.

Our first result is as follows.

Received August 29, 2013. Accepted February 5, 2014.
First published online January 26, 2015.

2010 Mathematics Subject Classification. Primary 13D45; Secondary 13N10.

© 2015 by The Editorial Board of the *Nagoya Mathematical Journal*

THEOREM 1. *Let I be a graded ideal in R . Set $\omega = \sum_{i=1}^n \omega_i$. Then the de Rham cohomology modules $H^*(\partial_1, \dots, \partial_n; H_I^*(R))$ are concentrated in degree $-\omega$; that is,*

$$H^*(\partial_1, \dots, \partial_n; H_I^*(R))_j = 0, \quad \text{for } j \neq -\omega.$$

We give an application of Theorem 1. Let f be a homogeneous polynomial in R , with $A = R/(f)$ an isolated singularity; that is, A_P is regular for all homogeneous prime ideals $P \neq \mathfrak{m}$. Let $H^i(\partial(f); A)$ be the i th Koszul cohomology of A with respect to $\partial_1(f), \dots, \partial_n(f)$. We show the following.

THEOREM 2 (with hypotheses as above). *There exists a filtration $\mathcal{F} = \{\mathcal{F}_\nu\}_{\nu \geq 0}$ consisting of K -subspaces of $H^{n-1}(\partial; H_{(f)}^1(R))$ with $\mathcal{F}_\nu = H^{n-1}(\partial; H_{(f)}^1(R))$ for $\nu \gg 0$, $\mathcal{F}_\nu \supseteq \mathcal{F}_{\nu-1}$, and $\mathcal{F}_0 = 0$ and injective K -linear maps*

$$\eta_\nu: \frac{\mathcal{F}_\nu}{\mathcal{F}_{\nu-1}} \rightarrow H^{n-1}(\partial(f); A)_{(\nu+1) \deg f - \omega}.$$

The techniques used in this theorem are generalized in [6] to show that $H^i(\partial; H_{(f)}^1(R)) = 0$ for $1 < i < n - 1$ and $H^1(\partial; H_{(f)}^1(R)) \cong K$. There is no software to compute de Rham cohomology of an $(A_n(K))$ -module M . As an application of Theorem 2, we prove the following.

EXAMPLE 0.1. Let $R = K[X_1, \dots, X_n]$, and let $f = X_1^2 + X_2^2 + \dots + X_{n-1}^2 + X_n^m$ with $m \geq 2$. Then

- (1) if m is odd, then $H^{n-1}(\partial; H_{(f)}^1(R)) = 0$;
- (2) if m is even, then
 - (a) if n is odd, then $H^{n-1}(\partial; H_{(f)}^1(R)) = 0$, and
 - (b) if n is even, then $\dim_K H^{n-1}(\partial; H_{(f)}^1(R)) \leq 1$.

We now describe in brief the contents of this article. In Section 1 we discuss a few preliminaries that we need. In Section 2 we introduce the concept of generalized Eulerian modules. In Section 3 we give a proof of Theorem 1. In Section 4 we give an outline of proof of Theorem 2. In Section 5 we prove Theorem 2. In Section 6 we give a proof of Example 0.1.

§1. Preliminaries

In this section we discuss a few preliminary results that we need.

REMARK 1.1. Although all the results are stated for de Rham cohomology of an $(A_n(K))$ -module M , we will actually work with de Rham homology.

Note that $H_i(\partial, M) = H^{n-i}(\partial, M)$ for any $(A_n(K))$ -module M . Let $S = K[\partial_1, \dots, \partial_n]$. Consider it as a subring of $A_n(K)$. Then note that $H_i(\partial, M)$ is the i th Koszul homology module of M with respect to ∂ .

1.2. Let M be a holonomic $(A_n(K))$ -module. Then for the case where $i = 0, 1$, the de Rham homology modules $H_i(\partial_n, M)$ are holonomic $(A_{n-1}(K))$ -modules (see [1, Theorem 6.2]).

The following result is well known (see [2, Corollary 1.6.13]).

LEMMA 1.3. *Let $\partial = \partial_r, \partial_{r+1}, \dots, \partial_n$, and let $\partial' = \partial_{r+1}, \dots, \partial_n$. Let M be a left $(A_n(K))$ -module. For each $i \geq 0$ there exists an exact sequence*

$$0 \rightarrow H_0(\partial_r; H_i(\partial'; M)) \rightarrow H_i(\partial; M) \rightarrow H_1(\partial_r; H_{i-1}(\partial'; M)) \rightarrow 0.$$

§2. Generalized Eulerian modules

Consider the Eulerian operator

$$\mathcal{E}_n = \omega_1 X_1 \partial_1 + \omega_2 X_2 \partial_2 + \dots + \omega_n X_n \partial_n.$$

If $r \in R$ is homogeneous, then recall that $\mathcal{E}_n r = (\deg r) \cdot r$. Note that degree of \mathcal{E}_n is zero.

Let M be a graded $(A_n(K))$ -module. If m is homogeneous, we set $|m| = \deg m$. We say that M is *Eulerian* $(A_n(K))$ -module if $\mathcal{E}_n m = |m| \cdot m$ for each homogeneous $m \in M$. This notion was discovered by Ma and Zhang (see their excellent paper [4]). They prove that local cohomology modules $H_i^*(R)$ are Eulerian $(A_n(K))$ -modules (see [4, Theorem 5.3]). In fact, they prove this when R is standard graded. The same proof can be adapted to prove the general case.

It can easily be seen that if M is an Eulerian $(A_n(K))$ -module, then so are each graded submodule and graded quotient of M . However, extensions of Eulerian modules need not be Eulerian (see [4, Remark 3.6]). To rectify this, we introduce the following notion. A graded $(A_n(K))$ -module M is said to be *generalized Eulerian* if for a homogeneous element m of M there exists a positive integer a (here a may depend on m) such that

$$(\mathcal{E}_n - |m|)^a m = 0.$$

We now prove that the class of generalized Eulerian modules is closed under extensions.

PROPOSITION 2.1. *Let $0 \rightarrow M_1 \xrightarrow{\alpha_1} M_2 \xrightarrow{\alpha_2} M_3 \rightarrow 0$ be a short exact sequence of graded $(A_n(K))$ -modules. Then the following are equivalent:*

- (1) M_2 is generalized Eulerian,
- (2) M_1 and M_3 are generalized Eulerian.

Proof. The assertion (1) \implies (2) is clear. We prove (2) \implies (1). Let $m \in M_2$ be homogeneous. Because M_3 is generalized Eulerian, we have

$$(\mathcal{E}_n - |m|)^b \alpha_2(m) = 0 \quad \text{for some } b \geq 1.$$

Set $v_2 = (\mathcal{E}_n - |m|)^b m \in M_2$. Because α_2 is $(A_n(K))$ -linear, we get $\alpha_2(v_2) = 0$. So $v_2 = \alpha_1(v_1)$ for some $v_1 \in M_1$. Note that $\deg v_1 = \deg v_2 = |m|$. Because M_1 is generalized Eulerian, we have

$$(\mathcal{E}_n - |m|)^a v_1 = 0 \quad \text{for some } a \geq 1.$$

Because α_1 is $(A_n(K))$ -linear, we get $(\mathcal{E}_n - |m|)^a v_2 = 0$. It follows that

$$(\mathcal{E}_n - |m|)^{a+b} m = 0. \quad \square$$

If M is a graded $(A_n(K))$ -module, then for $l \in \mathbb{Z}$ the module $M(l)$ denotes the shift of M by l ; that is, $M(l)_n = M_{n+l}$ for all $n \in \mathbb{Z}$. The following result was proved for Eulerian $(A_n(K))$ -modules in [4, Remark 2.5].

PROPOSITION 2.2. *Let M be a nonzero generalized Eulerian $(A_n(K))$ -module. Then for $l \neq 0$, the module $M(l)$ is not a generalized Eulerian $(A_n(K))$ -module.*

Proof. Suppose that $M(l)$ is a generalized Eulerian $(A_n(K))$ -module for some $l \neq 0$. Let $m \in M$ be homogeneous of degree r and nonzero. Because M is generalized Eulerian $(A_n(K))$ -module, we have

$$(\mathcal{E}_n - r)^a m = 0 \quad \text{for some } a \geq 1.$$

We may assume that $(\mathcal{E}_n - r)^{a-1} m \neq 0$. Now $m \in M(l)_{r-l}$. Because $M(l)$ is generalized Eulerian, we get

$$(\mathcal{E}_n - r + l)^b m = 0 \quad \text{for some } b \geq 1.$$

Notice that

$$0 = (\mathcal{E}_n - r + l)^b m = \left(l^b + \sum_{i=1}^b \binom{b}{i} l^{b-i} (\mathcal{E}_n - r)^i \right) m.$$

Multiply the term on the left by $(\mathcal{E}_n - r)^{a-1}$. We obtain

$$l^b(\mathcal{E}_n - r)^{a-1}m = 0.$$

Because $l \neq 0$, we get $(\mathcal{E}_n - r)^{a-1}m = 0$, a contradiction. □

§3. Proof of Theorem 1

In this section we prove Theorem 1. Notice that $H_i^j(R)$ are Eulerian $(A_n(K))$ -modules for all $i \geq 0$. Hence, Theorem 1 follows from the following more general result.

THEOREM 3.1. *Let M be a generalized Eulerian $(A_n(K))$ -module. Then $H_i(\partial; M)$ is concentrated in degree $-\omega = -\sum_{k=1}^n \omega_k$.*

Before proving Theorem 3.1, we need to prove a few preliminary results.

PROPOSITION 3.2. *Let M be a generalized Eulerian $(A_n(K))$ -module. Then for $i = 0, 1$, the $(A_{n-1}(K))$ -modules $H_i(\partial_n; M)(-\omega_n)$ are generalized Eulerian.*

Proof. Clearly, $H_i(\partial_n; M)(-\omega_n)$ are $(A_{n-1}(K))$ -modules for $i = 0, 1$. We have an exact sequence of $(A_{n-1}(K))$ -modules

$$0 \rightarrow H_1(\partial_n; M) \rightarrow M(\omega_n) \xrightarrow{\partial_n} M \rightarrow H_0(\partial_n; M) \rightarrow 0.$$

Note that $H_1(\partial_n; M)(-\omega_n) \subset M$. Let $\xi \in H_1(\partial_n; M)(-\omega_n)$ be homogeneous. As M is generalized Eulerian, we have

$$(\mathcal{E}_n - |\xi|)^a \xi = 0 \quad \text{for some } a \geq 1.$$

Notice that $\mathcal{E}_n = \mathcal{E}_{n-1} + \omega_n X_n \partial_n$. Also note that $X_n \partial_n$ commutes with \mathcal{E}_{n-1} . Thus,

$$0 = (\mathcal{E}_{n-1} - |\xi| + \omega_n X_n \partial_n)^a \xi = ((\mathcal{E}_{n-1} - |\xi|)^a + (*)X_n \partial_n) \xi.$$

Because $\partial_n \xi = 0$, we get $(\mathcal{E}_{n-1} - |\xi|)^a \xi = 0$. It follows that $H_1(\partial_n; M)(-\omega_n)$ is a generalized Eulerian $(A_{n-1}(K))$ -module.

Let $\xi \in H_0(\partial_n; M)(-\omega_n)$ be homogeneous of degree r . Then $\xi = \alpha + \partial_n M$, where $\alpha \in M_{r-\omega_n}$. Because M is generalized Eulerian, we get

$$(\mathcal{E}_n - r + \omega_n)^a \alpha = 0 \quad \text{for some } a \geq 1.$$

Notice that $\mathcal{E}_n = \mathcal{E}_{n-1} + \omega_n X_n \partial_n = \mathcal{E}_{n-1} + \omega_n \partial_n X_n - \omega_n$, so $\mathcal{E}_n - r + \omega_n = \mathcal{E}_{n-1} - r + \omega_n \partial_n X_n$. Notice that $\partial_n X_n$ commutes with \mathcal{E}_{n-1} . Thus,

$$0 = (\mathcal{E}_{n-1} - r + \omega_n \partial_n X_n)^a \alpha = (\mathcal{E}_{n-1} - r)^a \alpha + \partial_n \cdot * \alpha.$$

Going mod $\partial_n M$, we get

$$(\mathcal{E}_{n-1} - r)^a \xi = 0.$$

It follows that $H_0(\partial_n; M)(-\omega_n)$ is a generalized Eulerian $(A_{n-1}(K))$ -module. □

REMARK 3.3. If M is Eulerian, then the same proof shows that $H_i(\partial_n; M)(-\omega_n)$ is an Eulerian $(A_{n-1}(K))$ -module for $i = 0, 1$. However, as the proof of the following theorem shows, we can prove only that $H_1(\partial_{n-1}, \partial_n; M)(-\omega_{n-1} - \omega_n)$ is a generalized Eulerian $(A_{n-1}(K))$ -module.

PROPOSITION 3.4. *Let M be a generalized Eulerian $(A_n(K))$ -module. Let $\partial = \partial_i, \partial_{i+1}, \dots, \partial_n$; here $i \geq 2$. Then for each $j \geq 0$ the de Rham homology module*

$$H_j(\partial; M) \left(- \sum_{k=i}^n \omega_k \right)$$

is a generalized Eulerian $(A_{i-1}(K))$ -module.

Proof. We prove this result by descending induction on i . For $i = n$, the result holds by Proposition 3.2. Set $\partial' = \partial_{i+1}, \dots, \partial_n$. By induction hypothesis $H_j(\partial'; M)(-\sum_{k=i+1}^n \omega_k)$ is generalized Eulerian $(A_i(K))$ -module. By Proposition 3.2 again, for $l = 0, 1$ and for each $j \geq 0$,

$$H_l \left(\partial_i; H_j(\partial'; M) \left(- \sum_{k=i+1}^n \omega_k \right) \right) (-\omega_i) = H_l(\partial_i; H_j(\partial'; M)) \left(- \sum_{k=i}^n \omega_k \right)$$

is generalized Eulerian. By Lemma 1.3 we have the exact sequence

$$0 \rightarrow H_0(\partial_i; H_j(\partial'; M)) \rightarrow H_j(\partial; M) \rightarrow H_1(\partial_i; H_{j-1}(\partial'; M)) \rightarrow 0.$$

The modules at the left and right end become generalized Eulerian after shifting by $-\sum_{k=i}^n \omega_k$. By Proposition 2.1 it follows that for each $j \geq 0$ the de Rham homology module

$$H_j(\partial; M) \left(- \sum_{k=i}^n \omega_k \right)$$

is a generalized Eulerian $(A_{i-1}(K))$ -module. □

We now consider the case when $n = 1$.

PROPOSITION 3.5. *Let M be a generalized Eulerian $(A_1(K))$ -module. Then for $l = 0, 1$ the modules $H_l(\partial_1; M)$ are concentrated in degree $-\omega_1$.*

Proof. We have an exact sequence of K -vector spaces

$$0 \rightarrow H_1(\partial_1; M) \rightarrow M(\omega_1) \xrightarrow{\partial_1} M \rightarrow H_0(\partial_1; M) \rightarrow 0.$$

Let $\xi \in H_1(\partial_1; M)(-\omega_1)$ be homogeneous and nonzero. Because $\xi \in M$, we have

$$(\omega_1 X_1 \partial_1 - |\xi|)^a \xi = 0 \quad \text{for some } a \geq 1.$$

Notice that $(\omega_1 X_1 \partial_1 - |\xi|)^a = (*)\partial_1 + (-1)^a |\xi|^a$. Because $\partial_1 \xi = 0$, we get $(-1)^a |\xi|^a \xi = 0$. Because $\xi \neq 0$, we get $|\xi| = 0$. It follows that $H_1(\partial_1; M)$ is concentrated in degree $-\omega_1$.

Let $\xi \in H_0(\partial_1, M)$ be nonzero and homogeneous of degree r . Let $\xi = \alpha + \partial_1 M$, where $\alpha \in M_r$. Because M is generalized Eulerian, we get

$$(\omega_1 X_1 \partial_1 - r)^a \alpha = 0 \quad \text{for some } a \geq 1.$$

Notice that $\omega_1 X_1 \partial_1 = \omega_1 \partial_1 X_1 - \omega_1$, so we have

$$0 = (\omega_1 \partial_1 X_1 - (r + \omega_1))^a \alpha = (\partial_1 * + (-1)^a (r + \omega_1)^a) \alpha.$$

In $M/\partial_1 M$, we have $(-1)^a (r + \omega_1)^a \xi = 0$. Because $\xi \neq 0$, we get $r = -\omega_1$. It follows that $H_0(\partial_1; M)$ is concentrated in degree $-\omega_1$. □

We now give the following.

Proof of Theorem 3.1. Set $\partial' = \partial_2, \dots, \partial_n$. By Proposition 3.4, $N_j = H_j(\partial'; M)(-\sum_{k=2}^n \omega_k)$ is a generalized Eulerian $(A_1(K))$ -module, for each $j \geq 0$. We use exact sequence in Lemma 1.3 and shift it by $-\sum_{k=2}^n \omega_k$ to obtain an exact sequence

$$0 \rightarrow H_0(\partial_1, N_j) \rightarrow H_j(\partial; M) \left(-\sum_{k=2}^n \omega_k \right) \rightarrow H_1(\partial_1, N_{j-1}) \rightarrow 0$$

for each $j \geq 0$. By Proposition 3.5, the modules on the left and right of the above exact sequence are concentrated in degree $-\omega_1$. It follows that for each $j \geq 0$ the K -vector space $H_j(\partial; M)$ is concentrated in degree $-\omega = -\sum_{k=1}^n \omega_k$. □

§4. Outline of proof of Theorem 2

The proof of Theorem 2 is a bit long and has a lot of technical details. For the convenience of the reader, we give an outline of the proof.

4.1. By [5, Lemma 2.7], we have $H_1(\partial, R_f) \cong H_1(\partial, H^1_{(f)}(R))$. Thus, it is sufficient to work with $H_1(\partial, R_f)$ in order to prove Theorem 2. We consider elements of R_f^m as column vectors. For $x \in R_f^m$, we write $x = (x_1, \dots, x_m)'$; here $'$ indicates “transpose”.

4.2. Let $\xi \in R_f^m \setminus R^m$. The element $(a_1/f^i, a_2/f^i, \dots, a_m/f^i)'$, with $a_j \in R$ for all j , is said to be a *normal form* of ξ if

- (1) $\xi = (a_1/f^i, a_2/f^i, \dots, a_m/f^i)'$,
- (2) f does not divide a_j for some j , and
- (3) $i \geq 1$.

It can easily be shown that the normal form of ξ exists and is unique (see Proposition 5.1). Let $(a_1/f^i, a_2/f^i, \dots, a_m/f^i)'$ be the normal form of ξ . Set $L(\xi) = i$. Notice that $L(\xi) \geq 1$.

4.3. Construction of a function $\theta: Z_1(\partial, R_f) \setminus R^n \rightarrow H_1(\partial(f); A)$

Let $\xi \in Z_1(\partial, R_f) \setminus R^n$. Let $(a_1/f^i, a_2/f^i, \dots, a_n/f^i)'$ be the normal form of ξ . Thus, we have $\sum_{j=1}^n \partial/\partial X_j(a_j/f^i) = 0$, so we have

$$\frac{1}{f^i} \left(\sum_{j=1}^n \frac{\partial a_j}{\partial X_j} \right) - \frac{i}{f^{i+1}} \left(\sum_{j=1}^n a_j \frac{\partial f}{\partial X_j} \right) = 0.$$

It follows that

$$f \text{ divides } \sum_{j=1}^n a_j \frac{\partial f}{\partial X_j}.$$

Thus, $(\bar{a}_1, \dots, \bar{a}_n)' \in Z_1(\partial(f); A)$. We set

$$\theta(\xi) = [(\bar{a}_1, \dots, \bar{a}_n)'] \in H_1(\partial(f); A).$$

REMARK 4.4. It can be shown that if $\xi \in Z_1(\partial, R_f)_{-\omega}$ is nonzero, then $\xi \notin R^n$ (see Section 5.2). If $L(\xi) = i$, then by Section 5.3 we have

$$\theta(\xi) \in H_1(\partial(f); A)_{(i+1) \deg f - \omega}.$$

The next result uses the fact that A is an isolated singularity.

PROPOSITION 4.5. *If $\xi \in B_1(\partial, R_f)_{-\omega}$ is nonzero, then $\theta(\xi) = 0$.*

4.6. Let $\xi \in R_f^m$. We define $L(f)$ as follows.

Case 1: $\xi \in R_f^m \setminus R^m$. Let $(a_1/f^i, a_2/f^i, \dots, a_m/f^i)'$ be the normal form of ξ . Set $L(\xi) = i$. Notice that $L(\xi) \geq 1$ in this case.

Case 2: $\xi \in R^m \setminus \{0\}$. Set $L(\xi) = 0$.

Case 3: $\xi = 0$. Set $L(\xi) = -\infty$.

The following properties of the function L can be easily verified.

PROPOSITION 4.7 (with hypotheses as above). *Let $\xi, \xi_1, \xi_2 \in R_f^m$, and let $\alpha, \alpha_1, \alpha_2 \in K$. Then we have the following:*

- (1) *if $L(\xi_1) < L(\xi_2)$, then $L(\xi_1 + \xi_2) = L(\xi_2)$;*
- (2) *if $L(\xi_1) = L(\xi_2)$, then $L(\xi_1 + \xi_2) \leq L(\xi_2)$;*
- (3) *$L(\xi_1 + \xi_2) \leq \max\{L(\xi_1), L(\xi_2)\}$;*
- (4) *if $\alpha \in K^*$, then $L(\alpha\xi) = L(\xi)$;*
- (5) *$L(\alpha\xi) \leq L(\xi)$ for all $\alpha \in K$;*
- (6) *$L(\alpha_1\xi_1 + \alpha_2\xi_2) \leq \max\{L(\xi_1), L(\xi_2)\}$;*
- (7) *let $\xi_1, \dots, \xi_r \in R_f^m$, and let $\alpha_1, \dots, \alpha_r \in K$. Then*

$$L\left(\sum_{j=1}^r \alpha_j \xi_j\right) \leq \max\{L(\xi_1), L(\xi_2), \dots, L(\xi_r)\}.$$

4.8. We now use the fact that $H_1(\partial, R_f)$ is concentrated in degree $-\omega = -\sum_{k=1}^n \omega_k$. Thus,

$$H_1(\partial, R_f) = H_1(\partial, R_f)_{-\omega} = \frac{Z_1(\partial, R_f)_{-\omega}}{B_1(\partial, R_f)_{-\omega}}.$$

Let $x \in H_1(\partial, R_f)$ be nonzero. Define

$$L(x) = \min\{L(\xi) \mid x = [\xi], \text{ where } \xi \in Z_1(\partial, R_f)_{-\omega}\}.$$

It can be shown that $L(x) \geq 1$. If $x = 0$, then set

$$L(0) = -\infty.$$

We now define a function

$$\begin{aligned} \tilde{\theta}: H_1(\partial, R_f) &\rightarrow H_1(\partial(f); A) \\ x \mapsto &\begin{cases} \theta(\xi) & \text{if } x \neq 0, x = [\xi], \text{ and } L(x) = L(\xi), \\ 0 & \text{if } x = 0. \end{cases} \end{aligned}$$

It can be shown that $\tilde{\theta}(x)$ is independent of choice of ξ (see Proposition 5.6). Also note that if $L(x) = i$, then $\tilde{\theta}(x) \in H_1(\partial(f); A)_{(i+1)\deg f - \omega}$.

4.9. We now construct a filtration $\mathcal{F} = \{\mathcal{F}_\nu\}_{\nu \geq 0}$ of $H_1(\partial, R_f)$. Set

$$\mathcal{F}_\nu = \{x \in H_1(\partial, R_f) \mid L(x) \leq \nu\}.$$

In Section 5, we prove the following.

PROPOSITION 4.10. *We have the following:*

- (1) \mathcal{F}_ν is a K subspace of $H_1(\partial, R_f)$,
- (2) $\mathcal{F}_\nu \supseteq \mathcal{F}_{\nu-1}$ for all $\nu \geq 1$,
- (3) $\mathcal{F}_\nu = H_1(\partial, R_f)$ for all $\nu \gg 0$,
- (4) $\mathcal{F}_0 = 0$.

Let $\mathcal{G} = \bigoplus_{\nu \geq 1} \mathcal{F}_\nu / \mathcal{F}_{\nu-1}$. For $\nu \geq 1$, we define

$$\begin{aligned} \eta_\nu: \frac{\mathcal{F}_\nu}{\mathcal{F}_{\nu-1}} &\rightarrow H_1(\partial(f); A)_{(\nu+1)\deg f - \omega}, \\ \xi \mapsto &\begin{cases} 0 & \text{if } \xi = 0, \\ \tilde{\theta}(x) & \text{if } \xi = x + \mathcal{F}_{\nu-1} \text{ is nonzero.} \end{cases} \end{aligned}$$

It can be shown that $\eta_\nu(\xi)$ is independent of choice of x (see Proposition 5.10). Finally we prove the following result.

THEOREM 4.11 (with notation as above). *For all $\nu \geq 1$,*

- (1) η_ν is K -linear, and
- (2) η_ν is injective.

§5. Proof of Theorem 2

In this section we give a proof of Theorem 2 with all details. The reader is advised to read the preceding section before reading this section.

We first prove the following.

PROPOSITION 5.1. *Let $\xi \in R_f^m \setminus R^m$. Then a normal form of ξ exists and is unique.*

Proof. Existence: Let $\xi \in R_f^m \setminus R^m$. Let $\xi = (b_1/f^{i_1}, b_2/f^{i_2}, \dots, b_m/f^{i_m})'$ with $f \nmid b_j$ if $b_j \neq 0$. Note that $i_j \leq 0$ is possible. Let

$$i_r = \max\{i_j \mid i_j \geq 1 \text{ and } b_j \neq 0\}.$$

Notice that $i_r \geq 1$. Then

$$\xi = \left(\frac{b_1 f^{i_r - i_1}}{f^{i_r}}, \frac{b_2 f^{i_r - i_2}}{f^{i_r}}, \dots, \frac{b_m f^{i_r - i_m}}{f^{i_r}} \right)'.$$

Note that $f \nmid b_r$. Thus, the expression above is a normal form of ξ .

Uniqueness: Let $(a_1/f^i, \dots, a_m/f^i)'$ and $(b_1/f^r, \dots, b_m/f^r)'$ be two normal forms of ξ . We first assert that $i < r$ is not possible, for if this holds, then because $a_j/f^i = b_j/f^r$, we get $b_j = a_j f^{r-i}$, so $f \mid b_j$ for all j , a contradiction.

A similar argument shows that $i > r$ is not possible, so $i = r$. Thus, $a_j = b_j$ for all j . Thus, the normal form of ξ is unique. □

5.2. Let $\xi \in Z_1(\partial, R_f)_{-\omega}$ be nonzero. Let $\xi = (\xi_1, \dots, \xi_n)'$. Note that

$$\xi \in (R_f(\omega_1) \oplus R_f(\omega_2) \oplus \dots \oplus R_f(\omega_n))_{-\omega}.$$

It follows that

$$\xi_j \in (R_f)_{-\sum_{k \neq j} \omega_k}.$$

It follows that $\xi \in R_f^n \setminus R^n$.

5.3. Let $(a_1/f^i, \dots, a_n/f^i)'$ be the normal form of ξ . Then

$$\deg a_j = i \deg f - \sum_{k \neq j} \omega_k.$$

In particular, going mod f , we get

$$\overline{a_j} \in A(-\deg f + \omega_j)_{(i+1) \deg f - \omega}.$$

Notice that $\deg \partial f / \partial X_j = \deg f - \omega_j$. It follows that

$$(\overline{a_1}, \dots, \overline{a_n})' \in Z_1(\partial(f); A)_{(i+1) \deg f - \omega}.$$

Thus, $\theta(\xi) \in H_1(\partial(f); A)_{(i+1) \deg f - \omega}$.

5.4. Let $\mathbb{K} = \mathbb{K}(\partial; R_f)$ be the de Rham complex on R_f written homologically, so

$$\mathbb{K} = \cdots \rightarrow \mathbb{K}_3 \xrightarrow{\phi_3} \mathbb{K}_2 \xrightarrow{\phi_2} \mathbb{K}_1 \xrightarrow{\phi_1} \mathbb{K}_0 \rightarrow 0.$$

Here $\mathbb{K}_0 = R_f$, $\mathbb{K}_1 = \bigoplus_{k=1}^n R_f(\omega_k)$,

$$\mathbb{K}_2 = \bigoplus_{1 \leq i < j \leq n} R_f(\omega_i + \omega_j), \quad \text{and} \quad \mathbb{K}_3 = \bigoplus_{1 \leq i < j < l \leq n} R_f(\omega_i + \omega_j + \omega_l).$$

Let $\mathbb{K}' = \mathbb{K}(\partial(f); A)$ be the Koszul complex on A with respect to $\partial f / \partial X_1, \dots, \partial f / \partial X_n$. Thus,

$$\mathbb{K}' = \cdots \rightarrow \mathbb{K}'_3 \xrightarrow{\psi_3} \mathbb{K}'_2 \xrightarrow{\psi_2} \mathbb{K}'_1 \xrightarrow{\psi_1} \mathbb{K}'_0 \rightarrow 0.$$

Here $\mathbb{K}'_0 = A$, $\mathbb{K}'_1 = \bigoplus_{k=1}^n A(-\deg f + \omega_k)$,

$$\mathbb{K}'_2 = \bigoplus_{1 \leq i < j \leq n} A(-2 \deg f + \omega_i + \omega_j), \quad \text{and}$$

$$\mathbb{K}'_3 = \bigoplus_{1 \leq i < j < l \leq n} A(-3 \deg f + \omega_i + \omega_j + \omega_l).$$

We now prove Proposition 4.5.

Proof of Proposition 4.5. Let $u \in B_1(\partial; R_f)_{-\omega}$ be nonzero. Let $\xi \in (\mathbb{K}_2)_{-\omega}$ be homogeneous, with $\phi_2(\xi) = u$. Let $\xi = (\xi_{ij} \mid 1 \leq i < j \leq n)'$. Notice that

$$\xi_{ij} \in R_f(\omega_i + \omega_j)_{-\omega} = (R_f)_{-\sum_{k \neq i,j} \omega_k}.$$

It follows that $\xi \in R_f^{\binom{n}{2}} \setminus R^{\binom{n}{2}}$. Set

$$c = \min\{j \mid j = L(\xi) \text{ where } \phi_2(\xi) = u \text{ and } \xi \in (\mathbb{K}_2)_{-\omega} \text{ is homogeneous}\}.$$

Notice that $c \geq 1$. Let $\xi \in (\mathbb{K}_2)_{-\omega}$ be such that $L(\xi) = c$ and $\phi_2(\xi) = u$. Let $(b_{ij}/f^c \mid 1 \leq i < j \leq n)'$ be the normal form of ξ . Let $u = (u_1, \dots, u_n)'$. Then for $l = 1, \dots, n$,

$$u_l = \sum_{i < l} \frac{\partial}{\partial X_i} \left(\frac{b_{il}}{f^c} \right) - \sum_{j > l} \frac{\partial}{\partial X_j} \left(\frac{b_{lj}}{f^c} \right).$$

So

$$u_l = \frac{f}{f^{c+1}} \left(\sum_{i < l} \frac{\partial(b_{il})}{\partial X_i} - \sum_{j > l} \frac{\partial(b_{lj})}{\partial X_j} \right) + \frac{c}{f^{c+1}} \left(- \sum_{i < l} b_{il} \frac{\partial f}{\partial X_i} + \sum_{j > l} b_{lj} \frac{\partial f}{\partial X_j} \right).$$

Set

$$v_l = c \left(- \sum_{i < l} b_{il} \frac{\partial f}{\partial X_i} + \sum_{j > l} b_{lj} \frac{\partial f}{\partial X_j} \right).$$

Therefore,

$$u_l = \frac{f^* + v_l}{f^{c+1}}.$$

CLAIM. $f \nmid v_l$ for some l . First assume the claim. Then $((f^* + v_1)/f^{c+1}, \dots, (f^* + v_n)/f^{c+1})'$ is the normal form of u . Thus,

$$\theta(u) = [(\bar{v}_1, \dots, \bar{v}_n)'] = [\psi_2(-c\bar{b})] = 0.$$

We now prove our claim. Suppose, if possible, that $f \mid v_l$ for all l . Then

$$\psi_2(-c\bar{b}) = (\bar{v}_1, \dots, \bar{v}_l)' = 0,$$

so $-cb \in Z_2(\partial(f); A)$. Because $H_2(\partial(f); A) = 0$, we get $-cb \in B_2(\partial(f); A)$. Thus, $-c\bar{b} = \psi_3(\bar{\gamma})$. Here

$$\gamma = (\gamma_{ijl} \mid 1 \leq i < j < l \leq n)'.$$

Thus,

$$(5.4.1) \quad -cb_{ij} = \sum_{k < i < j} \gamma_{kij} \frac{\partial f}{\partial X_k} - \sum_{i < k < j} \gamma_{ikj} \frac{\partial f}{\partial X_k} + \sum_{i < j < k} \gamma_{ijk} \frac{\partial f}{\partial X_k} + \alpha_{ij} f.$$

We need to compute the degree of γ_{ijl} . Note that $\xi \in (\mathbb{K}_2)_{-\omega}$, so

$$\frac{b_{ij}}{f^c} \in (R_f(\omega_i + \omega_j))_{-\omega}.$$

It follows that

$$(5.4.2) \quad \deg b_{ij} = c \deg f - \omega + \omega_i + \omega_j.$$

It can be easily checked that

$$\bar{b} \in (\mathbb{K}'_2)_{(c+2) \deg f - \omega},$$

so

$$\gamma \in (\mathbb{K}'_3)_{(c+2) \deg f - \omega}.$$

It follows that

$$(5.4.3) \quad \deg \gamma_{ijl} = (c - 1) \deg f - \omega + \omega_i + \omega_j + \omega_l.$$

We first consider the case when $c = 1$. Then by (5.4.1), we have $\alpha_{ij} = 0$. Also,

$$\deg \gamma_{ijl} = -\omega + \omega_i + \omega_j + \omega_l < 0 \quad \text{if } n > 3,$$

so if $n > 3$, we get $\gamma_{ijl} = 0$. Thus, $b = 0$, so $\xi = 0$, a contradiction.

We now consider the case when $n = 3$. Note that $\gamma = \gamma_{123}$ is a constant. Thus,

$$b = \left(\gamma \frac{\partial f}{\partial X_3}, -\gamma \frac{\partial f}{\partial X_2}, \gamma \frac{\partial f}{\partial X_1} \right)'$$

A direct computation yields $u = 0$, a contradiction.

We now consider the case when $c \geq 2$. Notice that by (5.4.1), we have

$$\frac{-cb_{ij}}{f^c} = \frac{1}{f^c} \sum_{k < i < j} \gamma_{kij} \frac{\partial f}{\partial X_k} - \frac{1}{f^c} \sum_{i < k < j} \gamma_{ikj} \frac{\partial f}{\partial X_k} + \frac{1}{f^c} \sum_{i < j < k} \gamma_{ijk} \frac{\partial f}{\partial X_k} + \frac{\alpha_{ij}}{f^{c-1}}.$$

Notice that

$$\frac{\gamma_{kij} \partial f / \partial X_k}{f^c} = \frac{\partial}{\partial X_k} \left(\frac{\gamma_{kij}}{f^{c-1}} \right) - \frac{*}{f^{c-1}}.$$

Put

$$\tilde{\gamma}_* = \frac{1}{c(c-1)} \gamma_*.$$

Thus, we obtain

$$\frac{b_{ij}}{f^c} = \sum_{k < i < j} \frac{\partial}{\partial X_k} \left(\frac{\tilde{\gamma}_{kij}}{f^{c-1}} \right) - \sum_{i < k < j} \frac{\partial}{\partial X_k} \left(\frac{\tilde{\gamma}_{ikj}}{f^{c-1}} \right) + \sum_{i < j < k} \frac{\partial}{\partial X_k} \left(\frac{\tilde{\gamma}_{ijk}}{f^{c-1}} \right) + \frac{\tilde{b}_{ij}}{f^{c-1}}.$$

Set

$$\delta = \left(\frac{\tilde{\gamma}_{ijl}}{f^{c-1}} \mid 1 \leq i < j < l \leq n \right) \quad \text{and} \quad \tilde{\xi} = \left(\frac{\tilde{b}_{ij}}{f^{c-1}} \mid 1 \leq i < j \leq n \right).$$

Then

$$\xi = \phi_3(\delta) + \tilde{\xi},$$

so we have $u = \phi_2(\xi) = \phi_2(\tilde{\xi})$. This contradicts choice of c . □

5.5. By Theorem 3.1 we have

$$H_1(\partial; R_f) = H_1(\partial; R_f)_{-\omega} = \frac{Z_1(\partial; R_f)_{-\omega}}{B_1(\partial; R_f)_{-\omega}}.$$

Let $x \in H_1(\partial; R_f)$ be nonzero. Define

$$L(x) = \min\{L(\xi) \mid x = [\xi], \text{ where } \xi \in Z_1(\partial, R_f)_{-\omega}\}.$$

Let $\xi = (\xi_1, \dots, \xi_n)' \in Z_1(\partial, R_f)_{-\omega}$ be such that $x = [\xi]$, so $\xi \in (\mathbb{K}_1)_{-\omega}$. Thus, $\xi_i \in R_f(+\omega_i)_{-\omega}$, so if $\xi \neq 0$, then $\xi \in R_f^n \setminus R^n$. It follows that $L(\xi) \geq 1$. Thus, $L(x) \geq 1$.

We now define a function

$$\begin{aligned} \tilde{\theta}: H_1(\partial, R_f) &\rightarrow H_1(\partial(f); A), \\ x &\mapsto \begin{cases} \theta(\xi) & \text{if } x \neq 0, x = [\xi], \text{ and } L(x) = L(\xi), \\ 0 & \text{if } x = 0. \end{cases} \end{aligned}$$

PROPOSITION 5.6 (with hypotheses as above). *The element $\widetilde{\theta(x)}$ is independent of the choice of ξ .*

Proof. Suppose that $x = [\xi_1] = [\xi_2]$ is nonzero and that $L(x) = L(\xi_1) = L(\xi_2) = i$. Let $(a_1/f^i, \dots, a_n/f^i)'$ be the normal form of ξ_1 , and let $(b_1/f^i, \dots, b_n/f^i)'$ be the normal form of ξ_2 . It follows that $\xi_1 = \xi_2 + \delta$, where $\delta \in B_1(\partial; R_f)_{-\omega}$. By Proposition 4.7(1), we get $j = L(\delta) \leq i$. Let $(c_1/f^j, \dots, c_n/f^j)'$ be the normal form of δ . We consider two cases.

Case 1: $j < i$. Then note that $a_k = b_k + f^{i-j}c_k$ for $k = 1, \dots, n$. It follows that

$$\theta(\xi_1) = [(\overline{a_1}, \dots, \overline{a_n})] = [(\overline{b_1}, \dots, \overline{b_n})] = \theta(\xi_2).$$

Case 2: $j = i$. Then note that $a_k = b_k + c_k$ for $k = 1, \dots, n$. It follows that

$$\theta(\xi_1) = \theta(\xi_2) + \theta(\delta).$$

However, by Proposition 4.5, $\theta(\delta) = 0$, so $\theta(\xi_1) = \theta(\xi_2)$. Thus, $\widetilde{\theta(x)}$ is independent of choice of ξ . □

5.7. We now construct a filtration $\mathcal{F} = \{\mathcal{F}_\nu\}_{\nu \geq 0}$ of $H_1(\partial, R_f)$. Set

$$\mathcal{F}_\nu = \{x \in H_1(\partial, R_f) \mid L(x) \leq \nu\}.$$

We prove the following proposition.

PROPOSITION 5.8. *We have the following:*

- (1) \mathcal{F}_ν is a K subspace of $H_1(\partial; R_f)$,
- (2) $\mathcal{F}_\nu \supseteq \mathcal{F}_{\nu-1}$ for all $\nu \geq 1$,
- (3) $\mathcal{F}_\nu = H_1(\partial; R_f)$ for all $\nu \gg 0$,
- (4) $\mathcal{F}_0 = 0$.

Proof. (1) Let $x \in \mathcal{F}_\nu$, and let $\alpha \in K$. Then by Proposition 4.7,

$$L(\alpha x) \leq L(x) \leq \nu,$$

so $\alpha x \in \mathcal{F}_\nu$.

Let $x, x' \in \mathcal{F}_\nu$ be nonzero. Let $\xi, \xi' \in Z_1(\partial; R_f)$ be such that $x = [\xi], x' = [\xi']$ and $L(x) = L(\xi), L(x') = L(\xi')$. Then $x + x' = [\xi + \xi']$. It follows that

$$L(x + x') \leq L(\xi + \xi') \leq \max\{L(\xi), L(\xi')\} \leq \nu.$$

Note that the second inequality follows from Proposition 4.7. Thus, $x + x' \in \mathcal{F}_\nu$.

(2) This is clear.

(3) Let $\mathcal{B} = \{x_1, \dots, x_m\}$ be a K -basis of $H_1(\partial; R_f) = H_1(\partial; R_f)_{-\omega}$. Let

$$c = \max\{L(x_i) \mid i = 1, \dots, m\}.$$

We claim that

$$\mathcal{F}_\nu = H_1(\partial; R_f) \quad \text{for all } \nu \geq c.$$

Fix $\nu \geq c$. Let $\xi_i \in Z_1(\partial; R_f)_{-\omega}$ be such that $x_i = [\xi_i]$ and $L(x_i) = L(\xi_i)$ for $i = 1, \dots, m$.

Let $u \in H_1(\partial; R_f)$. Say that $u = \sum_{i=1}^m \alpha_i x_i$ for some $\alpha_1, \dots, \alpha_m \in K$. Then $u = [\sum_{i=1}^m \alpha_i \xi_i]$. It follows that

$$L(u) \leq L\left(\sum_{i=1}^m \alpha_i x_i\right) \leq \max\{L(\xi_i) \mid i = 1, \dots, m\} = c \leq \nu.$$

Here the second inequality follows from Proposition 4.7, so $u \in \mathcal{F}_\nu$. Thus, $\mathcal{F}_\nu = H_1(\partial; R_f)$.

(4) If $x \in H_1(\partial; R_f)$ is nonzero, then $L(x) \geq 1$. It follows that $\mathcal{F}_0 = 0$. \square

5.9. Let $\mathcal{G} = \bigoplus_{\nu \geq 1} \mathcal{F}_\nu / \mathcal{F}_{\nu-1}$. For $\nu \geq 1$ we define

$$\eta_\nu : \frac{\mathcal{F}_\nu}{\mathcal{F}_{\nu-1}} \rightarrow H_1(\partial(f); A)_{(\nu+1) \deg f - \omega},$$

$$u \mapsto \begin{cases} 0 & \text{if } u = 0, \\ \tilde{\theta}(x) & \text{if } u = x + \mathcal{F}_{\nu-1} \text{ is nonzero.} \end{cases}$$

PROPOSITION 5.10 (with hypotheses as above). *The element $\eta_\nu(u)$ is independent of choice of x .*

Proof. Suppose that $u = x + \mathcal{F}_{\nu-1} = x' + \mathcal{F}_{\nu-1}$ is nonzero. Then $x = x' + y$, where $y \in \mathcal{F}_{\nu-1}$. Because $u \neq 0$, we have $x, x' \in \mathcal{F}_\nu \setminus \mathcal{F}_{\nu-1}$, so $L(x) = L(x') = \nu$. Say that $x = [\xi]$, $x' = [\xi']$ and that $y = [\delta]$, where $\xi, \xi', \delta \in Z_1(\partial; R_f)$ with $L(\xi) = L(\xi') = \nu$ and $L(\delta) = L(y) = k \leq \nu - 1$. Thus, we have $\xi = \xi' + \delta + \alpha$, where $\alpha \in B_1(\partial; R_f)_{-\omega}$. Let $L(\alpha) = r$. Note that $r \leq \nu$.

Let $(a_1/f^\nu, \dots, a_n/f^\nu)'$, $(a'_1/f^\nu, \dots, a'_n/f^\nu)'$, $(b_1/f^k, \dots, b_n/f^k)'$, and $(c_1/f^r, \dots, c_n/f^r)'$ be normal forms of ξ , ξ' , δ , and α , respectively. Thus, we have

$$a_j = a'_j + f^{\nu-k} b_j + f^{\nu-r} c_j \quad \text{for } j = 1, \dots, n.$$

Case 1: $r < \nu$. In this case we have $\overline{a_j} = \overline{a'_j}$ in A for each $j = 1, \dots, n$, so $\theta(\xi) = \theta(\xi')$. Thus, $\tilde{\theta}(x) = \tilde{\theta}(x')$.

Case 2: $r = \nu$. In this case notice that $\overline{a_j} = \overline{a'_j} + \overline{c_j}$ in A for each $j = 1, \dots, n$, so $\theta(\xi) = \theta(\xi') + \theta(\alpha)$. However, $\theta(\alpha) = 0$ as $\alpha \in B_1(\partial; R_f)_{-\omega}$ (see Proposition 4.5). Thus, $\tilde{\theta}(x) = \tilde{\theta}(x')$. □

Note that neither θ nor $\tilde{\theta}$ is linear. However, we prove the following.

PROPOSITION 5.11 (with notation as above). *For all $\nu \geq 1$, η_ν is K -linear.*

Proof. Let $u, u' \in \mathcal{F}_\nu / \mathcal{F}_{\nu-1}$. We first show that $\eta_\nu(\alpha u) = \alpha \eta_\nu(u)$ for all $\alpha \in K$. We have nothing to show if $\alpha = 0$ or if $u = 0$, so assume that $\alpha \neq 0$ and that $u \neq 0$. Say that $u = x + \mathcal{F}_{\nu-1}$. Then $\alpha u = \alpha x + \mathcal{F}_{\nu-1}$. Because $\tilde{\theta}(\alpha x) = \alpha \tilde{\theta}(x)$, we get the result.

Next we show that $\eta_\nu(u + u') = \eta_\nu(u) + \eta_\nu(u')$. We have nothing to show if u or u' is zero. Next we consider the case when $u + u' = 0$. Then $u = -u'$,

so $\eta_\nu(u) = -\eta_\nu(u')$. Thus, in this case

$$\eta_\nu(u + u') = 0 = \eta_\nu(u) + \eta_\nu(u').$$

Now consider the case when u, u' are nonzero and $u + u'$ is nonzero. Say that $u = x + \mathcal{F}_{\nu-1}$ and that $u' = x' + \mathcal{F}_{\nu-1}$. Note that because $u + u'$ is nonzero, $x + x' \in \mathcal{F}_\nu \setminus \mathcal{F}_{\nu-1}$. Let $x = [\xi]$, and let $x' = [\xi']$, where $\xi, \xi' \in Z_1(\partial; R_f)_{-\omega}$ and $L(\xi) = L(\xi') = \nu$. Then $x + x' = [\xi + \xi']$. Note that $L(\xi + \xi') \leq \nu$ by Proposition 4.7. But $L(x + x') = \nu$, so $L(\xi + \xi') = \nu$. Let $(a_1/f^\nu, \dots, a_n/f^\nu)'$, $(a'_1/f^\nu, \dots, a'_n/f^\nu)'$ be normal forms of ξ and ξ' , respectively. Note that $((a_1 + a'_1)/f^\nu, \dots, (a_n + a'_n)/f^\nu)'$ is the normal form of $\xi + \xi'$. It follows that $\theta(\xi + \xi') = \theta(\xi) + \theta(\xi')$. Thus, $\theta(x + x') = \tilde{\theta}(x) + \tilde{\theta}(x')$. Therefore,

$$\eta_\nu(u + u') = \eta_\nu(u) + \eta_\nu(u'). \quad \square$$

Finally we have the main result of this section.

Proof of Theorem 2. Let $\nu \geq 1$. By Proposition 5.11, we know that η_ν is a linear map of K -vector spaces. We now prove that η_ν is injective.

Suppose, if possible, that η_ν is not injective. Then there exists nonzero $u \in \mathcal{F}_\nu/\mathcal{F}_{\nu-1}$ with $\eta_\nu(u) = 0$. Say that $u = x + \mathcal{F}_{\nu-1}$. Also, let $x = [\xi]$, where $\xi \in Z_1(\partial; R_f)_{-\omega}$ and $L(\xi) = L(x) = \nu$. Let $(a_1/f^\nu, \dots, a_n/f^\nu)'$ be the normal form of ξ . Thus, we have

$$0 = \eta_\nu(u) = \tilde{\theta}(x) = \theta(\xi) = [(\overline{a_1}, \dots, \overline{a_n})'].$$

It follows that $(\overline{a_1}, \dots, \overline{a_n})' = \psi_2(\overline{b})$, where $\overline{b} = (\overline{b_{ij}} \mid 1 \leq i < j \leq n)'$. It follows that, for $l = 1, \dots, n$,

$$\overline{a_l} = \sum_{i < l} \overline{b_{il}} \frac{\partial f}{\partial X_i} - \sum_{l > j} \overline{b_{lj}} \frac{\partial f}{\partial X_j}.$$

Then it follows that for $l = 1, \dots, n$ we have the following equation in R :

$$(5.11.1) \quad a_l = \sum_{i < l} b_{il} \frac{\partial f}{\partial X_i} - \sum_{l > j} b_{lj} \frac{\partial f}{\partial X_j} + d_l f,$$

for some $d_l \in R$. Note that (5.11.1) is of homogeneous elements in R . Thus, we have the following:

$$(5.11.2) \quad \frac{a_l}{f^\nu} = \frac{\sum_{i < l} b_{il} \frac{\partial f}{\partial X_i}}{f^\nu} - \frac{\sum_{l > j} b_{lj} \frac{\partial f}{\partial X_j}}{f^\nu} + \frac{d_l}{f^{\nu-1}}.$$

We consider two cases.

Case 1: $\nu \geq 2$. Set $\widetilde{b}_{ij} = -b_{il}/(c-1)$. Then note that

$$\frac{b_{il} \frac{\partial f}{\partial X_i}}{f^\nu} = \frac{\partial}{\partial X_i} \left(\frac{\widetilde{b}_{il}}{f^{\nu-1}} \right) - \frac{*}{f^{\nu-1}}.$$

By (5.11.2) we have, for $l = 1, \dots, n$,

$$\frac{a_l}{f^\nu} = \sum_{i < l} \frac{\partial}{\partial X_i} \left(\frac{\widetilde{b}_{il}}{f^{\nu-1}} \right) - \sum_{l < j} \frac{\partial}{\partial X_j} \left(\frac{\widetilde{b}_{lj}}{f^{\nu-1}} \right) + \frac{c_l}{f^{\nu-1}}.$$

Put $\xi' = (c_1/f^{\nu-1}, \dots, c_n/f^{\nu-1})'$, and put $\delta = (\widetilde{b}_{ij}/f^{\nu-1} \mid 1 \leq i < j \leq n)$. Then we have

$$\xi = \phi_2(\delta) + \xi',$$

so we have $x = [\xi] = [\xi']$. This yields $L(x) \leq L(\xi') \leq \nu - 1$. This is a contradiction.

Case 2: $\nu = 1$. Note that $\xi \in (\mathbb{K}_1)_{-\omega}$. Thus, for $l = 1, \dots, n$ we have

$$\frac{a_l}{f} \in (R_f(\omega_l))_{-\omega}.$$

It follows that

$$\deg a_l = \deg f - \sum_{k \neq l} \omega_k.$$

Also note that $\deg \partial f / \partial X_i = \deg f - \omega_i$. By comparing degrees in (5.11.1) we get $a_l = 0$ for all l . Thus, $\xi = 0$, so $x = 0$. Therefore, $u = 0$, a contradiction. □

§6. Example 0.1

Let $R = K[X_1, \dots, X_n]$, and let $f = X_1^2 + \dots + X_{n-1}^2 + X_n^m$, with $m \geq 2$. Set $A = R/(f)$. In this section we compute $H_1(\partial; H_{(f)}^1(R))$.

6.1. We give $\omega_i = \deg X_i = m$ for $i = 1, \dots, n - 1$, and we give $\omega_n = \deg X_n = 2$. Note that f is a homogeneous polynomial in R of degree $2m$. Also note that $\omega = \sum_{k=1}^n \omega_k = (n - 1)m + 2$.

6.2. First note that the Jacobian ideal J of f is primary to the unique graded maximal ideal of R . It follows that A is an isolated singularity. Note that $J = (X_1, \dots, X_{n-1}, X_n^{m-1})$. Let $H_i(J; A)$ be the i th Koszul homology of A with respect to J .

PROPOSITION 6.3. *The Hilbert series, $P(t)$, of $H_1(J; A)$ is*

$$P(t) = \sum_{k=0}^{m-2} t^{2m+2k}.$$

Proof. It is easily verified that X_1, \dots, X_{n-1} is an A -regular sequence. Set

$$B = A/(X_1, \dots, X_{n-1})A = \frac{K[X_n]}{(X_n^m)} = K \oplus KX_n \oplus X_n^2 \oplus \dots \oplus KX_n^{m-1}.$$

Note that we have an exact sequence

$$0 \rightarrow H_1(J; A) \rightarrow B(-2(m-1)) \xrightarrow{X_n^{m-1}} B.$$

It follows that $H_1(J; A) = X_n B(-2(m-1))$. The result follows. □

6.4. By Theorem 2 there exists a filtration $\mathcal{F} = \{\mathcal{F}_\nu\}_{\nu \geq 0}$ consisting of K -subspaces of $H_1(\partial; H_{(f)}^1(R))$ with $\mathcal{F}_\nu = H^{n-1}(\partial; H_{(f)}^1(R))$ for $\nu \gg 0$, $\mathcal{F}_\nu \supseteq \mathcal{F}_{\nu-1}$, and $\mathcal{F}_0 = 0$ and injective K -linear maps

$$\eta_\nu : \frac{\mathcal{F}_\nu}{\mathcal{F}_{\nu-1}} \longrightarrow H_1(\partial(f); A)_{(\nu+1) \deg f - \omega}.$$

Notice that

$$(\nu + 1) \deg f - \omega = (\nu + 1)2m - (n - 1)m - 2 = (2\nu - n + 3)m - 2.$$

If $\eta_\nu \neq 0$, then by Proposition 6.3 it follows that

$$(2\nu - n + 3)m - 2 = 2m + 2j \quad \text{for some } j = 0, \dots, m - 2.$$

Thus, we obtain

$$(6.4.1) \quad 2\nu m = (n - 1)m + 2(j + 1).$$

It follows that m divides $2(j + 1)$. Because $2(j + 1) \leq 2m - 2$, it follows that $2(j + 1) = m$. Thus, m is even.

6.5. Say that $m = 2r$. Then by (6.4.1) we have

$$2\nu r = (n - 1)r + r,$$

so $\nu = n/2$. It follows that n is even. Furthermore, note that $\eta_\nu = 0$ for $\nu \neq n/2$ and that if $\nu = n/2$ then by (6.3) $\dim \mathcal{F}_{n/2}/\mathcal{F}_{n/2-1} \leq 1$. It follows that in this case $\dim H_1(\partial; H_{(f)}^1(R)) \leq 1$.

6.6. In conclusion we have the following:

- (1) if m is odd, then $H^{n-1}(\partial; H_{(f)}^1(R)) = 0$;
- (2) if m is even, then
 - (a) if n is odd then $H^{n-1}(\partial; H_{(f)}^1(R)) = 0$,
 - (b) if n is even then $\dim_K H^{n-1}(\partial; H_{(f)}^1(R)) \leq 1$.

This proves Example 0.1.

REFERENCES

- [1] J.-E. Björk, *Rings of Differential Operators*, North-Holland Math. Library **21**, North-Holland, Amsterdam, 1979. [MR 0549189](#).
- [2] W. Bruns and J. Herzog, *Cohen–Macaulay Rings*, Cambridge Stud. Adv. Math. **39**, Cambridge University Press, Cambridge, 1993. [MR 1251956](#).
- [3] G. Lyubeznik, *Finiteness properties of local cohomology modules (an application of D-modules to commutative algebra)*, Invent. Math. **113** (1993), 41–55. [MR 1223223](#). [DOI 10.1007/BF01244301](#).
- [4] L. Ma and W. Zhang, *Eulerian graded D-modules*, Math. Res. Lett. **21** (2014), 149–167. [MR 3247047](#). [DOI 10.4310/MRL.2014.v21.n1.a13](#).
- [5] T. J. Puthenpurakal, *De Rham cohomology of local cohomology modules*, preprint, [arXiv:1302.0116v2](#) [math.AC].
- [6] T. J. Puthenpurakal and R. B. T. Reddy, *de Rham cohomology of $H_{(f)}^1(R)$, where $V(f)$ is a smooth hypersurface in \mathbb{P}^n* , preprint, [arXiv:1310.4654v1](#) [math.AC].

Department of Mathematics
Indian Institute of Technology Bombay
Powai, Mumbai 400 076
India
tputhen@math.iitb.ac.in