

Original Article

Cite this article: Visker JR, Ferguson DP. (2018) Postnatal undernutrition in mice causes cardiac arrhythmogenesis which is exacerbated when pharmacologically stressed. *Journal of Developmental Origins of Health and Disease* 9: 417–424. doi: 10.1017/S2040174418000156

Received: 6 September 2017
Revised: 20 February 2018
Accepted: 20 February 2018
First published online: 12 April 2018

Key words:

cardiovascular arrhythmias; Developmental Origins of Health and Disease; developmental programming

Address for correspondence:

J. R. Visker, Department of Kinesiology,
Michigan State University, 308 West Circle
Drive Room 27S, East Lansing, MI, 48824, USA.
E-mail: viskerjo@msu.edu

Postnatal undernutrition in mice causes cardiac arrhythmogenesis which is exacerbated when pharmacologically stressed

J. R. Visker and D. P. Ferguson

Department of Kinesiology, Michigan State University, East Lansing, MI, USA

Abstract

Growth restriction caused by postnatal undernutrition increases risk for cardiovascular disease in adulthood with the potential to induce arrhythmogenesis. Thus, the purpose was to determine if undernutrition during development produced arrhythmias at rest and when stressed with dobutamine in adulthood. Mouse dams were fed (CON: 20% protein), or low-protein (LP: 8%) diet before mating. A cross-fostering model was used where pups nursed by dams fed LP diet in early [EUN; postnatal day (PN) 1–10], late (LUN; PN11–21) and whole (PUN; 1–21) phases of postnatal life. Weaned pups were switched to CON diets for the remainder of the study (PN80). At PN80, body composition (magnetic resonance imaging), and quantitative electrocardiogram (ECG) measurements were obtained under 1% isoflurane anesthesia. After baseline ECG, an IP injection (1.5 µg/g body weight) of dobutamine was administered and ECG repeated. Undernutrition significantly ($P < 0.05$) reduced body weight in LUN (22.68 ± 0.88 g) and PUN (19.96 ± 0.32 g) but not in CON (25.05 ± 0.96 g) and EUN (25.28 ± 0.9207 g). Fat mass decreased in all groups compared with controls (CON: 8.00 ± 1.2 g, EUN: 6.32 ± 0.65 g, LUN: 5.11 ± 1.1 g, PUN: 3.90 ± 0.25 g). Lean mass was only significantly reduced in PUN (CON: 17.99 ± 0.26 g, EUN: 17.78 ± 0.39 g, LUN: 17.34 ± 0.33 g, PUN: 15.85 ± 0.28 g). Absolute heart weights were significantly less from CON, with PUN having the smallest. ECG showed LUN had occurrences of atrial fibrillation; EUN had increases of 1st degree atrioventricular block upon stimulation, and PUN had increased risk for ventricular depolarization arrhythmias. CON did not display arrhythmias. Undernutrition in early life resulted in ventricular arrhythmias under stressed conditions, but undernutrition occurring in later postnatal life there is an increased incidence of atrial arrhythmias.

Introduction

Cardiovascular disease is one of the leading causes of death in the United States and incidence rates have been increasing throughout the past decade.¹ The literature has routinely shown that cardiovascular disease is a result of genetic predisposition, lack of physical activity engagement and a diet high in saturated fats.^{2–5} However, an emerging risk factor for cardiovascular impairment is growth restriction caused by early life nutrition. Professor David Barker championed the Developmental Origins of Health and Disease (DOHaD) hypothesis which states that a brief period of malnutrition or growth restriction in early life will increase the likelihood of cardiovascular disease in adulthood.^{2–7} Yearly, 15 million babies worldwide will be born prematurely and/or experience growth delays due to poor postnatal nutrition. In fact, over 7,000,000 deaths per year is attributed to cardiovascular disease as a result of a suboptimal nutritional environment in early life. Yet, this topic is surprisingly under-researched in terms of mechanisms associated with cardiovascular disease development as a result of early life undernutrition.⁸

Evidence has shown that those who suffered from undernutrition in the early formative years have modified cardiovascular development. This phenotype is represented in humans, mice, rats, sheep, pigs and cows.^{2–5,9,10} Mice that are undernourished early (EUN) in development [postnatal day (PN) 1 to PN11] exhibit less cardiomyocyte number and with the majority being mononucleated cardiomyocytes. While those undernourished later (LUN, PN11 to PN21) in life show the effects of impaired cardiomyocyte hypertrophy.^{5,11–14} Thus, suboptimal nutrition during these different time points could have differential effects on the function of the heart. Specifically, any adverse conditions experienced in the postnatal environment that lead to cardiac reprogramming on heart size, heart structure, or Ca^{2+} management^{14–17} may be associated with increased likelihood of cardiac arrhythmias.^{11,18–20}

The use of an electrocardiogram (ECG) as an imaging tool for cardiac impairment has been widely used in the clinical and research setting, with recent advancements leading to the development of ECGs designed to be used in a murine model.^{21–23} Recent evidence from our lab and others^{24,25} has suggested that cardiac impairment as a result of early life

undernutrition is quiescent during rest and pathological during times of stress. Thus, in order to evaluate cardiac health as a result of early life nutrition the heart must be evaluated during rest and when stressed. Dobutamine is a common adrenergic agent used to increase cardiac output.^{26–29} Dobutamine acts on the β_1 receptors of the sympathetic nervous system to stimulate an increase in stroke volume by causing an inotropic effect on the heart, with a secondary response to increase heart rate (HR) and blood pressure.^{26,28,29} A growing use for dobutamine administration has been in the use of Pharmacological Stress Echocardiography to observe cardiac function and diagnose cardiovascular disease.^{26,28} Pharmacological stress testing is an alternative method to dynamic exercise testing that when combined with noninvasive imaging allows for the detection of myocardial impairment.³⁰ Thus, we used pharmacological stress testing coupled with ECG analysis in mice that were undernourished during the early, late and whole period of postnatal development as a means to determine whether early life nutrition predisposes mice for cardiac arrhythmogenesis.

Methods

All experiments were conducted according to the recommendations of the Guide for Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committee at Michigan State University. All animals were housed in the vivarium on wood-chip bedding in a single room maintained at 21°C with a 12 h light/dark cycle.

Nutritive model

To induce postnatal undernutrition we used methods previously described.³¹ In brief, second and third parity FVB (FVB/N; Charles River Laboratories, Wilmington, MA, USA) mouse dams were fed either a semi-purified control diet (20% protein; Research Diets, New Brunswick, NJ, USA) based on AIN93G, or a low-protein (LP) isocaloric diet (8% protein) beginning 1 week before mating. Table 1 displays the composition of the diets. These diets are casein, cornstarch and soybean oil and have been validated for rodent use during gestation, lactation and early postnatal life. For a detailed review on the theoretical framework for diet composition readers are directed to the work by Reeves *et al.*³² Mating was timed by introducing males for a 24-h period; only pups born within the same 24-h period (PN0) were studied. On PN1 all pups born to dams on the control diet were pooled, and redistributed to one of two experimental groups: (1) *Control (CON)*: pups born to and suckled by well-nourished control dams and weaned to a control diet at weaning (PN21); $n=16$; (2) *postnatal undernutrition (PUN)*: pups born to well-nourished dams and suckled by dams fed the LP diet, then fed the control diet at weaning (PN21). Mouse dams that are fed a LP diet produce 15–20% less milk – thus pups nursed by LP dams will experience a global nutrient deficit and possibly a slightly greater deficit in protein intake.^{33–38} This methodology closely mimics the nutrient intake of many infants who are born small for gestational age.³⁹ In order to distinguish cardiac differences at different stages of postnatal life we used a cross-fostering model at PN10 to segregated postnatal undernutrition to the early phase of undernutrition (EUN; PN1–10; $n=14$ litters) and late phase of undernutrition (LUN; PN11–21; $n=15$ litters). As displayed in Figure 1, at PN10 a subcohort pups suckled to control dams were cross-fostered to dams fed the LP diet (*LUN*). Additionally at

Table 1. Composition of control (CON) and low-protein (LP) diets

Diets	Protein (g)	Fat (g)	Carbohydrate (g)	Fiber (g)
CON	196.6	70	609.4	50
LP	80.6	70	728.8	60
	Protein (g%)	Fat (g%)	Carbohydrate (g%)	Fiber (g%)
CON	19.7	7	60.9	5
LP	8.1	7	73.1	6
	Protein (kcal)	Fat (kcal)	Carbohydrate (kcal)	Total (kcal)
CON	786.4	630	2437.5	3853.9
LP	322.2	630	2915.3	3867.5
	Protein (kcal%)	Fat (kcal%)	Carbohydrate (kcal%)	Total (kcal%)
CON	20	16	63	100
LP	8	16	75	100

Diets were casein and cornstarch based with soybean oil as the only source of fat. LP diet was achieved by replacing casein with additional cornstarch. The two diets contained the same amount of oil (70 g), and the fatty acid profile (weight percentage of total fat) consists of palmitic (10.3), oleic (22.8), linoleic (51) and linolenic (6.8) acids.




Fig. 1. Cross-foster model for early life nutrition. Control diet = 20% protein, low-protein (LP) diet = 8% protein. Both diets are isocaloric. Postnatal undernutrition (PUN) refers to mice suckled to dams fed the LP diet from postnatal day (PN) 1 to PN21. Early phase of undernutrition (EUN) refers to mice suckled to dams fed the LP diet from PN1–PN10, and then pups were cross-fostered to dams fed the control diet from PN11 to PN21. Late phase of undernutrition (LUN) refers to mice suckled to dams fed the CON diet from PN1–PN10; then cross-fostered to dams fed the LP from PN11 to PN21. Control group (CON) refers to mice suckled to dams fed the CON diet PN1–PN21. At PN21 all pups were weaned from dams and fed the control diet.

PN10 a subcohort of pups suckled to control dams were cross-fostered to dams fed the LP diet (*EUN*). At PN21 all pups (CON, PUN, EUN and LUN) were weaned and fed the control diet until ECG measurement at PN80. Thus, at the time of measurement all mice were consuming a nonrestrictive diet.

Each dam received seven pups (2–3 males and 4–5 females) and individual pups within a litter were identified with a unique tattoo. Females were only evaluated in this study, as work from our lab has shown that males did not display cardiac impairment as measured by treadmill testing, echocardiography and calcium flux of isolated cardiomyocytes and therefore it was hypothesized that they would not display cardiac arrhythmias.⁴⁰ Pups were allocated so that on PN1, average pup weight for all groups was

the same. Litter size was maintained constant throughout lactation by introducing 'donor' pups of similar body weight and age to replace any deaths; donor pups were not studied.

Body composition

At PN80 mice were weighed then fat mass and lean mass were measured by Quantitative Magnetic Resonance (EchoMRI, Houston, TX, USA).³¹ Body composition was determined using the supplied software. Mice were then evaluated for cardiac function via ECG.

ECG

All tests were done with mice placed under 1% isoflurane anesthesia, with the mice on an ECG/heated board (EMKA, Paris, France) and the limbs taped to the four electrodes. Leads I, II and III recordings were obtained for 5 min. The mouse was then administered an intraperitoneal injection of dobutamine (1.5 µg/g body weight; Hospira, Lake Forest, IL, USA) and the measurements were repeated.

ECG parameters that were obtained were HR (beats/min), R-R interval (ms), QRS duration (ms), ST deflection (V) and QT duration (ms). Only parameters found to be within established ranges were analyzed.⁴¹ Shape-based software (EMKA, Paris, France) was used to detect any change in morphology and presence of an arrhythmia. These parameters have been indicated and validated in studies previously analyzing various heart diseases.^{42–44}

After completion of the studies, the mice were allowed to recover for 24 h after which they were euthanized and the hearts were dissected quantitatively and weighed. Heart weights are reported as absolute weight and as normalized for body surface area as calculated using Meeh's formula.⁴⁵

$$\text{Body surface area} = 9.662 \times (\text{body weight, g})^{0.667}.$$

Evaluation of estrous cycle

A validated visual method to evaluate the stage of the estrous cycle was implemented as previously described.⁴⁶ In brief, each mouse was held by the tail with the forepaws resting on a cage lid. Vaginal openings were then visually inspected and classified based on previous literature for the stage of estrous cycle. Mice used in the study were within the follicular phase with low circulating estrogen levels.

Statistics

All data were analyzed in JMP v7.0 (SAS, Cary, NC, USA). Body composition and heart weight were analyzed by a one-way analysis of variance (ANOVA) with the main effects being diet (CON, EUN, LUN and PUN).

ECG parameters were analyzed with a two-way ANOVA with the main effects being diet (CON, EUN, LUN and PUN) and treatment (baseline or dobutamine) and a Tukey's honest significant difference *post-hoc* test. An α level of 0.05 was set *a priori*.

Results

Body composition and heart weight

Our nutritive model successfully induced growth restriction (Fig. 2), as evident by a reduction ($P < 0.05$) in body weight, in the PUN (19.96 ± 0.32 g) and LUN (22.68 ± 0.88 g) groups as compared with controls (25.05 ± 0.96 g) and EUN (25.28 ± 0.92 g). In addition to having the lowest absolute weight the PUN mice had a




Fig. 2. Body weight and composition at postnatal day (PN) 80. Mice were undernourished during early (EUN), late (LUN) and throughout (PUN) lactation phase alongside the control (CON) group. Weight was significantly decreased in the LUN and PUN groups compared with CON and EUN. Lean mass was significantly decreased in the PUN group only. Fat mass was significantly different between all groups. *Significantly different from control; #significantly different from EUN; ^significantly different from LUN; ^significantly different from PUN. $P < 0.05$ values are mean \pm SE.

significant ($P < 0.05$) reduction in lean mass (15.85 ± 0.28 g) as compared with CON, EUN and LUN. There was no difference in lean mass between CON (17.99 ± 0.26 g), EUN (17.78 ± 0.39 g) and LUN (17.34 ± 0.33 g). Fat mass was greatest in the CON (8.00 ± 1.2 g) group followed by EUN (6.32 ± 0.65 g), LUN (5.11 ± 1.1 g) and PUN (3.90 ± 0.25 g), each group was significantly different ($P < 0.05$) from the other.

The absolute heart weight of the CON group (0.118 ± 0.005 g) was significantly ($P < 0.001$) higher than the EUN (0.103 ± 0.002 g), LUN (0.103 ± 0.004 g) and the PUN group (0.094 ± 0.001 g); with the PUN group being significantly less than the three other groups. There was no difference between the EUN and LUN absolute heart weights (Fig. 3b).

When comparing heart weight (standardized to body surface area), amongst the four nutritive groups the only significant decrease in heart weight was the EUN group (1.288 ± 0.031 kg/cm²) as compared with CON (1.405 ± 0.089 kg/cm²), LUN (1.350 ± 0.040 kg/cm²) and PUN (1.342 ± 0.031 kg/cm²) ($P < 0.05$, Fig. 3a).

ECG

Table 2 displays the results obtained from the ECG analysis for mice that were undernourished during critical periods of development along with the response to dobutamine treatment.

Baseline measurements

The HR of the CON group was 534.3 ± 68.7 beats/min. The EUN group displayed HR values of 556.9 ± 29.4 beats/min, whereas the PUN and LUN baseline groups HR was 534.8 ± 68.4 beats/min and 551.6 ± 59.5 beats/min, respectively. There was no difference ($P > 0.05$) in baseline HR values between the groups. As RR interval is closely associated with HR there was also no difference in RR intervals ($P > 0.05$) between the groups (CON RR interval




Fig. 3. (a) Heart weight (standardized to body surface area): mice were undernourished during early (EUN), late (LUN) and throughout (PUN) lactation alongside the control (CON) group. *The EUN group had a significantly decreased heart weight as compared with the CON group ($P < 0.05$, values are mean \pm SE). (b) Absolute heart weight: the control group had a higher heart weight than the other treatment groups with the PUN group being significantly less than the CON, EUN and LUN group ($P < 0.001$). The EUN and LUN groups were not significantly different from each other (values are mean \pm SE; ^{A,B,C}different letters indicate significant difference).

Table 2. Electrocardiogram (ECG) parameters for mice that were undernourished during the early, late and whole phase of lactation along with a control group with and without stimulation of dobutamine

Group	Treatment	Heart rate (BPM)	RR interval (ms)	QRS duration (ms)	ST deflection (V)	QT duration (ms)
CON	Baseline	534.3 \pm 68.7	114.1 \pm 15.5	34.5 \pm 2.4	-0.085 \pm 0.1	62.4 \pm 4.2
	Dobutamine	488.7 \pm 30.4*	123.2 \pm 7.8*	37.7 \pm 3.5*	-0.085 \pm 0.04	70.3 \pm 5.3*
EUN	Baseline	556.9 \pm 29.4	108.1 \pm 6.2	32.0 \pm 2.6	-0.062 \pm 0.05	60.2 \pm 5.6
	Dobutamine	501.6 \pm 34.6*	120.4 \pm 8.6*	36.9 \pm 2.5*	-0.088 \pm 0.07	72.8 \pm 3.5*
LUN	Baseline	551.6 \pm 59.5	110.1 \pm 12.7	31.0 \pm 2.9	-0.09 \pm 0.06	62.8 \pm 5.1
	Dobutamine	526.3 \pm 54.3*	115.2 \pm 19.3*	35.3 \pm 4.2*	-0.12 \pm 0.04	77.2 \pm 5.3*
PUN	Baseline	534.8 \pm 68.4	114.2 \pm 15.2	33.0 \pm 3.4	-0.081 \pm 0.07	64.7 \pm 6.4
	Dobutamine	528.3 \pm 18.2	113.7 \pm 4.0*	39.1 \pm 3.4*	-0.11 \pm 0.04	80.1 \pm 6.3*

BPM, beats per minute.

ECG parameters obtained from lead II on control (CON), postnatal undernutrition (PUN), early phase of undernutrition (EUN) and late phase of undernutrition (LUN) groups at baseline and following dobutamine treatment.

*Significant ($P < 0.05$) change from baseline measurement to dobutamine treatment.

114.1 \pm 15.5 ms; EUN 108.1 \pm 6.2 ms; LUN 125.0 \pm 15.2 ms; PUN 114.2 \pm 15.2 ms). Electrical function of the ventricles at baseline showed that there was no difference ($P > 0.05$) in QRS duration between the four groups (CON 34.5 \pm 2.4 ms; EUN 32.0 \pm 2.6 ms, LUN 31.0 \pm 2.9 ms; PUN 33.0 \pm 3.4 ms). There was no significant ($P > 0.05$) difference in QT duration between any of the groups at baseline (CON QT duration 62.4 \pm 4.2 ms; EUN 60.2 \pm 5.6 ms; LUN 62.8 \pm 5.1 ms; PUN 64.7 \pm 6.4 ms). Additionally ST deflection at baseline was not significantly different ($P > 0.05$) between the diet groups (CON -0.085 \pm 0.1 V; EUN -0.062 \pm 0.05 V; LUN -0.090 \pm 0.06; PUN -0.081 \pm 0.07).

Dobutamine treatment

HR had a blunted decrease ($P = 0.0132$) in the PUN group with dobutamine treatment as compared with the other groups with a -1.2% change from baseline as compared with an -8.5% change in the CON mice, -9.9% change in the EUN and -4.6% change in the LUN. Corresponding to the HR the RR interval response with dobutamine treatment significantly increased ($P < 0.001$) in all groups. The PUN group had a 38% change in RR interval from baseline (120.9 \pm 25.9 ms) to dobutamine stimulation (166.9 \pm 22.6 ms). In comparison, the CON group had 11% change from baseline (123.9 \pm 6.1 ms) to dobutamine stimulation (137.6 \pm 5.4 ms). Treatment with dobutamine also significantly

increased the QRS duration ($P < 0.001$) for all groups. The PUN group had an 18.48% change from baseline (33.0 \pm 3.4 ms) to dobutamine stimulation (39.1 \pm 3.4 ms). In opposition, the CON group had 9.2% change from baseline (34.5 \pm 2.4 ms) to dobutamine stimulation (37.7 \pm 3.5 ms). The EUN and LUN groups recorded 15.3% (32.0 \pm 2.6 ms at baseline to 36.9 \pm 2.5 ms with dobutamine) and 13.8% change (31.0 \pm 2.9 ms at baseline to 35.3 \pm 4.2 ms with dobutamine) in QRS duration, respectively. As shown in Figure 4a trending interaction effect ($P = 0.09$) showed the PUN group had the longest QRS duration (39.1 \pm 3.4 ms), as compared with the other groups (CON 37.7 \pm 3.5 ms; EUN 36.9 \pm 2.5 ms; LUN 35.3 \pm 4.2).

Dobutamine treatment had a significant ($P = 0.01$) effect on QT duration (Fig. 5) with the PUN (80.1 \pm 6.3 ms) and LUN (77.2 \pm 5.3 ms) groups having a longer QT duration as compared with CON (70.3 \pm 5.3 ms) and EUN (72.8 \pm 3.5 ms) groups with dobutamine treatment. There was no difference in ST deflection following dobutamine treatment ($P > 0.05$).

Shape-based arrhythmia detection

There were three types of arrhythmias detected [wide QRS complex (25.0% of PUN baseline and 87.5% of PUN + dobutamine), 1st degree atrioventricular (AV) block (12.5% of EUN baseline and 12.5% of EUN + dobutamine) and atrial flutter




Fig. 4. QRS duration from the electrocardiogram at baseline and with administration of dobutamine. A trend ($P=0.09$) for increased QRS duration was observed in the postnatal undernutrition (PUN) group following dobutamine stimulation.




Fig. 5. QT duration from the electrocardiogram at baseline and following administration of dobutamine. *Significant ($P=0.01$) difference in QT duration in postnatal undernutrition (PUN) and late phase of undernutrition (LUN) groups following dobutamine stimulation as compared with the control (CON) and early phase of undernutrition (EUN) group.

(EUN baseline = 62.5%, EUN + dobutamine = 25.0%, LUN baseline = 62.5%, LUN + dobutamine = 62.5%, PUN baseline = 50.0% and PUN + dobutamine = 25.0%). The control group did not present any arrhythmias at baseline or with dobutamine stimulation (Fig. 6).

Discussion

The DOHaD hypothesis states that growth restriction in neonatal life permanently effects the body's structure and function, in ways that may lead to chronic disease in adult life.^{2–4,7} Professor David Barker first championed this hypothesis and the resulting influence on cardiovascular disease over 30 years ago. Since then the literature has documented the effect of neonatal nutrition on growth rate, organ size, body composition, along with clinical relevance on insulin sensitivity and hypertension.^{9,10,47} However, there is limited information on how undernutrition during various windows of postnatal life influences cardiac function. Thus, the purpose of this study was to determine whether early life undernutrition during key windows of development produced alterations in the depolarization and repolarization phases of the heart as measured by an ECG. Furthermore, we assessed the electrical activity of the heart at rest and when pharmacologically stressed with dobutamine, to determine if cardiac functional impairment was quiescent when the mouse was not in a stressed state.

We evaluated three-time points of postnatal undernutrition along with a control group. The EUN group experienced an initial deficit in nutrition from PN1 to PN11, the LUN group undergoing undernutrition from PN11 to PN21, and lastly, the PUN group enduring the lengthiest undernourishment period from PN1 to PN21, while the CON group did not experience undernutrition. Following the period of undernutrition the mice were nutritionally rehabilitated.

PN1–PN11 (EUN) is the cardiomyocyte proliferative phase of development in mice where mononucleated cells divide in order to increase cardiomyocyte number.^{11,48} During this period the electrical conductance system is continuing to mature as well.^{11,48} At the end of this period (PN7–10) cardiomyocytes undergo nuclear but not cellular division resulting in binucleated cardiomyocytes.^{11,48} The period of PN11–PN21 (LUN) is considered to be the hypertrophy phase of the cardiomyocytes, where the cells increase cross-sectional area in order to produce contractile force to overcome after load experienced by extra uterine life.⁴⁸ Thus, any nutritional insult, during these periods of development will have a lifelong effect, on the structure and function of the heart.

Postnatal growth failure occurs in a considerable majority of extremely low birth weight infants.^{33,34,49} Suboptimal nutrition during early life development has been shown to alter body composition including lean mass and increased adiposity.⁵⁰ Our nutritive model (Fig. 1) produced changes in fat and lean mass dependent on the window of development. LUN and PUN mice weighed less than CON and EUN, indicating that EUN mice experienced catch up growth, however the increase in body mass at PN80 was primarily due to an increase in fat mass (Fig. 2). The LUN group did not have increased fat deposition but displayed similar lean mass to the CON and EUN groups. The PUN group displayed a reduction in lean mass and fat mass as compared with the other groups. The PUN's inability to experience lean mass catch up growth could be due to a reduction in myogenic stem cell frequency⁵⁰ and a reduction in ribosomal abundance.³¹ The combination of these impairments would reduce muscle protein synthesis and reduce lean mass over the lifespan.

One aspect to the clinical significance of developmental programming is that in addition to a reduction in body mass there can be organ specific growth restriction,⁵¹ as evident by the fact that the longer the nutrient restriction the smaller the absolute heart weight at PN80 (Fig. 3b), however, when standardizing heart weight to body size the EUN group only displayed a significant impairment in heart weight (Fig. 3a). During PN1–10, when EUN was restricted, cardiomyocytes are proliferating thus undernutrition during that time period will reduce cell number and impair growth which cannot be recovered in the later phases of development with cardiomyocyte hypertrophy.^{24,25}

As the reduction in heart mass was a result of cardiomyocyte impairment it was hypothesized that there would be an associated functional impairment, which could manifest in the atria and the ventricles.

Atrial impairment

Under adrenergic stimulation the EUN group developed 1st degree AV block representing a slow conduction velocity in the electrical impulse being sent from the sinoatrial node to the atrioventricular node. Previous studies using a murine model have shown that 1st degree AV block will progress into 2nd degree AV block and left ventricular hypertrophy if hypertension is present.⁵² The DOHaD literature has consistently




Fig. 6. Electrocardiogram rhythm strips collect from EMKA Technologies Software. (a) Control (CON) group baseline measurement, with normal sinus rhythm. (b) CON group + dobutamine treatment, no arrhythmias detected. (c) Early phase of undernutrition (EUN) group baseline measurement, normal sinus rhythm with minimal artifact. (d) EUN group + dobutamine treatment, software detection of 1st degree atrioventricular Block. (e) late phase of undernutrition (LUN) group baseline measurement reveals atrial flutter. (f) LUN group + dobutamine treatment reveals atrial flutter. (g) postnatal undernutrition (PUN) group baseline measurement, no arrhythmias detected. (h) PUN group + dobutamine treatment, wide QRS complex detection.

shown that early life undernutrition leads to an increased incidence of hypertension.^{2–5,7}

When undernutrition was isolated to the hypertrophy phase of postnatal development (LUN) the ECG analysis revealed atrial flutter at rest and under beta-adrenergic stimulation. Atrial flutter is a condition in which the electrical and contractile properties of the atria are impaired and disorganized.⁵³ Previous literature has hypothesized that atrial flutter and fibrillation are caused by remodeling of the cardiac tissue, causing the cells to become frail and fibrotic,⁵⁴ with histological analysis of hearts from mice subjected to early life growth restriction having an increase in fibrotic tissue,^{24,25} while we did not perform histological analysis on these hearts the manifestation of atrial flutter could be a result of fibrotic tissue present. The development of atrial flutter if gone untreated can potentially lead to an increased risk of atrial fibrillation and stroke due to the increased risk of blood clot formation in the atria.^{55–59} It has been well documented in the literature that early life growth restriction increases the risk for stroke.¹² While the etiology of stroke risk from fetal origins may be independent of atrial fibrillation the fact that early life undernutrition also increases the risk of atrial flutter and fibrillation there is a further increase in the risk of stroke for individuals exposed to early life growth restriction.

It was observed in a small proportion of the EUN (three mice) and PUN (two mice) that with an injection of dobutamine there was atrial flutter withdrawal. A possible explanation for this is the dobutamine stimulation prolonged the refractory period and

slowed the conduction velocity thus preventing re-entry back into atrial flutter, acting as an anti-arrhythmic drug.⁶⁰

Ventricular impairment

Ventricular arrhythmias are generally accepted to have more clinical significance than atrial, due to the increased mortality risk.⁶¹ The PUN group presented with a healthy ECG tracing at baseline, however when stressed with dobutamine the PUN group displayed a widened QRS complex which represents a prolonged ventricular depolarization. We hypothesize that undernutrition in the late phase (PN11–21) of cardiomyocyte development impaired ventricular conduction as both the PUN and LUN groups had significantly longer QT durations when stimulated with dobutamine indicating a lengthened ventricular repolarization.

We did potentially observe a clinical condition in our PUN and LUN mice, termed long QT syndrome (Fig. 5). The pathology of long QT syndrome is characterized by impairment in ventricular repolarization leading to sudden cardiac death. The long QT syndrome is a family of arrhythmic disorders caused by dysfunction in one of 13 different ion channel complex proteins, including calcium, sodium and potassium channels.⁶² During the hypertrophy phase of cardiomyocyte development the sarcoplasmic reticulum matures in order to optimize calcium flux during depolarization,⁶³ thus early life nutrition during this phase could increase the risk of sudden cardiac death syndrome.

In conclusion, these findings suggest that undernourishment during the various phases (EUN, LUN and PUN) of cardiomyocyte

development causes functional and structural impairment. Most notably functional impairment only arose when the heart was stressed. Furthermore, our data indicate that undernutrition during these distinct windows of development results in specific arrhythmias present in adulthood. It was hypothesized that any arrhythmias present in the PUN group would be a combination of arrhythmias present in the EUN and LUN group. As this was not observed in our data, perhaps there is a mechanistic cascade associated with the timing of re-feed that result in the PUN group being different from EUN and LUN and not simply a combination of the two.

Acknowledgments. The authors thank Dr. Daniel Columbus from the Prairie Swine Research Center.

Author contributions. D.P.F. collected the data and co-wrote the manuscript. J.R.V. analyzed the data and co-wrote the manuscript.

Financial Support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflicts of Interest. None.

Ethical Standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national guidelines on the care and use of laboratory animals (Institutional Animal Care and Use Committee) and has been approved by Michigan State University.

References

- Mozaffarian D, Benjamin EJ, Go AS, *et al.* Heart disease and stroke statistics-2016 update: a report from the American Heart Association. *Circulation*. 2016; 133, e38–360.
- Barker DJ. The fetal origins of coronary heart disease. *Acta Paediatr Suppl*. 1997; 422, 78–82.
- Barker DJ. In utero programming of cardiovascular disease. *Theriogenology*. 2000; 53, 555–574.
- Barker DJ. The origins of the developmental origins theory. *J Intern Med*. 2007; 261, 412–417.
- Thornburg KL. The programming of cardiovascular disease. *J Dev Orig Health Dis*. 2015; 6, 366–376.
- Feltes BC, de Faria Poloni J, Bonatto D. The developmental aging and origins of health and disease hypotheses explained by different protein networks. *Biogerontology*. 2011; 12, 293–308.
- Barker DJ, Winter PD, Osmond C, Margetts B, Simmonds SJ. Weight in infancy and death from ischaemic heart disease. *Lancet*. 1989; 2, 984–985.
- Ritche MD, Loustalot F, Bowman BA, Hong Y. Trends in mortality rates by subtypes of heart disease in the United States, 2000–2010. *JAMA*. 2014; 312, 2037–2039.
- Fall CH, Barker DJ, Osmond C, *et al.* Relation of infant feeding to adult serum cholesterol concentration and death from ischaemic heart disease. *BMJ*. 1992; 304, 801–805.
- Wells JC. Environmental quality, developmental plasticity and the thrifty phenotype: a review of evolutionary models. *Evol Bioinform Online*. 2007; 3, 109–120.
- Louey S, Jonker SS, Giraud GD, Thornburg KL. Placental insufficiency decreases cell cycle activity and terminal maturation in fetal sheep cardiomyocytes. *J Physiol*. 2007; 580, 639–648.
- Morton M, Tsang H, Hohimer R, *et al.* Left ventricular size, output, and structure during guinea pig pregnancy. *Am J Physiol*. 1984; 246, R40–48.
- Black RE, Victoria CG, Walker SP, *et al.* Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013; 382, 427–451.
- Bubb KJ, Cock ML, Black MJ, *et al.* Intrauterine growth restriction delays cardiomyocyte maturation and alters coronary artery function in the fetal sheep. *J Physiol*. 2007; 578, 871–881.
- Ferguson DP, Dangott LJ, Schmitt EE, Vellers HL, Lightfoot JT. Differential skeletal muscle proteome of high- and low-active mice. *J Appl Physiol (1985)*. 2014; 116, 1057–1067.
- Taffet GE, Tate CA. CaATPase content is lower in cardiac sarcoplasmic reticulum isolated from old rats. *Am J Physiol*. 1993; 264, H1609–H1614.
- Taffet GE, Pham TT, Bick DL, *et al.* The calcium uptake of the rat heart sarcoplasmic reticulum is altered by dietary lipid. *J Membr Biol*. 1993; 131, 35–42.
- Baum M, Ortiz L, Quan A. Fetal origins of cardiovascular disease. *Curr Opin Pediatr*. 2003; 15, 166–170.
- Skilton MR, Phang M. From the α to the ω -3: breaking the link between impaired fetal growth and adult cardiovascular disease. *Nutrition*. 2016; 32, 725–731.
- Hannan RD, Jenkins A, Jenkins AK, Brandenburger Y. Cardiac hypertrophy: a matter of translation. *Clin Exp Pharmacol Physiol*. 2003; 30, 517–527.
- Fenske S, Probstle R, Auer F, *et al.* Comprehensive multilevel in vivo and in vitro analysis of heart rate fluctuations in mice by ECG telemetry and electrophysiology. *Nat Protoc*. 2016; 11, 61–86.
- SC W. A practical approach to using mice in atherosclerosis research. *Clin Biochem Rev*. 2004; 25, 81–93.
- Mongue-Din H, Salmon A, Fiszman MY, Fromes Y. Non-invasive restrained ECG recording in conscious small rodents: a new tool for cardiac electrical activity investigation. *Pflugers Arch*. 2007; 454, 165–171.
- Drenckhahn JD, Schwarz QP, Gray S, *et al.* Compensatory growth of healthy cardiac cells in the presence of diseased cells restores tissue homeostasis during heart development. *Dev Cell*. 2008; 15, 521–533.
- Drenckhahn JD, Strasen J, Heinecke K, *et al.* Impaired myocardial development resulting in neonatal cardiac hypoplasia alters postnatal growth and stress response in the heart. *Cardiovasc Res*. 2015; 106, 43–54.
- Boiten HJ, van Domburg RT, Geleijnse ML, *et al.* Cardiac stress imaging for the prediction of very long-term outcomes: dobutamine stress echocardiography or dobutamine (99m)Tc-sestamibi SPECT? *J Nucl Cardiol*. 2016; <https://doi.org/10.1007/s12350-016-0521-4>.
- Cortigiani L, Sorbo S, Miccoli M, *et al.* Prognostic value of cardiac power output to left ventricular mass in patients with left ventricular dysfunction and dobutamine stress echo negative by wall motion criteria. *Eur Heart J Cardiovasc Imaging*. 2016; 18, 153–158.
- Henri C, Piérard LA, Lancellotti P, *et al.* Exercise testing and stress imaging in valvular heart disease. *Can J Cardiol*. 2014; 30, 1012–1026.
- Kim MN, Kim SA, Kim YH, *et al.* Head to head comparison of stress echocardiography with exercise electrocardiography for the detection of coronary artery stenosis in women. *J Cardiovasc Ultrasound*. 2016; 24, 135–143.
- Standbridge K, Reyes E. The role of pharmacological stress testing in women. *J Nucl Cardiol*. 2016; 23, 997–1007.
- Fiorotto ML, Davis TA, Sosa HA, *et al.* Ribosome abundance regulates the recovery of skeletal muscle protein mass upon recuperation from postnatal undernutrition in mice. *J Physiol*. 2014; 592, 5269–5286.
- Reeves PG, Nielsen FH, Fahey GC Jr. AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J Nutr*. 1993; 123, 1939–1951.
- Sampson DA, Hunsaker HA, Jansen GR. Dietary protein quality, protein quantity and food intake: effects on lactation and on protein synthesis and tissue composition in mammary tissue and liver in rats. *J Nutr*. 1986; 116, 365–375.
- Grimble RF, Mansaray YK. Effects in rats of dietary protein inadequacy on lactose production, milk volume and components of the lactose synthetase complex (EC 2.4.1.22). *Ann Nutr Metab*. 1987; 31, 179–184.
- Crnich LS, Chase HP. Models of infantile undernutrition in rats: effects on milk. *J Nutr*. 1978; 108, 1755–1760.
- Grigor MR, Allan JE, Carrington JM, *et al.* Effect of dietary protein and food restriction on milk production and composition, maternal tissues and enzymes in lactating rats. *J Nutr*. 1987; 117, 1247–1258.
- Pine AP, Jessop NS, Oldham JD. Maternal protein reserves and their influence on lactational performance in rats. *J Nutr*. 1994; 71, 13–27.
- Mueller AJ, Cox WM Jr. The effect of changes in diet on the volume and composition of rat milk. *J Nutr*. 1946; 31, 249–259.

39. Lim K, Zimanyi MA, Black MJ. Effect of maternal protein restriction in rats on cardiac fibrosis and capillarization in adulthood. *Pediatr Res.* 2006; 60, 83–87.
40. Ferguson DP, Dangott LJ, Vellers HL, Schmitt EE, Lightfoot JT. Differential protein expression in the nucleus accumbens of high and low active mice. *Behav Brain Res.* 2015; 291, 283–288.
41. Berul CI, Aronovitz MJ, Wang PJ, Mendelsohn ME. In vivo cardiac electrophysiology studies in the mouse. *Circulation.* 1996; 94, 2641–2648.
42. Zwanenburg A, Jellema RK, Jennekens W, et al. Heart rate-mediated blood pressure control in preterm fetal sheep under normal and hypoxic-ischemic conditions. *Pediatr Res.* 2013; 73, 420–426.
43. Álvarez-García J, Vives-Borras M, Gomis P, et al. Electrophysiological effects of selective atrial coronary artery occlusion in humans. *Circulation.* 2016; 133, 2235–2242.
44. Han Y, Jing J, Tian F, et al. ST elevation acute myocardial infarction accelerates non-culprit coronary lesion atherosclerosis. *Int J Cardiovasc Imaging.* 2014; 30, 253–261.
45. Gouma E, Simos Y, Verginadis I, et al. A simple procedure for estimation of total body surface area and determination of a new value of Meeh's constant in rats. *Lab Anim.* 2012; 46, 40–45.
46. Byers SL, Wiles MV, Dunn SL, Taft RA. Mouse estrous cycle identification tool and images. *PLoS One.* 2012; 7, e35538.
47. Matthews PA, Sameulsson AM, Seed P, et al. Fostering in mice induces cardiovascular and metabolic dysfunction in adulthood. *J Physiol.* 2011; 589, 3869–3881.
48. Botting KJ, Wang KC, Padhee M, et al. Early origins of heart disease: low birth weight and determinants of cardiomyocyte endowment. *Clin Exp Pharmacol Physiol.* 2012; 39, 814–823.
49. Dusick AM, Poindexter BB, Ehrenkranz RA, Lemons JA. Growth failure in the preterm infant: can we catch up? *Semin Perinatol.* 2003; 27, 302–310.
50. Woo M, Isganaitis E, Fitzpatrick C, et al. Early life nutrition modulates muscle stem cell number: implications for muscle mass and repair. *Stem Cells Dev.* 2011; 20, 1763–1769.
51. Wang J, Li D, Dangott LJ, Wu G. Proteomics and its role in nutrition research. *J Nutr.* 2006; 136, 1759–1762.
52. Wei L, Taffet GE, Khoury DS, et al. Disruption of Rho signaling results in progressive atrioventricular conduction defects while ventricular function remains preserved. *FASEB.* 2004; 18, 857–859.
53. Yagi S, Akaike M, Aihara K, et al. Endothelial nitric oxide synthase-independent protective action of statin against angiotensin II-induced atrial remodeling via reduced oxidant injury. *Hypertension.* 2010; 55, 918–923.
54. Jansen HJ, Moghtadaei M, Mackasey M, et al. Atrial structure, function and arrhythmogenesis in aged and frail mice. *Sci Rep.* 2017; 7, 44336.
55. Allesie MA, Konings K, Kirchhof CJ, Wijffels M. Electrophysiologic mechanisms of perpetuation of atrial fibrillation. *Am J Cardiol.* 1996; 77, 10A–23A.
56. Devkota A, Bakhit A, Dufresne A, et al. Arrhythmias and electrocardiographic changes in systolic heart failure. *N Am J Med Sci.* 2017; 8, 171–174.
57. Friedrichs GS. Experimental models of atrial fibrillation/flutter. *J Pharmacol Toxicol Methods.* 2000; 43, 117–123.
58. Ozcan C, Battaglia E, Young R, Suzuki G. LKB1 knockout mouse develops spontaneous atrial fibrillation and provides mechanistic insights into human disease process. *J Am Heart Assoc.* 2015; 4, e001733.
59. Zhang Q, Timofeyev V, Lu L, et al. Functional roles of a Ca²⁺-activated K⁺ channel in atrioventricular nodes. *Circ Res.* 2008; 102, 465–471.
60. van der Hooft CS, Heeringa J, van Herpen G, et al. Drug-induced atrial fibrillation. *J Am Coll Cardiol.* 2004; 44, 2117–2124.
61. Velasco A, Stirrup Reyes E, Hage FG. Guidelines in review: Comparison between AHA/ACC and ESC guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *J Nucl Cardiol.* 2017; 24, 1893–1901.
62. Iyer V, Roman-Campos D, Sampson KJ, et al. Purkinje cells as sources of arrhythmias in long QT syndrome type 3. *Sci Rep.* 2017; 5, 13287.
63. Hamaguchi S, Kawakami Y, Honda Y, et al. Developmental changes in excitation-contraction mechanisms of the mouse ventricular myocardium as revealed by functional and confocal imaging analyses. *J Pharmacol Sci.* 2013; 123, 167–175.