

Original Article

*Joint First Authors.

Cite this article: Haining K *et al* (2020). Neuropsychological deficits in participants at clinical high risk for psychosis recruited from the community: relationships to functioning and clinical symptoms. *Psychological Medicine* 50, 77–85. <https://doi.org/10.1017/S0033291718003975>

Received: 17 August 2018
Revised: 9 October 2018
Accepted: 5 December 2018
First published online: 13 March 2019

Key words:

Clinical high-risk; functioning; neurocognition; prevention; psychosis

Author for correspondence:

Peter J. Uhlhaas, E-mail: peter.uhlhaas@glasgow.ac.uk

Neuropsychological deficits in participants at clinical high risk for psychosis recruited from the community: relationships to functioning and clinical symptoms

Kate Haining^{1,*}, Claire Matrunola^{2,*}, Lucy Mitchell¹, Ruchika Gajwani², Joachim Gross^{1,3}, Andrew I. Gumley³, Stephen M. Lawrie⁴, Matthias Schwannauer⁵, Frauke Schultze-Lutter⁶ and Peter J. Uhlhaas¹

¹Institute for Neuroscience and Psychology, University of Glasgow, Glasgow, UK; ²Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK; ³Institute of Biomagnetism and Biosignalanalysis, Westphalian Wilhelms University Muenster, Muenster, Germany; ⁴Department of Psychiatry, University of Edinburgh, Edinburgh, UK; ⁵Department of Clinical Psychology, University of Edinburgh, Edinburgh, UK and ⁶Department of Psychiatry and Psychotherapy, Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

Abstract

Background. The current study examined the pattern of neurocognitive impairments in a community-recruited sample of clinical high-risk (CHR) participants and established relationships with psychosocial functioning.

Methods. CHR-participants ($n = 108$), participants who did not fulfil CHR-criteria (CHR-negatives) ($n = 42$) as well as a group of healthy controls (HCs) ($n = 55$) were recruited. CHR-status was assessed using the Comprehensive Assessment of At-Risk Mental States (CAARMS) and the Schizophrenia Proneness Instrument, Adult Version (SPI-A). The Brief Assessment of Cognition in Schizophrenia Battery (BACS) as well as tests for emotion recognition, working memory and attention were administered. In addition, role and social functioning as well as premorbid adjustment were assessed.

Results. CHR-participants were significantly impaired on the Symbol-Coding and Token-Motor task and showed a reduction in total BACS-scores. Moreover, CHR-participants were characterised by prolonged response times (RTs) in emotion recognition as well as by reductions in both social and role functioning, GAF and premorbid adjustments compared with HCs. Neurocognitive impairments in emotion recognition accuracy, emotion recognition RT, processing speed and motor speed were associated with several aspects of functioning explaining between 4% and 12% of the variance.

Conclusion. The current data obtained from a community sample of CHR-participants highlight the importance of dysfunctions in motor and processing speed and emotion recognition RT. Moreover, these deficits were found to be related to global, social and role functioning, suggesting that neurocognitive impairments are an important aspect of sub-threshold psychotic experiences and a possible target for therapeutic interventions.

Background

Cognitive deficits are a core feature of schizophrenia (ScZ) and have been found in the domains of working memory (WM), verbal learning, motor abilities, attention, processing speed and social cognition (Green *et al.*, 2004). There is substantial evidence that neurocognitive and social cognitive impairments in ScZ are associated with poor occupational and social outcomes (Green *et al.*, 2000; Hooker and Park, 2002; Fett *et al.*, 2011), making them a potential target for therapeutic interventions.

More recently, one focus has been the identification of neurocognitive impairments in participants meeting clinical high-risk criteria (CHR) for the development of psychosis (Klosterkötter *et al.*, 2001; Yung *et al.*, 2005). These include ultra-high-risk (UHR) criteria that involve the presence of attenuated, psychotic symptoms (Miller *et al.*, 2003; Yung *et al.*, 2005). Moreover, UHR-criteria include a genetic risk plus functional deterioration syndrome as well as brief limited intermittent psychotic symptoms (BLIPs).

In addition, CHR-criteria have been developed based on the basic symptom (BS) concept proposed by Huber and colleagues (Schultze-Lutter, *et al.*, 2008). BS involve the presence of self-experienced perceptual and cognitive anomalies that are thought to represent the earliest manifestation of psychosis risk (Schultze-Lutter *et al.*, 2008). CHR-criteria confer a 10–30% risk of developing ScZ within a 2–5 year period (Fusar-Poli *et al.*, 2013, 2015a, 2015b).

More recent studies have shown that the combined presence of both BS- and UHR-criteria increases the predictive power significantly (Schultze-Lutter *et al.*, 2014).

There is extensive evidence on the presence of neurocognitive deficits in CHR-populations across a range of domains that mirror observations in established ScZ, including impairments in WM, attention, speed of processing, verbal memory, verbal fluency, executive functions and motor speed with small-to-medium effect sizes (Fusar-Poli *et al.*, 2012; Giuliano *et al.*, 2012; Bora *et al.*, 2014). Follow-up studies have suggested that certain deficits may indicate stable vulnerability markers, e.g. sustained attention (Francey *et al.*, 2005), whereas others may be predictive of transition to psychosis, such as verbal IQ, processing speed, verbal memory and WM (Brewer *et al.*, 2005; Lencz *et al.*, 2006; Pukrop and Klosterkötter, 2010; Seidman *et al.*, 2010; Michel *et al.*, 2014).

Moreover, previous studies have found deficits in emotion recognition, theory of mind and social perception in CHR-participants (Thompson *et al.*, 2011) in agreement with extensive evidence for dysfunctions in social cognition in ScZ-patients (Green *et al.*, 2015). More specifically, impaired facial emotion recognition in CHR-groups has been reported in several studies (Addington *et al.*, 2008a, 2008b; van Rijn *et al.*, 2011; Amminger *et al.*, 2012), suggesting that emotion recognition deficits may emerge before the onset of psychosis.

The current study aimed to extend these findings by examining the relationship between neurocognition, social cognition and current psychosocial functioning in a CHR-sample recruited from the general community. The large majority of studies investigating neurocognition in CHR-populations involve participants who are help-seeking and recruited through clinical pathways. Accordingly, it is unclear to what extent neurocognitive deficits generalise to more representative samples recruited outside clinical pathways. This is potentially an important question as there may be differences between clinically referred *v.* community CHR-samples, for example, regarding transition rates (Fusar-Poli *et al.*, 2015b).

To address this issue, we recruited a sample of $n = 108$ CHR-participants through an online-screening platform (McDonald *et al.*, 2018) as well as a group of $n = 42$ participants who did not fulfil CHR-criteria (CHR-negatives) but were characterised by psychiatric comorbidities, such as affective disorders and substance abuse, and a group of $n = 55$ healthy controls (HCs). Neurocognition was assessed with the Brief Assessment of Cognition in Schizophrenia Battery (BACS) (Keefe *et al.*, 2004) as well as tasks from the Penn Computerized Neurocognitive Battery (CNB) (Moore *et al.*, 2015). The Global Assessment of Functioning (GAF) as well as scales for role (GF: Role) and social (GF: Social) functioning (Cornblatt *et al.*, 2007) were used to assess psychosocial functioning.

A secondary objective was to examine the relationship between neurocognitive deficits and social and occupational functioning in community-recruited CHR-participants. Previous studies reported conflicting findings on this relationship in CHR-participants recruited from clinical pathways. Niendam *et al.* (2006) reported that impairments in verbal learning and memory were associated with current social functioning. A follow-up study found that improvements in social functioning predicted gains in processing speed and visual learning and memory (Niendam *et al.*, 2007). Similar findings were reported by Lin *et al.* (2011). However, findings by Jahshan *et al.* (2010) indicated that improvements in neurocognitive performance were not

significantly associated with functioning as measured by the GAF scale. Finally, Carrión *et al.* (2011) examined impairments in both social and role functioning in relation to neurocognitive performance and found that speed of processing was predictive of poorer social and role functioning.

Methods

Recruitment and participants

The YouR-Study is a longitudinal study to identify neurobiological and psychological mechanisms and predictors of psychosis-risk (Uhlhaas *et al.*, 2017) and is funded by the Medical Research Council (MRC).

CHR-participants were recruited through an online-screening approach (see <http://www.your-study.org.uk>) that identified CHR-participants from the general population through email invitations, posters and flyers over a 4-year period (see McDonald *et al.*, 2018). Specifically, email invitations were sent out to colleges and universities in Glasgow and Edinburgh through which the majority of study participants were identified. It is estimated that ~100 000 participants were invited to the study.

Approximately 2800 participants filled out the online versions of the (a) the 16-item Prodromal Questionnaire (PQ-16) (Ising *et al.*, 2012) and (b) a nine-item scale of perceptual and cognitive anomalies (PCA) that was developed to assess BS. Participants were invited for clinical interviews if they positively endorsed six or more items on the PQ-16 or three or more on the PCA.

Previous analysis (McDonald *et al.*, 2018) had shown that ~50% participants fulfilled the PQ-16 cut-off criteria while ~70% met criteria for the PCA. Out of the ~2800 of participants who met online cut-offs, ~20% took part in clinical assessments. Moreover, an additional sample of $n = 21$ participants meeting first-episode criteria were identified.

To establish CHR-criteria, the positive scale of the Comprehensive Assessment of At-Risk Mental States (CAARMS) (Yung *et al.*, 2005) and items of the Schizophrenia Proneness Instrument (SPI-A) (Schultze-Lutter, *et al.*, 2007) as defined by Cognitive-Perceptive Basic Symptoms (COPER) and Cognitive Disturbances (COGDIS) were administered through trained research assistants and M.Sc./Ph.D. level researchers. Inter-rater reliability of CHR-status as determined by the CAARMS and SPI-A ratings was assessed over 18 sessions, reaching good-to-excellent reliability (CAARMS: 92%; SPI-A: 95.7%).

CHR-participants were excluded for current or past diagnosis with Axis I psychotic disorders. Other co-morbid Axis I diagnoses, such as mood or anxiety disorders, were not exclusionary and all participants were between 16 and 35 years of age (for more details, see Uhlhaas *et al.*, 2017).

Participants were recruited into the CHR-group if they met (a) SPI-A COGDIS/COPER-criteria; (b) CAARMS criteria for the attenuated psychosis group (subthreshold psychotic syndrome present in the last year without a decline in functioning); (c) CAARMS criteria for genetic risk plus functional deterioration (family history of psychosis plus a 30% drop in GAF); and (d) CAARMS criteria for the BLIPs group (BLIPs).

Moreover, the M.I.N.I. International Neuropsychiatric Interview (M.I.N.I. 6.0) (Sheehan *et al.*, 1998), the scales for premorbid adjustment (Cannon-Spoor *et al.*, 1982) and social and role functioning scales (Cornblatt, *et al.*, 2007) were administered. Neuropsychological assessment consisted of the BACS (Keefe *et al.*, 2004) as well as three tasks from the CNB battery (Moore

et al., 2015): (a) the Continuous Performance Test, (b) the N-Back Task and (c) the Emotion Recognition Task.

In addition to CHR-participants, two samples were recruited consisting of (1) participants who entered the study similar to CHR-participants but who did not meet CHR-criteria (CHR-negative) and (2) a group of HCs without an Axis I diagnosis or family history of psychotic disorders. The former group was included to assess the impact of psychiatric comorbidity, such as affective disorders and substance abuse, on neurocognitive parameters.

Statistical analyses

All statistical analyses were performed using SPSS version 24. BACS and CNB raw test scores for each neurocognitive domain were standardised by creating *z*-scores using the means and standard deviations of HCs. BACS raw scores were additionally corrected for gender. When the homogeneity of variances assumption was violated in one-way analysis of variance (ANOVA) analyses, Welch's *F* was reported. Since the one-way ANOVA is considered a robust test against the normality assumption, no alternative tests were applied. The Hochberg's GT2 test was used as a post hoc test for ANOVA analyses whereas the Games–Howell test was used as a post hoc test for Welch analyses. For Kruskal–Wallis *H* tests, Dunn's pairwise tests were carried out post hoc.

All BACS and CNB neurocognitive domains were entered into stepwise multiple linear regressions in order to assess the relationship between functioning, neurocognition and psychopathology in the CHR group.

Results

Sample characteristics

Baseline demographic and clinical characteristics of the three groups are summarised in Table 1.

CHRs, CHR-negatives and HCs did not differ significantly on age, gender or years of education. The CHR-group had significantly higher CAARMS-positive severity scores, poorer premorbid adjustment, lower GAF scores as well as reduced role and social functioning compared with HCs and CHR-negatives. Significant differences between groups were also found for medication status with 49.1% of CHR-participants receiving current medication. The CHR-group was also characterised by extensive psychiatric comorbidity, in particular with affective disorders. Moreover, differences in CHR-subgroups [UHR ($n = 34$), BS ($n = 29$), UHR/BS ($n = 45$)] were explored (online Supplementary Table S1). The BS group had significantly higher GAF scores and lower CAARMS-positive severity scores than the UHR/BS group.

Neuropsychology

Table 2 summarises the neurocognitive performance for CHRs, CHR-negatives and HCs. Due to incorrect task performance, one CHR participant was removed from the CNB WM accuracy and WM RT analysis, and one CHR-negative participant was removed from the CNB attention accuracy analysis.

Significant group effects were demonstrated for motor speed [$F_{(2,202)} = 8.48$, $p < 0.001$], BACS composite [$F_{(2,105)} = 3.44$, $p < 0.05$], emotion recognition RT [$F_{(2,105)} = 3.74$, $p < 0.05$] and processing speed [$F_{(2,202)} = 4.23$, $p < 0.05$]. These effects were

observed between CHRs and controls for all domains apart from processing speed where CHRs significantly differed only from CHR-negatives. Fig. 1 displays the effect sizes for each neurocognitive domain for both CHRs and CHR-negatives.

In the CHR-group, motor speed had the largest effect size (Cohen's $d = 0.63$). A small-to-medium effect size was found for emotion recognition RT ($d = 0.37$), processing speed ($d = 0.35$), BACS composite ($d = 0.35$), attention accuracy ($d = 0.28$) and WM accuracy ($d = 0.23$). In the CHR-negative group, a small-to-medium effect size was found for motor speed ($d = 0.43$), verbal fluency ($d = 0.29$) and attention RT ($d = 0.24$).

Furthermore, analysis was carried out to explore recognition of specific emotion categories (online Supplementary Table S2). CHR-participants were significantly slower in their RTs compared with HCs for recognising happy faces [$F_{(2,102)} = 6.90$, $p < 0.01$; $d = 0.46$]. No additional emotion recognition deficits emerged.

We also examined differences in neurocognition in relation to CHR-subgroups (online Supplementary Table S1). There was a significant difference between groups on motor speed [$F_{(3,159)} = 5.47$, $p < 0.01$], while a trend was observed for emotion recognition RT [$F_{(3,74)} = 2.72$, $p = 0.05$], BACS composite [$F_{(3,72)} = 2.30$, $p = 0.09$] and attention RT [$F_{(3,159)} = 2.28$, $p = 0.08$]. CHR-participants in the UHR and UHR/BS groups had significantly slower motor speed than HCs. Individuals in the UHR/BS groups also had significantly slower emotion recognition RTs than HCs ($p = 0.046$). No post-hoc differences were found for BACS composite or attention RT. CHR subgroup effect sizes for each neurocognitive domain are reported in online Supplementary Fig. S1.

Cognition, psychopathology and functioning

Stepwise multiple linear regressions were performed to assess the relationship between functioning, neurocognition and psychopathology in the CHR-group (Tables 3–4). All BACS and CNB neurocognitive domains were included in the regression. Motor speed significantly predicted GAF, accounting for 4% of the variance while emotion recognition RT explained 5% of the variance in CAARMS-positive severity scores. Emotion recognition RT together with emotion recognition accuracy and processing speed significantly predicted social functioning, accounting for 11% of the variance while processing speed alone significantly predicted role functioning, explaining 5% of the variance.

Fear RT was found to be a significant predictor for both GAF and social functioning, accounting for 4% and 10% of the variance, respectively, and together with anger RT, fear RT significantly predicted role functioning, accounting for 12% of the variance. Happy RT significantly predicted CAARMS-positive severity scores, accounting for 10% of the variance.

Discussion

The current study examined neurocognition and its relationship to functioning in a sample of CHR-participants recruited from the general community. Deficits in neurocognition are a hallmark of ScZ (Heinrichs and Zakzanis, 1998; Rajji *et al.*, 2009) and have been observed in CHR-participants across a number of domains with small-to-medium effect sizes (Fusar-Poli *et al.*, 2012; Giuliano *et al.*, 2012; Bora *et al.*, 2014). Importantly, there is evidence to suggest that impairments in neurocognition impact on psychosocial functioning in CHR-participants (Niendam *et al.*,

Table 1. Baseline demographic and clinical characteristics of CHR, HC and CHR-negative participants

Characteristic	CHRs (<i>N</i> = 108)	HCs (<i>N</i> = 55)	CHR-Ns (<i>N</i> = 42)	df	<i>F</i> / χ^2 / <i>H</i>	<i>p</i>	Post hoc contrasts
Age (years), <i>M</i> ± s.d.	21.85 ± 4.33	22.31 ± 3.39	23.24 ± 5.00	2, 97	<i>F</i> = 1.27	0.29	
Gender, <i>N</i> female (%)	82 (75.9)	37 (63.7)	28 (66.7)	2	χ^2 = 2.01	0.37	
Years of education, <i>M</i> ± s.d.	15.50 ± 3.13	16.38 ± 2.84	16.57 ± 3.62	2, 202	<i>F</i> = 2.29	0.10	
GAF, median (range)	59.50 (21–95)	88 (67–97)	70 (43–94)	2	<i>H</i> = 105.13	<0.001	CHR v. CHR-N v. HC
CAARMS-positive severity, median (range)	28.50 (0–72)	0 (0–12)	5 (0–24)	2	<i>H</i> = 129.41	<0.001	CHR v. CHR-N v. HC
GF: social, median (range)	8 (5–10)	9 (8–10)	8 (6–9)	2	<i>H</i> = 64.44	<0.001	CHR v. HC v. CHR-N
GF: role, median (range)	8 (5–9)	9 (5–9)	8 (5–9)	2	<i>H</i> = 45.05	<0.001	CHR v. HC v. CHR-N
PAS, median (range)							
Childhood	0.11 (0–0.57)	0.04 (0–0.21)	0.07 (0–0.46)	2	<i>H</i> = 25.92	<0.001	CHR v. HC
Early adolescence	0.17 (0–0.54)	0.06 (0–0.23)	0.11 (0–0.46)	2	<i>H</i> = 42.51	<0.001	HC v. CHR, CHR-N
Late adolescence	0.14 (0–0.57)	0.06 (0–0.29)	0.11 (0–0.71)	2	<i>H</i> = 27.41	<0.001	HC v. CHR, CHR-N
Medication, <i>N</i> (%)	53 (49.1)	1 (1.8)	19 (45.2)	10	χ^2 = 45.49	<0.001	
Anti-psychotic	0 (0)	0 (0)	1 (2.4)				
Mood stabiliser	1 (0.9)	0 (0)	0 (0)				
Anti-depressant	23 (21.3)	0 (0)	10 (23.8)				
Other	13 (12.0)	1 (1.8)	6 (14.3)				
Multiple	16 (14.8)	0 (0)	2 (4.7)				
Diagnosis, <i>N</i> (%)	97 (89.8)	3 (5.45)	26 (61.9)	2	χ^2 = 109.5	<0.001	
Anxiety disorders	80 (74.1)	0 (0)	19 (45.2)				
Mood disorders	67 (62.0)	0 (0)	12 (28.6)				
Eating disorders	11 (10.2)	0 (0)	1 (2.4)				
Suicide risk	57 (52.8)	1 (1.8)	10 (23.8)				
Alcohol dependence/abuse	31 (28.7)	2 (3.6)	9 (21.4)				
Substance dependence/abuse	13 (12.0)	0 (0)	1 (2.4)				

CHR, clinical high-risk; HC, healthy control; CHR-N, clinical high-risk-negative.

2006, 2007; Carrión *et al.*, 2011; Lin *et al.*, 2011). However, it is unclear to what extent these findings generalise to CHR-samples recruited from the general community.

Recent evidence has highlighted the importance of studying CHR-populations outside clinical referral pathways to identify the similarities and differences in clinical characteristics, demographic variables and neurocognition (Mills *et al.*, 2017; Schultze-Lutter *et al.*, 2018). Overall, our sample of CHR-participants recruited through a novel online screening platform (McDonald *et al.*, 2018) was characterised by similar levels of functioning and psychiatric comorbidity as previously observed in cohorts recruited through early intervention centres.

However, with regard to the pattern of neurocognitive deficits, there were differences and similarities with previous studies. We observed neurocognitive impairments that are consistent with a large body of work that has highlighted neurocognitive deficits in CHR-samples with mild-to-moderate effect sizes (Fusar-Poli *et al.*, 2012; Giuliano *et al.*, 2012; Bora and Murray, 2014). However, there were also certain differences to previous data, particularly with regard to the extent of dysfunctions in neuropsychological variables (see online Supplementary Fig. S2). Specifically, we observed that the neurocognitive domains that

were most prominently impaired were processing and motor speed.

The symbol-coding task has been consistently shown to be impaired in ScZ-patients with large effect sizes (Dickinson *et al.*, 2007). Moreover, it discriminates between CHR and controls (Seidman *et al.*, 2010; Fusar-Poli *et al.*, 2012) and predicts psychosis onset in CHR-individuals (Pukrop and Klosterkötter, 2010; Michel *et al.*, 2014). In the current study, we observed that CHR-participants showed a similar deficit that was associated with an effect size of $d = 0.35$. Interestingly, processing speed was largely intact in the CHR-negative group (effect size: $d < 0.1$), highlighting that the symbol-coding task may delineate specific cognitive impairments associated with psychosis risk.

In addition, CHR-participants were characterised by pronounced impairments in motor speed. While abnormalities in the motor system that involve psychomotor slowing are considered a core feature of ScZ (Morrens *et al.*, 2006), alterations in the motor system in CHR-participants are only recently being investigated. Evidence suggests that youths who later develop a ScZ-spectrum disorder have been reported to show poorer motor function in childhood (Dickson *et al.*, 2012) and abnormal involuntary movements were linked to CHR symptoms in a child

Table 2. Neurocognitive performance of CHR, HC and CHR-N participants

Domain	CHR (N = 108)		HCs (N = 55)		CHR-Ns (N = 42)		df	F	p	Cohen's d	Post hoc contrasts
	M	s.d.	M	s.d.	M	s.d.					
BACS											
Verbal memory	-0.07	1.19	0	1	0.18	1.07	2, 202	0.77	0.47	0.11	
Motor speed	-0.71	1.13	0	1	-0.40	0.92	2, 202	8.48	<0.001	0.63	HC v. CHR
Processing speed	-0.39	1.11	0	1	0.11	1.16	2, 202	4.23	<0.05	0.35	CHR v. CHR-N
Verbal fluency	-0.11	0.96	0	1	-0.26	0.81	2, 202	0.94	0.39	0.11	
Executive function	-0.07	1.26	0	1	-0.03	1.19	2, 202	0.05	0.95	0.06	
Working memory	0.01	1.39	0	1	0.27	1.18	2, 105	0.73	0.48	0.00	
BACS composite	-0.49	1.58	0	1	-0.05	1.33	2, 105	3.56	<0.05	0.35	HC v. CHR
CNB											
Emotion recognition accuracy	-0.07	0.99	0	1	-0.16	0.94	2, 202	0.31	0.73	0.06	
Emotion recognition RT	0.57	1.65	0	1	0.21	1.39	2, 105	3.74	<0.05	0.37	HC v. CHR
Attention accuracy	-0.69	2.83	0	1	-0.28	2.82	2, 201	1.50	0.23	0.28	
Attention RT	-0.16	0.84	0	1	-0.24	1.00	2, 202	0.90	0.41	0.17	
Working memory accuracy	-0.31	1.45	0	1	-0.21	1.28	2, 104	1.32	0.27	0.23	
Working memory RT	-0.07	0.76	0	1	-0.09	1.03	2, 201	0.14	0.87	0.06	

CHR, clinical high-risk; HC, healthy control; CHR-N, clinical high-risk-negative; RT, response times. CHR effect sizes, measured by Cohen's d, are classified as small (0.2), medium (0.5) and large (0.8).




Fig. 1. CHR and CHR-negative effect sizes, as measured by Cohen's d, for BACS and CNB data: classified as small (0.2), medium (0.5) and large (0.8). Error bars indicate standard errors of the mean. Positive values indicate impaired performance while negative values indicate better performance compared with HCs.

and adolescent community sample (Kindler *et al.*, 2016). These findings are consistent with reduced motor speed, dexterity and movement abnormalities in CHR-populations (e.g. Niendam *et al.*, 2006; Carrion *et al.*, 2011; Fusar-Poli *et al.*, 2012; Bora *et al.*, 2014; Dean and Mittal, 2015; Dean *et al.*, 2016). However, in contrast to the symbol-coding task, impairments in motor speed were also present in the CHR-negative group (effect size: $d = 0.4$), suggesting that psychomotor-slowness may be related to aspects of general psychopathology rather than psychosis risk *per se*.

In addition to impaired motor and processing speed, we also observed slower RTs during emotion recognition, while the

accuracy of emotion recognition was intact, highlighting the importance of reduced processing speed across different domains of functioning. Emotion recognition deficits have been reported in some CHR studies (Addington *et al.*, 2008a, 2008b; van Rijn *et al.*, 2011; Amminger *et al.*, 2012) while others have found emotion recognition to be intact (Pinkham *et al.*, 2007; Seiferth *et al.*, 2008; Gee *et al.*, 2012). There is also preliminary evidence for the possibility of emotion recognition deficits as a predictor for transition to psychosis (Allott *et al.*, 2014).

Interestingly, other domains of neurocognition that were found to be impaired in previous studies were not replicated in our community-recruited CHR-group. Verbal memory, for

Table 3. Linear regression results for the effects of neurocognitive performance on clinical characteristics at baseline in CHR participants

Variable	<i>B</i>	s.e.	β	R^2	<i>F</i>	<i>p</i>
GAF						
Motor speed	3.04	1.04	0.20	0.04	8.56	<0.01
CAARMS-positive severity						
Emotion recognition RT	2.74	0.86	0.22	0.05	10.10	<0.01
Social functioning						
Emotion recognition RT	-0.14	0.05	-0.20			
Emotion recognition accuracy	0.14	0.07	0.13	0.11	8.38	<0.001
Processing speed	0.14	0.06	0.15			
Role functioning						
Processing speed	0.20	0.06	0.22	0.05	10.05	<0.01

RT, response times.

Table 4. Linear regression results for the effects of emotion specific RTs on clinical characteristics at baseline in CHR participants

Variable	<i>B</i>	s.e.	β	R^2	<i>F</i>	<i>p</i>
GAF						
Fear RT	-2.01	0.73	-0.19	0.04	7.68	<0.01
CAARMS-positive severity						
Happy RT	4.24	0.92	0.31	0.10	21.38	<0.001
Social functioning						
Fear RT	-0.21	0.04	-0.32	0.10	23.55	<0.001
Role functioning						
Fear RT	-0.26	0.05	-0.40	0.12	13.31	<0.001
Anger RT	0.18	0.07	0.19			

RT, response times.

example, which has been associated with medium effect sizes in CHR-populations (Fusar-Poli *et al.*, 2012), was relatively intact in the current study. Previous reports have found verbal fluency and memory to be associated with subsequent transition to psychosis (Fusar-Poli *et al.*, 2012). Moreover, there is evidence to suggest that poorer verbal memory predicts more rapid transitioning (Seidman *et al.*, 2010).

The current study could not replicate impaired memory, executive function and attention in our CHR-group. Evidence is emerging of deficits in declarative memory in FEP (Mesholam-Gately *et al.*, 2009) and in CHR populations (Seidman *et al.*, 2016). The domain of attention has been argued to represent a stable vulnerability marker in CHR-populations (e.g. Francey *et al.*, 2005). More recent data from the NAPLS-2 cohort has demonstrated impairments in WM and attention in CHR-participants who later transitioned to psychosis relative to CHR-participants who did not transition (Seidman *et al.*, 2016).

Finally, executive functions have been found to be impaired in CHR samples (Lencz *et al.*, 2006; Carrión *et al.*, 2011; Fusar-Poli *et al.*, 2012; Seidman *et al.*, 2016). A meta-analysis found executive functioning, along with domains of memory and attention, to be the most consistently impaired and already established at the time of the FEP (Mesholam-Gately *et al.*, 2009).

Our data show that there are subtle differences between neurocognition and functioning levels in CHR-subgroups. Current evidence suggests that self-experienced BS represent the earliest manifestation of psychosis risk or an early prodromal state (EPS) while positive symptoms constitute coping mechanisms that emerge later during development (late prodromal state, LPS) (Fusar-Poli *et al.*, 2013). Consistent with this notion, we observed that CHR-participants who met UHR-criteria and UHR/BS-criteria had more pronounced cognitive impairments, in particular in motor speed, compared with the BS only group. This is consistent with previous findings that neurocognitive impairments differentiate EPS from the LPS-participants. Frommann *et al.* (2010) found individuals in the LPS to be impaired across all domains, while those in the EPS showed a specific deficit in the executive control/processing speed domain, raising the question of potentially progressive impairments in cognition across the at-risk phase. Alternatively, it has been suggested that BS criteria help to identify a more homogenous group with respect to neurocognitive profiles (Simon *et al.*, 2006).

Our data also support previous findings that deficits in neurocognition impact on functioning parameters in CHR-participants (Niendam *et al.*, 2006, 2007; Carrión *et al.*, 2011; Lin, *et al.*, 2011). Consistent with previous findings that highlighted that reduced

processing speed is an important determinant of functioning (Carrion *et al.*, 2011), our data suggest that impaired processing speed significantly correlates with role and social functioning. Emotion recognition RT, emotion recognition accuracy and processing speed combined explained 11% of the variance in social functioning while processing speed alone accounted for 5% of the variance in role functioning in our CHR-sample. In addition, we found that emotion recognition RT explained 5% of the variance in CAARMS-positive severity scores, while motor speed alone explains 4% of the variance in global functioning.

While these data replicate previous findings (e.g. Carrion *et al.*, 2011) and highlight the importance of processing speed for explaining psychosocial functioning, the relatively low amount of variance that is being accounted for also suggests that other factors are involved in contributing towards impaired functioning in CHR-participants. Given the importance of psychosocial functioning as an outcome parameter in CHR-populations, further studies need to address the contribution of other factors that could potentially allow insights into origin and mechanism(s) of impaired role and social functioning in CHR-participants.

Limitations

The current study has several limitations. With regard to the sample characteristics, the number of female CHR-participants in the current study exceeded previous studies in the field. The reason of the higher number of self-referrals is not completely clear but may be in part explained by the greater willingness of female participants to engage in studies and perhaps increased awareness of mental health issues. If the latter is correct, different strategies may have to be employed to engage male participants in early intervention. Secondly, we did not assess negative symptoms in CHR-participants that have been shown to mediate the relationship between neurocognitive deficits and functioning in previous studies (Meyer *et al.*, 2014; Glenthøj *et al.*, 2017). Finally, it is currently unclear whether neurocognitive deficits in our community recruited CHR-sample are predictive for the persistence of sub-threshold psychosis symptoms and/or conversion to psychosis as has been suggested by previous findings (Seidman *et al.*, 2016; Lam *et al.*, 2018).

Summary and conclusions

The current data support the view that neurocognitive deficits are a core feature of the CHR-participants recruited from the general community, replicating previous findings from CHR-cohorts recruited from clinical referral pathways. This is also supported by the fact that cognitive impairments were largely specific to the CHR-group. Thus, participants who did not meet CHR-criteria but who were characterised by affective disorders and substance abuse did not show neurocognitive impairments to the same extent as observed in the CHR-group, supporting the view that dysfunctional cognition is related to an extended psychosis phenotype.

Follow-up data need to confirm whether such deficits are also predictive for clinical outcomes and transitioning to psychosis in community-recruited CHR-participants. If this is the case, neurocognitive testing could potentially be used to stratify young people with subthreshold psychotic symptoms and support targeted interventions for improving cognitive processes. This approach is furthermore motivated by the finding that neurocognitive

deficits were related to aspects of psychosocial functioning, replicating existing data from clinically identified CHR-groups.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291718003975>.

Author ORCIDs.  Peter J. Uhlhaas, 0000-0002-0892-2224

Acknowledgement. We acknowledge the support of the Scottish Mental Health Research Network (<http://www.smhrn.org.uk>) now called the NHS Research Scotland Mental Health Network (NRS MHN: <http://www.nhs.researchscotland.org.uk/research-areas/mental-health>) for providing assistance with participant recruitment, interviews and cognitive assessments. We would like to thank both the participants and patients who took part in the study and the research assistants of the YouR-study for supporting the recruitment and assessment of CHR-participants.

Financial support. This study was supported by the project MR/L011689/1 from the Medical Research Council (MRC).

References

- Addington J, Penn D, Woods SW, Addington D and Perkins DO (2008a) Facial affect recognition in individuals at clinical high risk for psychosis. *The British Journal of Psychiatry* **192**, 67–68.
- Addington J, Penn D, Woods SW, Addington D and Perkins DO (2008b) Social functioning in individuals at clinical high risk for psychosis. *Schizophrenia Research* **99**, 119–124.
- Allott KA, Schäfer MR, Thompson A, Nelson B, Bendall S, Bartholomeusz CF, Yuen HP, McGorry PD, Schlögelhofer M, Bechdolf A and Amminger GP (2014) Emotion recognition as a predictor of transition to a psychotic disorder in ultra-high risk participants. *Schizophrenia Research* **153**, 25–31.
- Amminger GP, Schäfer MR, Klier CM, Schlögelhofer M, Mossaheb N, Thompson A, Bechdolf A, Allott K, McGorry PD and Nelson B (2012) Facial and vocal affect perception in people at ultra-high risk of psychosis, first-episode schizophrenia and healthy controls. *Early Intervention in Psychiatry* **6**, 450–454.
- Bora E, Lin A, Wood SJ, Yung AR, McGorry PD and Pantelis C (2014) Cognitive deficits in youth with familial and clinical high risk to psychosis: a systematic review and meta-analysis. *Acta Psychiatrica Scandinavica* **130**, 1–15.
- Bora E and Murray RM (2014) Meta-analysis of cognitive deficits in ultra-high risk to psychosis and first-episode psychosis: do the cognitive deficits progress over, or after, the onset of psychosis? *Schizophr Bull* **40**, 744–755. <https://doi.org/10.1093/schbul/sbt085>.
- Brewer WJ, Francey SM, Wood SJ, Jackson HJ, Pantelis C, Phillips LJ, Yung AR, Anderson VA and McGorry PD (2005) Memory impairments identified in people at ultra-high risk for psychosis who later develop first-episode psychosis. *American Journal of Psychiatry* **162**, 71–78.
- Cannon-Spoor HE, Potkin SG and Wyatt RJ (1982) Measurement of pre-morbid adjustment in chronic schizophrenia. *Schizophrenia Bulletin* **8**, 470–484.
- Carrion RE, Goldberg TE, McLaughlin D, Auther AM, Correll CU and Cornblatt BA (2011) Impact of neurocognition on social and role functioning in individuals at clinical high risk for psychosis. *American Journal of Psychiatry* **168**, 806–813.
- Cornblatt BA, Auther AM, Niendam T, Smith CW, Zinberg J, Bearden CE and Cannon TD (2007) Preliminary findings for two new measures of social and role functioning in the prodromal phase of schizophrenia. *Schizophrenia Bulletin* **33**, 688–702.
- Dean DJ and Mittal VA (2015) Spontaneous parkinsonisms and striatal impairment in neuroleptic free youth at ultrahigh risk for psychosis. *NPJ Schizophrenia* **1**, 14006.
- Dean DJ, Orr JM, Newberry RE and Mittal VA (2016) Motor behavior reflects reduced hemispheric asymmetry in the psychosis risk period. *Schizophrenia Research* **170**, 137–142.

- Dickinson D, Ramsey ME and Gold JM (2007) Overlooking the obvious: a meta-analytic comparison of digit symbol coding tasks and other cognitive measures in schizophrenia. *Archives of General Psychiatry* **64**, 532–542.
- Dickson H, Laurens KR, Cullen AE and Hodgins S (2012) Meta-analyses of cognitive and motor function in youth aged 16 years and younger who subsequently develop schizophrenia. *Psychological Medicine* **42**, 743–755.
- Fett AKJ, Viechtbauer W, Penn DL, van Os J and Krabbendam L (2011) The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: a meta-analysis. *Neuroscience & Biobehavioral Reviews* **35**, 573–588.
- Francey SM, Jackson HJ, Phillips LJ, Wood SJ, Yung AR and McGorry PD (2005) Sustained attention in young people at high risk of psychosis does not predict transition to psychosis. *Schizophrenia Research* **79**, 127–136.
- Frommann I, Pukrop R, Brinkmeyer J, Bechdorf A, Ruhrmann S, Berning J, Decker P, Riedel M, Möller HJ, Wölwer W and Gaebel W (2010) Neuropsychological profiles in different at-risk states of psychosis: executive control impairment in the early – and additional memory dysfunction in the late – prodromal state. *Schizophrenia Bulletin* **37**, 861–873.
- Fusar-Poli P, Deste G, Smieskova R, Barlati S, Yung AR, Howes O, Stieglitz RD, Vita A, McGuire P and Borgwardt S (2012) Cognitive functioning in prodromal psychosis: a meta-analysis. *Archives of General Psychiatry* **69**, 562–571.
- Fusar-Poli P, Borgwardt S, Bechdorf A, Addington J, Riecher-Rössler A, Schultze-Lutter F, Keshavan M, Wood S, Ruhrmann S, Seidman LJ and Valmaggia L (2013) The psychosis high-risk state: a comprehensive state-of-the-art review. *JAMA Psychiatry* **70**, 107–120.
- Fusar-Poli P, Cappucciati M, Rutigliano G, Schultze-Lutter F, Bonoldi I, Borgwardt S, Riecher-Rössler A, Addington J, Perkins D, Woods SW and McGlashan TH (2015a) At risk or not at risk? A meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. *World Psychiatry* **14**, 322–332.
- Fusar-Poli P, Schultze-Lutter F, Cappucciati M, Rutigliano G, Bonoldi I, Stahl D, Borgwardt S, Riecher-Rössler A, Addington J, Perkins DO and Woods SW (2015b) The dark side of the moon: meta-analytical impact of recruitment strategies on risk enrichment in the clinical high risk state for psychosis. *Schizophrenia Bulletin* **42**, 732–743.
- Gee DG, Karlsgodt KH, van Erp TG, Bearden CE, Lieberman MD, Belger A, Perkins DO, Olvet DM, Cornblatt BA, Constable T and Woods SW (2012) Altered age-related trajectories of amygdala-prefrontal circuitry in adolescents at clinical high risk for psychosis: a preliminary study. *Schizophrenia Research* **134**, 1–9.
- Giuliano AJ, Li H, Mesholam-Gately RI, Sorenson SM, Woodberry KA and Seidman LJ (2012) Neurocognition in the psychosis risk syndrome: a quantitative and qualitative review. *Current Pharmaceutical Design* **18**, 399–415.
- Glenthøj LB, Jepsen JRM, Hjorthøj C, Bak N, Kristensen TD, Wenneberg C, Krakauer K, Nordentoft M and Fagerlund B (2017) Negative symptoms mediate the relationship between neurocognition and function in individuals at ultrahigh risk for psychosis. *Acta Psychiatrica Scandinavica* **135**, 250–258.
- Green MF, Kern RS, Braff DL and Mintz J (2000) Neurocognitive deficits and functional outcome in schizophrenia: are we measuring the ‘right stuff’? *Schizophrenia Bulletin* **26**, 119–136.
- Green MF, Nuechterlein KH, Gold JM, Barch DM, Cohen J, Essock S, Fenton WS, Frese F, Goldberg TE, Heaton RK and Keefe RS (2004) Approaching a consensus cognitive battery for clinical trials in schizophrenia: the NIMH-MATRICES conference to select cognitive domains and test criteria. *Biological Psychiatry* **56**, 301–307.
- Green MF, Horan WP and Lee J (2015) Social cognition in schizophrenia. *Nature Reviews Neuroscience* **16**, 620.
- Heinrichs RW and Zakzanis KK (1998) Neurocognitive deficit in schizophrenia: a quantitative review of the evidence. *Neuropsychology* **12**, 426.
- Hooker C and Park S (2002) Emotion processing and its relationship to social functioning in schizophrenia patients. *Psychiatry Research* **112**, 41–50.
- Ising HK, Veling W, Loewy RL, Rietveld MW, Rietdijk J, Dragt S, Klaassen RM, Nieman DH, Wunderink L, Linszen DH and van der Gaag M (2012) The validity of the 16-item version of the Prodromal Questionnaire (PQ-16) to screen for ultra high risk of developing psychosis in the general help-seeking population. *Schizophrenia Bulletin* **38**, 1288–1296.
- Jahshan C, Heaton RK, Golshan S and Cadenhead KS (2010) Course of neurocognitive deficits in the prodrome and first episode of schizophrenia. *Neuropsychology* **24**, 109.
- Keefe RS, Goldberg TE, Harvey PD, Gold JM, Poe MP and Coughenour L (2004) The Brief Assessment of Cognition in Schizophrenia: reliability, sensitivity, and comparison with a standard neurocognitive battery. *Schizophrenia Research* **68**, 283–297.
- Keefe RS, Harvey PD, Goldberg TE, Gold JM, Walker TM, Kennel C and Hawkins K (2008) Norms and standardization of the Brief Assessment of Cognition in Schizophrenia (BACS). *Schizophrenia Research* **102**, 108–115.
- Kindler J, Schultze-Lutter F, Michel C, Martz-Iringartinger A, Linder C, Schmidt SJ, Stegmayer K, Schimmelmann BG and Walther S (2016) Abnormal involuntary movements are linked to psychosis-risk in children and adolescents: results of a population-based study. *Schizophrenia Research* **174**, 58–64.
- Klosterkötter J, Hellmich M, Steinmeyer EM and Schultze-Lutter F (2001) Diagnosing schizophrenia in the initial prodromal phase. *Archives of General Psychiatry* **58**, 158–164.
- Lam M, Lee J, Rapisarda A, See YM, Yang Z, Lee SA, Abdul-Rashid NA, Kraus M, Subramaniam M, Chong SA and Keefe RS (2018) Longitudinal cognitive changes in young individuals at ultrahigh risk for psychosis. *JAMA Psychiatry* **75**, 929–939.
- Lencz T, Smith CW, McLaughlin D, Auther A, Nakayama E, Hovey L and Cornblatt BA (2006) Generalized and specific neurocognitive deficits in prodromal schizophrenia. *Biological Psychiatry* **59**, 863–871.
- Lin A, Wood SJ, Nelson B, Brewer WJ, Spiliotacopoulos D, Bruxner A, Broussard C, Pantelis C and Yung AR (2011) Neurocognitive predictors of functional outcome two to 13 years after identification as ultra-high risk for psychosis. *Schizophrenia Research* **132**, 1–7.
- McDonald M, Christoforidou E, Van Rijsbergen N, Gajwani R, Gross J, Gumley AI, Lawrie SM, Schwannauer M, Schultze-Lutter F and Uhlhaas PJ (2018) Using online screening in the general population to detect participants at clinical high-risk for psychosis. *Schizophrenia Bulletin*. doi: 10.1093/schbul/sby069.
- Mesholam-Gately RI, Giuliano AJ, Goff KP, Faraone SV and Seidman LJ (2009) Neurocognition in first-episode schizophrenia: a meta-analytic review. *Neuropsychology* **23**, 315.
- Meyer EC, Carrión RE, Cornblatt BA, Addington J, Cadenhead KS, Cannon TD, McGlashan TH, Perkins DO, Tsuang MT, Walker EF and Woods SW (2014) The relationship of neurocognition and negative symptoms to social and role functioning over time in individuals at clinical high risk in the first phase of the North American Prodrome Longitudinal Study. *Schizophrenia Bulletin* **40**, 1452–1461.
- Michel C, Ruhrmann S, Schimmelmann BG, Klosterkötter J and Schultze-Lutter F (2014) A stratified model for psychosis prediction in clinical practice. *Schizophrenia Bulletin* **40**, 1533–1542.
- Miller TJ, McGlashan TH, Rosen JL, Cadenhead K, Ventura J, McFarlane W, Perkins DO, Pearlson GD and Woods SW (2003) Prodromal assessment with the structured interview for prodromal syndromes and the scale of prodromal symptoms: predictive validity, interrater reliability, and training to reliability. *Schizophrenia Bulletin* **29**, 703–715.
- Mills JG, Fusar-Poli P, Morgan C, Azis M and McGuire P (2017) People meeting ultra high risk for psychosis criteria in the community. *World Psychiatry* **16**, 322–323.
- Moore TM, Reise SP, Gur RE, Hakonarson H and Gur RC (2015) Psychometric properties of the Penn computerized neurocognitive battery. *Neuropsychology* **29**, 235.
- Morrens M, Hulstijn W and Sabbe B (2006) Psychomotor slowing in schizophrenia. *Schizophrenia Bulletin* **33**, 1038–1053.
- Niendam TA, Bearden CE, Johnson JK, McKinley M, Loewy R, O’Brien M, Nuechterlein KH, Green MF and Cannon TD (2006) Neurocognitive performance and functional disability in the psychosis prodrome. *Schizophrenia Research* **84**, 100–111.
- Niendam TA, Bearden CE, Zinberg J, Johnson JK, O’Brien M and Cannon TD (2007) The course of neurocognition and social functioning in individuals at ultra high risk for psychosis. *Schizophrenia Bulletin* **33**, 772–781.

- Pinkham AE, Penn DL, Perkins DO, Graham KA and Siegel M** (2007) Emotion perception and social skill over the course of psychosis: a comparison of individuals 'at-risk' for psychosis and individuals with early and chronic schizophrenia spectrum illness. *Cognitive Neuropsychiatry* **12**, 198–212.
- Pukrop R and Klosterkötter J** (2010) Neurocognitive indicators of clinical high-risk states for psychosis: a critical review of the evidence. *Neurotoxicity Research* **18**, 272–286.
- Rajji TK, Ismail Z and Mulsant BH** (2009) Age at onset and cognition in schizophrenia: meta-analysis. *The British Journal of Psychiatry* **195**, 286–293.
- Schultze-Lutter F, Addington J, Ruhrmann S and Klosterkötter J** (2007) Schizophrenia proneness instrument, adult version (SPI-A). *Rome: Giovanni Fioriti*.
- Schultze-Lutter F, Ruhrmann S, Berning J, Maier W and Klosterkötter J** (2008) Basic symptoms and ultrahigh risk criteria: symptom development in the initial prodromal state. *Schizophrenia Bulletin* **36**, 182–191.
- Schultze-Lutter F, Klosterkötter J and Ruhrmann S** (2014) Improving the clinical prediction of psychosis by combining ultra-high risk criteria and cognitive basic symptoms. *Schizophrenia Research* **154**, 100–106.
- Schultze-Lutter F, Michel C, Ruhrmann S and Schimmelmann BG** (2018) Prevalence and clinical relevance of interview-assessed psychosis-risk symptoms in the young adult community. *Psychological Medicine* **48**, 1167–1178.
- Seidman LJ, Giuliano AJ, Meyer EC, Addington J, Cadenhead KS, Cannon TD, McGlashan TH, Perkins DO, Tsuang MT, Walker EF and Woods SW** (2010) Neuropsychology of the prodrome to psychosis in the NAPLS consortium: relationship to family history and conversion to psychosis. *Archives of General Psychiatry* **67**, 578–588.
- Seidman LJ, Shapiro DI, Stone WS, Woodberry KA, Ronzio A, Cornblatt BA, Addington J, Bearden CE, Cadenhead KS, Cannon TD and Mathalon DH** (2016) Association of neurocognition with transition to psychosis: baseline functioning in the second phase of the North American prodrome longitudinal study. *JAMA Psychiatry* **73**, 1239–1248.
- Seiferth NY, Pauly K, Habel U, Kellermann T, Shah NJ, Ruhrmann S, Klosterkötter J, Schneider F and Kircher T** (2008) Increased neural response related to neutral faces in individuals at risk for psychosis. *Neuroimage* **40**, 289–297.
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R and Dunbar GC** (1998) The Mini-International Neuropsychiatric Interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry* **59**, 22.
- Simon AE, Dvorsky DN, Boesch J, Roth B, Isler E, Schueler P, Petralli C and Umbricht D** (2006) Defining subjects at risk for psychosis: a comparison of two approaches. *Schizophrenia Research* **81**, 83–90.
- Thompson AD, Bartholomeusz C and Yung AR** (2011) Social cognition deficits and the 'ultra high risk' for psychosis population: a review of literature. *Early Intervention in Psychiatry* **5**, 192–202.
- Uhlhaas PJ, Gajwani R, Gross J, Gumley AI, Lawrie SM and Schwannauer M** (2017) The youth mental health risk and resilience study (YouR-Study). *BMC Psychiatry* **17**, 43.
- Van Rijn S, Aleman A, de Sonneville L, Sprong M, Ziermans T, Schothorst P, Van Engeland H and Swaab H** (2011) Misattribution of facial expressions of emotion in adolescents at increased risk of psychosis: the role of inhibitory control. *Psychological Medicine* **41**, 499–508.
- Velthorst E, Nieman DH, Linszen D, Becker H, de Haan L, Dingemans PM, Birchwood M, Patterson P, Salokangas RK, Heinimaa M and Heinz A** (2010) Disability in people clinically at high risk of psychosis. *The British Journal of Psychiatry* **197**, 278–284.
- Yung AR, Yung AR, Pan Yuen H, McGorry PD, Phillips LJ, Kelly D, Dell'olio M, Francey SM, Cosgrave EM, Killackey E and Stanford C** (2005) Mapping the onset of psychosis: the comprehensive assessment of at-risk mental states. *Australian and New Zealand Journal of Psychiatry* **39**, 964–971.