

Review Article

Dr N C-W Tan takes responsibility for the integrity of the content of the paper

Cite this article: Ashman A, Psaltis AJ, Wormald PJ, Tan NC-W. Extended endoscopic approaches to the maxillary sinus. *J Laryngol Otol* 2020;**134**:473–480. <https://doi.org/10.1017/S0022215120000882>

Accepted: 17 March 2020
First published online: 8 June 2020

Key words:

Endoscopic Surgical Procedure;
Minimally Invasive Surgery; Maxillary Sinus;
Otorhinolaryngology;
Paranasal Sinus Diseases

Author for correspondence:

Mr Neil C-W Tan,
Department of Otolaryngology,
Royal Cornwall Hospital, Treliske,
Truro TR1 3LQ, UK
E-mail: n.tan@exeter.ac.uk

Extended endoscopic approaches to the maxillary sinus

A Ashman¹, A J Psaltis², P J Wormald² and N C-W Tan³

¹Department of Otolaryngology, Oxford University Hospitals, UK, ²Department of Otolaryngology, University of Adelaide and Queen Elizabeth Hospital, Adelaide, Australia and ³Department of Otolaryngology, Royal Cornwall Hospital, Truro, UK

Abstract

Objectives. Treatment of inflammatory and neoplastic disease in the maxillary sinus, pterygopalatine and infratemporal fossae requires appropriate surgical exposure. As modern rhinology evolves, so do the techniques available. This paper reviews extended endoscopic approaches to the maxillary sinus and the evidence supporting each technique.

Methods. A literature search of the Ovid Medline and PubMed databases was performed using appropriate key words relating to endoscopic approaches to the maxillary sinus.

Results. Mega-antrostomy and medial maxillectomy have a role in the surgical treatment of refractory inflammatory disease and sinonasal neoplasms. The pre-lacrimal fossa approach provides excellent access but can be limited because of anatomical variations. Both the trans-septal and endoscopic Denker's approaches were reviewed; these appear to be associated with morbidity, without any significant increase in exposure over the afore-described approaches.

Conclusion. A range of extended endoscopic approaches to the maxillary sinus exist, each with its own anatomical limitations and potential complications.

Introduction

A range of surgical approaches to the maxillary sinus can be considered in the treatment of inflammatory or neoplastic rhinological disease. Historically, the maxillary sinus was accessed by external approaches such as lateral rhinotomy, even in cases of benign disease. The endoscopic approach has now replaced external approaches as the standard of care, because of the reduced morbidity, improved visualisation and lower recurrence rates of benign tumours.¹ However, standard middle meatal antrostomy offers limited access to the maxillary sinus.

In cases where the disease is extensive, originates from the anterior wall, or involves the pterygopalatine or infratemporal fossa, a range of extended endoscopic approaches are available to allow improved access of both endoscopes and instruments. These are listed in [Table 1](#) alongside the traditional external and open approaches. This review aimed to discuss the benefits and drawbacks of each of the endoscopic modalities available to the modern rhinologist.

Materials and methods

A literature search of the Ovid Medline and PubMed databases was performed using appropriate key words relating to endoscopic approaches to the maxillary sinus. Relevant case series, and cadaveric, imaging and interventional studies, were identified and reviewed for relevance.

Anatomy

The largest of the paranasal sinuses, the maxillary sinus, has a final volume of around 10 ml.² It is the first sinus to develop embryologically, starting at the 10th week, and is almost always present at birth.³ Subsequent pneumatization occurs as the facial skeleton grows and matures, with the sinus reaching 25 per cent of its final size by age 2 years, up to 50 per cent by age 8 years, and by the 16th year the sinus will have reached its final adult size.⁴

The sinuses are pyramidal voids within the maxilla, the base being the lateral border of the nasal cavity and the apex lying towards the zygomatic process. The roof of the sinus forms the orbital floor, and the base of the sinus may extend into the alveolar part of the maxilla to the extent that dental roots may project into the sinus.

Drainage

Drainage is via the natural ostium in the superomedial part of the sinus that opens into the hiatus semilunaris, posterior to the uncinate process. The cilia lining the mucosa beat

Table 1. Endoscopic, and external or open approaches to maxillary sinus

Endoscopic approaches
– Middle meatal antrostomy
– Mega-antrostomy
– Modified or radical medial maxillectomy
– Pre-lacrimal approach
– Transseptal approach
– Endoscopic canine fossa trephine
– Endoscopic Denker's approach
External & open approaches
– Caldwell–Luc
– External ethmoidectomy
– Lateral rhinotomy
– Midfacial degloving

towards the natural ostium, and thus clear maxillary secretions out of the sinus through this mechanism.⁵

Accessory ostia are present in around one-fifth of patients, and are associated with a higher likelihood of mucosal disease, recurrent acute sinusitis and retention cysts.⁶ At surgery, an accessory ostium may be mistaken for the natural ostium, leading to the 'missed ostium' sequence whereby the natural ostium remains obstructed by the unciniate and is not included within the antrostomy.⁷ This may have the functional effect known as mucus circulation, whereby secretions cleared via the natural ostium are drawn back into the sinus through the accessory ostium, thus contributing to rhinorrhoea and post-nasal drip.

Indications for surgery

Inflammatory disease of the maxillary sinus may occur in the context of allergic or infective rhinosinusitis, with or without the formation of inflammatory polyps. Unilateral maxillary disease is often related to odontogenic causes such as periapical abscesses of the upper molars.⁸ When such cases fail to respond to maximal medical therapy, uncinectomy and middle meatal antrostomy is performed to ventilate the sinus, reduce the inflammatory burden by removal of polyps and debris, and, possibly most importantly, improve access for topical medications.

Benign, pre-malignant and malignant neoplasms of the maxillary sinus may be amenable to endoscopic resection. Inverted papilloma is the most common benign neoplasm of the sinonasal cavity; it is managed by meticulous surgical resection because of its risk of malignant transformation, its locally invasive nature and its tendency to recur. Malignant neoplasms of the paranasal sinuses can originate from or involve the maxillary sinus, and thus suitable access must be offered to allow for an appropriate oncological resection. Finally, lesions involving the pterygopalatine and infratemporal fossae may be approached endoscopically through the posterior wall of the maxillary sinus, requiring extended endoscopic approaches.

Uncinectomy and middle meatal antrostomy

At endoscopic sinus surgery, the maxillary sinus is commonly accessed via an uncinectomy and middle meatal antrostomy (Figure 1 and Table 2). However, this allows only limited




Fig. 1. Endoscopic images of middle meatal antrostomy. (a) Natural ostium (O) and inferior unciniate process (UP). (b) Widened middle meatal antrostomy demonstrating access to superior maxillary sinus (MS) only.

access to the sinus. A 2010 study using cadaveric heads found that a middle meatal antrostomy only provided access to 24–34 per cent of the total sinus volume, and rarely offered access to the anterior wall and sinus floor, regardless of the angled instruments used.⁹ This approach is adequate to drain mucopus and treat minor inflammatory disease or postero-medially based tumours; however, it is not always adequate for more significant disease.

Endoscopic maxillary mega-antrostomy

Endoscopic maxillary mega-antrostomy is typically used as a revision procedure in patients with maxillary sinusitis refractory to surgery.^{10,11} After a standard middle meatal antrostomy, the antrostomy is extended posteriorly to the back wall of the sinus and inferiorly to the level of the nasal floor, thereby requiring a posterior partial inferior turbinectomy (Figure 2). The resulting antrostomy is greatly enlarged and allows the sinus to drain more easily by gravity, as well as admitting topical therapy more readily (Table 2). This is potentially useful in patients with impaired mucociliary clearance such as cystic fibrosis or primary

Table 2. Pros and cons of endoscopic techniques to access maxillary sinus

Technique	Benefits	Drawbacks
Uncinectomy & middle meatal antrostomy	<ul style="list-style-type: none"> – Simple & fast to perform – Excellent access for ventilation & washout of mucopus – Good access to posterior area of sinus 	<ul style="list-style-type: none"> – Limited access to anterior & inferior portions of sinus
Endoscopic maxillary mega-antrostomy or modified medial maxillectomy	<ul style="list-style-type: none"> – Allows for dependant drainage after high-volume saline rinses in conditions such as cystic fibrosis & primary ciliary disorders – Good access to inferior portions of sinus – Maintains head of inferior turbinate, to reduce risk of nasal dryness & crusting 	<ul style="list-style-type: none"> – May still have limited access to anterior face of sinus or antero-inferior sump of sinus
Radical medial maxillectomy	<ul style="list-style-type: none"> – Widest endoscopic surgical access achievable for tumours of maxillary sinus & pterygopalatine or infratemporal fossa – Allows use of 0° endoscopes & straight instrumentation 	<ul style="list-style-type: none"> – Potentially more morbid – increased risk of crusting, nasal dryness & epiphora
Pre-lacrimal recess	<ul style="list-style-type: none"> – Wide access with straight & angled endoscopes directly into entire maxillary sinus – Maintains & reconstitutes head of inferior turbinate, to reduce risk of nasal dryness & crusting – Does not violate nasolacrimal system 	<ul style="list-style-type: none"> – Potential for minor crusting in anterior nasal cavity from lateral nasal wall incision

ciliary dyskinesia,¹² as well as in cases where there is a high inflammatory burden in the maxillary sinus such as severe polyposis or fungal sinusitis.

Inadequate resection of fungal debris or polyps, especially in areas typically inaccessible through a standard middle meatal antrostomy such as the pre-lacrimal recess, will leave




Fig. 2. Endoscopic images of maxillary mega-antrostomy. (a) Sharp resection of mid-portion of inferior turbinate (IT). (b) Exposed inferior meatus. (c) Mucosal flap (MF) medialised to expose bone of inferior meatus. (d) View into maxillary sinus following bony removal. MT = middle turbinate




Fig. 3. Endoscopic images of radical medial maxillectomy. (a) Frontal process of maxilla (FP) exposed and prepared for osteotomies. (b) Lacrimal sac (LS) exposed with sharp transection. (c) View into maxillary sinus, demonstrating multifocal inverted papilloma (IP) with anterior and superior attachments. (d) Wide surgical access to entire maxillary sinus following tumour resection.

residual pro-inflammatory material and may be a cause of surgical failure.¹³

Costa *et al.* published the long-term outcomes for a cohort of 122 patients.¹⁴ The authors reported that all patients with recalcitrant maxillary sinusitis experienced symptomatic improvement, with almost 75 per cent describing complete symptom resolution, with no late complications.

Endoscopic medial maxillectomy

In its radical form, endoscopic medial maxillectomy involves the removal of the entire inferior turbinate, uncinate process, ethmoid bulla, and the medial maxillary wall containing the distal portion of the nasolacrimal duct (Figure 3). Radical endoscopic medial maxillectomy is indicated for the resection of benign and malignant sinonasal neoplasms where wide surgical access and tumour clearance is required (Table 2).^{15–17} However, resection of the inferior turbinate head may

contribute to nasal crusting, dryness and empty nose syndrome.^{18,19}

The modified endoscopic medial maxillectomy can be considered akin to endoscopic maxillary mega-antrostomy, in which the head of the inferior turbinate and nasolacrimal duct are preserved (Figure 4), but with additional bone of the inferomedial maxillary wall resected. Its use has been described in treating refractory sinus disease.²⁰ Furthermore, similar to endoscopic maxillary mega-antrostomy, modified endoscopic medial maxillectomy gives superior access to the maxillary sinus (compared to middle meatal antrostomy), whilst preserving the head of the inferior turbinate. This maintains the humidification function of the inferior turbinate, resulting in less crusting when compared to radical endoscopic medial maxillectomy.¹⁸ Several authors have described modifications of this technique in an attempt to preserve even more inferior turbinate, through en bloc excision and re-implantation of the entire turbinate,²¹ or through




Fig. 4. Endoscopic images of modified medial maxillectomy. (a) Boundaries of mucosal flap raised after sharp resection of inferior turbinate (IT), demonstrating inverted papilloma (IP). (b) Back-biting forceps used to resect bone inferior to head of inferior turbinate. (c) View of maxillary sinus with 70-degree endoscope, demonstrating anterior attachment of inverted papilloma and access to anterior and inferior maxillary sinus. (d) Anterior maxillary face following tumour resection. S = septum; MT = middle turbinate

medialisation of the turbinate and nasolacrimal duct during the procedure.²²

To date, no studies have directly compared modified endoscopic medial maxillectomy with endoscopic maxillary mega-antrostomy in terms of surgical access, especially of the anterior maxillary face. The additional morbidity of the modified endoscopic medial maxillectomy compared with the endoscopic maxillary mega-antrostomy is minimal, however, and if the surgical pathology dictates a wider approach it should certainly be considered. In the context of recalcitrant inflammatory disease, both the modified endoscopic medial maxillectomy and endoscopic maxillary mega-antrostomy procedures have been reported to offer excellent disease control, with no requirement for further revision surgery on short- and long-term follow up.^{10,14,23,24}

Pre-lacrimal recess approach

An approach to the maxillary sinus via the pre-lacrimal recess allows for direct access to the sinus with 0-degree endoscopes

and instrumentation, and thus improved visualisation of its contents, whilst preserving the inferior turbinate and nasolacrimal duct (Figure 5).²⁵ This approach involves a vertical incision along the lateral nasal wall, just behind the piriform aperture, and medialisation of a mucosal flap with exposure of the bone of the antero-medial maxillary wall. Using sharp osteotomies or a drill, bone overlying the pre-lacrimal recess is removed and the lacrimal apparatus identified, thus allowing direct access into the maxillary sinus. After the operation, the mucosal flap is replaced and sutured with absorbable material. The pre-lacrimal recess approach is of value in the removal of recurrent antrochoanal polyps, benign tumours and refractory fungal infections,^{26–29} as well as being useful in approaches to the pterygopalatine and infratemporal fossae (Table 2).^{30,31}

The technique was initially described by Zhou *et al.* in 2013.³² These authors went on to describe the long-term results of this approach, including those of 71 patients with inverted papilloma.³³ The group reported complications such as upper lid and alar numbness, in 7 per cent of cases, and




Fig. 5. Endoscopic images of pre-lacrimal recess approach. (a) Sharp incision of mucosa (blue line) of head of inferior turbinate (IT). (b) Mucosal flap (MF) medialisated with exposure of bone overlying pre-lacrimal fossa prepared for osteotomy. (c) View into maxillary sinus with well-exposed anterior maxillary face (Ant Max). (d) Head of inferior turbinate reconstituted and sutured in place with absorbable material.

mild alar collapse, in 5.6 per cent of cases. A further series of 51 patients with sinonasal inverted papilloma was reported by Suzuki and colleagues.³⁴ They reported no significant morbidity resulting from this technique, although partial osteotomy of the piriform aperture was required for access in eight cases, which can lead to alar collapse or notching at the alar margin.³⁵ We would not advocate resection of the piriform aperture because of the above risks.

Two studies of computed tomograms demonstrated great variation in the anatomy of the pre-lacrimal recess. Kashlan and Craig found that the anteroposterior dimension of the pre-lacrimal recess was widest inferiorly, with a mean of 8.4 mm but a range of 1.9–14.2 mm, with the height having a range of 18.5–39.9 mm.³⁶ Sieskiewicz *et al.* found that the width of the pre-lacrimal recess had a range of 0–15.2 mm at the level of the inferior turbinate, and in 30 per cent of patients studied the recess was too narrow to accept a 4 mm endoscope.³⁷ However, even if the pre-lacrimal recess itself is narrow, the nasolacrimal apparatus can be mobilised and pushed medially in order to admit suitable instrumentation, without affecting its long-term function.

Trans-septal approach

The trans-septal approach was used to access the anterior maxillary face through asymmetrical septal incisions and an inferior cartilaginous septal resection. This approach was found to offer a 15–20 degree improvement in access in the axial plane.^{38,39} However, an imaging study of six cadaveric heads suggested that, in some cases, access to the anterior-inferior part of the maxillary sinus might be limited by the height of the nasal floor.⁴⁰ This technique has fallen out of favour because of the excellent access that the pre-lacrimal recess approach offers, with no risk of the morbidities associated with the transseptal approach such as septal perforation; thus, we would not advocate its use unless access is limited through alternative approaches.

Endoscopic canine fossa trephine

Access may also be improved by performing canine fossa trephination, with the potential for better outcomes when treating severely diseased maxillary sinuses. A cadaveric study comparing canine fossa trephination with exclusively

transnasal endoscopic access found that a greater amount of debris remained in the maxillary sinus following the transnasal approach (3.88 cm³ vs 2.88 cm³).⁴¹ A single-blind, randomised study of 29 patients with unilateral chronic rhinosinusitis, by Byun and Lee, reported improved Sino-Nasal Outcome Test 20 (SNOT-20) outcome scores at 12 months in patients undergoing canine fossa trephination compared to those undergoing middle meatal antrostomy alone.⁴² A similar study by Seiberling and colleagues showed improved post-operative nasal endoscopy scores following canine fossa trephine compared with middle meatal antrostomy in 42 patients.⁴³

There is an associated risk of damage to branches of the infraorbital nerve, namely the anterior superior alveolar nerve and the middle superior alveolar nerve, with the potential for adverse effects including facial paraesthesia and dental numbness. Traditional canine fossa approaches have been reported to result in adverse effects in 76 per cent of cases.⁴⁴ A cadaveric study by Robinson and Wormald found that the anterior superior alveolar nerve emerged from its foramen as a double trunk in 25 per cent of the 40 sides dissected. One or more branches from the anterior superior alveolar nerve trunks were identified in 60 per cent, and a middle superior alveolar nerve was identified in 23 per cent. The authors concluded that the safest point to perform canine fossa puncture is on the mid-pupillary line, at the level of the floor of the pyriform aperture.⁴⁵ A case series of 99 canine fossa approaches by Singhal *et al.* found that the use of anatomical landmarks reduced the frequency of adverse effects to 53 per cent, and the addition of endoscopic guidance reduced this further, to 40 per cent.⁴⁴

The high initial adverse effects are very minor and largely soft tissue related, and are secondary to the excessive raising of soft tissue from the anterior maxillary face. Early studies reported a moderate rate of persistent paraesthesia of 28.6 per cent.⁴⁵ However, with the use of anatomical landmarks with endoscopy, thereby allowing for refined trephination of the anterior maxillary face, the incidence of complications was reduced to 3.1 per cent (3 out of 97) at one month and 1 per cent (1 out of 97) at three months.⁴⁶

Endoscopic Denker's approach

Tumours involving the pterygopalatine and infratemporal fossae may, in some cases, prove difficult to access via a trans-septal approach or radical medial maxillectomy. An endoscopic Denker's (Sturmann–Canfield) approach can provide improved access.^{47,48} This technique involves a soft tissue approach similar to the radical endoscopic medial maxillectomy, with a mucosal flap raised anterior to the inferior turbinate head and posteriorly along the floor of the nose, with exposure of the frontal process of the maxilla. Rather than staying medial to the piriform aperture, as in the endoscopic medial maxillectomy, the bony dissection is taken laterally over the face of the maxilla, whilst staying below the infraorbital nerve. The bony window therefore includes the medial wall, as in a radical endoscopic medial maxillectomy, but also the piriform aperture and anterior maxillary face.⁴⁹

This approach may be appropriate in the management of extensive sinonasal tumours, but is associated with risks such as infraorbital nerve damage, alar collapse and nasolacrimal duct stenosis.⁴⁹ Another potential application is in cases where tumours involve the nasolacrimal duct or ascending process of the maxilla, when the approach can be used as a means of obtaining an anterior margin. As the pre-lacrimal

recess approach offers almost complete access to the entirety of the maxillary sinus, pterygopalatine and infratemporal fossae, we cannot recommend Denker's approach.

Conclusion

A variety of extended endoscopic approaches to the maxillary sinus, pterygopalatine and infratemporal fossae exist, each with its own anatomical limitations and potential complications. The pre-lacrimal recess approach offers excellent visualisation of the entire sinus in most cases of benign disease. The medial maxillectomy should be a technique held within the armament of the modern endoscopic rhinologist for when the surgical pathology necessitates wider access. Careful pre-operative review of the anatomy and disease burden is mandated prior to selecting an appropriate operative strategy.

Competing interests. None declared

References

- Goudakos JK, Blioukas S, Nikolaou A, Vlachtsis K, Karkos P, Markou KD. Endoscopic resection of sinonasal inverted papilloma: systematic review and meta-analysis. *Am J Rhinol Allergy* 2018;**32**:167–74
- Wagenmann M, Naclerio RM. Anatomic and physiologic considerations in sinusitis. *J Allergy Clin Immunol* 1992;**90**:419–23
- Nunez-Castruita A, Lopez-Serna N, Guzman-Lopez S. Prenatal development of the maxillary sinus: a perspective for paranasal sinus surgery. *Otolaryngol Head Neck Surg* 2012;**146**:997–1003
- Lorkiewicz-Muszynska D, Kociemba W, Rewekant A, Sroka A, Jonczyk-Potoczna K, Patelska-Banaszewska M *et al.* Development of the maxillary sinus from birth to age 18. Postnatal growth pattern. *Int J Pediatr Otorhinolaryngol* 2015;**79**:1393–400
- Gosau M, Rink D, Driemel O, Draenert FG. Maxillary sinus anatomy: a cadaveric study with clinical implications. *Anat Rec (Hoboken)* 2009;**292**:352–4
- Yenigun A, Fazliogullari Z, Gun C, Uysal II, Nayman A, Karabulut AK. The effect of the presence of the accessory maxillary ostium on the maxillary sinus. *Eur Arch Otorhinolaryngol* 2016;**273**:4315–19
- Parsons DS, Stivers FE, Talbot AR. The missed ostium sequence and the surgical approach to revision functional endoscopic sinus surgery. *Otolaryngol Clin North Am* 1996;**29**:169–83
- Little RE, Long CM, Loehrl TA, Poetker DM. Odontogenic sinusitis: a review of the current literature. *Laryngoscope Investig Otolaryngol* 2018;**3**:110–14
- Robey A, O'Brien EK, Leopold DA. Assessing current technical limitations in the small-hole endoscopic approach to the maxillary sinus. *Am J Rhinol Allergy* 2010;**24**:396–401
- Cho DY, Hwang PH. Results of endoscopic maxillary mega-antrostomy in recalcitrant maxillary sinusitis. *Am J Rhinol* 2008;**22**:658–62
- Woodworth BA, Parker RO, Schlosser RJ. Modified endoscopic medial maxillectomy for chronic maxillary sinusitis. *Am J Rhinol* 2006;**20**:317–19
- Thompson C, Conley DB. What is the optimal maxillary antrostomy size during sinus surgery? *Curr Opin Otolaryngol Head Neck Surg* 2015;**23**:34–8
- Bassiouni A, Naidoo Y, Wormald PJ. When FESS fails: the inflammatory load hypothesis in refractory chronic rhinosinusitis. *Laryngoscope* 2012;**122**:460–6
- Costa ML, Psaltis AJ, Nayak JV, Hwang PH. Long-term outcomes of endoscopic maxillary mega-antrostomy for refractory chronic maxillary sinusitis. *Int Forum Allergy Rhinol* 2015;**5**:60–5
- Sadeghi N, Al-Dhahri S, Manoukian JJ. Transnasal endoscopic medial maxillectomy for inverting papilloma. *Laryngoscope* 2003;**113**:749–53
- Wormald PJ, Ooi E, van Hasselt CA, Nair S. Endoscopic removal of sinonasal inverted papilloma including endoscopic medial maxillectomy. *Laryngoscope* 2003;**113**:867–73
- Luong A, Citardi MJ, Batra PS. Management of sinonasal malignant neoplasms: defining the role of endoscopy. *Am J Rhinol Allergy* 2010;**24**:150–5
- Kastl KG, Rettinger G, Keck T. The impact of nasal surgery on air-conditioning of the nasal airways. *Rhinology* 2009;**47**:237–41

- 19 Dayal A, Rhee JS, Garcia GJ. Impact of middle versus inferior total turbinectomy on nasal aerodynamics. *Otolaryngol Head Neck Surg* 2016;**155**:518–25
- 20 Durr ML, Goldberg AN. Endoscopic partial medial maxillectomy with mucosal flap for maxillary sinus mucoceles. *Am J Otolaryngol* 2014;**35**:115–19
- 21 Weber RK, Werner JA, Hildenbrand T. Endonasal endoscopic medial maxillectomy with preservation of the inferior turbinate. *Am J Rhinol Allergy* 2010;**24**:132–5
- 22 Suzuki M, Nakamura Y, Nakayama M, Inagaki A, Murakami S, Takemura K *et al.* Modified transnasal endoscopic medial maxillectomy with medial shift of preserved inferior turbinate and nasolacrimal duct. *Laryngoscope* 2011;**121**:2399–401
- 23 Wang EW, Gullung JL, Schlosser RJ. Modified endoscopic medial maxillectomy for recalcitrant chronic maxillary sinusitis. *Int Forum Allergy Rhinol* 2011;**1**:493–7
- 24 Thulasidas P, Vaidyanathan V. Role of modified endoscopic medial maxillectomy in persistent chronic maxillary sinusitis. *Int Arch Otorhinolaryngol* 2014;**18**:159–64
- 25 Morrissey DK, Wormald PJ, Psaltis AJ. Prelacrimal approach to the maxillary sinus. *Int Forum Allergy Rhinol* 2016;**6**:214–18
- 26 Lin YT, Lin CF, Yeh TH. Application of the endoscopic prelacrimal recess approach to the maxillary sinus in unilateral maxillary diseases. *Int Forum Allergy Rhinol* 2018;**8**:530–6
- 27 Comoglu S, Celik M, Enver N, Sen C, Polat B, Deger K. Transnasal prelacrimal recess approach for recurrent antrchoanal polyp. *J Craniofac Surg* 2016;**27**:1025–7
- 28 Lee JJ, Ahmad ZA, Kim D, Ryu G, Kim HY, Dhong HJ *et al.* Comparison between endoscopic prelacrimal medial maxillectomy and Caldwell-Luc approach for benign maxillary sinus tumors. *Clin Exp Otorhinolaryngol* 2019;**12**:287–93
- 29 Yu QQ, Guan G, Zhang NK, Zhang XW, Jiang Y, Lian YY *et al.* Intranasal endoscopic prelacrimal recess approach for maxillary sinus inverted papilloma. *Eur Arch Otorhinolaryngol* 2018;**275**:2297–302
- 30 Gao L, Zhou L, Dai Z, Huang X. The endoscopic prelacrimal recess approach to the pterygopalatine fossa and infratemporal fossa. *J Craniofac Surg* 2017;**28**:1589–93
- 31 Zhou B, Huang Q, Shen PH, Cui SJ, Wang CS, Li YC *et al.* The intranasal endoscopic removal of schwannoma of the pterygopalatine and infratemporal fossae via the prelacrimal recess approach. *J Neurosurg* 2016;**124**:1068–73
- 32 Zhou B, Han DM, Cui SJ, Huang Q, Wang CS. Intranasal endoscopic prelacrimal recess approach to maxillary sinus. *Chin Med J (Engl)* 2013;**126**:1276–80
- 33 Zhou B, Huang Q, Sun J, Li X, Zhang W, Cui S *et al.* Resection of inverted papilloma of the maxillary sinus via a prelacrimal recess approach: a multicenter retrospective analysis of surgical efficacy. *Am J Rhinol Allergy* 2018;**32**:518–25
- 34 Suzuki M, Nakamura Y, Yokota M, Ozaki S, Murakami S. Modified transnasal endoscopic medial maxillectomy through prelacrimal duct approach. *Laryngoscope* 2017;**127**:2205–9
- 35 Papesch E, Papesch M. The nasal pyriform aperture and its importance. *Otorhinolaryngol Head Neck Surg* 2016;**1**:89–91
- 36 Khashan K, Craig J. Dimensions of the medial wall of the prelacrimal recess. *Int Forum Allergy Rhinol* 2018;**8**:751–5
- 37 Sieskiewicz A, Buczko K, Janica J, Lukasiewicz A, Lebkowska U, Piszczatowski B *et al.* Minimally invasive medial maxillectomy and the position of nasolacrimal duct: the CT study. *Eur Arch Otorhinolaryngol* 2017;**274**:1515–19
- 38 Harvey RJ, Sheehan PO, Debnath NI, Schlosser RJ. Transseptal approach for extended endoscopic resections of the maxilla and infratemporal fossa. *Am J Rhinol Allergy* 2009;**23**:426–32
- 39 Ramakrishnan VR, Suh JD, Chiu AG, Palmer JN. Septal dislocation for endoscopic access of the anterolateral maxillary sinus and infratemporal fossa. *Am J Rhinol Allergy* 2011;**25**:128–30
- 40 Schreiber A, Ferrari M, Rampinelli V, Doglietto F, Belotti F, Lancini D *et al.* Modular endoscopic medial maxillectomies: quantitative analysis of surgical exposure in a preclinical setting. *World Neurosurg* 2017;**100**:44–55
- 41 Feldt BA, McMains KC, Weitzel EK. Cadaveric comparison of canine fossa vs transnasal maxillary sinus access. *Int Forum Allergy Rhinol* 2011;**1**:183–6
- 42 Byun JY, Lee JY. Canine fossa puncture for severe maxillary disease in unilateral chronic sinusitis with nasal polyp. *Laryngoscope* 2013;**123**:E79–84
- 43 Seiberling KA, Church CA, Tewfik M, Foreman A, Chang D, Ghostine M *et al.* Canine fossa trephine is a beneficial procedure in patients with Samter's triad. *Rhinology* 2012;**50**:104–8
- 44 Singhal D, Douglas R, Robinson S, Wormald PJ. The incidence of complications using new landmarks and a modified technique of canine fossa puncture. *Am J Rhinol* 2007;**21**:316–19
- 45 Robinson S, Wormald PJ. Patterns of innervation of the anterior maxilla: a cadaver study with relevance to canine fossa puncture of the maxillary sinus. *Laryngoscope* 2005;**115**:1785–8
- 46 Seiberling K, Ooi E, MiinYip J, Wormald PJ. Canine fossa trephine for the severely diseased maxillary sinus. *Am J Rhinol Allergy* 2009;**23**:615–18
- 47 Upadhyay S, Dolci RL, Buohliqah L, Fiore ME, Ditzel Filho LF, Prevedello DM *et al.* Effect of incremental endoscopic maxillectomy on surgical exposure of the pterygopalatine and infratemporal fossae. *J Neurol Surg B Skull Base* 2016;**77**:66–74
- 48 Prosser JD, Figueroa R, Carrau RI, Ong YK, Solares CA. Quantitative analysis of endoscopic endonasal approaches to the infratemporal fossa. *Laryngoscope* 2011;**121**:1601–5
- 49 Lee JT, Suh JD, Carrau RL, Chu MW, Chiu AG. Endoscopic Denker's approach for resection of lesions involving the anteroinferior maxillary sinus and infratemporal fossa. *Laryngoscope* 2017;**127**:556–60