

Coxeter Diagrams and the Köthe's Problem

Ziba Fazelpour and Alireza Nasr-Isfahani

Abstract. A ring Λ is called right Köthe if every right Λ -module is a direct sum of cyclic modules. In this paper, we give a characterization of basic hereditary right Köthe rings in terms of their Coxeter valued quivers. We also give a characterization of basic right Köthe rings with radical square zero. Therefore, we give a solution to Köthe's problem in these two cases.

1 Introduction

It is known that every finitely generated \mathbb{Z} -module is a direct sum of cyclic modules. The idea of this important property of abelian groups go back to Prüfer [30]. Köthe showed that artinian principal ideal rings have this property. He also proved that if a commutative artinian ring Λ has the property that each of its Λ -modules is a direct sum of cyclic modules, then it is a principal ideal ring. He posed the question to classify the noncommutative rings with this property [28]. Köthe's problem is one of the old problems in rings and modules theory that has not yet been solved. A ring for which any right module is a direct sum of cyclic modules, is now called a *right Köthe ring*. Nakayama gave an example of a right Köthe ring Λ that is not a principal right ideal ring (see [29, p. 289]). Later, Cohen and Kaplansky proved that if a commutative ring Λ is Köthe, then Λ is an artinian principal ideal ring [10]. Combining the results of Cohen and Kaplansky [10] and Köthe [28], one obtains that a commutative ring Λ is Köthe if and only if Λ is an artinian principal ideal ring. A right artinian ring Λ is called *representation-finite* provided Λ has, up to isomorphism, only finitely many finitely generated indecomposable right Λ -modules. Following [41], we call the ring Λ *right pure semisimple if every right Λ -module is a direct sum of finitely generated right Λ -modules*. It is known that a commutative ring Λ is pure semisimple, if and only if, Λ is a representation-finite ring, if and only if, Λ is a Köthe ring [22]. A ring Λ is a representation-finite ring if and only if Λ is right pure semisimple and left pure semisimple [3]. The problem of whether right pure semisimple rings are representation-finite, known as the pure semisimplicity conjecture, remains open (see [3, 41, 42]). It seems that there is a strong connection between pure semisimplicity conjecture and Köthe's problem.

Kawada completely solved Köthe's problem for the basic finite-dimensional K -algebras [25–27] (see also [31]). Kawada's papers contain a set of 19 conditions that

Received by the editors March 4, 2019; revised November 28, 2019.

Published online on Cambridge Core.

The research of the first author was in part supported by a grant from IPM. Also, the research of the second author was in part supported by a grant from IPM (No. 98170412).

AMS subject classification: 16G10, 16D70, 16G60.


Keywords: right Köthe ring, representation-finite ring, species, partial Coxeter functor, Coxeter functor, Coxeter valued quiver, separated diagram.

characterize Kawada algebras as well as the list of all possible finitely generated indecomposable modules. Using the multiplicity-free of top and soc of finitely generated indecomposable modules, Ringel gave a characterization of Kawada algebras [31]. Behboodi et al. proved that if Λ is a right Köthe ring in which all idempotents are central, then Λ is an artinian principal left ideal ring [6]. Recently the authors have studied Köthe's problem [17]. In fact, all known results related to the characterization of right Köthe rings follow from [17, Corollary 3.2].

In representation theory, representation-finite algebras are of particular importance, since in this case, one has a complete combinatorial description of the module category in terms of the Auslander-Reiten quiver. By [9, Theorem 4.4], any right Köthe ring Λ is right artinian; then there is a finite upper bound for the lengths of the finitely generated indecomposable right Λ -modules. Thus by [43, Proposition 54.3], any right Köthe ring is an artinian representation-finite ring. It seems that a solution of Köthe's problem needs a classification of all representation-finite rings and some further information about the structure of the modules over representation-finite rings, which is a rather difficult problem. In this paper, by using the representation theory techniques and classifications of representation-finite hereditary rings [11, 13, 14], we solve Köthe's problem in this case. As a consequence, we solve Köthe's problem for the class of rings with radical square zero.

We recall that a unitary ring Λ is defined to be *right hereditary* if every right ideal of Λ is projective. In [20], Gabriel proved that a hereditary finite-dimensional K -algebra Λ , over an algebraically closed field K , is representation-finite if and only if the underlying graph of its quiver Q_Λ (see [2]) is a disjoint union of the Dynkin diagrams \mathbb{A}_n , \mathbb{D}_n , \mathbb{E}_6 , \mathbb{E}_7 , and \mathbb{E}_8 presented in Table A, that also appear in Lie theory. It is also shown in [20] that there is a bijection between the isomorphism classes of finite-dimensional indecomposable representations and the positive roots of the corresponding Dynkin diagrams; see also [2, Ch. VII].

Table A. Dynkin diagrams



We recall from [19] that a *species* $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ is a finite set of division rings F_i and F_i - F_j -bimodules ${}_iM_j$, $i \neq j$. To an arbitrary basic ring Λ , one attaches its species \mathcal{M}_Λ as follows. Let $\Lambda/J \cong \bigoplus_{i=1}^n F_i$, where $n \in \mathbb{N}$; each F_i is a division ring

and J is the Jacobson radical of Λ . We can write $J/J^2 = \bigoplus_{1 \leq i, j \leq n} {}_iM_j$, where each ${}_iM_j = F_i(J/J^2)F_j$ is an F_i - F_j -bimodule. Then $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i, j \in I}$ is called the species of Λ . Let Λ be a basic hereditary ring and let $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i, j \in I}$ be the species of Λ . We recall from [37] that a Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is a quiver with vertices $1, 2, \dots, n$ corresponding to the division rings F_1, F_2, \dots, F_n . There exists a valued arrow


$$\begin{array}{ccc} & m_{ij} & \\ \circ & \xrightarrow{\quad} & \circ \\ i & & j \end{array}$$

in $(\mathcal{C}_\Lambda, \mathbf{m})$ if and only if the F_i - F_j -bimodule ${}_iM_j$ is non-zero and there exists exactly $m_{ij} \geq 3$ pairwise non-isomorphic indecomposable right $\begin{pmatrix} F_i & {}_iM_j \\ 0 & F_j \end{pmatrix}$ -modules. If $m_{ij} = 3$, we write simply

$$\begin{array}{ccc} \circ & \xrightarrow{\quad} & \circ \\ i & & j \end{array}$$

In [13], Dowbor, Ringel, and Simson proved that a hereditary artinian ring Λ is representation-finite if and only if the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is a disjoint union of the Coxeter diagrams presented in Table B, where the valued edge $\circ \xrightarrow{3} \circ$ is identified with $\circ \text{---} \circ$ (see [24]).

Table B. Coxeter diagrams



Let Λ be a basic hereditary ring and let \mathcal{D} be the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ . The ring Λ is called of the Dynkin type \mathcal{D} if \mathcal{D} is one of the Coxeter diagrams $A_n, B_n, D_n, E_6, E_7, E_8, F_4,$ and G_2 presented

in Table B. Also Λ is called of the Coxeter type \mathcal{D} if \mathcal{D} is one of the Coxeter diagrams \mathbb{H}_3 , \mathbb{H}_4 , and $\mathbb{I}_2(p)$ presented in Table B.

It is proved by Schofield [34, 35] that there exist hereditary bimodule rings of the form $\Lambda = \begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ of the Coxeter type $\mathbb{I}_2(5)$. However, the existence of such hereditary rings of the Coxeter type $\mathbb{I}_2(p)$ with $p \geq 7$ remains open. It depends on rather difficult questions concerning division rings extensions and leads to a generalized Artin problems; see [37, 39].

One of the main aims of this paper is to get a diagrammatic characterization of right Köthe rings Λ that are basic hereditary, or the square of the Jacobson radical of Λ is zero.

The paper is organized as follows. In Section 2, we prove some preliminary results that will be needed later in the paper. In Section 3, we collect some results of hereditary representation-finite rings that we need in the rest of the paper. In Section 4, we give a characterization of basic hereditary right Köthe rings of Dynkin type. In Section 5, we give a characterization of basic hereditary right Köthe rings in terms of their Coxeter valued quivers. Finally in Section 6, we give a characterization of basic right Köthe rings with radical square zero in terms of their separated quivers.

1.1 Notation

Throughout this paper, Λ is an associative ring with unit and all modules are unital. We denote by $\text{Mod-}\Lambda$ (resp. $\text{mod-}\Lambda$) the category of all right Λ -modules (resp. finitely generated right Λ -modules) and by J the Jacobson radical of Λ . A ring Λ is said to be *basic* if Λ/J is a direct product of division rings. For a right Λ -module M , we denote by $\text{top}(M)$ and $\text{rad}(M)$ its top and radical, respectively. Let X be a representation of a species \mathcal{M} (see Section 2) and let V be a right module over a division ring G . We denote by $\text{dim } X$ and $\text{dim}(V)_G$ the dimension vector of X and dimension of V , respectively. Let F and G be division rings and M be an F - G -bimodule. We denote $\text{dim}_F(M)$ by $\text{l.dim } M$ and $\text{dim}(M)_G$ by $\text{r.dim } M$. Also, we denote $M^L := \text{Hom}_F(M, F)$ and $M^R := \text{Hom}_G(M, G)$. We denote the left and right dualisations of the F - G -bimodule M by setting $M^{(0)} = M$, $M^{(j)} = (M^{(j-1)})^L$ for $j \geq 1$ and $M^{(j)} = (M^{(j+1)})^R$ for $j \leq -1$, respectively. Moreover, for each $m \geq 1$, we denote by $d_m(M)$ the sequence $(d_0^M, d_1^M, \dots, d_{m-1}^M)$, where $d_j^M = \text{r.dim } M^{(j)}$ for each j . Let X and Y be two right Λ -modules and $f: X \rightarrow Y$ be a homomorphism. We write $X \twoheadrightarrow Y$ (resp. $X \hookrightarrow Y$) when f is an epimorphism (resp. f is a monomorphism). Let A be a Λ -submodule of X . We denote by $f|_A$ the restriction of f to A . Let M be a right Λ -module and $n \in \mathbb{N}$. Then M^n denotes the direct sum of n copies of M . Let Q be a quiver and let i be a vertex of Q . We denote by i^+ and i^- the set of direct successors of i and the set of direct predecessors of i , respectively. Also, we denote by $|i^+|$ and $|i^-|$ the cardinal number of i^+ and i^- , respectively. Let K be a field, $n \in \mathbb{N}$ and $1 \leq i \leq n$. We denote by e_i the vector in K^n with a 1 in the i -th coordinate and zero in the j -th coordinate, for each $j \neq i$. Throughout the paper, we use standard quiver representation and path algebra terminology as applied in the monographs [2, 5]. We also refer to [2, Ch. VII] for a detailed explanation of the reflection functors technique introduced by Bernstein, Gelfand, and Ponomarev in [7], later developed by Dlab and Ringel in [11] and Auslander, Platzeck, and Reiten in [4].

2 Preliminaries

We recall from [11,13,14] that a *valued quiver* (Γ, \mathbf{d}) of a species $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ is a finite quiver $\Gamma = (\Gamma_0, \Gamma_1)$, with the finite set Γ_0 of vertices corresponding to the division rings F_i together with non-negative integers $d_{ij} = r.\dim {}_iM_j$ and $d_{ji} = l.\dim {}_iM_j$ for $i \neq j$ and the set Γ_1 of valued arrows defined as follows. There exists a valued arrow

$$\begin{array}{ccc} & (d_{ij}, d_{ji}) & \\ \circ & \xrightarrow{\quad} & \circ \\ i & & j \end{array}$$

if and only if ${}_iM_j \neq 0$. If $d_{ij} = d_{ji} = 1$, we write simply

$$\begin{array}{ccc} \circ & \xrightarrow{\quad} & \circ \\ i & & j \end{array}$$

The valued quiver of the species of a basic ring Λ is denoted by $(\Gamma_\Lambda, \mathbf{d})$. A valued quiver is called *connected* if the underlying valued graph of the valued quiver is connected. A valued quiver is said to be *acyclic* if it has no cycles. In this paper, unless otherwise stated, we assume that $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ is a species with the property “ ${}_iM_j \neq 0$ implies that ${}_jM_i = 0$ ”. Note that the species of any basic hereditary right artinian ring has this property; see [16]. Following [11, 13, 14, 33, 40], a *representation* of \mathcal{M} is a family $(X_i, {}_j\varphi_i)_{i,j \in I}$ of right F_i -modules X_i and F_j -linear maps ${}_j\varphi_i: X_i \otimes_{F_i} {}_iM_j \rightarrow X_j$ for each arrow $i \rightarrow j$ of Γ . A representation $(X_i, {}_j\varphi_i)$ is called *finite-dimensional* provided that all X_i are finite-dimensional right F_i -modules. A morphism $\alpha: (X_i, {}_j\varphi_i) \rightarrow (Y_i, {}_j\psi_i)$ is given by right F_i -linear maps $\alpha_i: X_i \rightarrow Y_i$ such that ${}_j\psi_i(\alpha_i \otimes id_{{}_iM_j}) = \alpha_{jj}{}_j\varphi_i$ for each arrow $i \rightarrow j$ of Γ . We denote by $\text{Rep}(\mathcal{M})$ (resp. by $\text{rep}(\mathcal{M})$) the category of all representations of \mathcal{M} (resp. finite-dimensional representations of \mathcal{M}) [19]. Obviously representations of species are a generalization of representations of quivers.

Following [1], a submodule K of a right Λ -module N is called *small submodule* if for every submodule L of N , $K + L = N$ implies $L = N$. Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a species with the valued quiver (Γ, \mathbf{d}) and $(X_i, {}_j\varphi_i)$ be a representation of \mathcal{M} . Suppose that $(Y_i, {}_j\psi_i)$ is a subrepresentation of the representation $(X_i, {}_j\varphi_i)$ of \mathcal{M} . Then $(X_i, {}_j\varphi_i)/(Y_i, {}_j\psi_i) = (X_i/Y_i, {}_j\gamma_i)$, where for each arrow $i \rightarrow j$ of Γ the maps ${}_j\gamma_i$ are defined by the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & Y_i \otimes_{F_i} {}_iM_j & \xrightarrow{\iota_i \otimes id_{{}_iM_j}} & X_i \otimes_{F_i} {}_iM_j & \xrightarrow{\pi_i \otimes id_{{}_iM_j}} & X_i/Y_i \otimes_{F_i} {}_iM_j \longrightarrow 0 \\ & & \downarrow {}_j\psi_i & & \downarrow {}_j\varphi_i & & \downarrow {}_j\gamma_i \\ 0 & \longrightarrow & Y_j & \xrightarrow{\iota_j} & X_j & \xrightarrow{\pi_j} & X_j/Y_j \longrightarrow 0 \end{array}$$

in which $\iota_i: Y_i \rightarrow X_i$ denotes the inclusion map and $\pi_i: X_i \rightarrow X_i/Y_i$ projection. Set $\pi = (\pi_i)_i: (X_i, {}_j\varphi_i) \rightarrow (X_i/Y_i, {}_j\gamma_i)$. The representation $(Y_i, {}_j\psi_i)$ is called a *small subrepresentation* of the representation $(X_i, {}_j\varphi_i)$ if every morphism $\alpha: (Z_i, {}_j\phi_i) \rightarrow (X_i, {}_j\varphi_i)$ in $\text{rep}(\mathcal{M})$ with $\pi\alpha$ epic is epic.

The following proposition gives the necessary and sufficient conditions for a representation $(Y_i, {}_j\psi_i)$ to be a small subrepresentation of $(X_i, {}_j\varphi_i)$.

Proposition 2.1 Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a species. Suppose that the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is connected and acyclic and let $(Y_i, {}_j\psi_i)$ be a subrepresentation of a representation $(X_i, {}_j\varphi_i)$ of \mathcal{M} . Then $(Y_i, {}_j\psi_i)$ is a small subrepresentation of $(X_i, {}_j\varphi_i)$ if and only if the following conditions hold:

- (i) If k is a source vertex of Γ , then $Y_k = 0$.
- (ii) If k is not a source vertex of Γ , then $Y_k \subseteq \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$, where the sum is over all arrows with the target k .

Proof (\Rightarrow). Assume that k is a source vertex of Γ and $X_k \neq 0$. Assume that $Y_k = X_k$. We define a representation $(Z_i, {}_j\theta_i)$ of \mathcal{M} by taking $Z_k = 0$, $Z_i = X_i$ for each $i \neq k$ and all ${}_j\theta_i = {}_j\varphi_i|_{Z_i \otimes_{F_i} M_j}$. Then $\pi\alpha$ is an epimorphism, where $\alpha = (\alpha_i)_i: (Z_i, {}_j\theta_i) \rightarrow (X_i, {}_j\varphi_i)$ and each $\alpha_i: Z_i \rightarrow X_i$ is the inclusion map. Since α is not an epimorphism, $(Y_i, {}_j\psi_i)$ is not a small subrepresentation of $(X_i, {}_j\varphi_i)$, which gives a contradiction. Hence, $Y_k \neq X_k$. Assume that H_k is an F_k -submodule of X_k such that $H_k + Y_k = X_k$. We define a subrepresentation $(X'_i, {}_j\varphi'_i)$ of $(X_i, {}_j\varphi_i)$ by taking $X'_k = H_k$, $X'_i = X_i$ for each $i \neq k$ and all ${}_j\varphi'_i = {}_j\varphi_i|_{X'_i \otimes_{F_i} M_j}$. Then $\pi\ell$ is an epimorphism, where $\ell = (\ell_i)_i: (X'_i, {}_j\varphi'_i) \rightarrow (X_i, {}_j\varphi_i)$ and each $\ell_i: X'_i \rightarrow X_i$ is the inclusion map. The assumption $(Y_i, {}_i\psi_j)$ is a small subrepresentation of $(X_i, {}_i\varphi_j)$ yields the map ℓ is an epimorphism and so $H_k = X_k$. This proves that Y_k is a small submodule of X_k . Since X_k is a right F_k -module and F_k is a division ring, Y_k is a direct summand of X_k . Therefore, $Y_k = 0$. Assume that k is not a source vertex of Γ and $\eta_k: X_k \rightarrow X_k / \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$ is the canonical quotient map, where the sum is over all arrows with the target k . Assume that $\eta_k(X_k) \neq 0$ and $\eta_k(Y_k) = \eta_k(X_k)$. Then $X_k = Y_k + \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$. We define a subrepresentation $(V_i, {}_j\varepsilon_i)$ of $(X_i, {}_j\varphi_i)$ by taking $V_k = \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$, $V_i = X_i$ for each $i \neq k$ and ${}_j\varepsilon_i = {}_j\varphi_i|_{V_i \otimes_{F_i} M_j}$ for each i and j . Therefore, $\pi\beta$ is an epimorphism, where $\beta = (\beta_i)_i: (V_i, {}_j\varepsilon_i) \rightarrow (X_i, {}_j\varphi_i)$ and each $\beta_i: V_i \rightarrow X_i$ is the inclusion map. Thus, $(Y_i, {}_j\psi_i)$ is not a small subrepresentation of $(X_i, {}_j\varphi_i)$, which gives a contradiction. It follows that $\eta_k(Y_k) \neq \eta_k(X_k)$. Assume that E_k is an F_k -submodule of $\eta_k(X_k)$ such that $E_k + \eta_k(Y_k) = \eta_k(X_k)$. Set $D_k = \eta_k^{-1}(E_k)$. Then ${}_k\varphi_j(X_j \otimes_{F_j} M_k) \subset D_k$ for each arrow $j \rightarrow k$ of Γ . Hence, $D_k + Y_k = X_k$ and $(X''_i, {}_j\varphi''_i)$ is a subrepresentation of $(X_i, {}_j\varphi_i)$, where $X''_k = D_k$, $X''_i = X_i$ for each $i \neq k$ and all ${}_j\varphi''_i = {}_j\varphi_i|_{X''_i \otimes_{F_i} M_j}$. Therefore, $\pi\xi$ is an epimorphism, where $\xi = (\xi_i)_i: (X''_i, {}_j\varphi''_i) \rightarrow (X_i, {}_j\varphi_i)$ and each $\xi_i: X''_i \rightarrow X_i$ is the inclusion map. The assumption that $(Y_i, {}_i\psi_j)$ is a small subrepresentation of $(X_i, {}_i\varphi_j)$ yields $D_k = X_k$. It follows that $E_k = \eta_k(X_k)$. Consequently, $\eta_k(Y_k)$ is a small submodule of $\eta_k(X_k)$. Therefore, $Y_k \subseteq \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$.

(\Leftarrow). Let $(Z_i, {}_j\phi_i)$ be a representation of \mathcal{M} and $f = (f_i)_i: (Z_i, {}_j\phi_i) \rightarrow (X_i, {}_j\varphi_i)$ be a morphism in $\text{Rep}(\mathcal{M})$ such that πf is an epimorphism. Then for each vertex i of Γ , $\text{Im } f_i + Y_i = X_i$. If k is a source vertex of Γ , then the assumption (i) yields $X_k = \text{Im } f_k$. Assume that k is not a source vertex of Γ . Let $\eta_k: X_k \rightarrow X_k / \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$ be the canonical quotient map. Since $\text{Im } f_k + Y_k = X_k$, $\eta_k(\text{Im } f_k) + \eta_k(Y_k) = \eta_k(X_k)$. Hence, the assumption (ii) yields $\eta_k(\text{Im } f_k) = \eta_k(X_k)$. Therefore, it is sufficient to show that for each arrow $j \rightarrow k$ of Γ , ${}_k\varphi_j(X_j \otimes_{F_j} M_k) \subseteq \text{Im } f_k$. Assume that there exists an arrow $j_1 \rightarrow k$ of Γ such that ${}_k\varphi_{j_1}(X_{j_1} \otimes_{F_{j_1}} M_k) \not\subseteq \text{Im } f_k$. Since ${}_j\varphi_i(f_i \otimes id_{i,M_j}) = f_{jj}\varphi_i$ for each arrow $i \rightarrow j$ of Γ , we have a representation $(\text{Im } f_i, {}_j\delta_i)$, where ${}_j\delta_i = {}_j\varphi_i|_{\text{Im } f_i \otimes_{F_i} M_j}$.

Thus, for each arrow $i \rightarrow j$ of Γ , we have the following commutative diagram

$$\begin{array}{ccc} \text{Im } f_i \otimes_{F_i} {}_iM_j & \xrightarrow{j\delta_i} & \text{Im } f_j \\ \ell_i \otimes 1 \downarrow & & \downarrow \ell_j \\ X_i \otimes_{F_i} {}_iM_j & \xrightarrow{j\varphi_i} & X_j, \end{array}$$

where each $\ell_i: \text{Im } f_i \rightarrow X_i$ is the inclusion map. Therefore, $\text{Im } f_{j_1} \neq X_{j_1}$. Since $\text{Im } f_{j_1} + Y_{j_1} = X_{j_1}$, Y_{j_1} is not a small submodule of X_{j_1} . By the above argument, j_1 is not a source vertex of Γ and by (ii), $\eta_{j_1}(Y_{j_1}) = 0$. Since $\text{Im } f_{j_1} + Y_{j_1} = X_{j_1}$, $\eta_{j_1}(\text{Im } f_{j_1}) + \eta_{j_1}(Y_{j_1}) = \eta_{j_1}(X_{j_1})$. It follows that $\eta_{j_1}(\text{Im } f_{j_1}) = \eta_{j_1}(X_{j_1})$. Therefore, there exists an arrow $j_2 \rightarrow j_1$ of Γ such that ${}_{j_1}\varphi_{j_2}(X_{j_2} \otimes_{F_{j_2}} {}_{j_2}M_{j_1}) \not\subseteq \text{Im } f_{j_1}$. It follows that j_2 is not a source vertex of Γ and the same argument shows that there exists an arrow $j_3 \rightarrow j_2$ of Γ such that ${}_{j_2}\varphi_{j_3}(X_{j_3} \otimes_{F_{j_3}} {}_{j_3}M_{j_2}) \not\subseteq \text{Im } f_{j_2}$. Continuing in this way, we get a cycle in (Γ, \mathbf{d}) which gives a contradiction. Then ${}_k\varphi_j(X_j \otimes_{F_j} {}_jM_k) \subseteq \text{Im } f_k$ for each arrow $j \rightarrow k$ of Γ . It follows that f is an epimorphism. Consequently, $(Y_i, {}_j\psi_i)$ is a small subrepresentation of $(X_i, {}_j\varphi_i)$. ■

In the following proposition, we compute the radical and top of a representation of species.

Proposition 2.2 *Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a species. Suppose that the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is connected and acyclic and let $(X_i, {}_j\varphi_i)$ be a finite-dimensional representation of \mathcal{M} . Then*

- (i) $\text{rad}((X_i, {}_j\varphi_i)) = (Y_i, {}_j\psi_i)$, where $Y_k = \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$ if k is not a source vertex of Γ , $Y_k = 0$ if k is a source vertex of Γ , and ${}_j\psi_i = {}_j\varphi_i|_{Y_i \otimes_{F_i} {}_iM_j}$ for each i and j .
- (ii) $\text{top}((X_i, {}_j\varphi_i)) = (Z_i, {}_j\gamma_i)$, where $Z_k = X_k / \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$ if k is not a source vertex of Γ , $Z_k = X_k$ if k is a source vertex of Γ and ${}_j\gamma_i = 0$ for each i and j .

Proof For each vertex k of Γ , define the representation $F_k = (W_i, {}_j\chi_i)$, where $W_i = 0$ for $i \neq k$, $W_k = F_k$ and all ${}_j\chi_i = 0$. Since the species \mathcal{M} is acyclic, by [40, Proposition 1.1], the simple representations F_k form a complete list of all non-isomorphic simple objects in $\text{rep}(\mathcal{M})$, where k is a vertex of Γ . Therefore $(V_i, {}_j\phi_i)$ is a maximal subrepresentation of $(X_i, {}_j\varphi_i)$ if and only if there exists a vertex k of Γ such that $(V_i, {}_j\phi_i)$ satisfies one of the following conditions:

- (i') k is a source vertex of Γ , V_k is a maximal submodule of X_k , $V_i = X_i$ for each $i \neq k$, and ${}_j\phi_i = {}_j\varphi_i|_{V_i \otimes_{F_i} {}_iM_j}$ for each i and j ,
- (ii') k is not a source vertex of Γ , V_k is a maximal submodule of X_k which contains $\sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$, $V_i = X_i$ for each $i \neq k$, and ${}_j\phi_i = {}_j\varphi_i|_{V_i \otimes_{F_i} {}_iM_j}$ for each i and j .

It follows that $\text{rad}((X_i, {}_j\varphi_i)) = (Y_i, {}_j\psi_i)$, where $Y_k = \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$ if k is not a source vertex of Γ , whereas $Y_k = 0$ if k is a source vertex of Γ and ${}_j\psi_i = {}_j\varphi_i|_{Y_i \otimes_{F_i} {}_iM_j}$ for each i and j . Consequently, $\text{top}((X_i, {}_j\varphi_i)) = (Z_i, {}_j\gamma_i)$, where $Z_k = X_k / \sum_{j \rightarrow k} \text{Im}({}_k\varphi_j)$ if k is not a source vertex of Γ , whereas $Z_k = X_k$ if k is a source vertex of Γ and ${}_j\gamma_i = 0$ for each i and j . ■

As an immediate consequence of Propositions 2.1 and 2.2, we obtain the following corollary.

Corollary 2.3 Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a species. Suppose that the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is connected and acyclic and let $(X_i, {}_i\varphi_j)$ be a finite-dimensional representation of \mathcal{M} . Then $\text{rad}((X_i, {}_j\varphi_i))$ is the biggest small subrepresentation of $(X_i, {}_i\varphi_j)$.

It is well known that, for a quiver $Q = (Q_0, Q_1)$, the category of representations of Q over a field K is equivalent to $\text{Mod-}KQ$, where KQ is the path K -algebra of Q ; see [2]. This fact was generalized nicely for species. For a species $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$, one can form a tensor ring $R_{\mathcal{M}} = \bigoplus_{t \geq 0} N^{(t)}$, where $A = N^{(0)} = \bigoplus_{i \in I} F_i$, $N^{(1)} = \bigoplus_{i,j \in I} {}_iM_j$ and $N^{(t)} = N^{(t-1)} \otimes_A N^{(1)}$ for $t \geq 2$, with the component-wise addition and the multiplication induced by taking tensor products. The ring $R_{\mathcal{M}}$ is called the *tensor ring of \mathcal{M}* (see [19]). Following [14, 40], a species \mathcal{M} is called *right (resp. left) finite-dimensional* if the dimensions d_{ij} (resp. d_{ji}) are finite for all $i \neq j$. The species \mathcal{M} is called *finite-dimensional* if \mathcal{M} is left and right finite-dimensional. Assume that $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ is a right finite-dimensional species and the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is acyclic. We now define as in [12] a functor

$$(2.1) \quad F: \text{rep}(\mathcal{M}) \longrightarrow \text{mod-}R_{\mathcal{M}}$$

as follows. For each object $X = (X_i, {}_j\varphi_i)$ in $\text{rep}(\mathcal{M})$, we set $F(X) = \bigoplus_{i \in I} X_i$. The reader can easily verify that $F(X)$ is a finitely generated right $R_{\mathcal{M}}$ -module. If $(\alpha_i)_{i \in I}: (X_i, {}_j\varphi_i) \rightarrow (Y_i, {}_j\psi_i)$ is a morphism in $\text{rep}(\mathcal{M})$, we define $F((\alpha_i)_{i \in I})$ to be $\bigoplus_{i \in I} \alpha_i: \bigoplus_{i \in I} X_i \rightarrow \bigoplus_{i \in I} Y_i$. It is easy to verify that $F((\alpha_i)_{i \in I})$ is an $R_{\mathcal{M}}$ -homomorphism. It is well known that the functor F is an equivalence (see [40, Proposition 1.1]). From now on, we fix the functor F . Note that when the species \mathcal{M} is right finite-dimensional, the tensor ring $R_{\mathcal{M}}$ is semiprimary. In this case, Propositions 2.1 and 2.2 and Corollary 2.3 are an automatic translation of categorical properties from modules to representations. In particular, we have the following fact.

Proposition 2.4 Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a right finite-dimensional species. Suppose that the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is connected and acyclic and let $(X_i, {}_j\varphi_i)$ be a finite-dimensional representation of \mathcal{M} . Then $\text{top}(F(X_i, {}_j\varphi_i)) \cong F(\text{top}((X_i, {}_j\varphi_i)))$.

3 Representations of Species and Modules over Hereditary Artinian Rings

Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a finite-dimensional species and suppose that the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is acyclic and connected. Let k be a sink (resp. source) vertex of Γ and let $\mathcal{M}^k = (F_i, {}_iN_j)_{i,j \in I}$ be the new species, where

$${}_iN_j = \begin{cases} {}_jM_k^L & \text{if } i = k, \\ {}_iM_j & \text{if } i \neq k \text{ and } j \neq k, \\ 0 & \text{if } j = k. \end{cases}$$

(resp. ${}_iN_k = {}_kM_i^R, {}_kN_i = 0$ for each i and ${}_iN_j = {}_iM_j$ for each $i \neq k, j \neq k$) (see [14]). We recall from [11, 33] (see also [4] and [13, 14, 40]) that a pair of *partial Coxeter functors* (or *reflection functors*, see [2, Ch. VII])

$$\text{rep}(\mathcal{M}) \begin{matrix} \xrightarrow{S_k^+} \\ \xleftarrow{S_k^-} \end{matrix} \text{rep}(\mathcal{M}^k)$$

is defined as follows. Let $X = (X_i, {}_j\varphi_i)$ be a finite-dimensional representation of \mathcal{M} . Define $S_k^+X = (Y_i, {}_j\psi_i)$, where $Y_i = X_i$ for each $i \neq k$ and Y_k is the kernel of the morphism $({}_k\varphi_j)_j$,

$$0 \longrightarrow Y_k \xrightarrow{({}_j\sigma_k)_j} \bigoplus_{j \in \Gamma_0} X_j \otimes_{F_j} {}_jM_k \xrightarrow{({}_k\varphi_j)_j} X_k.$$

By using the natural isomorphism

$$\text{Hom}_{F_j}(Y_k \otimes_{F_k} {}_jM_k^L, X_j) \cong \text{Hom}_{F_k}(Y_k, X_j \otimes_{F_j} {}_jM_k),$$

we get ${}_j\psi_k: Y_k \otimes_{F_k} {}_jM_k^L \rightarrow Y_j$ and ${}_j\psi_i = {}_j\varphi_i$ for $i \neq k$. Also, if $\alpha = (\alpha_i)_i: X \rightarrow X'$ is a morphism in $\text{rep}(\mathcal{M})$, then $S_k^+\alpha = (\beta_i)_i$ is defined by $\beta_i = \alpha_i$ for $i \neq k$ and $\beta_k: Y_k \rightarrow Y'_k$ as the restriction of

$$\bigoplus_{j \in \Gamma_0} (\alpha_j \otimes 1): \bigoplus_{j \in \Gamma_0} X_j \otimes {}_jM_k \longrightarrow \bigoplus_{j \in \Gamma_0} X'_j \otimes {}_jM_k.$$

Also, for each sink vertex k of Γ , define the linear transformation $s_k^+ : \mathbb{Z}^n \rightarrow \mathbb{Z}^n$ by $s_k^+x = y$, where $|\Gamma_0| = n$, $y_i = x_i$ for $i \neq k$ and $y_k = -x_k + \sum_{i \in \Gamma_0} d_{ik}x_i$. For each finite-dimensional indecomposable representation $X = (X_i, {}_j\varphi_i)$ of \mathcal{M} , we can see that each ${}_k\varphi_i$ is an epimorphism. It follows that $\dim S_k^+X = s_k^+(\dim X)$, where $\dim X = (\dim(X_i)_{F_i})_{i \in I}$. The functor S_k^- is defined analogously. The functors S_k^+ and S_k^- induce quasi-inverse equivalences between the full subcategory of $\text{rep}(\mathcal{M})$ of the representations having no direct summand isomorphic to the simple projective representation \mathbf{F}_k , and the full subcategory of $\text{rep}(\mathcal{M}^k)$ of the representations having no direct summand isomorphic to the simple injective representation \mathbf{F}_k (see [11, 14]).

Let k be a vertex of Γ and let $s_k\Gamma$ be the quiver obtained from Γ by reversing the direction of all arrows starting or end in k ; see [11] and [2, Ch. VII]. An admissible sequence of sinks in a quiver Γ is a sequence k_1, \dots, k_n of all vertices in Γ such that each vertex k_t is a sink in $s_{k_{t-1}} \cdots s_{k_1}\Gamma$ for all $1 \leq t \leq n$ [2, 11]. Let k_1, \dots, k_n be an admissible sequence in Γ and let $(k'_j)_j$ be a sequence of vertices of Γ , where $j \in \mathbb{Z}$,

$$k'_j = \begin{cases} k_{r+1} & \text{if } j \geq 0, j = tn + r, 0 \leq r < n; \\ k_{n-r+1} & \text{if } j < 0, -j = tn + r, 0 \leq r < n, \end{cases}$$

and t is a positive integer. For any $m \in \mathbb{Z}$, the species $\mathcal{M}^{(m)}$ is defined in [14] as follows:

$$\mathcal{M}^{(m)} = \begin{cases} (\mathcal{M}^{(m-1)})^{k'_{m-1}} & \text{if } m \geq 1; \\ (\mathcal{M}^{(m+1)})^{k'_m} & \text{if } m \leq -1, \end{cases}$$

where $\mathcal{M}^{(0)} = \mathcal{M}$. The species \mathcal{M} has the right (resp. left) finite-dimensional property if the species $\mathcal{M}^{(m)}$ are finite-dimensional for all $m \geq 0$ (resp. $m \leq 0$). \mathcal{M} has the finite-dimensional property if it has both the left and the right finite-dimensional property. If \mathcal{M} has the right (resp. left) finite-dimensional property, then the following sequence is a right (resp. left) sequence of partial Coxeter functors of \mathcal{M} :

$$\text{rep}(\mathcal{M}) \begin{matrix} \xleftarrow{s_1^+} \\ \xrightarrow{s_1^-} \end{matrix} \text{rep}(\mathcal{M}^{(1)}) \rightleftharpoons \dots \rightleftharpoons \text{rep}(\mathcal{M}^{(m-1)}) \begin{matrix} \xleftarrow{s_m^+} \\ \xrightarrow{s_m^-} \end{matrix} \text{rep}(\mathcal{M}^{(m)}) \rightleftharpoons \dots$$

$$(\text{resp.}, \dots \rightleftharpoons \text{rep}(\mathcal{M}^{(-m-1)}) \begin{matrix} \xleftarrow{s_m^+} \\ \xrightarrow{s_m^-} \end{matrix} \text{rep}(\mathcal{M}^{(-m)}) \rightleftharpoons \dots \rightleftharpoons \text{rep}(\mathcal{M}^{(-1)}) \begin{matrix} \xleftarrow{s_0^+} \\ \xrightarrow{s_0^-} \end{matrix} \text{rep}(\mathcal{M})),$$

where S_j^+ and S_j^- are the pair of partial Coxeter functors corresponding to the sink k'_{j-1} in the valued quiver of $\mathcal{M}^{(j-1)}$. We denote by $F_{k'_j}^{(j)}$ the simple projective representation in $\mathcal{M}^{(j)}$ corresponding to the sink k'_j [14].

We need the following result proved in [14, Theorem 1].

Theorem 3.1 *Let \mathcal{M} be a finite-dimensional species and suppose that its valued quiver (Γ, \mathbf{d}) is connected and acyclic. Then \mathcal{M} is representation-finite if and only if \mathcal{M} has the finite-dimensional property and there exists $m > 0$ such that $s_m^+ \cdots s_1^+(e_i) \not\cong 0$ for any source i of Γ . Moreover, if m is minimal with the above property and $|\Gamma_0| = n$, then the mapping $\text{dim}: \text{rep}(\mathcal{M}) \rightarrow \mathbb{Z}^n$ is a one-one correspondence between isomorphism classes of finite-dimensional indecomposable representations of \mathcal{M} and vectors in \mathbb{Z}^n of the form $s_1^- \cdots s_t^-(e_{k'_t})$, where $t < m$ and k'_t is a sink in the valued quiver of $\mathcal{M}^{(t)}$. In other words, any finite-dimensional indecomposable representation X of \mathcal{M} has the form $X \cong P_i$ for some $0 \leq i < m$, where $P_0 = F_{k_1}$ and $P_i = S_1^- \cdots S_i^- F_{k'_i}^{(i)}$.*

We recall from [13] that a sequence $a = (a_1, \dots, a_m)$ of length $m \geq 2$ with $a_i \in \mathbb{N}$ is called a *dimension sequence* provided there exist $x_i, y_i \in \mathbb{N}$ ($1 \leq i \leq m$), with

$$a_i x_i = x_{i-1} + x_{i+1} \quad \text{and} \quad a_i y_i = y_{i-1} + y_{i+1} \quad (1 \leq i < m),$$

where $x_0 = -1, y_0 = y_m = x_1 = 0$ and $x_m = y_1 = 1$.

Lemma 3.2 *Let F and G be division rings, M be an F - G -bimodule and $\Lambda = \begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ be an artinian ring. Then there exist precisely 3 pairwise non-isomorphic finitely generated indecomposable right Λ -modules if and only if $\text{r.dim } M = \text{l.dim } M = 1$. Moreover, in this case, $F \cong G$ as division rings and $M^L \cong M^R$ as G - F -bimodules.*

Proof Assume that $\Lambda = \begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ is an artinian ring such that there exist precisely 3 pairwise non-isomorphic, finitely generated, indecomposable right Λ -modules. Then by [39, Corollary 3.5], $d_3(M)$ is a dimension sequence. It follows that by [39, Lemma 3.1] and [13, Proposition 2], $\text{r.dim } M = \text{l.dim } M = 1$.

Conversely, assume that $\text{r.dim } M = \text{l.dim } M = 1$. Let $M = xG$ for some $0 \neq x \in M$. Define a division ring embedding $\alpha: F \rightarrow G$ by the formula $fx = x\alpha(f)$ for any $f \in F$. So we have an F - G -isomorphism $\varphi: M \rightarrow {}_{\alpha(F)}G_G$ by the formula $\varphi(xg) = g$. Since $\text{l.dim } M = 1$, we can assume that $M = Fx$. Define a division ring embedding $\beta: G \rightarrow F$ by the formula $xg = \beta(g)x$. Thus, we have an F - G -isomorphism $\psi: M \rightarrow {}_F F_{\beta(G)}$ by the formula $\psi(fx) = f$. Therefore, ${}_{\alpha(F)}G_G \cong M \cong {}_F F_{\beta(G)}$ as F - G -bimodules. It follows that $M^R \cong {}_G G_{\alpha(F)}$ and $M^L \cong {}_{\beta(G)}F_F$ as G - F -bimodules. Clearly, $\beta = \alpha^{-1}$ and the isomorphism $\psi\varphi^{-1}$ is just equal β . Notice that the same β yields also the isomorphism ${}_G G_{\alpha(F)} \cong {}_{\beta(G)}F_F$, so $M^L \cong M^R$ as G - F -bimodules. Therefore by [11, Proposition 2.6], there exist precisely 3 pairwise non-isomorphic finitely generated indecomposable right Λ -modules. ■


Let F and G be division rings and let M be an F - G -bimodule. We say that the F - G -bimodule M is *trivial* if there exist precisely 3 pairwise non-isomorphic finitely generated indecomposable right $\binom{F}{0} \binom{M}{G}$ -modules.

Let Λ be a basic hereditary ring and suppose that the number of vertices of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the natural number n . A vector $x \in \mathbb{N}^n$ is called *branch system* of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ if there exists an admissible sequence k_1, \dots, k_t in the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ such that $s_t^+ \cdots s_1^+ x = e_j$ for some $1 \leq j \leq n$. This generalizes the positive part of the usual rank 2 root systems (see [13]). The following well-known result is playing a key role in our study in this paper.

Proposition 3.3 *Let Λ be a basic hereditary artinian ring. Then the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is a disjoint union of the Coxeter diagrams presented in Table B if and only if Λ is a representation-finite ring. Moreover in this case:*

- (i) *there exists a bijection between the indecomposable finite-dimensional representations of \mathcal{M}_Λ and the branch system of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ ;*
- (ii) *the ring Λ is isomorphic with the tensor ring $R_{\mathcal{M}_\Lambda}$ of the species \mathcal{M}_Λ of Λ ;*
- (iii) *the functor $F: \text{rep}(\mathcal{M}_\Lambda) \rightarrow \text{mod-}\Lambda$ in (2.1) with $\Lambda = R_{\mathcal{M}_\Lambda}$ is an equivalence of categories;*
- (iv) *if the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is one of the Coxeter diagrams $\mathbb{A}_n, \mathbb{B}_n, \mathbb{D}_n, \mathbb{E}_6, \mathbb{E}_7, \mathbb{E}_8$ and \mathbb{F}_4 presented in Table B, then the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is one of the valued Dynkin diagrams presented in Table C.*

Table C. Valued Dynkin diagrams (consult also [37])



Proof This follows from [13, Theorem 2].

(i) This follows from Theorem 3.1, see also [13, Theorem 2].

(ii) The proof in the case when Λ is a basic hereditary finite-dimensional algebra over a field is given in [12]. The proof in the general case follows from [38, Theorem 4.5 and Corollary 4.6] together with [37, Lemma 3.3]; see also [16, Theorem 3].

(iii) By [39, Lemma 3.1 and Corollary 3.5] and [13, Proposition 2'], the species \mathcal{M}_Λ of Λ is finite-dimensional. Since the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is acyclic and connected, by [40, Proposition 1.1], the functor $F: \text{rep}(\mathcal{M}_\Lambda) \rightarrow \text{mod-}\Lambda$ in (2.1) with $\Lambda = R\mathcal{M}_\Lambda$ is an equivalence of categories.

(iv) Assume that the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is one of the Coxeter diagrams $\mathbb{A}_n, \mathbb{D}_n, \mathbb{E}_6, \mathbb{E}_7$, and \mathbb{E}_8 presented in Table B. Then by Lemma 3.2, the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is one of the valued Dynkin diagrams $\mathbb{A}_n, \mathbb{D}_n, \mathbb{E}_6, \mathbb{E}_7$ and \mathbb{E}_8 presented in Table C, respectively. If the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the Coxeter diagram \mathbb{B}_n presented in Table B, then by [39, Lemma 3.1 and Corollary 3.5] and [13, Proposition 2'], the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is one of the valued Dynkin diagrams \mathbb{B}_n and \mathbb{C}_n presented in Table C. If the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the Coxeter diagram \mathbb{F}_4 presented in Table B, by the same arguments the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the valued Dynkin diagram \mathbb{F}_4 presented in Table C. ■

Let Λ be a basic hereditary ring such that the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is one of the valued Dynkin diagrams presented in Table C. Therefore, there exist non-zero natural numbers f_i satisfying $d_{ij}f_j = d_{ji}f_i$ for each vertex i, j of Γ_Λ . Assume that the number of vertices of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the natural number n . We denote by \mathbb{Q}^n the vector space of all $x = (x_1, \dots, x_n)$ over the field of rational numbers. Then we define a symmetric positive definite bilinear form $B: \mathbb{Q}^n \times \mathbb{Q}^n \rightarrow \mathbb{Q}$ as follows. For each $x, y \in \mathbb{Q}^n$, $B(x, y) = \sum_i f_i x_i y_i - \frac{1}{2} \sum_{i,j} d_{ij} f_j x_i y_j$. For each vertex k of Γ_Λ , we have a reflection s_k , where $s_k : \mathbb{Q}^n \rightarrow \mathbb{Q}^n$ is a linear transformation given by $s_k x = x - (2B(x, e_k)/B(e_k, e_k))e_k$. A group of all linear transformations of \mathbb{Q}^n generated by the reflections s_k , k is a vertex of Γ_Λ , is called *Weyl group* and is denoted by \mathcal{W} . It is well known that the set $R = \{x \in \mathbb{Q}^n \mid x = we_k \text{ for some } w \in \mathcal{W} \text{ and } k \text{ is a vertex of } \Gamma_\Lambda\}$ is a reduced root system such that the set $\{e_k \mid k \text{ is a vertex of } \Gamma_\Lambda\}$ is a base for R ; see [8, 11, 24]. If the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is one of the Coxeter diagrams $\mathbb{A}_n, \mathbb{B}_n, \mathbb{D}_n, \mathbb{E}_6, \mathbb{E}_7, \mathbb{E}_8$, and \mathbb{F}_4 presented in Table B; then by Theorem 3.1, [33, Theorem 6.5] and [13, Theorem 1] (see also [11, Proposition 1.9]), there exists a bijection between the isomorphism classes of finite-dimensional indecomposable representations of \mathcal{M}_Λ and the positive roots of $(\Gamma_\Lambda, \mathbf{d})$. Now we assume that the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the Coxeter diagram \mathbb{G}_2 presented in Table B, then the category $\text{mod-}\Lambda$ has exactly 6 non-isomorphic indecomposable modules. By [39, Corollary 3.5], $d_6(M)$ is a dimension sequence. Therefore by [13, Proposition 2'], $d_6(M)$ is one of the following sequences up to cyclic permutation and reversion:

$$(1, 2, 2, 2, 1, 4), (1, 2, 3, 1, 2, 3), (1, 3, 1, 3, 1, 3).$$

If $d_6(M) = (1, 3, 1, 3, 1, 3)$, then by [39, Lemma 3.1 and Corollary 3.5], the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is \mathbb{G}_2 presented in Table C. Thus by Proposition 3.3 and [11, Proposition 1.9], the branch system of the $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is exactly the positive roots of $(\Gamma_\Lambda, \mathbf{d})$. But if $d_6(M)$ is one of the sequences (up to cyclic permutation and reversion) $(1, 2, 2, 2, 1, 4)$ and $(1, 2, 3, 1, 2, 3)$, then the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is one of the valued diagrams

$$\overset{(1,2)}{\circ} \text{---} \overset{(2,2)}{\circ} \text{---} \overset{(1,4)}{\circ} \text{---} \overset{(2,3)}{\circ} \text{---} \overset{(2,1)}{\circ} \text{---} \overset{(4,1)}{\circ} \text{---} \overset{(3,1)}{\circ}.$$

It follows that by Proposition 3.3, the branch system of its Coxeter valued quiver is different from the positive roots of the corresponding Dynkin diagram. Therefore in Section 5, by using reflection functors, we study the Köthe property for basic hereditary rings Λ of the Dynkin type \mathbb{G}_2 and of the Coxeter types $\mathbb{H}_3, \mathbb{H}_4,$ and $\mathbb{I}_2(p)$ with $p = 5$ or $7 \leq p < \infty$.

4 Right Köthe Rings of Dynkin Type

Following [31], we say that a finitely generated indecomposable right Λ -module M is *multiplicity-free top* if composition factors of $\text{top}(M)$ are pairwise non-isomorphic. The species \mathcal{M}_Λ has the *multiplicity-free top property* if every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top.

We start this section with the following fact that is frequently used in our study of right Köthe rings of Dynkin type.

Proposition 4.1 *The following two conditions are equivalent for a basic ring Λ .*

- (i) Λ is a right Köthe ring.
- (ii) Λ is artinian and every indecomposable right Λ -module of finite length is multiplicity-free top.

If, in addition, Λ is hereditary, then (i) is equivalent with the following statement:

- (iii) Λ is a representation-finite ring and the species \mathcal{M}_Λ has the multiplicity-free top property.

Proof The equivalence of (i) \Leftrightarrow (ii) is a consequence of [17, Corollary 3.3].


(ii) \Leftrightarrow (iii) Since Λ is assumed to be hereditary, the equivalence of (ii) and (iii) follows from Propositions 2.4 and 3.3, and the equivalence (i) \Leftrightarrow (ii) proved earlier. ■

Proposition 4.2 *If Λ is a basic hereditary artinian ring of the Dynkin type \mathbb{A}_n , then Λ is a right Köthe ring.*

Proof Assume that Λ is a basic hereditary artinian ring of the Dynkin type \mathbb{A}_n . Then there exists a bijection between the finite-dimensional indecomposable representations of \mathcal{M}_Λ and the positive roots of $(\Gamma_\Lambda, \mathbf{d})$. Since by [8, p. 265], the positive roots of $(\Gamma_\Lambda, \mathbf{d})$ are $\sum_{i \leq k < j} e_k$, where $1 \leq i < j \leq n + 1$, by Proposition 2.2, every finite-dimensional indecomposable representation of the species \mathcal{M}_Λ is multiplicity-free top. Therefore by Proposition 4.1, Λ is a right Köthe ring. ■

Proposition 4.3 *Let Λ be a basic hereditary ring of the Dynkin type \mathbb{D}_n . Then Λ is a right Köthe ring if and only if Λ is an artinian ring such that the following*

conditions hold:



- (i) $|(n - 2)^+| \leq 2;$
- (ii) For each $i \leq n - 3$, there exists at most one arrow with the source i .

Proof (\Rightarrow). Assume that $|(n - 2)^+| > 2$. Since by Proposition 3.3, the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the valued Dynkin diagram \mathbb{D}_n presented in Table C, by using [8, p. 271], there exists a finite-dimensional indecomposable representation X of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} X = {}_{0 \dots 0121}$. Therefore by Proposition 2.2, there exists a finite-dimensional indecomposable representation X of \mathcal{M}_Λ such that $\text{top}(X) \cong \mathbf{F}_{n-2} \oplus \mathbf{F}_{n-2}$, which is a contradiction by Proposition 4.1. Now, we show that for each $i \leq n - 3$, there exists at most one arrow with the source i . Assume that there exists $i \leq n - 3$ such that $|i^+| = 2$. Since by using [8, p. 271], there exists a finite-dimensional indecomposable representation Y of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} Y = {}_{0 \dots 012 \dots 2221}$, where the first 2 is in the i -th coordinate, by Proposition 2.2, there exists a finite-dimensional indecomposable representation Y of \mathcal{M}_Λ that is not multiplicity-free top. It follows that by Proposition 4.1, Λ is not right Köthe, which is a contradiction.

(\Leftarrow). Assume that $X = (X_i, j\varphi_i)$ is a finite-dimensional indecomposable representation of \mathcal{M}_Λ and there exists $1 \leq t \leq n$ such that $\mathbf{F}_t \oplus \mathbf{F}_t \subseteq \text{top}(X)$. Since by Proposition 3.3, the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the valued Dynkin diagram \mathbb{D}_n presented in Table C, by [8, p. 271], the positive roots of $(\Gamma_\Lambda, \mathbf{d})$ can be expressed as combinations of simple roots as follows:

- (1) $e_i + \dots + e_k$ $1 \leq i \leq k \leq n - 1;$
- (2) $e_i + \dots + e_k + 2e_{k+1} + \dots + 2e_{n-2} + e_{n-1} + e_n$ $1 \leq i \leq k \leq n - 3;$
- (3) $e_i + \dots + e_{n-2} + e_n$ $1 \leq i \leq n - 2;$
- (4) $e_i + \dots + e_{n-2} + e_{n-1} + e_n$ $1 \leq i \leq n - 2.$

By Proposition 2.2, $\dim(X_t)_{F_t} = 2$ and the vertex t is source. It follows that by (ii), $t = n - 2$. Therefore, $|(n - 2)^+| = 3$, which is a contradiction. Thus, every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Therefore, by Propositions 3.3 and 4.1, Λ is a right Köthe ring. ■

An *arm of length t* is a pair (Q', k) consisting of a quiver Q' of type \mathbb{A}_n presented in Table A and the vertex k of Q' , which has at most one neighbor in Q' . We say that a quiver Q has an arm (Q', k) if Q' is a full subquiver of Q and there are no arrows between the vertices of Q outside of Q' and the vertices of Q' different from k . Let c be a vertex of a quiver Δ of tree type. We say that an arrow $\alpha: x \rightarrow y$ points to c provided y and c belong to the same connected component of the quiver obtained from Δ by deleting α . Let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a species such that for any $i, j \in I$, $F_i \cong F_j$ as division rings and $\text{r.dim } {}_iM_j = \text{l.dim } {}_iM_j = 1$ and suppose that the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is of tree type. Let $X = (X_i, j\varphi_i)$ be a finite-dimensional representation of \mathcal{M}

and (Q', k) be an arm of (Γ, \mathbf{d}) . We say that X is *conical* on (Q', k) provided ${}_j\varphi_i$ is injective for any arrow $i \rightarrow j$ of Q' that points to k and for the remaining arrows $l \rightarrow t$ of Q' , ${}_t\varphi_l$ is surjective. The representation X is said to be *thin*, provided $\dim (X_i)_{F_i} \leq 1$ for all vertices i . The support of X is the set of vertices i with $X_i \neq 0$ [32].


The following lemma is a generalization of [32, Corollary 1.4].

Lemma 4.4 *Let F be a division ring and let $\mathcal{M} = (F_i, {}_iM_j)_{i,j \in I}$ be a species such that for each i and j , $F_i \cong F$ as division rings and ${}_iM_j \cong {}_F F_F$ as bimodule and the valued quiver (Γ, \mathbf{d}) of \mathcal{M} is of tree type. Let (Q', k) be an arm of (Γ, \mathbf{d}) . Then any finite-dimensional representation X of \mathcal{M} has a decomposition as $X = X' \oplus X''$, where X' is conical on (Q', k) and the support of X'' is contained in $Q' \setminus \{k\}$. In particular, if X is a finite-dimensional indecomposable representation of \mathcal{M} such that $X_k \neq 0$, then X is conical on (Q', k) .*

Proof Let $X = (X_i, {}_j\varphi_i)$ be a finite-dimensional representation of \mathcal{M} . Let \mathcal{M}' be a subspecies of \mathcal{M} with the valued quiver Q' . Then by [11, Theorem], there exists a bijection between the finite-dimensional indecomposable representations of \mathcal{M}' and the positive roots of Q' . Moreover, \mathcal{M}' is representation-finite. Therefore by using [8, p. 265], every finite-dimensional representation of \mathcal{M}' is a direct sum of thin indecomposable representations. Thus, the restriction $X|_{Q'}$ of X to Q' is a direct sum of thin indecomposable representations $X(j)$ with $j \in J$. Let J' be the set of indices $j \in J$ such that $X(j)_k \neq 0$ and let J'' be the set of indices $j \in J$ such that $X(j)_k = 0$. For each vertex t of Q' , we set $X'_t = \bigoplus_{j \in J'} X(j)_t$ and $X''_t = \bigoplus_{j \in J''} X(j)_t$. Moreover, if t is a vertex in $\Gamma \setminus Q'$, then we set $X'_t = X_t$ and $X''_t = 0$. Thus, $X = X' \oplus X''$, where $X' = (X'_i, {}_j\varphi_i)$ and $X'' = (X''_i, {}_j\varphi_i)$. Since the representations $X(j)$ with $j \in J'$ are conical on (Q', k) , X' is conical on (Q', k) . Also, since the representations $X(j)$ with $j \in J''$ satisfy $X(j)_k = 0$, the support of X'' is contained in $Q' \setminus \{k\}$. Therefore, the proof is complete. ■

Let \mathcal{M} be a species and assume that \mathcal{M}' is a subspecies of \mathcal{M} . If the species \mathcal{M} has the multiplicity-free top property, then clearly the species \mathcal{M}' has the multiplicity-free top property. This fact is frequently used in the rest of the paper.


Proposition 4.5 *Let Λ be a basic hereditary ring of the Dynkin type E_6 . Then Λ is a right Köthe ring if and only if Λ is an artinian ring such that the following conditions hold:*



- (i) $1 \leq |3^+| \leq 2, |4^+| \leq 1$ and $|2^+| \leq 1$;
- (ii) For each $y \in 3^-$, there exists at least one arrow with the target y .


Proof (\Rightarrow). Assume that Λ is a right Köthe ring. Let \mathcal{M}' be a subspecies of \mathcal{M}_Λ such that the underlying valued graph of the valued quiver (Γ', \mathbf{d}) of \mathcal{M}' is the valued

Dynkin diagram \mathbb{D}_5 presented in Table C. Then by using Proposition 4.3, $|3^+| \leq 2$ and there exists at most one arrow with the source 2 and one arrow with the source 4. Consequently, $|4^+| \leq 1$ and $|2^+| \leq 1$. If $|3^+| = 0$, then by Proposition 3.3, the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ has the orientation




Hence by using [8, p. 275], there exists a finite-dimensional indecomposable representation X of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} X = {}_{12321}^2$. It follows that by Proposition 2.2, there exists a finite-dimensional indecomposable representation X of \mathcal{M}_Λ such that $\mathbf{F}_6 \oplus \mathbf{F}_6 \subseteq \text{top}(X)$, which is a contradiction, by Proposition 4.1. Therefore, $1 \leq |3^+| \leq 2$. Now we show that for each $y \in 3^-$, there exists at least one arrow with the target y . Assume that there exists $y \in 3^-$ such that there is no arrow with the target y . Since $|4^+| \leq 1$ and $|2^+| \leq 1$, $y = 6$. Therefore by Proposition 2.2, $\mathbf{F}_6 \oplus \mathbf{F}_6 \subseteq \text{top}(X)$, where X is a finite-dimensional indecomposable representation of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} X = {}_{12321}^2$. Hence, X is not multiplicity-free top, which is a contradiction, by Proposition 4.1.

(\Leftarrow). Let $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i,j \in I}$ be the species of Λ . Assume that $X = (X_i, {}_j\varphi_i)$ is a finite-dimensional indecomposable representation of \mathcal{M}_Λ such that $\mathbf{F}_i \oplus \mathbf{F}_i \subseteq \text{top}(X)$ for some vertex i of Γ_Λ . Since by Lemma 3.2, there exists a division ring F such that each $F_i \cong F$ as division rings, by using [8, p. 275], $\dim (X_1)_F \leq 1$, $\dim (X_2)_F \leq 2$, $\dim (X_3)_F \leq 3$, $\dim (X_4)_F \leq 2$, $\dim (X_5)_F \leq 1$, and $\dim (X_6)_F \leq 2$. Thus by assumptions (i) and(ii) and Proposition 2.2, $i = 3$ and $\dim (X_3)_F = 3$. It follows that by Lemma 4.4, X is one of the following representations




where $\dim(\text{Im}({}_3\varphi_2))_F = 1$ and $\dim(\text{Im}({}_3\varphi_4))_F = 1$. Since there exists a bijection between the finite-dimensional indecomposable representations of \mathcal{M}_Λ and the positive roots of $(\Gamma_\Lambda, \mathbf{d})$, by using [8, p. 275], there are no indecomposable representations of \mathcal{M}_Λ with the dimension vectors ${}^*13^*{}^*$ or ${}^*{}^*31^*$. It follows that X is not indecomposable, which is a contradiction. Thus, every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Therefore by Propositions 3.3 and 4.1, Λ is a right Köthe ring. ■

Proposition 4.6 *Let Λ be a basic hereditary ring of the Dynkin type \mathbb{E}_7 . Then Λ is a right Köthe ring if and only if Λ is an artinian ring such that the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ has the orientation*



Proof (\Rightarrow). Assume that Λ is a right Köthe ring. Then by the same argument as in the proof of Propositions 4.3 and 4.5, we can see that $1 \leq |3^+| \leq 2$, $|4^+| \leq 1$, $|5^+| \leq 1$, and $|2^+| \leq 1$ and for each $y \in 3^-$, there exists at least one arrow with the target y . Since by using [8, p. 279], there exists a finite-dimensional indecomposable representation Y of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} Y = {}_234321$, by Propositions 2.2 and 4.1, $(\mathcal{C}_\Lambda, \mathbf{m})$ has the orientation



(\Leftarrow). Let $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i,j \in I}$ be the species of Λ . Assume that $X = (X_i, {}_j\psi_i)$ is a finite-dimensional indecomposable representation of \mathcal{M}_Λ such that $F_j \oplus F_j \subseteq \text{top}(X)$ for some vertex j of Γ_Λ . Since Λ is of the Dynkin type \mathbb{E}_7 , by Lemma 3.2, there exists a division ring F such that for each $i \in I$, $F_i \cong F$ as division rings. If $X_3 = 0$, then by Proposition 4.2, X is multiplicity-free top, which is a contradiction. Hence $X_3 \neq 0$. Therefore by Lemma 4.4, X is the representation

$$\begin{array}{ccccccc}
 & & X_6 & & & & \\
 & & \uparrow & & & & \\
 X_1 & \longleftarrow & X_2 & \longleftarrow & X_3 & \longleftarrow & X_4 & \longleftarrow & X_5 & \longleftarrow & X_7.
 \end{array}$$

If $X_7 = 0$, by the proof of Proposition 4.5, X is not indecomposable, which is a contradiction. Thus, $X_7 \neq 0$. If $\dim(X_5)_F = 2$, then by Lemma 4.4, $\overline{X} = \overline{Z}' \oplus \overline{Z}''$, where

$$\begin{aligned}
 \overline{X} &:= X_1 \longleftarrow X_2 \oplus X_6 \longleftarrow X_3 \longleftarrow X_4 \longleftarrow X_5 \longleftarrow X_7, \\
 \overline{Z}' &:= X'_1 \longleftarrow X'_2 \oplus X'_6 \longleftarrow X'_3 \longleftarrow X'_4 \longleftarrow X'_5 \longleftarrow 0, \\
 \overline{Z}'' &:= X''_1 \longleftarrow X''_2 \oplus X''_6 \longleftarrow X''_3 \longleftarrow X''_4 \longleftarrow X''_5 \xleftarrow{5\psi_7} X_7
 \end{aligned}$$

and ${}_5\psi_7: X_7 \rightarrow X_5''$ is an isomorphism. Thus, $X = Z' \oplus Z''$, where

$$\begin{array}{ccccccc}
 & & & & X'_6 & & \\
 & & & & \uparrow & & \\
 Z' := & X'_1 & \longleftarrow & X'_2 & \longleftarrow & X'_3 & \longleftarrow & X'_4 & \longleftarrow & X'_5 & \longleftarrow & 0 \\
 & & & & & & & & & & & \\
 & & & & X''_6 & & \\
 & & & & \uparrow & & \\
 Z'' := & X''_1 & \longleftarrow & X''_2 & \longleftarrow & X''_3 & \longleftarrow & X''_4 & \longleftarrow & X''_5 & \xleftarrow{{}_5\psi_7} & X_7.
 \end{array}$$

Since by using [8, p. 278], $\dim(X_7)_F = 1, X_5' \neq 0$. It follows that X is not indecomposable, which is a contradiction. Therefore, $\dim(X_5)_F \neq 2$. Since by using [8, p. 278], $\dim(X_1)_F \leq 2, \dim(X_2)_F \leq 3, \dim(X_3)_F \leq 4, \dim(X_4)_F \leq 3, \dim(X_5)_F \leq 2, \dim(X_6)_F \leq 2$ and $\dim(X_7)_F \leq 1$, so $\dim(X_7)_F = \dim(X_5)_F = 1$. Hence by Proposition 2.2, $j = 3$ or $j = 4$. If $j = 4$, then $\dim(X_4)_F = 3$. Set

$$\begin{array}{ccccccc}
 & & & & X_6 & & \\
 & & & & \uparrow & & \\
 T := & X_1 & \longleftarrow & X_2 & \longleftarrow & X_3 & \longleftarrow & X_4 & \xleftarrow{{}_4\psi_5} & X_5.
 \end{array}$$

Thus, by Lemma 4.4, $\bar{T} = \bar{T}' \oplus \bar{T}''$, where

$$\begin{array}{l}
 \bar{T} := X_1 \longleftarrow X_2 \oplus X_6 \longleftarrow X_3 \longleftarrow X_4 \longleftarrow X_5, \\
 \bar{T}' := X'_1 \longleftarrow X'_2 \oplus X'_6 \longleftarrow X'_3 \longleftarrow X'_4 \longleftarrow 0, \\
 \bar{T}'' := X''_1 \longleftarrow X''_2 \oplus X''_6 \longleftarrow X''_3 \longleftarrow X''_4 \xleftarrow{{}_4\psi_5} X_5
 \end{array}$$

and ${}_4\psi_5: X_5 \rightarrow X_4''$ is an isomorphism. So $X = T' \oplus T''$, where

$$\begin{array}{ccccccc}
 & & & & X'_6 & & \\
 & & & & \uparrow & & \\
 T' := & X'_1 & \longleftarrow & X'_2 & \longleftarrow & X'_3 & \longleftarrow & X'_4 & \longleftarrow & 0 & \longleftarrow & 0, \\
 & & & & & & & & & & & \\
 & & & & X''_6 & & \\
 & & & & \uparrow & & \\
 T'' := & X''_1 & \longleftarrow & X''_2 & \longleftarrow & X''_3 & \longleftarrow & X''_4 & \xleftarrow{{}_4\psi_5} & X_5 & \longleftarrow & X_7.
 \end{array}$$

Since $\dim(X_5)_F = 1, X_4' \neq 0$ and $T'' \neq 0$. Consequently, X is not indecomposable, which is a contradiction. Now assume that $j = 3$. Therefore, either $\dim(X_3)_F = 3$ and $\dim(\text{Im}({}_3\psi_4))_F = 1$ or $\dim(X_3)_F = 4$ and $1 \leq \dim(\text{Im}({}_3\psi_4))_F \leq 2$. Set

$$\begin{array}{ccccccc}
 & & & & X_6 & & \\
 & & & & \uparrow & & \\
 H := X_1 & \longleftarrow & X_2 & \longleftarrow & X_3 & \longleftarrow & X_4.
 \end{array}$$

Then by Lemma 4.4, $\overline{H} = \overline{H'} \oplus \overline{H''}$, where

$$\begin{aligned}
 \overline{H} &:= X_1 \longleftarrow X_2 \oplus X_6 \longleftarrow X_3 \longleftarrow X_4, \\
 \overline{H'} &:= X'_1 \longleftarrow X'_2 \oplus X'_6 \longleftarrow X'_3 \longleftarrow 0, \\
 \overline{H''} &:= X''_1 \longleftarrow X''_2 \oplus X''_6 \longleftarrow X''_3 \xleftarrow{{}_3\psi_4} X_4,
 \end{aligned}$$


and ${}_3\psi_4: X_4 \rightarrow X''_3$ is an isomorphism. It follows that $X = H' \oplus H''$, where

$$\begin{array}{ccccccc}
 & & & & X'_6 & & \\
 & & & & \uparrow & & \\
 H' := X'_1 & \longleftarrow & X'_2 & \longleftarrow & X'_3 & \longleftarrow & 0 \longleftarrow 0 \longleftarrow 0, \\
 & & & & X''_6 & & \\
 & & & & \uparrow & & \\
 H'' := X''_1 & \longleftarrow & X''_2 & \longleftarrow & X''_3 & \xleftarrow{{}_3\psi_4} & X_4 \longleftarrow X_5 \longleftarrow X_7,
 \end{array}$$

and $X'_3 \neq 0$. Consequently, X is not indecomposable, which is a contradiction. Hence every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Therefore, by Propositions 3.3 and 4.1, Λ is a right Köthe ring. ■

Lemma 4.7 *If Λ is a basic hereditary artinian ring of the Dynkin type \mathbb{E}_8 , then Λ is not a right Köthe ring.*

Proof Assume that Λ is a right Köthe ring. Then by Proposition 4.1, every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Thus by using Proposition 4.6, $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is one of the following quivers:



Since by using [8, p. 284], there exists a finite-dimensional indecomposable representation Z of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} Z = {}^3_{2465432}$, by Proposition 2.2,

either $F_8 \oplus F_8 \subseteq \text{top}(Z)$ or $F_7 \oplus F_7 \oplus F_7 \subseteq \text{top}(Z)$, which is a contradiction. Therefore, Λ is not right Köthe. ■

A Characterization of Right Köthe Hereditary Rings

In this section we give a characterization of right Köthe hereditary rings.

Theorem 5.1 *Let F and G be division rings and M be an F - G -bimodule. Then $\Lambda = \begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ is a right Köthe ring if and only if there exists $m \geq 3$ such that $d_m(M) = (m - 2, 1, 2, \dots, 2, 1)$ is a dimension sequence.*

Proof We start by an observation that is a direct consequence of Proposition 2.2 (see also Proposition 2.4). For a basic hereditary ring $\Lambda (\cong R_{\mathcal{M}_\Lambda})$ as above being right artinian, a finitely generated indecomposable right Λ -module $N = (X_F, Y_G, \varphi: X \otimes_F M \rightarrow Y)$ is multiplicity-free top if and only if either N is a simple projective module and then $(\dim(X)_F, \dim(Y)_G) = (0, 1)$ or $\dim(X)_F = 1$. This follows from the fact that φ is surjective and $X_F \neq 0$ in case N is not simple projective.

(\Rightarrow). Assume that $\Lambda = \begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ is a right Köthe ring. Then there exists $m \geq 3$ such that Λ has only m pairwise non-isomorphic finitely generated indecomposable right Λ -modules of finite length. Thus, by [40, Proposition 1.1], \mathcal{M}_Λ is a finite-dimensional species. Hence by Theorem 3.1, \mathcal{M}_Λ has the finite-dimensional property and $\{P_0, P_1, \dots, P_{m-1}\}$ is the set of all finite-dimensional indecomposable representations (up to isomorphism) of \mathcal{M}_Λ , where P_0 is a simple projective representation and $P_i = S_1^- \cdots S_i^- F_{k_i}^{(i)}$, for each $1 \leq i \leq m - 1$. By using [39, Proposition 3.2] and [40, Proposition 1.1] (see also [13, Proposition 1]), $M^{(m-2)L} \cong M^{2R}$ as bimodules and by using of [40, Lemma 1.3] (see also [13, Proposition 1]), for each $1 \leq t \leq m - 2$, $\dim P_{t+1} = d_t^M \dim P_t - \dim P_{t-1}$. It follows that $d_m(M) = (d_0^M, d_1^M, \dots, d_{m-1}^M)$ is a dimension sequence. Since Λ is right Köthe, by Proposition 4.1, every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Assume that $m > 3$ and $\dim P_i = (x_i, y_i)$, for each $0 \leq i \leq m - 1$ (comparing to the formula before Lemma 3.2 for the sequence $a := d_m(M)$, the indexing of x_i 's and y_i 's are shifted by 1). Then by the entrance observation $x_i = 1$, for every $i = 1, \dots, m - 1$. Since $\dim P_0 = (0, 1)$ and $\dim P_1 = (1, d_0^M)$, we have $d_1^M = x_2 = 1$ and consequently $d_0^M > 1$, since otherwise $m = 3$ by Lemma 3.2. Also, we have $x_3 = d_2^M - 1$, so $d_2^M = 2$. Proceeding in a similar way, by induction we obtain

$$d_m(M) = (d_0^M, 1, 2, \dots, 2, d_{m-1}^M).$$

Consequently, by [13, Proposition 2'] and Lemma 3.2, $(d_0^M - (m - 3), 1, d_{m-1}^M)$ is a dimension sequence. Hence, we infer that $d_0^M = m - 2$ and $d_{m-1}^M = 1$. Therefore, $d_m(M) = (m - 2, 1, 2, \dots, 2, 1)$.

(\Leftarrow). Assume that there exists $m \geq 3$ such that $d_m(M) = (m - 2, 1, 2, \dots, 2, 1)$ is a dimension sequence. Then by [39, Corollary 3.5], Λ has only m pairwise non-isomorphic finitely generated indecomposable right Λ -modules of finite length. Thus, Λ is representation-finite. It follows from [40, Proposition 1.1] that Λ is a basic hereditary artinian ring and \mathcal{M}_Λ is representation-finite. Therefore by Theorem 3.1,

$\{\mathbf{P}_0, \mathbf{P}_1, \dots, \mathbf{P}_{m-1}\}$ is the set of all finite-dimensional indecomposable representations (up to isomorphism) of \mathcal{M}_Λ , where \mathbf{P}_0 is a simple projective representation and $\mathbf{P}_i = S_1^- \cdots S_i^- F_{k_i}^{(i)}$, for each $1 \leq i \leq m - 1$. Since $d_m(M) = (m - 2, 1, 2, \dots, 2, 1)$ and by using [40, Lemma 1.3] (see also [13, Proposition 1]), $\dim \mathbf{P}_{t+1} = d_t^M \dim \mathbf{P}_t - \dim \mathbf{P}_{t-1}$ for each $1 \leq t \leq m - 2$, so computing inductively, the consecutive dimension vectors, with the starting values $\dim \mathbf{P}_0 = (0, 1)$ and $\dim \mathbf{P}_1 = (1, m - 2)$, we infer that $x_i = 1$, for every $i = 1, \dots, m - 1$, where (x_i, y_i) are as above. Thus by entrance observation and Proposition 2.2, every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Therefore by Proposition 4.1, Λ is a right Köthe ring. ■


Example 5.2 Let $\Lambda = \begin{pmatrix} \mathbb{C} & \mathbb{C} \\ 0 & \mathbb{R} \end{pmatrix}$, where \mathbb{C} is the field of complex numbers and \mathbb{R} is the field of real numbers. Then Λ is a basic hereditary artinian ring and the category $\text{mod-}\Lambda$ has exactly 4 non-isomorphic indecomposable modules, since the underlying valued graph of $(\Gamma_\Lambda, \mathbf{d})$ is equal to \mathbb{B}_2 from Table C and Λ is a finite-dimensional algebra over its center $\mathbb{R}.I_2$, in particular $(\mathbb{C}\mathbb{C}_\mathbb{R})^L \cong (\mathbb{C}\mathbb{C}_\mathbb{R})^R$; see [11]. Thus, the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is \mathbb{B}_2 presented in Table B (i.e., $\circ \xrightarrow{4} \circ$) and by the very definition $d_4(\mathbb{C}\mathbb{C}_\mathbb{R}) = (2, 1, 2, 1)$. Therefore by Theorem 5.1, Λ is a right Köthe ring.

Now let $\Lambda = \begin{pmatrix} \mathbb{R} & \mathbb{C} \\ 0 & \mathbb{C} \end{pmatrix}$. Then Λ has the analogous properties as in the previous case with the one exception; namely, $d_4(\mathbb{R}\mathbb{C}_\mathbb{C}) = (1, 2, 1, 2)$, hence Λ is not a right Köthe ring. Notice that in this case, $\dim \mathbf{P}_2 = (2, 1)$, so the representation \mathbf{P}_2 of \mathcal{M}_Λ is not multiplicity-free top.


Let Λ be a basic hereditary artinian ring such that the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is one of the Coxeter diagrams $\mathbb{B}_n, \mathbb{F}_4, \mathbb{H}_3, \mathbb{H}_4, \mathbb{G}_2$, and $\mathbb{I}_2(p)$ presented in Table B. Then by Proposition 3.3, $\Lambda \cong R_{\mathcal{M}_\Lambda}$. Assume that Λ is a right Köthe ring. Then by Proposition 4.1, \mathcal{M}_Λ has the multiplicity-free top property. Let $\mathcal{M}' = (F, G, M)$ be a subspecies of \mathcal{M}_Λ such that $\begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ has only $m \geq 3$ pairwise non-isomorphic indecomposable right modules. Therefore by Theorem 5.1, $d_m(M) = (m - 2, 1, 2, \dots, 2, 1)$ is a dimension sequence.

Proposition 5.3 Let Λ be a basic hereditary ring of the Dynkin type \mathbb{B}_n . Assume that $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i,j \in I}$ is the species of Λ . Then Λ is a right Köthe ring if and only if Λ is an artinian ring such that one of the following conditions holds:

- (i) $d_4({}_{n-1}M_n) = (2, 1, 2, 1)$ is a dimension sequence and the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the quiver




- (ii) $d_4({}_nM_{n-1}) = (2, 1, 2, 1)$ is a dimension sequence and the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the quiver



where $1 \leq t \leq n - 1$.


Proof (\Rightarrow). Assume that Λ is a right Köthe ring. Then by Proposition 4.1, every finite-dimensional indecomposable representation of the species \mathcal{M}_Λ has multiplicity-free top. The underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the Coxeter diagram



thus by Theorem 5.1, either $d_4({}_{n-1}M_n) = (2, 1, 2, 1)$ is a dimension sequence or $d_4({}_nM_{n-1}) = (2, 1, 2, 1)$ is a dimension sequence. If $d_4({}_{n-1}M_n) = (2, 1, 2, 1)$ is a dimension sequence, then by Proposition 3.3, the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ is the valued Dynkin diagram \mathbb{B}_n presented in Table C. Since there exists a bijection between the isomorphism classes of finite-dimensional indecomposable representations of \mathcal{M}_Λ and the positive roots of $(\Gamma_\Lambda, \mathbf{d})$ and since by [8, pp. 267–268], the positive roots of $(\Gamma_\Lambda, \mathbf{d})$ can be expressed as combinations of simple roots as follows:

- (1) $\sum_{i \leq k \leq n} e_k \qquad (1 \leq i \leq n),$
- (2) $\sum_{i \leq k \leq j} e_k \qquad (1 \leq i < j \leq n),$
- (3) $\sum_{i \leq k < j} e_k + 2 \sum_{j \leq k \leq n} e_k \qquad (1 \leq i < j \leq n),$


thus by Proposition 2.2, the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is



If $d_4({}_nM_{n-1}) = (2, 1, 2, 1)$ is a dimension sequence, then by Proposition 3.3, the underlying valued graph of the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ is the valued Dynkin diagram \mathbb{C}_n presented in Table C. Since there exists a bijection between the isomorphism classes of finite-dimensional indecomposable representations of \mathcal{M}_Λ and the positive roots of $(\Gamma_\Lambda, \mathbf{d})$ and also since by using [8, pp. 269–270], the positive roots of $(\Gamma_\Lambda, \mathbf{d})$ can be expressed as combinations of simple roots as follows:

- (1) $\sum_{i \leq k < j} e_k \qquad (1 \leq i < j \leq n),$
- (2) $\sum_{i \leq k < j} e_k + 2 \sum_{j \leq k < n} e_k + e_n \qquad (1 \leq i < j \leq n),$
- (3) $\sum_{i \leq k < n} e_k + e_n \qquad (1 \leq i \leq n);$

thus, by Proposition 2.2, the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is



where $1 \leq t \leq n - 1$.

(\Leftarrow). Assume that $d_4(n_{-1}M_n) = (2, 1, 2, 1)$ is a dimension sequence and the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the quiver

$$\circ \xleftarrow{4} \circ \text{-----} \circ \xleftarrow{2} \circ \text{-----} \circ$$

$n \quad n-1 \quad 2 \quad 1$

Then by Proposition 3.3, the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the valued Dynkin quiver

$$\mathbb{B}_n : \quad \circ \xleftarrow{(1,2)} \circ \text{-----} \circ \xleftarrow{2} \circ \text{-----} \circ$$

$n \quad n-1 \quad 2 \quad 1$

Thus by using [8, pp. 267–268] and Proposition 2.2, \mathcal{M}_Λ has the multiplicity-free top property. Therefore by Propositions 3.3 and 4.1, Λ is a right Köthe ring.

Now we assume that $d_4(nM_{n-1}) = (2, 1, 2, 1)$ is a dimension sequence and the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is the quiver

$$\circ \xrightarrow{4} \circ \text{-----} \circ \xrightarrow{2} \circ \text{-----} \circ \xleftarrow{2} \circ \text{-----} \circ \xleftarrow{2} \circ$$

$n \quad n-1 \quad t-1 \quad t \quad t+1 \quad 2 \quad 1$

where $1 \leq t \leq n - 1$. Thus by Proposition 3.3, the valued quiver $(\Gamma_\Lambda, \mathbf{d})$ of Λ is the valued Dynkin quiver

$$\mathbb{C}_n : \quad \circ \xrightarrow{(2,1)} \circ \text{-----} \circ \xrightarrow{2} \circ \text{-----} \circ \xleftarrow{2} \circ \text{-----} \circ \xleftarrow{2} \circ$$

$n \quad n-1 \quad t-1 \quad t \quad t+1 \quad 2 \quad 1$

where $1 \leq t \leq n - 1$. It follows that by using [8, pp. 269–270] and Proposition 2.2, \mathcal{M}_Λ has the multiplicity-free top property. Consequently by Propositions 3.3 and 4.1, Λ is a right Köthe ring. ■

Lemma 5.4 *If Λ is a basic hereditary artinian ring of the Dynkin type \mathbb{F}_4 , then Λ is not a right Köthe ring.*


Proof Let F and G be division rings and let M be an F - G -bimodule such that $\mathcal{M}' = (F, G, M)$ is a subspecies of \mathcal{M}_Λ and $\begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ has only 4 pairwise non-isomorphic indecomposable right modules. Assume to the contrary that Λ is a right Köthe ring. Then by Theorem 5.1, $d_4(M) = (2, 1, 2, 1)$. It follow from Proposition 3.3 that the valued quiver (Γ, \mathbf{d}) of Λ is one of the valued Dynkin quivers

$$\begin{aligned} \mathbb{F}_4 : & \quad \circ \text{-----} \circ \xrightarrow{(1,2)} \circ \text{-----} \circ \text{-----} \circ \\ \mathbb{F}_4 : & \quad \circ \text{-----} \circ \xrightarrow{(1,2)} \circ \text{-----} \circ \text{-----} \circ \\ \mathbb{F}_4 : & \quad \circ \text{-----} \circ \xleftarrow{(1,2)} \circ \text{-----} \circ \text{-----} \circ \\ \mathbb{F}_4 : & \quad \circ \text{-----} \circ \xleftarrow{(1,2)} \circ \text{-----} \circ \text{-----} \circ \end{aligned}$$

Since by using [8, p. 287], there exists a finite-dimensional indecomposable representation X of \mathcal{M}_Λ with the dimension vector $\mathbf{dim} X = 2342$, by Proposition 2.2, \mathcal{M}_Λ has no the multiplicity-free top property, which is a contradiction, by Proposition 4.1. Therefore, Λ is not right Köthe. ■

Lemma 5.5 *If Λ is a basic hereditary artinian ring of the Coxeter type \mathbb{H}_3 , then Λ is not a right Köthe ring.*


Proof Let F and G be division rings and let M be an F - G -bimodule such that $\mathcal{M}' = (F, G, M)$ is a subspecies of \mathcal{M}_Λ and $\begin{pmatrix} F & M \\ 0 & G \end{pmatrix}$ has only 5 pairwise non-isomorphic indecomposable right modules. Assume to the contrary that Λ is a right Köthe ring. Then by Proposition 4.1, every finite-dimensional indecomposable representation of the species \mathcal{M}_Λ has multiplicity-free top. Thus by Theorem 5.1, $d_5(M) = (3, 1, 2, 2, 1)$. If $(\mathcal{C}_\Lambda, \mathbf{m})$ is the quiver



then

$$\begin{aligned} \dim \mathbf{P}_5 &= s_1^- s_2^- s_3^- s_4^- s_5^- (0, 0, 1) = s_1^- s_2^- s_3^- s_4^- (0, 2, 1) = s_1^- s_2^- s_3^- (0, 2, 3) \\ &= s_1^- s_2^- (2, 2, 3) = s_1^- (2, 3, 3) = (2, 3, 6). \end{aligned}$$


Hence by Proposition 2.2, $\mathbf{F}_1 \oplus \mathbf{F}_1 \subseteq \text{top}(\mathbf{P}_5)$, which is a contradiction. If $(\mathcal{C}_\Lambda, \mathbf{m})$ is the quiver




then

$$\begin{aligned} \dim \mathbf{P}_4 &= s_1^- s_2^- s_3^- s_4^- (0, 0, 1) = s_1^- s_2^- s_3^- (0, 2, 1) = s_1^- s_2^- (0, 2, 1) \\ &= s_1^- (2, 2, 1) = (2, 3, 1). \end{aligned}$$

Thus, $\mathbf{F}_1 \oplus \mathbf{F}_1 \subseteq \text{top}(\mathbf{P}_4)$, which is a contradiction. Suppose that $(\mathcal{C}_\Lambda, \mathbf{m})$ is the quiver



Then $\dim \mathbf{P}_5 = s_1^- s_2^- s_3^- s_4^- s_5^- (0, 1, 0) = s_1^- s_2^- s_3^- s_4^- (0, 1, 2) = s_1^- s_2^- s_3^- (1, 1, 2) = s_1^- s_2^- (1, 2, 2) = s_1^- (1, 2, 4) = (1, 2, 4)$. Therefore $\mathbf{F}_2 \oplus \mathbf{F}_2 \subseteq \text{top}(\mathbf{P}_5)$, which is a contradiction. If $(\mathcal{C}_\Lambda, \mathbf{m})$ is the quiver



then $\dim \mathbf{P}_7 = s_1^- s_2^- s_3^- s_4^- s_5^- s_6^- s_7^- (0, 1, 0) = s_1^- s_2^- s_3^- s_4^- s_5^- s_6^- (1, 1, 0) = s_1^- s_2^- s_3^- s_4^- s_5^- (1, 1, 2) = s_1^- s_2^- s_3^- s_4^- (1, 4, 2) = s_1^- s_2^- s_3^- (3, 4, 2) = s_1^- s_2^- (3, 4, 2) = s_1^- (3, 5, 2) = (2, 5, 2)$. Thus by Proposition 2.2, $\mathbf{F}_3 \oplus \mathbf{F}_3 \subseteq \text{top}(\mathbf{P}_7)$, which is a contradiction. Therefore Λ is not a right Köthe ring. ■

As an immediate consequence of Lemma 5.5, we have the following corollary.

Corollary 5.6 *If Λ is a basic hereditary artinian ring of the Coxeter type \mathbb{H}_4 , then Λ is not a right Köthe ring.*


The pair (Γ, \mathbf{m}) is called a (general) Coxeter valued quiver if $\Gamma = (\Gamma_0, \Gamma_1)$ is a finite quiver and $\mathbf{m}: \Gamma_1 \rightarrow \mathbb{N} \cup \{\infty\}$ is a function such that $\mathbf{m}(\alpha) \geq 3$, for any arrow

$\alpha \in \Gamma_1$. Notice that each Coxeter valued quiver is uniquely determined by the underlying Coxeter valued graph and some selection of its orientation.


Let Λ be a basic hereditary ring and let $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i,j \in I}$ be the species of Λ . Note that the species \mathcal{M}_Λ does not necessarily have the property “ ${}_iM_j \neq 0$ implies that ${}_jM_i = 0$ ”. We say that the bimodule ${}_iM_j$ belongs to a fixed connected component (Γ, \mathbf{m}) of the Coxeter valued quiver of Λ , provided we have $i, j \in \Gamma_0$. Recall that a ring Λ is called *indecomposable* if Λ is not a direct product of two non-zero rings. Now, we are ready to give a characterization of basic hereditary right Köthe rings in terms of their Coxeter valued quivers.

Theorem 5.7 *Let Λ be a basic hereditary ring. Then Λ is right Köthe if and only if Λ is an artinian ring such that the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is a finite disjoint union of the following Coxeter valued quivers:*

- (i) \mathbb{A}_n with any orientation;
- (ii) \mathbb{B}_n with the orientation




- (iii) \mathbb{B}_n with the orientation




with $1 \leq t \leq n - 1$;

- (iv) \mathbb{D}_n with the following conditions:




- (a) $|(n - 2)^+| \leq 2$;
- (b) for each $i \leq n - 3$, there exists at most one arrow with the source i ;

- (v) \mathbb{E}_6 with the following conditions:



- (a) $1 \leq |3^+| \leq 2, |4^+| \leq 1$ and $|2^+| \leq 1$;
- (b) For each $y \in 3^-$, there exists at least one arrow with the target y ;

- (vi) \mathbb{E}_7 with the orientation



(vii) \mathbb{G}_2 with the orientation



(viii) $\mathbb{I}_2(p)$ with the orientation



with $p = 5$ or $7 \leq p < \infty$;

where additionally in the cases (ii), (iii), (vii), and (viii) the dimension sequences of the unique nontrivial bimodules ${}_iM_j$ belong to these components have very restrictive shapes given respectively as follows:

- (ii) $d_4({}_{n-1}M_n) = (2, 1, 2, 1)$;
- (iii) $d_4({}_nM_{n-1}) = (2, 1, 2, 1)$;
- (vii) $d_6({}_1M_2) = (4, 1, 2, 2, 2, 1)$;
- (viii) $d_p({}_1M_2) = (p - 2, 1, 2, \dots, 2, 1)$.

Proof Since Λ is a basic hereditary ring, it has a ring product decomposition $\Lambda = \Lambda_1 \oplus \dots \oplus \Lambda_s$, where each Λ_i is an indecomposable basic hereditary ring. Then the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is a disjoint union of the Coxeter valued quivers $(\mathcal{C}_{\Lambda_i}, \mathbf{m})$ of Λ_i .

(\Rightarrow). Assume that Λ is a right Köthe ring. Then Λ_i is an indecomposable basic hereditary right Köthe ring, for each $1 \leq i \leq s$. Thus, without loss of generality, we can assume that Λ is an indecomposable basic hereditary right Köthe ring. Then by Proposition 4.1, Λ is a representation-finite ring and every finite-dimensional indecomposable representation of \mathcal{M}_Λ is multiplicity-free top. Thus by Proposition 3.3 and Lemmas 4.7, 5.4, 5.5, and Corollary 5.6, the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_\Lambda, \mathbf{m})$ of Λ is one of the Coxeter diagrams $\mathbb{A}_n, \mathbb{B}_n, \mathbb{D}_n, \mathbb{E}_6, \mathbb{E}_7, \mathbb{G}_2$ and $\mathbb{I}_2(p)$ presented in Table B. Therefore by Propositions 4.2, 4.3, 4.5, 4.6, 5.3, and Theorem 5.1, the proof complete.

(\Leftarrow). By Propositions 4.2, 4.3, 4.5, 4.6, 5.3, and Theorem 5.1, for each $1 \leq i \leq s$, every right Λ_i -module is a direct sum of cyclic modules. Therefore, Λ is a right Köthe ring. ■

6 A Characterization of Right Köthe Rings with Radical Square Zero

Let U and T be two rings, let M be a T - U -bimodule and let $R = \begin{pmatrix} T & M \\ 0 & U \end{pmatrix}$. Let \mathcal{D}_R be the category whose objects are triples (X, Y, f) , where X is a right T -module, Y is a right U -module and $f \in \text{Hom}_U(X \otimes_T M, Y)$. If $\alpha \in \text{Hom}_{\mathcal{D}_R}((X, Y, f), (X', Y', f'))$, then $\alpha = (\alpha_1, \alpha_2)$, where $\alpha_1 \in \text{Hom}_T(X, X')$ and $\alpha_2 \in \text{Hom}_U(Y, Y')$ such that $\alpha_2 f = f'(\alpha_1 \otimes id_M)$. The functor $G: \mathcal{D}_R \rightarrow \text{Mod-}R$ is defined in [18] (see also [19]) as follows. Let (X, Y, f) be an object in the category \mathcal{D}_R . For $(x, y) \in X \oplus Y$ and $\begin{pmatrix} t & m \\ 0 & u \end{pmatrix} \in R$, define

$$(x, y) \begin{pmatrix} t & m \\ 0 & u \end{pmatrix} = (xt, f(x \otimes m) + yu).$$

It is easy to see that $X \oplus Y$ is a right R -module. We define $G((A, B, f))$ to be $A \oplus B$. Let $\alpha = (\alpha_1, \alpha_2) \in \text{Hom}_{\mathcal{D}_R}((X, Y, f), (X', Y', f'))$. We set $G(\alpha) = \alpha_1 \oplus \alpha_2$. The reader can easily verify that $G(\alpha)$ is an R -homomorphism. It is well known that the functor G is an equivalence. From now on, we will identify the categories \mathcal{D}_R and $\text{Mod-}R$.

Let Λ be a basic artinian ring with radical square zero and $R = \begin{pmatrix} \Lambda/J & J \\ 0 & \Lambda/J \end{pmatrix}$. Let \mathcal{A} denote the full subcategory of $\text{Mod-}R$ whose objects are (X, Y, f) , where X and Y are two right Λ/J -modules and $f \in \text{Hom}_{\Lambda/J}(X \otimes_{\Lambda/J} J, Y)$ is an epimorphism. Then we have the natural functor $H: \text{Mod-}\Lambda \rightarrow \mathcal{A}$ which is defined in [18] (see also [19]) as follows. Let M be a right Λ -module. Then $H(M) = (M/MJ, MJ, f_M)$, where $f_M: M/MJ \otimes_{\Lambda/J} J \rightarrow MJ$ is induced from the multiplication map $M \otimes_{\Lambda} J \rightarrow MJ$. It is well known that the functor H is full and dense and $M \in \text{Mod-}\Lambda$ is indecomposable if and only if $H(M)$ in \mathcal{A} is indecomposable.

Proposition 6.1 *Let Λ be a basic artinian ring with radical square zero. Then Λ is a right Köthe ring if and only if $R = \begin{pmatrix} \Lambda/J & J \\ 0 & \Lambda/J \end{pmatrix}$ is a right Köthe ring.*


Proof (\Rightarrow). Assume that Λ is a right Köthe ring. Since Λ is a basic artinian ring, by [21, Proposition 1.8], R is a basic artinian ring. Let (A, B, f) be a finitely generated indecomposable right R -module. If $(A, B, f) \notin \mathcal{A}$, then f is not an epimorphism. Since B is semisimple, $B = B' \oplus \text{Im } f$ for some Λ/J -submodule B' of B . Thus $(A, B, f) \cong (A, \text{Im } f, f) \oplus (0, B', f)$. Since (A, B, f) is indecomposable, $(A, B, f) \cong (0, B', f)$ and hence B' is a simple right Λ -module. Therefore by [23, Corollary 2.2], (A, B, f) is multiplicity-free top. Now assume that $(A, B, f) \in \mathcal{A}$. Then there exists an indecomposable right Λ -module M such that $H(M) \cong (A, B, f)$. It follows that $(A, B, f) \cong (M/MJ, MJ, f_M)$, where $f_M: M/MJ \otimes_{\Lambda/J} J \rightarrow MJ$ is induced from the multiplication map $M \otimes_{\Lambda} J \rightarrow MJ$. Since Λ is right Köthe, by Proposition 4.1, $M/MJ = S_1 \oplus \dots \oplus S_t$, where $t \in \mathbb{N}$, each S_i is a simple right Λ -module and $S_i \not\cong S_j$ for each $i \neq j$. Therefore by [23, Corollary 2.2], $\text{top}((A, B, f)) = (S_1, 0, 0) \oplus \dots \oplus (S_t, 0, 0)$, where for $i \neq j$, $(S_i, 0, 0) \not\cong (S_j, 0, 0)$. This proves that every finitely generated indecomposable right R -module is multiplicity-free top. Therefore by Proposition 4.1, R is a right Köthe ring.

(\Leftarrow). Let N be a finitely generated indecomposable right Λ -module. Then by [21, Exercise 1C], $H(N) = (N/NJ, NJ, f_N)$ is a finitely generated indecomposable right R -module. Since by [23, Corollary 2.2], $\text{top}(H(N)) = (S_1, 0, 0) \oplus \dots \oplus (S_t, 0, 0)$, where $t \in \mathbb{N}$ and $N/NJ = S_1 \oplus \dots \oplus S_t$, by Proposition 4.1, for each $i \neq j$, $S_i \not\cong S_j$. Therefore, Λ is a right Köthe ring. ■

Let Λ be a basic artinian ring with radical square zero and let $\mathcal{M}_{\Lambda} = (F_i, {}_iM_j)_{i,j \in I}$ be the species of Λ . Note that the species \mathcal{M}_{Λ} does not necessarily have the property “ ${}_iM_j \neq 0$ implies that ${}_jM_i = 0$ ”. Let $(\Gamma_{\Lambda}, \mathbf{d})$ be the valued quiver of Λ . We recall from [15] that a *separated quiver* of Λ is the valued graph $(\Gamma_{\Lambda}^s, \mathbf{d}^s)$ with the vertex set $\{(i, l) : i \in I, l = 0, 1\}$ and the arrows

$$\begin{matrix} & (d_{ij}, d_{ji}) & \\ & \circ \xrightarrow{\quad} \circ & \\ (i, 0) & & (j, 1) \end{matrix}$$

precisely when ${}_iM_j \neq 0$. If $d_{ij} = d_{ji} = 1$, we write simply




It is easy to see that the separated quiver of Λ coincides with the valued quiver of R (see [15]). Moreover, it is well known that $R = \begin{pmatrix} \Lambda/J & J \\ 0 & \Lambda/J \end{pmatrix}$ is a basic hereditary artinian ring (see [15] and also [21]).

We conclude this section with the following result, which is a characterization of basic right Köthe rings with radical square zero in terms of their separated quivers.


Theorem 6.2 *Let Λ be a basic ring with radical square zero and let $\mathcal{M}_\Lambda = (F_i, {}_iM_j)_{i,j \in I}$ be the species of Λ (note that \mathcal{M}_Λ does not necessarily have the property “ ${}_iM_j \neq 0$ implies that ${}_jM_i = 0$ ”). Then the following statements are equivalent:*

- (i) Λ is a right Köthe ring;
- (ii) $R = \begin{pmatrix} \Lambda/J & J \\ 0 & \Lambda/J \end{pmatrix}$ is an artinian ring such that the Coxeter valued quiver $(\mathcal{C}_R, \mathbf{m})$ of R is a finite disjoint union of the Coxeter valued quivers presented in Theorem 5.7;
- (iii) Λ is a representation-finite ring and the separated quiver $(\Gamma_\Lambda^s, \mathbf{d})$ of Λ is a finite disjoint union of the following valued Dynkin quivers:
 - (a) \mathbb{A}_n with any orientation;
 - (b) \mathbb{B}_n with the orientation




where there exist precisely 4 pairwise non-isomorphic finitely generated indecomposable right $\begin{pmatrix} F_{n-1} & n-1M_n \\ 0 & F_n \end{pmatrix}$ -modules;

- (c) \mathbb{C}_n with the orientation




where $1 \leq i \leq n-1$ and there exist precisely 4 pairwise non-isomorphic finitely generated indecomposable right $\begin{pmatrix} F_n & nM_{n-1} \\ 0 & F_{n-1} \end{pmatrix}$ -modules;

- (d) \mathbb{D}_n with the following conditions:




- (1) $|(n-2)^+| \leq 2$;
- (2) For each $i \leq n-3$, there exists at most one arrow with the source i ;

- (e) \mathbb{E}_6 with the following conditions:



- (1) $1 \leq |3^+| \leq 2, |4^+| \leq 1$ and $|2^+| \leq 1$;
- (2) For each $y \in 3^-$, there exists at least one arrow with the target y ;

(f) \mathbb{E}_7 with the orientation



(g) the valued quiver



where there exist precisely 6 pairwise non-isomorphic finitely generated indecomposable right $\begin{pmatrix} F_1 & {}_1M_2 \\ 0 & F_2 \end{pmatrix}$ -modules;

(h) the valued quiver



where $p = 5$ or $7 \leq p < \infty$ and there exist precisely p pairwise non-isomorphic finitely generated indecomposable right $\begin{pmatrix} F_1 & {}_1M_2 \\ 0 & F_2 \end{pmatrix}$ -modules.

Proof

(i) \iff (ii) follows from Theorems 5.7 and 6.1.

(ii) \implies (iii) follows from Proposition 3.3 and [39, Lemma 3.1].

(iii) \implies (i) Since Λ is a basic artinian ring with radical square zero, R is a basic hereditary artinian ring. Thus, $R = R_1 \oplus \dots \oplus R_m$, where each R_i is an indecomposable basic hereditary artinian ring. Since R is a right Köthe ring if and only if each R_i is a right Köthe ring and the valued quiver of R is a disjoint union of the valued quivers of R_i , without loss of generality, we can assume that R is an indecomposable basic hereditary artinian ring. Let the separated quiver $(\Gamma_\Lambda^s, \mathbf{d})$ of Λ be one of the valued Dynkin quivers $\mathbb{A}_n, \mathbb{B}_n, \mathbb{C}_n, \mathbb{D}_n, \mathbb{E}_6$, and \mathbb{E}_7 presented in (iii). Since R is a representation-finite ring, by Proposition 3.3, the underlying Coxeter valued graph of the Coxeter valued quiver $(\mathcal{C}_R, \mathbf{m})$ of R is one of the Coxeter diagrams $\mathbb{A}_n, \mathbb{B}_n, \mathbb{D}_n, \mathbb{E}_6$, and \mathbb{E}_7 presented in Table B. Hence, by assumption and Theorem 5.7, R is a right Köthe ring. Now assume that the separated quiver $(\Gamma_\Lambda^s, \mathbf{d})$ of Λ is the quiver



and there exist precisely 6 pairwise non-isomorphic finitely generated indecomposable right $\begin{pmatrix} F_1 & {}_1M_2 \\ 0 & F_2 \end{pmatrix}$ -modules. Then by [39, Corollary 3.5], $d_6({}_1M_2)$ is a dimension sequence and hence by [13, Proposition 2'], $d_6({}_1M_2) = (4, 1, 2, 2, 2, 1)$. Thus by Theorem 5.7, R is a right Köthe ring. If the separated quiver $(\Gamma_\Lambda^s, \mathbf{d})$ of Λ is the quiver presented in (h), by the similar argument we can see that R is a right Köthe ring. Therefore by Proposition 6.1, Λ is a right Köthe ring. ■

Acknowledgments The authors would like to thank Daniel Simson for his comments on the earlier version of this paper. Also, we are indebted to the anonymous

reviewers' insightful comments, suggestions and feedback on this paper and their time and effort spent on it.

References

- [1] F. W. Anderson and K. R. Fuller, *Rings and categories of modules*, Second ed., Graduate Texts in Mathematics, 13, Springer-Verlag, New York, 1992. <https://doi.org/10.1007/978-1-4612-4418-9>
- [2] I. Assem, D. Simson, and A. Skowronski, *Elements of the representation theory of associative algebras*. Vol. 1, Techniques of representation theory, London Mathematical Society Student Texts, 65, Cambridge University Press, Cambridge, 2006. <https://doi.org/10.1017/CBO9780511614309>
- [3] M. Auslander, *Large modules over Artin algebras*. In: *Algebra, topology and category theory*. Academic Press, New York, 1976, pp. 1–17.
- [4] M. Auslander, M. I. Platzeck, and I. Reiten, *Coxeter functors without diagrams*. *Trans. Amer. Math. Soc.* 250(1979), 1–46. <https://doi.org/10.2307/1998978>
- [5] M. Auslander, I. Reiten, and S. O. Smalø, *Representation theory of Artin algebras*. Cambridge Studies in Advanced Mathematics, 36, Cambridge University Press, Cambridge, 1995.
- [6] M. Behboodi, A. Ghorbani, A. Morazadeh-Dehkordi, and S. H. Shojaei, *On left Köthe rings and a generalization of a Köthe–Cohen–Kaplansky theorem*. *Proc. Amer. Math. Soc.* 142(2014), no. 8, 2625–2631. <https://doi.org/10.1090/S0002-9939-2014-11158-0>
- [7] I. N. Bernstein, I. M. Gelfand, and V. A. Ponomarev, *Coxeter functors, and Gabriel's theorem*. (Russian) *Uspehi Mat. Nauk* 28(1973), no. 2(170), 19–33.
- [8] N. Bourbaki, *Lie groups and Lie algebras*. Chapters 4–6. Translated from the 1968 French original by Andrew Pressley. *Elements of Mathematics* (Berlin), Springer-Verlag, Berlin, 2002. <https://doi.org/10.1007/978-3-540-89394-3>
- [9] S. U. Chase, *Direct products of modules*. *Trans. Amer. Math. Soc.* 97(1960), 457–473. <https://doi.org/10.2307/1993382>
- [10] I. S. Cohen and I. Kaplansky, *Rings for which every module is a direct sum of cyclic modules*. *Math. Z.* 54(1951), 97–101. <https://doi.org/10.1007/BF01179851>
- [11] V. Dlab and C. M. Ringel, *Indecomposable representations of graphs and algebras*. *Mem. Amer. Math. Soc.* 6(1976), no. 173. <https://doi.org/10.1090/memo/0173>
- [12] V. Dlab and C. M. Ringel, *On algebras of finite representation type*. *J. Algebra* 33(1975), 306–394. [https://doi.org/10.1016/0021-8693\(75\)90125-8](https://doi.org/10.1016/0021-8693(75)90125-8)
- [13] P. Dowbor, C. M. Ringel, and D. Simson, *Hereditary Artinian rings of finite representation type*. In: *Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979)*. Lecture Notes in Math., 832, Springer, Berlin, 1980, pp. 232–241.
- [14] P. Dowbor and D. Simson, *A characterization of hereditary rings of finite representation type*. *Bull. Amer. Math. Soc. (N.S.)* 2(1980), no. 2, 300–302. <https://doi.org/10.1090/S0273-0979-1980-14741-2>
- [15] P. Dowbor and D. Simson, *Quasi-Artin species and rings of finite representation type*. *J. Algebra* 63(1980), no. 2, 435–443. [https://doi.org/10.1016/0021-8693\(80\)90082-4](https://doi.org/10.1016/0021-8693(80)90082-4)
- [16] J. A. Drozd, *The structure of hereditary rings*. (Russian) *Mat. Sb. (N.S.)* 113(155)(1980), no. 1(9), 161–172, 176.
- [17] Z. Fazelpour and A. Nasr-Isfahani, *Connection between representation-finite and Köthe rings*. *J. Algebra* 514(2018), 25–39. <https://doi.org/10.1016/j.jalgebra.2018.08.006>
- [18] R. M. Fossum, P. A. Griffith, and I. Reiten, *Trivial extensions of Abelian categories. Homological algebra of trivial extensions of abelian categories with applications to ring theory*. Lecture Notes in Mathematics, 456, Springer-Verlag, Berlin–New York, 1975.
- [19] P. Gabriel, *Indecomposable representation. II*. *Symposia Mathematica*, Vol. XI (Convegno di Algebra Commutativa, INDAM, Rome, 1971), Academic Press, London, 1973, pp. 81–104.
- [20] P. Gabriel, *Unzerlegbare Darstellungen I*. *Manuscripta Math.* 6(1972), 71–103; correction, *ibid.* 6(1972), 309. <https://doi.org/10.1007/BF01298413>
- [21] K. R. Goodearl and R. B. Warfield Jr., *An introduction to noncommutative Noetherian rings*, Second ed., London Mathematical Society Student Texts, 61, Cambridge University Press, Cambridge, 2004. <https://doi.org/10.1017/CBO9780511841699>
- [22] P. Griffith, *On the decomposition of modules and generalized left uniserial rings*. *Math. Ann.* 184(1969/1970), 300–308. <https://doi.org/10.1007/BF01350858>
- [23] A. Haghany and K. Varadarajan, *Study of modules over formal triangular matrix rings*. *J. Pure Appl. Algebra* 147(2000), no. 1, 41–58. [https://doi.org/10.1016/S0022-4049\(98\)00129-7](https://doi.org/10.1016/S0022-4049(98)00129-7)

- [24] J. E. Humphreys, *Reflection groups and Coxeter groups*. Cambridge studies in Advanced Mathematics, 29, Cambridge University Press, Cambridge, 1990.
<https://doi.org/10.1017/CBO9780511623646>
- [25] Y. Kawada, *On Köthe's problem concerning algebras for which every indecomposable module is cyclic I*. Sci. Rep. Tokyo Kyoiku Daigaku Sect. A 7(1962), 154–230.
- [26] Y. Kawada, *On Köthe's problem concerning algebras for which every indecomposable module is cyclic II*. Sci. Rep. Tokyo Kyoiku Daigaku Sect. A 8(1963), 1–62.
- [27] Y. Kawada, *On Köthe's problem concerning algebras for which every indecomposable module is cyclic III*. Sci. Rep. Tokyo Kyoiku Daigaku Sect. A 8(1965), 165–250.
- [28] G. Köthe, *Verallgemeinerte Abelsche Gruppen mit hyperkomplexem Operatorenring*. (German) Math. Z. 39(1935), no. 1, 31–44. <https://doi.org/10.1007/BF01201343>
- [29] T. Nakayama, *Note on uni-serial and generalized uni-serial rings*. Proc. Imp. Acad. Tokyo 16(1940), 285–289.
- [30] H. Prüfer, *Untersuchungen über die Zerlegbarkeit der Abzählbaren primären abelschen Gruppen*. (German) Math. Z. 17(1923), no. 1, 35–61. <https://doi.org/10.1007/BF01504333>
- [31] C. M. Ringel, *Kawada's theorem*. In: *Abelian group theory (Oberwolfach, 1981)*. Lecture Notes in Math., 874, Springer, Berlin–New York, 1981, pp. 431–447.
- [32] C. M. Ringel, *Representation theory of Dynkin quivers, Three contributions*. Front. Math. China II(2016), no. 4, 765–814. <https://doi.org/10.1007/s11464-016-0548-5>
- [33] C. M. Ringel, *Representations of K -species and bimodules*. J. Algebra 41(1976), no. 2, 269–302.
[https://doi.org/10.1016/0021-8693\(76\)90184-8](https://doi.org/10.1016/0021-8693(76)90184-8)
- [34] A. H. Schofield, *Hereditary artinian rings of finite representation type and extensions of simple artinian rings*. Math. Proc. Cambridge Philos. Soc. 102(1987), no. 3, 411–420.
<https://doi.org/10.1017/S0305004100067463>
- [35] A. H. Schofield, *Representations of rings over skew fields*. London Mathematical Society Lecture Note Series, 92, Cambridge University Press, Cambridge, 1985.
<https://doi.org/10.1017/CBO9780511661914>
- [36] D. Simson, *A class of potential counterexamples to the pure semisimplicity conjecture*. In: *Advances in algebra and model theory (Essen, 1994; Dresden, 1995)*. Algebra Logic Appl., 9, Gordon and Breach, Amsterdam, 1997, pp. 345–373.
- [37] D. Simson, *An Artin problem for division ring extensions and the pure semisimplicity conjecture II*. J. Algebra 227(2000), no. 2, 670–705. <https://doi.org/10.1006/jabr.1999.8245>
- [38] D. Simson, *Categories of representations of species*. J. Pure Appl. Algebra 14(1979), no. 1, 101–114.
[https://doi.org/10.1016/0022-4049\(79\)90015-X](https://doi.org/10.1016/0022-4049(79)90015-X)
- [39] D. Simson, *On right pure semisimple hereditary rings and an Artin problem*. J. Pure Appl. Algebra 104(1995), no. 3, 313–332. [https://doi.org/10.1016/0022-4049\(94\)00068-X](https://doi.org/10.1016/0022-4049(94)00068-X)
- [40] D. Simson, *Partial Coxeter functors and right pure semisimple hereditary rings*. J. Algebra 71(1981), no. 1, 195–218. [https://doi.org/10.1016/0021-8693\(81\)90115-0](https://doi.org/10.1016/0021-8693(81)90115-0)
- [41] D. Simson, *Pure semisimple categories and rings of finite representation type*. J. Algebra 48(1977), no. 2, 290–296. [https://doi.org/10.1016/0021-8693\(77\)90307-6](https://doi.org/10.1016/0021-8693(77)90307-6)
- [42] D. Simson, *Pure semisimple categories and rings of finite representation type, Corrigendum*. J. Algebra 67(1980), no. 1, 254–256. [https://doi.org/10.1016/0021-8693\(80\)90320-8](https://doi.org/10.1016/0021-8693(80)90320-8)
- [43] R. Wisbauer, *Foundations of module and ring theory. A handbook for study and research*. Revised and translated from the 1988 German edition, Algebra, Logic and Applications, 3, Gordon and Breach Science Publishers, Philadelphia, PA, 1991.

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran, Iran

e-mail: z.fazelpour@ipm.ir

Department of Pure Mathematics, Faculty of Mathematics and Statistics, University of Isfahan, P.O. Box: 81746-73441, Isfahan, Iran

and

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746, Tehran, Iran

e-mail: nasr_a@sci.ui.ac.ir nasr@ipm.ir