

## Research Article

**Cite this article:** Mendes BP, da Silva Jr. IA, DaMata JP, Castro-Gomes T, Vieira LQ, Ribeiro-Dias F, Horta MF (2019). Metacyclogenesis of *Leishmania (Viannia) guyanensis*: a comprehensive study of the main transformation features in axenic culture and purification of metacyclic promastigotes by negative selection with *Bauhinia purpurea* lectin. *Parasitology* **146**, 716–727. <https://doi.org/10.1017/S0031182018002111>

Received: 29 May 2018

Revised: 9 October 2018

Accepted: 18 November 2018

First published online: 27 December 2018

**Key words:**

*Bauhinia purpurea*; *Leishmania guyanensis*; metacyclogenesis

**Author for correspondence:**

M. Fátima Horta, E-mail: [phorta@icb.ufmg.br](mailto:phorta@icb.ufmg.br)

# Metacyclogenesis of *Leishmania (Viannia) guyanensis*: a comprehensive study of the main transformation features in axenic culture and purification of metacyclic promastigotes by negative selection with *Bauhinia purpurea* lectin

Bárbara Pinheiro Mendes<sup>1</sup>, Ildefonso Alves da Silva Jr.<sup>2</sup>, Jarina Pena DaMata<sup>1</sup>, Thiago Castro-Gomes<sup>3</sup>, Leda Quercia Vieira<sup>1</sup>, Fátima Ribeiro-Dias<sup>2</sup> and Maria Fátima Horta<sup>1</sup>

<sup>1</sup>Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, 31.270-901, Belo Horizonte, MG, Brazil; <sup>2</sup>Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás, 74605-050, Goiânia, GO, Brazil and <sup>3</sup>Departamento de Parasitologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, 31.270-901, Belo Horizonte, MG, Brazil

**Abstract**

*Leishmania (Viannia) guyanensis* is one species that causes cutaneous leishmaniasis in the New World. The incidence of infections with this parasite is probably underestimated and few studies exist on this species, despite its epidemiological importance. In particular, there are no studies concerning *L. guyanensis* metacyclogenesis and no technique for obtaining metacyclic promastigotes for this species is presently available. Here, we have studied *L. guyanensis* metacyclogenesis in axenic culture, describing the main changes that occur during this process, namely, in morphology and size, sensitivity to complement-mediated lysis, surface carbohydrates and infectivity to macrophages. We have shown that metacyclogenesis in *L. guyanensis* promastigotes is basically complete on the 4<sup>th</sup> day of culture, as determined by decreased body size, increased flagellum length, resistance to complement-mediated lysis and infectivity. We have also found that only a fraction of the parasites is agglutinated by *Bauhinia purpurea* lectin. The non-agglutinated parasites, which also peaked on the 4<sup>th</sup> day of culture, had all morphological traits typical of the metacyclic stage. This is the first report describing metacyclogenesis in *L. guyanensis* axenic promastigotes and a simple and efficient method for the purification of metacyclic forms. Furthermore, a model of human macrophage infection with *L. guyanensis* was established.

**Introduction**

Protozoan of the genus *Leishmania* are parasites whose developmental stages alternate between female sand flies and mammalian hosts. Promastigotes, which are flagellated forms, are found in the midgut of the sand flies, while amastigotes, lacking the external flagella, live inside mammalian macrophages causing a spectrum of disease manifestations known as leishmaniasis. Infection of mammalian hosts starts with the inoculation of promastigotes by sand flies during bloodmeal. Promastigotes development in sand flies is associated with an increase of virulence, a process known as metacyclogenesis (Sacks and Perkins, 1985). A proportion of parasites undergo a sequential development transition from uninfected (procyclic) to infective (metacyclic) promastigotes after a period of replication, which terminates approximately upon the complete digestion of the bloodmeal (Sacks and Perkins, 1984). The metacyclogenesis process can also occur in *in vitro* axenic cultures, where parasites from the log-phase of growth resemble procyclic promastigotes and, entering into stationary phase, a fraction of them differentiates into promastigotes with properties reminiscent of authentic sand fly metacyclic promastigotes (Giannini, 1974; Sacks and Perkins, 1984; Sacks and Perkins, 1985; Da Silva and Sacks, 1987; Howard *et al.*, 1987; Sacks *et al.*, 1995; Lira *et al.*, 1998). *In vitro* metacyclogenesis occurs in different species of the genus *Leishmania*, as observed by several groups (Sacks and Da Silva, 1987; Louassini *et al.*, 1998; Zakai *et al.*, 1998; Pinto-da-Silva *et al.*, 2002; Almeida *et al.*, 2003; Gamboa *et al.*, 2008).

The infective metacyclic stage of all *Leishmania* species is morphologically distinct from the procyclic non-infective promastigotes, with a shorter, narrower body and a longer flagellum, which usually reach more than twice the body size in length (Sacks and Perkins, 1984; Zakai *et al.*, 1998). Metacyclogenesis, both in the sand fly gut and in *in vitro* axenic cultures, is also associated with changes in gene expression and structural modifications, particularly in the composition of the parasite surface glycocalyx, which is composed of a variety of

glycosylphosphatidylinositol-anchored molecules. These molecules include the abundant glycolipid lipophosphoglycan (LPG) and the protease gp63 (Sacks and Da Silva, 1987; McConville *et al.*, 1992; Turco and Descoteaux, 1992), known to play important roles in infectivity (Sacks *et al.*, 1990; Brodin *et al.*, 1992; Marín *et al.*, 2000; Ramos *et al.*, 2004, 2011). Some molecules, such as the META proteins (Uliana *et al.*, 1999; Ramos *et al.*, 2011; Santos *et al.*, 2011) and the MAT-1 transcript (Brodin *et al.*, 1992; Saraiva *et al.*, 1995; Marín *et al.*, 2000) are expressed only in metacyclic forms and are considered virulence factors. Differential regulation of genes also results in expression of proteins known as HASP and SHERP, essential for differentiation of *Leishmania major* in its sand fly vector (Sádlová *et al.*, 2000), only in infective stage parasites (McKean *et al.*, 2001).

The mechanism accounting for the differentiation into metacyclic forms is poorly understood. It is known that, *in vitro*, acidification of culture media and the scarcity of nutrients, typical of the stationary phase of growth, seem to be the main stimuli that determine the differentiation of promastigotes from procyclic to metacyclic forms (Bates, 2008). Moreover, Cunningham *et al.* (2001) showed that pteridin metabolism is relevant to metacyclogenesis in *L. major*. Likewise, Serafim *et al.* (2012) demonstrated that this process may be controlled by purines, particularly adenosine.

*Leishmania* promastigotes inoculated in mammals by the sand fly must survive microbicidal defence mechanisms to which they are exposed within vertebrate hosts. These defences include the lytic effects of alternative complement pathway activation (Jokiranta *et al.*, 1995) as well as the microbicidal mechanisms of the phagocytic cells (Camus *et al.*, 1995; Bogdan *et al.*, 1996). In fact, metacyclic promastigotes are more resistant to lysis by fresh normal serum than procyclic forms of several species of *Leishmania*, including *L. donovani*, *L. panamensis*, *L. major*, *L. mexicana*, *L. amazonensis* and *L. braziliensis* (Franke *et al.*, 1985; Puentes *et al.*, 1988; Louassini *et al.*, 1998). Resistance to complement-mediated lysis was related to the elongation of LPG chain in metacyclic promastigotes surface, which seems to protect the parasite from lysis acting like a physical barrier that block membrane attack complex insertion into parasite plasma membrane (Franke *et al.*, 1985; Puentes *et al.*, 1988). The increase in the surface metalloprotease gp63 expression in metacyclic forms also seems to play a role in the resistance to complement-mediated lysis (Brittingham *et al.*, 1995) due to the proteolytical inactivation of C3b bound to parasite surface (Mosser and Brittingham, 1997).

As described for *L. major* and *L. donovani* (Da Silva and Sacks, 1987; McConville *et al.*, 1992; Sacks *et al.*, 1995), developmental changes in LPG structure result in downregulation of lectin-binding sites. In both species, metacyclic promastigotes can be purified by negative selection, due to loss of the ability of being agglutinated by peanut lectin and concanavalin A. It was also shown that *L. tropica* (Lira *et al.*, 1998) and *L. amazonensis* (Courret *et al.*, 1999) metacyclic promastigotes can be purified from axenic cultures using antilipophosphoglycan monoclonal antibodies. *Leishmania braziliensis* metacyclic promastigotes also can be purified by negative selection using *Bauhinia purpurea* lectin (BPL), based on the changes promastigotes undergo on outer membrane carbohydrates during metacyclogenesis (Pinto-da-Silva *et al.*, 2002).

Although *L. guyanensis* is the causative agent of cutaneous leishmaniasis in the New World and one of the most important species in Brazil and several other countries in South America, there are no studies concerning *L. guyanensis* metacyclogenesis. Moreover, as a consequence, no technique for obtaining metacyclic promastigotes for the study of infections by this species is presently available.

The present work aimed to study the main features of metacyclogenesis transformation of *L. guyanensis* in axenic culture. Our results describe a series of characteristics, such as changes in morphology and size, sensitivity to complement-mediated lysis, surface carbohydrates and infectivity, associated with promastigotes metacyclogenesis transformation. Importantly, we were able to purify *L. guyanensis* metacyclic promastigotes using the methodology described by Pinto-da-Silva *et al.* (2002) for *L. braziliensis* using BPL, showing that *L. guyanensis* promastigotes not agglutinated by BPL exhibit the same morphology, resistance to complement and infectivity *in vitro* of metacyclic promastigotes.

## Material and methods

### Parasites and growth assessment

*Leishmania* (*Viannia*) *guyanensis* [M4147 strain (MHOM/BR/75/M4147) and M1176 strain (MHOM/BR/1975/M1176)] promastigotes were cultured in Schneider's insect medium (Sigma Chemical Co., St Louis, MD, USA) supplemented with 10% heat-inactivated fetal calf serum (Cripion, Andradina, SP, Brazil), and 50 mg L<sup>-1</sup> gentamicin, pH 7.2, at 25 °C. Parasites used herein are well-characterized strains of *L. guyanensis* kindly provided by Dr Maria Norma Melo (Departamento de Parasitologia, Universidade Federal de Minas Gerais), formerly typed by their isoenzymes profile differences (Lainson *et al.* (1982) and by antibodies (Jennings *et al.*, 2014) and later by ITS1 RFLP, using Hae III, according to Schönian *et al.* (2003). Parasite cultures were initiated from frozen stocks at 1 × 10<sup>6</sup> parasites mL<sup>-1</sup>. Promastigotes were harvested mainly in the 2<sup>nd</sup>, 4<sup>th</sup>, 7<sup>th</sup> and 10<sup>th</sup> days of culture throughout this work, maintained in the above conditions for a maximum of ten passages to prevent loss of metacyclic characteristics, such as virulence. Growth assessments were carried out by daily haemocytometer counting of promastigotes from an initial load of 1 × 10<sup>6</sup> parasites mL<sup>-1</sup>. M4147 strain of *L. guyanensis* was used throughout this work, except for a couple of experiments performed with the M1176 strain.

### Morphological analysis

Parasites were fixed in glass slides after spinning of 100 µL of the culture at 3500×g for 30 s in a cytospin centrifuge (Fanem, São Paulo, Brazil). Parasites were then stained with Instant Prov Kit (Laborclin, Pinhais, PR, Brazil). Morphological evaluation of metacyclogenesis was performed under light microscopy (1000×) by digital photography. The images were analysed and measurements of promastigote cell body size and flagellum lengths were made using Image J software (National Institute of Health Bethesda, MD, USA).

### Agglutination with BPL

Negative selection of *L. guyanensis* promastigotes was performed, using BPL (Vector Laboratories, Burlingame, CA, USA), as previously described for *L. braziliensis* (Pinto-da-Silva *et al.*, 2002). Briefly, non-agglutinated promastigotes from different phases of culture were washed in Hank's balanced salt solution (HBSS), pH 7.4, supplemented with 50 mg L<sup>-1</sup> gentamicin and quantified in a haemocytometer. Promastigotes (10<sup>8</sup> parasites mL<sup>-1</sup>) were washed and incubated with 50 µg mL<sup>-1</sup> of BPL at room temperature. After 30 min, the suspension was centrifuged at 40×g for 5 min to sediment agglutinated forms. Non-agglutinated promastigotes, obtained in the supernatant, were washed in HBSS twice at 3000×g for 15 min. The percentage of metacyclic forms was

determined in relation to total parasites and used in the experiments.

### Complement-mediated lysis assay

Total or BPL non-agglutinated promastigotes ( $5 \times 10^5$  parasites) from different days of culture (2<sup>nd</sup>, 4<sup>th</sup>, 7<sup>th</sup> or 10<sup>th</sup> day) were incubated with 10% of rabbit serum (Low-tox-M Rabbit Complement – Cerdalane Laboratories, Burlington, Canada) in 200  $\mu$ l of HBSS, pH 7.4, at 34 °C. After 1 h, the tubes were placed on ice and the number of surviving parasites determined by light microscopy using a haemocytometer, as described (Howard *et al.*, 1987; Lincoln *et al.*, 2004; da Silva *et al.*, 2015). Live promastigotes (intact, motile and translucent) are readily distinguished from dead parasites (with loss of morphological integrity, non-motile and opaque or granular) ('ghost cells'). Results were expressed as the percentage of parasite survival compared with controls that were not exposed to serum.

### Infection of macrophages

Peripheral blood monocytes were obtained by venipuncture of healthy donors ( $n = 12$ ) from Blood Bank of Instituto Goiano de Hematologia e Oncologia (Goiânia, Goiás, Brazil) in EDTA-vacuum tubes (Greiner bio-one, Vacuette, Americana, SP, Brazil). Blood sample was layered on Ficoll gradient (GE Healthcare Bio-Sciences AB, Uppsala, Uppsala län, Sweden) and centrifuged at 1000 $\times$ g, at room temperature for 20 min. Mononuclear cells ( $5 \times 10^5$  cells) were sedimented in 24 wells plate onto round coverslips in 500  $\mu$ L RPMI 1640 medium supplemented with 2% of heat-inactivated human serum, 11 mM sodium bicarbonate, 2 mM L-glutamine, 100 U mL<sup>-1</sup> penicillin and 100  $\mu$ g mL<sup>-1</sup> streptomycin (all reagents from Sigma). Mononuclear cells were then derived to macrophages by 7 days in culture at 37 °C and 5% CO<sub>2</sub>, changing the culture medium completely each 48 h. Macrophages were then incubated with promastigotes (1:1 parasite:macrophage ratio) for 4 h, washed three times with warmed RPMI medium to remove extracellular parasites, and incubated until 24 or 48 h at 34 °C and 5% CO<sub>2</sub>. Cells were then fixed and stained with Instant Prov kit (Newprov) and analysed under a light microscope (1000 $\times$ ). Number of infected and uninfected cells and number of parasites per cell were determined in at least 300 macrophages per coverslip. Infection index represents the percentage of infected cells  $\times$  mean number of parasites per infected cell.

### Flow cytometry analysis

Parasites ( $1 \times 10^6$  mL<sup>-1</sup>) were analysed for light scatter by flow cytometry (FACScan, Becton Dickinson) in order to evaluate the size of parasites as described by Saraiva *et al.* (2005). Size of parasites was analysed after plotting FSC vs SSC using the software FlowJo (TreeStar).

### MTT assay for analysis of parasites viability

The viability of promastigotes during *in vitro* culture was accessed by the MTT colorimetric assay based on the ability of viable parasites to reduce the tetrazolium salt to an insoluble formazan product, as described by Dutta *et al.* (2005). Briefly,  $1 \times 10^5$  parasites were incubated with 2 mg mL<sup>-1</sup> of MTT in phenol red-free RPMI 1640 medium for 4 h in the dark at 34 °C. The microplate were centrifuged 3000 $\times$ g for 5 min, the supernatant was removed, the pellet was dissolved with DMSO and absorbance was measured as 492 nm in a spectrophotometer (Molecular Devices, Sunnyvale, CA, USA).

## Results

### Analysis of metacyclogenesis of *L. guyanensis* promastigotes in axenic culture

#### Growth

Initially, promastigote growth was verified by counting the parasites daily under light microscope. Parasites proliferate exponentially until the 3<sup>rd</sup> day in culture, characterizing the logarithmic (log) phase, and continue to grow until the 4<sup>th</sup> or 5<sup>th</sup> day after which proliferation stops, characterizing the stationary phase (Fig. 1). During *in vitro* promastigote growth, parasites are around 70–80% viable up to the 5<sup>th</sup> day of culture, but viability decreases from the 6<sup>th</sup> day on (Fig. 1). The following analyses on the main changes that occur during promastigote metacyclogenesis were performed in parasites harvested on 2<sup>nd</sup> (log phase), 4<sup>th</sup> (early stationary phase), 7<sup>th</sup> and, occasionally, on 10<sup>th</sup> (late stationary phase) days, indicated by the arrows (Fig. 1).


#### Size

(1) *Flow cytometry*: Parasites were analysed for light scatter by flow cytometry to determine changes in size during *in vitro* growth. We analysed forward scatter (FSC), which reflects the size and side scatter (SSC), which reflects the granularity of parasites. Figure 2A–C shows a sample analysis of 2<sup>nd</sup>, 4<sup>th</sup> and 7<sup>th</sup> day of culture. Analysis were performed inside the gate depicted in the figure, which excluded a defined population of very small size (which is frequent on the 2<sup>nd</sup> day) and intense granularity (which appears on the 7<sup>th</sup> day), that we assume to be dead parasites, due to the results on viability (Fig. 1). We can observe that the population of promastigotes gradually decreases in size during *in vitro* growth. The histogram of frequency of cells of different sizes shows that the majority of parasites on the 2<sup>nd</sup> day are larger than those on the 4<sup>th</sup> or 7<sup>th</sup> day (Fig. 2D). By comparison with other descriptions, metacyclic promastigotes should be the low FSC and SSC parasites. (2) *Morphology and morphometry*: We also evaluated the morphological alterations by visualization in light microscopy (Fig. 2L–N) and by measuring (Fig. 3) stained promastigotes during *in vitro* growth to identify metacyclic forms in cultures, according to their body and flagellum sizes. Total promastigotes from the 2<sup>nd</sup> day of culture were mostly slender and fusiform with the flagellum size equivalent to the body length (Fig. 2L). On the 4<sup>th</sup> day, parasites start losing their fusiform shape, becoming smaller (Fig. 2M) and rounded, in a mix population. On the 7<sup>th</sup> day, they completely assume a round/balloon shape (Fig. 2N). Parasite measures (Fig. 3) show a progressive decrease in the cellular body size (length and width) and an increase in the flagellum length during time, much like in other species of *Leishmania*, mainly from 2<sup>nd</sup> to 4<sup>th</sup> day, indicating that metacyclogenesis is occurring in axenic *L. guyanensis* culture. These results corroborate what we observed in flow cytometry. It is interesting to notice, looking at the median, that cellular body width decrease is more gradual than cellular body length decrease or flagellum length increase, which seems to change more quickly.

#### Resistance to complement-mediated lysis

In other species of *Leishmania*, transformation of procyclic into metacyclic form in promastigotes is accompanied by an increase in the capacity of parasites to resist to lysis mediated by the complement system. We performed this assay utilizing fresh rabbit serum as source of complement proteins to evaluate the degree of resistance of axenic promastigotes to complement-mediated lysis *in vitro*. We verified that parasites from the stationary phase (4<sup>th</sup> and 7<sup>th</sup> day of culture) are more resistant to lysis than those from log phase (2<sup>nd</sup> day of culture) (Fig. 4, clear boxes). Parasite resistance reaches up to 90% at the 10<sup>th</sup> day





**Fig. 1.** Growth curve and viability of *Leishmania guyanensis* promastigotes in axenic culture. Parasites ( $1 \times 10^6 \text{ mL}^{-1}$  initial load) were cultured in Schneider's insect medium and counted daily in a haemocytometer. Arrows mark the days when parasites were harvested for analysis, corresponding log, early stationary and late stationary phases. Viability was determined by the MTT assay, as described in 'Material and methods' section. Each point represents the mean  $\pm$  s.e. of three independent experiments carried out in duplicates.

(not shown) as compared with around 20% at the 2<sup>nd</sup> day. Total parasites from the 2<sup>nd</sup> day incubated with heat-inactivated rabbit serum have an average survival rate of 97% (not shown). These results corroborate the occurrence of metacyclogenesis in axenic culture of *L. guyanensis*.

#### Infectivity

Since metacyclic forms are more infective to macrophages, promastigotes were evaluated for their infectivity to human macrophages. Figure 5 shows that, at 24 h, total promastigotes from log phase (2<sup>nd</sup> day) infect around 25% of macrophages (A) which have less than two parasites average per cell (B). This rate of infection does not vary in 48 h. On the other hand, parasites from 4<sup>th</sup> day infect around 65% of macrophages (A), each macrophage containing more than three parasites average (B) at 24 h, and somewhat decreased at 48 h to 55% with less than three parasites average. When the population of promastigotes from the 7<sup>th</sup> day of culture is used, the percentage of infected macrophages did not vary much (55%) at 24 h but the number of parasites inside cells dropped to more than two parasites average (B). At 48 h the percentage of infected macrophages decreased to 40% and the number of parasites per cell to less than two, average. The infection index (C) reflects more accurately the infectivity of these populations of promastigotes, showing that promastigotes from the 4<sup>th</sup> and 7<sup>th</sup> day of culture are far more infective than promastigotes from the 2<sup>nd</sup> day of culture, although the population of promastigotes from the 7<sup>th</sup> day of culture is less infective than the population from the 4<sup>th</sup> day.

#### Agglutination mediated by BPL

In an attempt to purify the metacyclic promastigotes from total culture, parasites were incubated with BPL, which is known to selectively agglutinate procyclic forms of *L. braziliensis* (Pinto-da-Silva *et al.*, 2002). It had already been shown that *L. guyanensis* binds BPL (Gazola *et al.*, 2001), but there are no studies about selective agglutination of promastigotes during the different phases of growth. In Fig. 6, we show that incubation of promastigotes from the different days of axenic culture with BPL led to selective agglutination of parasites (Fig. 6A). In this experiment, we included parasites from the 3<sup>rd</sup> day to verify

whether the changes in the ability to bind BPL occurred from 2<sup>nd</sup> to 3<sup>rd</sup> or from 3<sup>rd</sup> to 4<sup>th</sup> day and from the 10<sup>th</sup> day to verify whether metacyclics persisted at a late phase growth. We observed that this procedure leaves only 25–30% of non-agglutinated parasites on the 2<sup>nd</sup> and 3<sup>rd</sup> day, while 60–80% of parasites did not agglutinate on the 4<sup>th</sup> and 7<sup>th</sup> days, reaching almost 100% on the 10<sup>th</sup> day, showing a remarkable increase of forms that do not bind BPL from the log phase of culture. Purified parasites keep their viability similarly to that of parasites from 4-day cultures, but are somewhat less viable than the parasites from 7<sup>th</sup> or 10<sup>th</sup> day cultures (not shown). These data confirm that forms that bind BPL, which must be the procyclic forms, predominate at log phase of growth of parasites in axenic culture, while those that do not bind BPL, which must be the metacyclic forms, predominate at the beginning of stationary phase in the culture up to the 3<sup>rd</sup> day. Agglutination by BPL is disrupted by lactose, a BPL-binding inhibitor, confirming the specificity of the binding (not shown). BPL differential promastigote agglutination is not a feature unique for the M4147 strain of *L. guyanensis* used throughout this work, since promastigotes from the M1176 strain are also differentially agglutinated by BPL, following the same proportions of agglutination, as evaluated on days 2, 4 and 7 of *in vitro* growth (Fig. 6B).


#### Analysis of *L. guyanensis* promastigotes negatively selected with BPL

##### Size as evaluated by flow cytometry

Negatively selected parasites from 2<sup>nd</sup>, 4<sup>th</sup> and 7<sup>th</sup> day cultures were analysed by flow cytometry (Fig. 2E–H). We observed that, on the 2<sup>nd</sup> day, the population that does not bind BPL (Fig. 2E) displays much smaller size when compared with populations of total culture (Fig. 2A). It is possible that either they are metacyclics from the previous culture or parasites that precociously became metacyclics. When we look at the population of negatively selected promastigotes from the 4<sup>th</sup> or 7<sup>th</sup> day (Fig. 2F and G), there is little or no difference in size, respectively, when compared with total culture (Fig. 2B and C), showing that these are stages in which most or all promastigotes had already become metacyclic in total culture. The differences in the appearance of the BPL-non-agglutinated and BPL-agglutinated parasites are striking when analysed by light microscopy. Non-agglutinated (BPL–) promastigotes (Fig. 2O) are round with long flagella, whereas agglutinated (BPL+) parasites (Fig. 2P) are much thinner and fusiform. We can see that on the 2<sup>nd</sup> day, BPL-negative population is still somewhat heterogeneous, but most parasites are smaller when compared with the whole population. BPL-negative promastigotes from days 4 and 7 are practically all small and homogenous in size. Figure 2I–J shows a comparison between the frequency of sizes in total culture and negatively selected culture. These data are compatible with those found by morphometry, which show a decrease of the cellular size, concomitant with the appearance of metacyclic forms in culture. BPL-negative population of M1176 strain also shows the same size and appearance as M4147 strain, used throughout this work, as accessed by flow cytometry and microscopy, respectively (not shown), also showing that the selected parasites have the typical features of metacyclic promastigotes.

##### Resistance to complement-mediated lysis

BPL-negative promastigote forms were also evaluated as to their resistance to complement-mediated lysis. Our results show that BPL-negative parasites (Fig. 4, shaded boxes) are indeed enriched in forms more resistant to complement-mediated lysis. Even with parasites from the 2<sup>nd</sup> day, we already may observe 70–80% of survival of purified parasites in comparison to around 20% in



**Fig. 2.** Analysis of *Leishmania guyanensis* promastigote sizes by morphometry. Parasites were fixed in glass slides and stained. Morphological evaluation of metacyclogenesis was performed under light microscopy (1000 $\times$ ) by digital photography. Measurements of promastigote cell body lengths (A), body widths (B) and flagellum lengths (C) were made using Image J software. Values represent the median, superior and inferior quartiles, and maximal and minimal sizes of four independent experiments in which the size of 60 parasites/experiment was determined (\* $P < 0.0001$  compared with 2<sup>nd</sup> day; # $P < 0.0001$  compared with 4<sup>th</sup> day; ° $P < 0.0001$  compared with each other).




Fig. 2. Continued.

the total culture, reaching more than 80% in parasites from the 4<sup>th</sup> or 7<sup>th</sup> day. These data show that we were able to enrich in complement-resistant promastigotes since the beginning of the stationary phase, corroborating that the parasites not agglutinated by BPL are the metacyclic forms.

#### Infectivity

The results of this experiment, carried out with parasites on the 4<sup>th</sup> day of culture, also corroborate that agglutination with BPL causes an enrichment of metacyclic forms at the culture, since BPL-negative parasites proved to be more infective than the parasites of total culture (Fig. 5D-F). Purified metacyclic forms were also shown to be more resistant to elimination by macrophages, since the rate of infection and parasites per cell are maintained from 24 to 48 h, as opposed to parasites of total culture at 4<sup>th</sup> day (Fig. 5A-C, repeated in Fig. 5D-E for better comparison) or 7<sup>th</sup> day (Fig. 5A-C).


#### Discussion

In a recent study, Pires *et al.* (2015), reporting the characterization of an isolate of *L. guyanensis* from a patient in Goiás, Brazil, emphasized the crucial importance of correct identification and characterization of species of the subgenus *Leishmania Viannia*,

suggesting that the presence of *L. guyanensis* may be underestimated in endemic regions. Despite the epidemiologic importance of *L. guyanensis*, in PubMed web site (<https://www.ncbi.nlm.nih.gov/pubmed>), as accessed 12 December 2018, of the 13 134 hits containing the keyword *Leishmania* in the title of the article, only 93 (0.71%) contains the words *Leishmania* and *guyanensis*, demonstrating the remarkable lack of data concerning this important species. No articles were found on *L. guyanensis* metacyclogenesis and no technique is described to isolate metacyclic promastigotes from axenic cultures, an important step to study infection *in vitro* and *in vivo* in animal models. Here, we have thoroughly studied the metacyclogenesis process in *L. guyanensis* in axenic cultures, and tested and validated a method currently used to purify *L. braziliensis* metacyclic promastigote (Pinto-da-Silva *et al.*, 2002).


Like other *Leishmania* species (da Silva and Sacks, 1987), *L. guyanensis* grows *in vitro*, presenting a standard growth curve with a log phase, which lasts up to the 4<sup>th</sup> day, from where a stationary phase starts (Fig. 1). Due to morphological and biochemical differences among the various developmental forms, it is possible to identify metacyclic and procyclic promastigotes in different species of *Leishmania spp* (Sacks and Da Silva, 1987; Louassini *et al.*, 1998; Zakai *et al.*, 1998; Pinto-da-Silva *et al.*, 2002; Almeida *et al.*, 2003; Gamboa *et al.*, 2008). Here, we found that, during axenic culture, *L. guyanensis* undergoes the





**Fig. 3.** Analysis of total culture or BPL-negative *Leishmania guyanensis* promastigote sizes by flow cytometry. Total (A–C) or BPL-negative (E–G) promastigotes from 2<sup>nd</sup>, 4<sup>th</sup> or 7<sup>th</sup> day of culture were analysed as to their size (FSC) and granulosity (SSC). The frequency of events inside defined gates are plotted together (D, H and I–K) for better comparison.

same types of changes that characterize metacyclogenesis. One is the morphological transformations regarding cell and flagellum sizes. These changes were gradual, with a decrease in cell body size and increased flagellum length (Figs 2 and 3). Promastigotes are considered to have a metacyclic morphology when presenting small body cell size and long flagellum (twice or more the body length) (Zakai *et al.*, 1998) and these forms were mainly found from the 4<sup>th</sup> day on. By flow cytometry, the changes in parasite size were readily and more accurately observable than by morphometry. Interestingly, however, the morphology of metacyclic *L. guyanensis* promastigotes are rather different from other species, although all have small bodies, *L.*




**Fig. 4.** Complement-mediated lysis of total or BPL-negative *Leishmania guyanensis* promastigotes. Total (T) or BPL-negative (BPL–) promastigotes from 2<sup>nd</sup>, 4<sup>th</sup> or 7<sup>th</sup> day of culture were incubated with 10% fresh or heat-inactivated rabbit serum for 1 h at 37 °C, and parasite viability was evaluated. Parasites incubated in culture medium only were used as controls. Values represent the median, superior and inferior quartiles, and maximal and minimal percentages of resistant parasites of three independent experiments (\* $P < 0.05$ ; # $P < 0.05$  compared with total 2<sup>nd</sup> day; \*\* $P < 0.005$ ).

*guyanensis* metacyclics have a balloon-shaped body (Fig. 2), whereas metacyclics from other species exhibit a slender appearance (Saraiva *et al.*, 2005). *Leishmania braziliensis*, a closely related species, have been described as having either slender (Pinto-da-Silva *et al.*, 2002) or round/oval shapes (Almeida *et al.*, 1993). Our results demonstrated that, like in other species, the size of the parasites decreases over time in axenic culture, indicating that metacyclogenesis is taking place.


Just as in the digestive tract of the vector, *Leishmania* promastigotes in axenic culture, transform from a non-infective to a highly infective stage, a central feature of metacyclogenesis. During this process, parasites become pre-adapted to survive within vertebrates, as they need to escape host cells microbicidal mechanisms. Complement-mediated lysis is one such mechanism. Metacyclic promastigotes, abundant during stationary phase of growth, have been proven to be much more resistant to complement-mediated lysis than procyclic forms mostly present during log phase (Sacks, 1989). Corroborating these findings, we demonstrated that stationary phase *L. guyanensis* promastigotes are far more resistant to complement-mediated lysis than log phase parasites (Fig. 4), confirming our previous findings that the frequency of *L. guyanensis* metacyclic promastigotes is much higher in stationary phase (Sousa-Franco *et al.*, 2006). It is noteworthy that, as from the 4<sup>th</sup> day (early stationary phase), the frequency of small parasites and parasites resistant to complement-mediated lysis is comparable to the frequency on the 7<sup>th</sup> or 10<sup>th</sup> day (late stationary phase), indicating that *L. guyanensis* metacyclogenesis is mostly completed in the early stationary phase, also confirmed by infectivity and BPL binding, as discussed later.

Infectivity of the parasite *in vitro* or *in vivo* is also a manner of evaluating metacyclogenesis, since metacyclic promastigote forms are more infective to their host cells than procyclic forms (Louassini *et al.*, 1998; Pinto-da-Silva *et al.*, 2002; Yao *et al.*, 2008; Serafim *et al.*, 2012; Da Silva *et al.*, 2015). Our experiments demonstrated that *L. guyanensis* early (4<sup>th</sup> day) or late (7<sup>th</sup> day) stationary phase promastigotes are more infective to human macrophages *in vitro* than those in the log phase (Fig. 5), as observed both from the number of infected macrophages and the number of parasites inside macrophages, reflected in the great difference between the infection index on days 2 and 4. Parasites from the early stationary phase (4<sup>th</sup> day) are more



**Fig. 5.** Infectivity of total and BPL-negative *Leishmania guyanensis* promastigotes. Total (T) or BPL-negative (BPL-) parasites from 2<sup>nd</sup>, 4<sup>th</sup> or 7<sup>th</sup> day were incubated with human macrophages. Infectivity was evaluated after 24 (triangles) and 48 (squares) h after infection. Values represent the mean of infectivity of promastigotes to 12 or six different donors in five independent experiments (\**P* < 0.05; \*\**P* < 0.005; \*\*\**P* < 0.0005). Infectivity of total promastigotes on the 4<sup>th</sup> day (A–C) is repeated on the second column (D–E) for easier comparison.





**Fig. 6.** *Leishmania guyanensis* promastigotes profile of agglutination by *Bauhinia purpurea* lectin (BPL). Parasite cultures of M4147 (A) or M1176 (B) strain were washed and incubated with  $50 \mu\text{g mL}^{-1}$  BPL. Values of experiment in (A) represent the median, superior and inferior quartiles, and maximal and minimal percentages of non-agglutinated BPL-negative parasites of three independent experiments ( $*P < 0.005$ ). Experiment in (B) is a unique experiment, performed merely to validate the method for other strain. ND, not determined.

successful in macrophage infection than parasites from the late stationary phase (7<sup>th</sup> day) (Fig. 5), possibly due to the higher number of viable parasites in the early stationary phase (Fig. 1). One interesting fact is that, at 48 h, the infection index is lower than at 24 h, suggesting that a number of invading parasites are being eliminated by the macrophages. These parasites may correspond to procyclic forms still in culture (observe that this does not occur with the purified metacyclic forms).

Another expressive shift that occurs during differentiation into metacyclic form in several species of *Leishmania* is the modification in LPG molecules. Changes in the composition of LPG carbohydrates associated with metacyclogenesis have been demonstrated in experiments where metacyclic forms lose their ability to bind to insect gut (Pinto-da Silva *et al.*, 2002; Serafim *et al.*, 2012). In *L. major*, LPG almost doubles the mean number of repeating units per molecule and arabinose (Ara) covers the galactose (Gal) side chains (Sacks *et al.*, 1990). Metacyclic forms of *L. braziliensis* also undergo modifications in the LPG molecule by adding glucose residues and increasing the galactose-mannose (Gal-Man) repeating units (Soares *et al.*, 2005). These changes may be identified by differential binding of parasites to lectins (Sacks and Da Silva, 1987; Sacks *et al.*, 1995) or anti-LPG antibodies (Lira *et al.*, 1998; Mahoney *et al.*, 1999; Chaves *et al.*,

2003; Pinto-da-Silva *et al.*, 2005), which allows parasites to be purified or enriched by negative selection.

Here, we defined a method for purification of *L. guyanensis* metacyclic forms, employing the method used with *L. braziliensis*, first shown by Pinto-da-Silva *et al.* (2002). Gazola *et al.* (2001), in a study with several plant lectins, had already shown that BPL, which attaches to galactosyl ( $\beta$ -1, 3) N-acetylgalactosamine [ $\beta$ -gal (1-3)-GalNAc] residues, binds to *L. guyanensis* promastigotes, without, however, analysing whether this feature is shared by all promastigote forms during axenic growth of promastigotes. Our results corroborate these data and here, we further followed this feature during metacyclogenesis. At any point of the growth curve, we could see that BPL selectively agglutinated *L. guyanensis* promastigotes allowing the separation of two populations. The fact that, in the log phase of growth, more parasites are agglutinated and the opposite occurred from the early stationary phase (Fig. 6) indicated that BPL selectively binds to procyclic promastigotes. Indeed, analysis of the non-agglutinated forms, to compare their features with those of metacyclic forms, showed that these forms shared all the characteristics of metacyclic promastigotes studied here, namely morphology, resistance to complement-mediated lysis and *in vitro* macrophage infectivity (Figs 2-5).

In *Leishmania*, LPG is an interspecies polymorphic polysaccharide with variations in carbohydrates branching from the conserved column of Gal ( $\beta$ 1, 4) Man ( $\alpha$ 1) -PO<sub>4</sub> repeat units and cap oligosaccharides. In *L. braziliensis*, metacyclic forms also undergo modifications in the LPG molecule with the addition of glucose residues and increasing in mannose-galactose repeating units (Soares *et al.*, 2005). In *L. major*, during metacyclogenesis, there is also an increase in the number of repeating units per molecule and a decrease of repetitive units with  $\beta$  Gal or Gal  $\beta$  1-3 Gal  $\beta$  1-side chains and an increase of both repeating units without side chains or side chains containing Ara  $\alpha$  1-2 Gal  $\beta$  1-, and other structural changes (McConville *et al.*, 1992). In *L. guyanensis*, there are no reports of which are the changes, but the differential agglutination of promastigotes by BPL shows that there must also be such modifications. Since BPL binds  $\beta$ -gal (1-3)-GalNAc residues, our results show that, like in other species, these modifications are most likely the decrease in mannose-galactose residues. This corroborates the results obtained by Muskus *et al.* (1997), which showed that lectins and antibodies specific for the disaccharide units of *L. donovani* are able to distinguish *L. guyanensis*, *L. panamensis* and *L. braziliensis* promastigotes from the log and the stationary phases.

The results discussed so far show evidence that, from early stationary phase (4<sup>th</sup> day), the vast majority of promastigotes already have metacyclic characteristics, such as small size, resistance to complement-mediated lysis, loss of ability to be agglutinated by BPL and infectivity. Here, we could enrich the population of metacyclic promastigotes of *L. guyanensis* in culture, by negative selection of promastigotes by agglutination by BPL, using both log phase promastigotes (2<sup>nd</sup> day) and stationary phase parasites from early (4<sup>th</sup> day) or late phase (7<sup>th</sup> day). The efficacy of this method was verified by the same parameters described here. Purified parasites stay viable during the purification process, following the same pattern of the total culture (not shown). BPL (-) promastigotes viability is also patent given their strong ability to infect macrophages (Fig. 5).

Regarding the size, the frequency of non-agglutinated parasites obtained from log phase (2<sup>nd</sup> day) differs considerably from the total parasites in culture. On the other hand, in cultures from early stationary phase (4<sup>th</sup> day), we can see that almost all parasites have the same size as observed in total culture and 100% of the late stationary phase parasites (7<sup>th</sup> day) already have the same size as parasites from total culture (Fig. 2).


Non-agglutinated parasites, from both log phase and stationary phases, were more resistant to complement-mediated lysis than the total parasite culture (Fig. 4). Likewise, non-agglutinated parasites from initial stationary phase proved to be more infective to macrophages than those from total culture (Fig. 5). Furthermore, macrophages were not able to eliminate intracellular amastigotes derived from infection with parasites not agglutinated by BPL, contrary to what happens with the total population of promastigotes in culture whose infection rate is lower at 48 h (Fig. 5). It is likely that the increased resistance to complement-mediated lysis of promastigotes not agglutinated by BPL is due to changes of LPG, as it has been shown for *L. major* (Späth *et al.*, 2003; Gaur *et al.*, 2009), *L. mexicana* and *L. donovani* (Gaur *et al.*, 2009). For *L. major* and *L. donovani* (Gaur *et al.*, 2009), it has been shown that LPG in metacyclic forms is an important structure for the macrophage–parasite binding/infection of these cells. Here, we found a correlation between infectivity and agglutination by BPL, and it is also possible that the change in *L. guyanensis* LPG contributes for its infectivity.

Evaluation of parasites not agglutinated by BPL by methods well established for metacyclogenesis analysis of various species of *Leishmania* shows that, also in *L. guyanensis*, they display characteristics defined as metacyclogenesis markers, such as reduced cell body and flagellum longer than cell body, increased resistance to complement-mediated lysis and increased infectivity for macrophages, in agreement with the various works cited here. Therefore, we can conclude from the present data that parasites that are not agglutinated by BPL are the metacyclic forms of *L. guyanensis*.

It is expected that, even in cultures at stationary phase, heterogeneous populations of promastigote can be found in axenic cultures of *L. major* (Sacks, 1989) and in sandflies infected with *L. braziliensis* and *L. amazonensis* (Nieves and Pimenta, 2000). The presence of metacyclic forms in all stages of culture (log and early or late stationary) and all times evaluated were noted in our experiments. This indicates that metacyclogenesis can occur in an unsynchronized way, considering the entire population of parasites in culture. However, we did find an enrichment of metacyclic forms in the stationary phase, as expected. The results found in experiments using flow cytometry and morphology showed a more gradual appearance pattern of morphological changes, even in stationary phase (Figs 2 and 3). Conversely, data obtained from complement-mediated lysis assay and by agglutination with BPL show that in stationary phase the population of metacyclic forms is more homogeneous as far as the changes in surface molecules are concerned (Figs 4 and 6). We can see that the size of total parasites and BPL-enriched parasites is almost the same on day 4 and the same on day 7. Despite the similar size, around 30% of promastigotes are still agglutinated by BPL (Fig. 6) or sensitive to complement-mediated death (Fig. 4). This indicates that two different populations may still co-exist at the stationary phase. It is possible however that the minor population of the 4<sup>th</sup> day that still did not reach the final smallest size of BPL-enriched parasites could be the population that also binds BPL (Fig. 6). Metacyclogenesis is actually not a single step for all features and metacyclic promastigotes are characterized by all of them together. In this case, it looks like that the size is one of the first features to change. Thus, the size of the parasites, at least in this species, may not be a good metacyclogenesis marker. That is why we believe BPL enrichment is important, since BPL-enriched parasites are also complement-resistance-enriched parasites, more infective-enriched parasites and also have the smallest size.

Approaches for metacyclic purification have been described, exploring these morphological features. Späth and Beverley

(2001) purified *L. major* metacyclics by density gradient centrifugation, obtaining cells with typical metacyclic features at the 8–10% Ficoll interface. These distinct morphologies also allowed a description of a method to analyse metacyclic infective forms of *Leishmania* spp within the whole stationary population by non-fluorimetric parameters of flow cytometry (Saraiva *et al.*, 2005), which was used here. In conclusion, this is the first report describing metacyclogenesis in *L. guyanensis* axenic promastigotes, based on morphology and size, sensitivity to complement-mediated lysis, surface carbohydrates and infectivity. We have also described a simple and efficient method for the purification of metacyclic forms of *L. guyanensis* by negative selection using BPL. The method was proved suitable for two strains of *L. guyanensis*, suggesting that it is applicable for all parasites from this species. Furthermore, we have established a model of human macrophage infection with *L. guyanensis*, important for the study of other infection features.

**Author ORCIDs.**  Ildelfonso Alves da Silva Jr. 0000-0001-8184-9863; Thiago Castro-Gomes 0000-0003-1564-4645; Leda Quêrcia Vieira 0000-0001-5481-9618; Fátima Ribeiro-Dias 0000-0002-7950-3414; Maria Fátima Horta 0000-0002-9009-2716.

**Acknowledgements.** We thank Elimar Faria for the valuable technical assistance.

**Financial support.** This work was funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

**Conflict of interest.** None.

**Ethical standards.** All procedures for macrophage derivation were approved by local Ethical Committee (Hospital das Clínicas/UFG, prot. 132/2012).

## References

- Almeida MC, Cuba CA, de Sá CM, Pharoah MM, Howard KM and Miles MA (1993) Metacyclogenesis of *Leishmania (viannia) braziliensis in vitro*: evidence that lentil lectin is a marker of complement resistance and enhanced infectivity. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **87**, 325–329.
- Almeida MC, Vilhena V, Barral A and Barral-Netto M (2003) Leishmanial infection: analysis of its first steps: a review. *Memórias do Instituto Oswaldo Cruz* **98**, 861–870.
- Bates PA (2008) *Leishmania* sand fly interaction: progress and challenges. *Current Opinion in Microbiology* **11**, 340–344.
- Bogdan C, Gessner A, Solbach W and Rölinghoff M (1996) Invasion, control and persistence of *Leishmania* parasites. *Current Opinion in Immunology* **8**, 517–525.
- Brittingham A, Morrison CJ, McMaster WR, McGwire BS, Chang KP and Mosser DM (1995) Role of the *Leishmania* surface protease gp63 in complement fixation, cell adhesion, and resistance to complement-mediated lysis. *Journal of Immunology* **155**, 3102–3111.
- Brodin TN, Heath S and Sacks DL (1992) Genes selectively expressed in the infectious (metacyclic) stage of *Leishmania major* promastigotes encode a potential basic-zipper structural motif. *Molecular and Biochemical Parasitology* **52**, 241–250.
- Camus D, Zalis MG, Vannier-Santos MA and Banic DM (1995) The art of parasite survival. *Brazilian Journal of Medical and Biological Research* **28**, 399–413.
- Chaves CS, Soares DC, Silva RP and Saraiva EM (2003) Characterization of the species- and stage-specificity of two monoclonal antibodies against *Leishmania amazonensis*. *Experimental Parasitology* **103**, 152–159.
- Courret N, Prina E, Mougneau E, Saraiva EM, Sacks DL, Glaichenhaus N and Antoine JC (1999) Presentation of the *Leishmania* antigen LACK by infected macrophages is dependent upon the virulence of the phagocytosed parasites. *European Journal of Immunology* **29**, 762–773.
- Cunningham ML, Titus RG, Turco SJ and Beverley SM (2001) Regulation of differentiation to the infective stage of the protozoan parasite *Leishmania major* by tetrahydrobiopterin. *Science* **292**, 285–287.

- Da Silva R and Sacks DL (1987) Metacyclogenesis is a major determinant of *Leishmania* promastigote virulence and attenuation. *Infection and Immunity* 55, 2802–2806.
- Da Silva Jr IA, Morato CI, Quixabeira VB, Pereira LI, Dorta ML, de Oliveira MA, Horta MF and Ribeiro-Dias F (2015) *In vitro* metacyclogenesis of *Leishmania (viannia) braziliensis* and *Leishmania (leishmania) amazonensis* clinical field isolates, as evaluated by morphology, complement-resistance, and infectivity to human macrophages. *Biomedical Research International* 2015, 393049.
- Dutta A, Bandyopadhyay S, Mandal C and Chatterjee M (2005) Development of a modified MTT assay for screening antimonial resistant field isolates of Indian visceral leishmaniasis. *Parasitology International* 54, 119–122.
- Franke ED, McGreevy PB, Katz SP and Sacks DL (1985) Growth cycle-dependent generation of complement-resistant *Leishmania* promastigotes. *Journal of Immunology* 134, 2713–2718.
- Gamboa D, Torres K, De Doncker S, Zimic M, Arevalo J and Dujardin JC (2008) Evaluation of an *in vitro* and *in vivo* model for experimental infection with *Leishmania (viannia) braziliensis* and *L. (V.) peruviana*. *Parasitology* 135, 319–326.
- Gazola KC, Ferreira AV, Anacleto C, Michalick MS, Andrade AF and Moreira ES (2001) Cell surface carbohydrates and *in vivo* infectivity of glucantime-sensitive and resistant *Leishmania (viannia) guyanensis* cell lines. *Parasitology Research* 87, 935–940.
- Gaur U, Showalter M, Hickerson S, Dalvi R, Turco SJ, Wilson ME and Beverley SM (2009) *Leishmania donovani* lacking the Golgi GDP-Man transporter LPG2 exhibit attenuated virulence in mammalian hosts. *Experimental Parasitology* 122, 182–191.
- Giannini MS (1974) Effects of promastigote growth phase, frequency of subculture, and host age on promastigote-initiated infections with *Leishmania donovani* in the golden hamster. *Journal of Protozoology* 21, 521–527.
- Howard MK, Sayers G and Miles MA (1987) *Leishmania donovani* metacyclic promastigotes: transformation *in vitro*, lectin agglutination, complement-resistance, and infectivity. *Experimental Parasitology* 64, 147–156.
- Jennings YL, de Souza AA, Ishikawa EA, Shaw JJ, Lainson R and Silveira F (2014) Phenotypic characterization of *Leishmania* spp. causing cutaneous leishmaniasis in the lower Amazon region, western Pará state, Brazil, reveals a putative hybrid parasite, *Leishmania (viannia) guyanensis* x *Leishmania (viannia) shawi shawi*. *Parasite* 21, 39.
- Jokiranta TS, Jokipii L and Meri S (1995) Complement-resistance of parasites. *Scandinavian Journal of Immunology* 42, 9–20.
- Lainson R, Shaw JJ, Miles MA and Póvoa M (1982) Leishmaniasis in Brazil: XVII enzymic characterization of a *Leishmania* from the armadillo, *Dasypus novemcinctus* (Edentata), from Pará State. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 76, 810–811.
- Lincoln LM, Ozaki M, Donelson JE and Beetham JK (2004) Genetic complementation of *Leishmania* deficient in PSA (GP46) restores their resistance to lysis by complement. *Molecular and Biochemical Parasitology* 137, 185–189.
- Lira R, Méndez S, Carrera L, Jaffe C, Neva F and Sacks DL (1998) *Leishmania tropica*: the identification and purification of metacyclic promastigotes and use in establishing mouse and hamster models of cutaneous and visceral disease. *Experimental Parasitology* 89, 331–342.
- Louassini M, Adroher FJ, Foulquié MR and Benítez R (1998) Investigations on the *in vitro* metacyclogenesis of a visceral and a cutaneous human strain of *Leishmania infantum*. *Acta Tropica* 70, 355–368.
- Mahoney AB, Sacks DL, Saraiva E, Modi G and Turco SJ (1999) Intraspecific and stage-specific polymorphisms in lipophosphoglycan structure control *Leishmania donovani*-sand fly interactions. *Biochemistry* 38, 9813–9823.
- Marín M, Muskus C, Ramírez JR, Arbelaez LF, Alzate JF and Berberich C (2000) The gene encoding the metacyclogenesis-associated transcript Mat-1 is conserved in the genus *Leishmania* and shows a tendency to form dimers upon protein expression. *Parasitology Research* 86, 431–435.
- McConville MJ, Turco SJ, Ferguson MA and Sacks DL (1992) Developmental modification of lipophosphoglycan during the differentiation of *Leishmania major* promastigotes to an infectious stage. *EMBO Journal* 11, 3593–3600.
- McKean PG, Denny PW, Knuepfer E, Keen JK and Smith DF (2001) Phenotypic changes associated with deletion and overexpression of a stage-regulated gene family in *Leishmania*. *Cellular Microbiology* 3, 511–523.
- Mosser DM and Brittingham A (1997) *Leishmania*, macrophages and complement: a tale of subversion and exploitation. *Parasitology* 115, 9–23.
- Muskus C, Segura I, Oddone R, Turco SJ, Leiby DA, Toro L, Robledos S and Saraiva NG (1997) Carbohydrate and LPG expression in *Leishmania viannia* subgenus. *Journal of Parasitology* 83, 671–678.
- Nieves E and Pimenta PFP (2000) Development of *Leishmania (viannia) braziliensis* and *Leishmania (leishmania) amazonensis* in the sand fly *Lutzomyia migonei* (Diptera: Psychodidae). *Journal of Medical Entomology* 37, 134–140.
- Pinto-da-Silva LH, Camurate M, Costa KA, Oliveira SM, Da Cunha-e-Silva NL and Saraiva EM (2002) *Leishmania (Viannia) braziliensis* metacyclic promastigotes purified using *Bauhinia purpurea* lectin are complement-resistant and highly infective form macrophages *in vitro* and hamsters *in vivo*. *International Journal for Parasitology* 32, 1371–1377.
- Pinto-da-Silva LH, Fampa P, Soares DC, Oliveira SM, Souto-Pradon T and Saraiva EM (2005) The 3A1-La monoclonal antibody reveals key features of *Leishmania (l) amazonensis* metacyclic promastigotes and inhibits procyclic attachment to the sand fly midgut. *International Journal for Parasitology* 35, 757–764.
- Pires AS, Borges AF, Cappellazzo-Coelho A, Dorta ML, Lino-Junior R de S, Pereira LI, Pinto SA, Pelli-de-Oliveira MA, de Matos GG, Abrahamssohn IA, Uliana SR, Lima GM and Ribeiro-Dias F (2015) Identification and biological characterization of *Leishmania (viannia) guyanensis* isolated from a patient with tegumentary leishmaniasis in Goiás, a no endemic area for this species in Brazil. *Biomedical Research International* 2015, 350764.
- PubMed. <https://www.ncbi.nlm.nih.gov/pubmed> (Accessed 8 May 2018).
- Puentes SM, Sacks DL, Silva RP and Joiner KA (1988) Complement binding by two developmental stages of *Leishmania major* promastigotes varying in expression of a surface lipophosphoglycan. *Journal of Experimental Medicine* 167, 887–902.
- Ramos CS, Franco FAL, Smith DF and Uliana SRB (2004) Characterization of a new *Leishmania* META gene and genomic analysis of the META cluster. *FEMS Microbiology Letters* 238, 213–219.
- Ramos CS, Yokoyama-Yasunaka JKU, Guerra-Giraldez C, Price HP, Mortara RA, Smith DF and Uliana SRB (2011) *Leishmania amazonensis* META2 protein confers protection against heat shock and oxidative stress. *Experimental Parasitology* 127, 228–237.
- Sacks DL (1989) Metacyclogenesis in *Leishmania* promastigotes. *Experimental Parasitology* 69, 100–103.
- Sacks DL and Da Silva RP (1987) The generation of infective stage *Leishmania major* promastigotes is associated with the cell surface expression and release of a developmentally regulated glycolipid. *Journal of Immunology* 139, 3099–3106.
- Sacks DL and Perkins PV (1984) Identification of an infective stage of *Leishmania* promastigotes. *Science* 223, 1417–1419.
- Sacks DL and Perkins PV (1985) Development of infective stage *Leishmania* promastigotes within phlebotomine sandflies. *American Journal of Tropical Medicine and Hygiene* 34, 456–459.
- Sacks DL, Brodin TN and Turco SJ (1990) Developmental modification of the lipophosphoglycan from *Leishmania major* promastigotes during metacyclogenesis. *Molecular and Biochemical Parasitology* 42, 225–233.
- Sacks DL, Pimenta PF, McConville MJ, Schneider P and Turco SJ (1995) Stage-specific binding of *Leishmania donovani* to the sand fly vector midgut is regulated by conformational changes in the abundant surface lipophosphoglycan. *Journal of Experimental Medicine* 181, 685–697.
- Sádlová J, Price HP, Smith BA, Votýpka J, Volf P and Smith DF (2000) The stage-regulated HASPB and SHERP proteins are essential for differentiation of the protozoan parasite *Leishmania major* in its sand fly vector, *Phlebotomus papatasi*. *Cellular Microbiology* 12, 1765–1779.
- Santos MG, Silva ME, Zampieri RA, Lafraia RM and Floeter-Winte LM (2011) Correlation of meta 1 expression with culture stage, cell morphology and infectivity in *Leishmania (leishmania) amazonensis*. *Memórias do Instituto Oswaldo Cruz* 106, 190–193.
- Saraiva EM, Pimenta PF, Brodin TN, Rowton E, Modi GB and Sacks DL (1995) Changes in lipophosphoglycan and gene expression associated with the development of *Leishmania major* in *Phlebotomus papatasi*. *Parasitology* 111, 275–287.
- Saraiva EM, Pinto-da-Silva LH, Wanderley JLM, Bonomo AC, Barcinski MA and Moreira MEC (2005) Flow cytometric assessment of *Leishmania* spp metacyclic differentiation: validation by morphological features and specific markers. *Experimental Parasitology* 110, 39–47.



- Schönian G, Nasereddin A, Dinse N, Schweynoch C, Schallig HD, Presber W and Jaffe CL (2003) PCR diagnosis and characterization of *Leishmania* in local and imported clinical samples. *Diagnostic Microbiology and Infectious Disease* **47**, 349–358.
- Serafim TD, Figueiredo AB, Costa PAC, Marques-da-Silva EA, Gonçalves R, de Moura SA, Gontijo NF, da Silva SM, Michakick MS, Meyer-Fernandes JR, de Carvalho RP, Uliana SR, Fietto JL and Afonso LC (2012) *Leishmania* metacyclogenesis is promoted in the absence of purines. *PLoS Neglected Tropical Diseases* **6**, e1833.
- Soares RP, Cardoso TL, Barron T, Araujo MS and Turco SJ (2005) *Leishmania braziliensis*: a novel mechanism in the lipophosphoglycan regulation during metacyclogenesis. *International Journal for Parasitology* **35**, 245–253.
- Sousa-Franco J, Araújo-Mendes E, Silva-Jardim I, Lima-Santos J, Faria DR, Dutra WO and Horta MF (2006) Infection-induced respiratory burst in BALB/c macrophages kills *Leishmania guyanensis* amastigotes through apoptosis: possible involvement in resistance to cutaneous leishmaniasis. *Microbes and Infection* **8**, 390–400.
- Späth GF and Beverley SM (2001) A lipophosphoglycan-independent method for isolation of infective *Leishmania* metacyclic promastigotes by density gradient centrifugation. *Experimental Parasitology* **99**, 97–103.
- Späth GF, Garraway LA, Turco SJ and Beverley SM (2003) The role(s) of lipophosphoglycan (LPG) in the establishment of *Leishmania major* infections in mammalian hosts. *Proceedings of the National Academy of Sciences USA* **100**, 9536–9541.
- Turco SJ and Descoteaux A (1992) The lipophosphoglycan of *Leishmania* parasites. *Annual Review of Microbiology* **46**, 65–94.
- Uliana SRB, Goyal N, Freymüller E and Smith DF (1999) *Leishmania*: over-expression and comparative structural analysis of the stage-regulated meta 1 gene. *Experimental Parasitology* **92**, 1983–1191.
- Yao C, Chen Y, Sudan B, Donelson JE and Wilson ME (2008) *Leishmania chagasi*: homogenous metacyclic promastigotes isolated by buoyant density are highly virulent in a mouse model. *Experimental Parasitology* **118**, 129–133.
- Zakai HA, Chance ML and Bates PA (1998) *In vitro* stimulation of metacyclogenesis in *Leishmania braziliensis*, *L. donovani*, *L. major* and *L. mexicana*. *Parasitology* **116**, 305–309.