

# Altered social and non-social decision-making in recreational and dependent cocaine users

L. M. Hulka<sup>1</sup>, C. Eisenegger<sup>2</sup>, K. H. Preller<sup>1</sup>, M. Vonmoos<sup>1</sup>, D. Jenni<sup>1</sup>, K. Bendrick<sup>3</sup>,  
M. R. Baumgartner<sup>4</sup>, E. Seifritz<sup>1</sup> and B. B. Quednow<sup>1\*</sup>

<sup>1</sup>Department of Psychiatry, Psychotherapy and Psychosomatics, University Hospital of Psychiatry Zurich, Zurich, Switzerland

<sup>2</sup>Behavioural and Clinical Neuroscience Institute, Department of Experimental Psychology, University of Cambridge, Cambridge, UK

<sup>3</sup>Department of Economics, University of Konstanz, Konstanz, Germany

<sup>4</sup>Institute of Legal Medicine, University of Zurich, Zurich, Switzerland

**Background.** Maladaptive decision-making is assumed to be a core feature of cocaine addiction. Indeed, numerous studies have reported deficits in non-social decision-making tasks and reward-related impulsivity in dependent cocaine users. However, social decision-making has not been examined in cocaine users yet. Moreover, it is unknown if even recreational and non-dependent cocaine use is linked to decision-making deficits. Therefore, we investigated whether recreational and dependent cocaine users exhibit alterations in social and non-social decision-making.

**Method.** The performance of healthy controls ( $n=68$ ), recreational cocaine users ( $n=68$ ) and dependent cocaine users ( $n=30$ ) in classical decision-making paradigms (Iowa Gambling Task, Delay Discounting) and in social interaction paradigms (Distribution Game, Dictator Game) was assessed.

**Results.** Decisions in the social interaction tasks of both cocaine user groups were more self-serving compared with controls as cocaine users preferred higher monetary payoffs for themselves. In the Iowa Gambling Task, only dependent cocaine users were more likely to choose disadvantageous card decks, reflecting worse decision-making. They were also more likely to choose immediate smaller rewards over larger delayed rewards in the Delay Discounting task.

**Conclusions.** Our results imply that both recreational and dependent cocaine users are more concerned with their own monetary gain when interacting with another person. Furthermore, primarily dependent cocaine users are less foresighted and more impulsive regarding immediate reward. Overall, social interaction deficits are already present in recreational users, while non-social decision-making deficits occur predominantly in dependent cocaine users. Thus, social interaction training and cognitive remediation strategies may improve treatment success and quality of life in cocaine dependence.

Received 1 November 2012; Revised 7 June 2013; Accepted 25 June 2013; First published online 22 July 2013

**Key words:** Delay of gratification, dependence, intertemporal discounting, neuroeconomics, prefrontal cortex.

## Introduction

Cocaine is the second most used illegal drug in Europe after cannabis and it is estimated that 15.5 million Europeans have tried cocaine at least once in their life, amounting to a lifetime prevalence of 4.6%. Moreover, cocaine is the primary illegal drug responsible (European Monitoring Centre for Drugs and Drug Addiction, 2012) for drug-dependence treatment in North and South America (United Nations Office on Drugs and Crime, 2011). Cocaine is classified as a highly addictive drug (Nutt *et al.* 2007) and it is

estimated that 5–6% of users will meet dependence criteria within the first year of use and around 21% by the age of 45 years (Wagner & Anthony, 2007). The health risks associated with cocaine abuse include severe medical complications, such as cardiovascular or respiratory incidences, and a number of psychiatric disorders (Buttner, 2012). Because drug addiction results in high economic and societal costs (Olesen *et al.* 2012), and effective pharmacological treatment options are currently lacking (O'Brien, 2005), an adequate characterization of the core feature of cocaine addiction, maladaptive decision-making, is crucial for the development of effective prevention and treatment strategies.

The term 'decision-making' describes the ability to select an optimal course of action from multiple alternatives. Selecting an adequate choice entails constant updating and integrating of information about

\* Address for correspondence: B. B. Quednow, Ph.D., Experimental and Clinical Pharmacopsychology, Department of Psychiatry, Psychotherapy and Psychosomatics, University Hospital of Psychiatry Zurich, Lengstrasse 31, CH-8032 Zurich, Switzerland.  
(Email: quednow@bli.uzh.ch)

the value of present and potential actions as well as future states pertaining to current needs (Fellows, 2004; Lucantonio et al. 2012). Decision-making deficits in dependent cocaine users (DCU) are well captured by the paradox that they compulsively seek and take the drug despite encountering adverse legal, financial, health-related and social consequences (Koob, 2009). In line with these everyday life difficulties, experimental studies in DCU have provided evidence that chronic cocaine use is linked to deficits in the processing of reward and punishment contingencies, as measured by the Iowa Gambling Task (IGT) (Bechara et al. 2002; Verdejo-Garcia et al. 2007a; Kjome et al. 2010), and a preference for smaller immediate over larger delayed rewards, henceforth referred to as Delay Discounting (DD) (Kirby & Petry, 2004; Heil et al. 2006; Bickel et al. 2011a). Consequently, it has been suggested that DCU experience 'myopia for the future'; they thus fail to incorporate ongoing feedback to guide future behaviours and instead make impulsive decisions that are based on immediate reward availability (Rachlin & Green, 1972; Ainslie, 1975; Bechara et al. 2002). An elevated reward impulsivity as measured by DD has been associated with negative outcomes in the financial, academic and health domains (Mischel et al. 2011) and poor treatment response in DCU (Washio et al. 2011).

Because we live in complex, continuously changing social environments, decision-making often takes place in the form of social interaction and is strongly influenced by self and others regarding preferences and social cognitive abilities (e.g. emotion recognition, theory of mind, empathy) (Couture et al. 2006; Fehr & Camerer, 2007). Social decision-making (SDM) encompasses multiple facets including trust, cooperation, fairness, altruism, norm-abiding decision-making, punishment, social learning, and competitive social interactions (Rilling & Sanfey, 2011). To date SDM has not been investigated in an experimental setting in cocaine users; however, a growing number of findings imply that cocaine users may exhibit deficits during social interactions. For instance, decision-making deficits in crack cocaine-dependent individuals were associated with self-reported social dysfunction (Cunha et al. 2011). Moreover, cocaine users feature a 22-fold increased risk for an antisocial personality disorder (ASPD) and clinical reports have given account of egocentrism and blunted emotion in cocaine-dependent individuals (Rounsaville, 2004). Notably, adequate socio-cognitive abilities are known to have a strong impact on the development, course and outcome of psychiatric diseases (Couture et al. 2006) and may also affect the course of dependence and treatment success in stimulant abusers (Homer et al. 2008). Thus, understanding how cocaine addiction

may be associated with maladaptive social interaction is fundamentally important. Recent advances in game-theoretic approaches have provided the unique opportunity to quantify SDM in psychiatric disorders (Kishida et al. 2010). Therefore, we relied on economic decision-making tasks to investigate social preferences such as fairness and efficiency preferences in cocaine users in comparison with controls. In these tasks, participants are considered fair if they distribute money evenly between themselves and their interaction partner, whereas they are deemed unfair if they allocate money in a more self-serving manner as reflected by a higher monetary payoff for themselves and a smaller payoff for their interaction partner. Furthermore, the design also allows an assessment of individuals' efficiency preferences, i.e. any subject motivated by efficiency concerns values the total monetary payoff for the group positively.

Given that the transition to dependence is not dichotomous but rather gradual, advancing from habitual to compulsive use (Haber, 2008), recreational cocaine use can be thought of as an intermediate step in addiction. Recreational cocaine users (RCU) are not (yet) addicted but administer the drug regularly for personal pleasure. A growing number of studies suggest that the recreational use of cocaine or prescription stimulants is associated with subtle cognitive impairments in attention, memory and components of executive functions (Rahman & Clarke, 2005; Colzato et al. 2007, 2009b; Reske et al. 2010, 2011; Soar et al. 2012; Vonmoos et al. 2013) that are similar but less pronounced compared with DCU (Jovanovski et al. 2005; Vonmoos et al. 2013). Additionally, we recently reported changes in early information processing and blue-yellow colour vision deficits in RCU, suggesting putative alterations of catecholamine neurotransmission at an early stage of cocaine abuse (Hulka et al. 2013; Preller et al. 2013). Moreover, young adults with recreational stimulant use showed more pronounced risk-taking behaviour (Leland & Paulus, 2005) and a subgroup of cocaine-preferring occasional stimulant users exhibited altered neural activity during a reinforcement-based decision-making task (Stewart et al. 2013). However, decision-making has not systematically been investigated in relatively pure RCU.

In this cross-sectional study, we investigate SDM and non-SDM (NSDM) behaviour of RCU and DCU in comparison with an age-, sex- and verbal intelligence-matched healthy control group. We report on effects of cocaine abuse on measures of fairness and efficiency preferences and extend on previous reports in the domain of risk taking and discounting of delayed rewards by incorporating a group of RCU. Based on prior studies demonstrating more subtle cognitive and decision-making deficits in recreational

stimulant users compared with dependent users (Colzato *et al.* 2009a; Reske *et al.* 2010, 2011; Soar *et al.* 2012; Stewart *et al.* 2013; Vonmoos *et al.* 2013) and a very clear dose-dependent association of cognitive dysfunctions and cumulative cocaine use (Vonmoos *et al.* 2013), we hypothesize that RCU exhibit similar but less pronounced behavioural changes compared with DCU.

## Method

### Participants

The present sample represents the cross-sectional part of the longitudinal Zurich Cocaine Cognition Study (ZuCo<sup>2</sup>St) and consists of 68 RCU, 30 DCU and 68 healthy control subjects (total of 166 subjects). Details regarding recruitment, selection process and study procedure are provided in the online Supplementary text. Inclusion criteria for the two cocaine user groups were cocaine as the primary drug (>2-fold higher cocaine concentrations in the hair samples than any other drug), cocaine use of >0.5 g per month (over the past 6 months), and an abstinence duration of <6 months. Cocaine dependence was diagnosed according to the Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) criteria, with only DCU meeting these criteria. All participants had to be aged between 18 and 60 years and proficient in German. Exclusion criteria were use of opioids, excessive MDMA intake (>50 pills lifetime in RCU, >200 pills in DCU, >5 pills in controls), excessive cannabis use (>5 g per week or daily use), intake of prescription drugs affecting the CNS, presence of a current or previous Axis I DSM-IV psychiatric disorder [other than cocaine and alcohol abuse/dependence, attention deficit hyperactivity disorder (ADHD), and a former affective disorder], neurological disorders or head injury, and a family history of a severe DSM-IV psychiatric disorder such as schizophrenia, bipolar disorder or obsessive-compulsive disorder. Participants were instructed to abstain from illegal drugs for a minimum of 3 days and from alcohol for at least 24 h prior to study completion.

Polytoxic drug abuse is one of the major confounding factors in addiction research and the reliability of self-reported data has been questioned (Hser, 1997). Therefore, urine samples were collected on the day of testing to control for recent drug use. To objectively characterize drug use over the last 6 months, hair samples were collected on the day of testing and analysed with liquid chromatography–mass spectrometry (see online Supplementary text).

Participants received financial compensation of 170–225 Swiss Francs (CHF), depending on their

decisions in some of the tasks. The study was approved by the Cantonal Ethics Committee of Zurich and all participants provided written informed consent.

### Clinical interviews and questionnaires

The Structured Clinical Interview for DSM-IV Disorders (SCID-I) was carried out by trained psychologists and all participants completed the SCID-II personality questionnaire to evaluate the severity of ASPD symptoms (Wittchen *et al.* 1997a,b). To estimate pre-morbid verbal intelligence the Mehrfachwahl-Wortschatz-Intelligenztest (MWT-B; multiple choice vocabulary intelligence test) was applied (Lehrl, 1999). Drug use was assessed by means of the Interview for Psychotropic Drug Consumption, which has been described in detail elsewhere (Quednow *et al.* 2004). The brief version of the Cocaine Craving Questionnaire was used to assess current cocaine craving (Tiffany *et al.* 1993; Sussner *et al.* 2006). As psychiatric co-morbidities such as ADHD and depression are frequently present among addicted individuals (Rounsaville, 2004; Ivanov *et al.* 2008; Perez de Los Cobos *et al.* 2011), we used the Attention Deficit Hyperactivity Disorder Self-Rating Scale (ADHD-SR; Rosler *et al.* 2004) and the Beck Depression Inventory (BDI; Beck *et al.* 1961). In the ADHD-SR, clinically relevant ADHD symptoms were diagnosed if at least six of items 1–9 (inattention) were affirmed, at least three of the items 10–14 (hyperactivity), and at least one of the items 15–18 (function).

### Behavioural tasks

#### SDM

Participants' social preferences were assessed in a Distribution Game followed by a Dictator Game (Charness & Rabin, 2002; Engelmann & Strobel, 2004) implemented in z-Tree (Fischbacher, 2007). Participants were informed that each of the games only had one trial and that each game was played with a different interaction partner. The Distribution Game involves two players, player A and B. Player A chooses one of 10 possible point distributions ranging from a fair distribution where both players would receive 25 points each to the most opportunistic distribution where player A would receive 40 points and player B one point. Player B is a passive recipient and is merely informed about which distribution player A chose and how many points both players receive. In addition, the Distribution Game also allows classification of subjects according to their efficiency preferences. A subject that is motivated by efficiency concerns values the total monetary payoff for the dyad positively.

Participants were classified as 'fair' when they chose the first distribution, as 'unfair and efficient' when they chose distributions two to five (yielding a higher total payoff), and as 'unfair and inefficient' when they selected distributions six to 10 (resulting in lower overall payoff for the dyad; online Supplementary Fig. S1). The Dictator Game always followed the Distribution Game, and the participants were told that they would play with another player B. Player A receives an endowment of 50 points and can give any amount from 0 to 50 points to player B. All subjects received a payment according to the points earned in the tasks. In both tasks, each point earned was worth CHF 0.25. At the end of the experiment subjects received payment in cash or via online banking. In order to guarantee anonymity of the cocaine users, interaction partners were simulated by the computer (details in the online Supplementary material). After study completion participants were asked whether they had doubts about the realness of their interaction partners by means of a five-point Likert scale (1=not at all to 5=very much).

#### NSDM

We tested participants' risk-taking preferences and planning abilities using the IGT, which has been described in detail before (Bechara *et al.* 2002; Quednow *et al.* 2007). Intertemporal choice was measured using the DD according to Kirby *et al.* (1999). Further details of the tasks are given in the online Supplementary material.

#### Statistical analysis

Statistical analyses were performed with PASW 19.0 (SPSS Inc., USA). Demographic data and drug use patterns of the three groups were analysed by means of analysis of variance (ANOVA) with Sidak-corrected *post-hoc* analyses and by means of frequency analyses (Pearson's  $\chi^2$  test). We conducted correlation analyses (Pearson's product-moment) to examine if performance in SDM and NSDM tasks was related. Moreover, we conducted multiple regression analyses to examine the association of pre-selected predictors including age, sex, years of education, and two dummy coded group contrasts (controls *v.* RCU, controls *v.* DCU) with SDM and NSDM. Further multiple regression analyses were conducted to investigate how drug use patterns are related to performance in the SDM and NSDM tasks. Finally, the potential effects of psychiatric symptoms, cocaine craving and recent drug use (positive urine toxicology) on SDM and NSDM were explored by means of multiple regression analyses. As the assumptions of homoscedasticity and parametric

distribution were not met by some variables, the drug use variables grams per week and lifetime use in grams were log-transformed ( $\log_{10}$ ) and the constant 1 was added because the data contained 0 values. To reduce data quantity, to obtain a measure of how strongly cocaine users deviate from controls' SDM preferences, and because of a significant intercorrelation of the Distribution and Dictator Games ( $r=0.61, p<0.001$ ) we computed a composite SDM score. The composite score was derived by averaging z-transformed measures of the Distribution and Dictator Games (payoffs B) according to means and standard deviations of the control group.

## Results

### Demographic variables

Groups did not differ regarding socio-economic status (online Supplementary Table S1) and demographic variables except for years of education (Table 1). RCU and DCU did not differ from controls regarding age, but DCU were by trend slightly older than RCU. Moreover, there were marginally, but not significantly, more males in the cocaine user groups compared with controls and a previous study has reported sex differences in IGT performance (Bolla *et al.* 2004). Therefore, we introduced years of education, age and sex as covariates in all statistical models. As RCU and DCU both reported more symptoms of ADHD (14 RCU and eight DCU exhibited clinically relevant ADHD symptoms) and depression than controls (Table 1), additional analyses were conducted to examine a potential association of these factors with decision-making.

### Self-reported and objective drug use

Self-reported drug use showed that RCU used cocaine on a regular basis, with a mean weekly consumption of about 1 g cocaine. Several participants tested positive for cocaine and cannabis in the urine toxicology analyses but we decided not to exclude them in order to investigate potential effects of recent drug use (Table 2).

Results from the hair toxicology analyses revealed that self-reported cocaine use (g/week, cumulative dose, duration of use) corresponded with concentrations of cocaine and its metabolites in the hair samples ( $r=0.29-0.41$ , all  $p<0.01$ ). Importantly, hair toxicology provided evidence that the RCU and DCU enrolled in the present study are unique with regard to three crucial aspects (Table 3). (1) For both drug user groups, cocaine had been the main drug of use over the past 6 months and concentrations of cocaine

**Table 1.** Demographic data

	Controls (n=68)	Recreational cocaine users (n=68)	Dependent cocaine users (n=30)	Value <sup>a</sup>	p <sup>a</sup>	df
Mean age, years (s.d.) [range]	30.63 (9.15) [19–57]	28.71 (6.19) [20–46]	32.80 (9.54) [19–56]	2.77	0.07	2
Sex, n (%)						
Male	46 (68)	50 (73)	22 (73)	0.66	0.72	2
Female	22 (32)	18 (27)	8 (27)			
Mean years of education (s.d.)	10.61 (1.77)	10.50 (1.96)*	9.48 (1.19)*	4.6	0.01	2
Mean estimated verbal IQ (s.d.)	104.66 (10.41)	103.21 (9.58)	100.93 (12.01)	1.36	0.26	2
Smoking, n (%)						
Smokers	54 (79)	61 (90)	27 (90)	3.5	0.17	2
Non-smokers	14 (21)	7 (10)	3 (10)			
Mean ADHD-SR (s.d.)	7.84 (4.71)	13.16 (8.98)**	17.00 (8.85)**	17.6	0.00	2
Mean Beck Depression Inventory (s.d.)	4.41 (4.38)	7.35 (6.14)*	11.8 (8.58)***	15.85	0.00	2
Mean ASPD symptoms, SCID-II (s.d.)	16.62 (13.75)	28.28 (21.15)**	31.43 (26.90)**	8.29	0.00	2
Mean doubts about interaction partner realness (s.d.)	2.74 (1.48)	2.12 (1.33)*	2.07 (1.46)	4.01	0.02	2
Mean Cocaine Craving Questionnaire (s.d.)	–	19.04 (9.10)	20.90 (11.68)	–0.85 <sup>b</sup>	0.40	96

df, Degrees of freedom; s.d., standard deviation; IQ, intelligence quotient; ADHD-SR, Attention Deficit Hyperactivity Disorder-Self-Rating; ASPD, Antisocial Personality Disorder; SCID-II, Structured Clinical Interview for DSM-IV Axis II.

<sup>a</sup> Analysis of variance (all groups) or  $\chi^2$  test (all groups) for frequency data.

<sup>b</sup> *t* test (only cocaine user groups).

Mean value was significantly different from that of the control group: \*  $p < 0.05$ , \*\*  $p < 0.01$  (*post-hoc* test, Sidak).

†† Mean value was significantly different from that of the recreational cocaine user group ( $p < 0.01$ , *post-hoc* test, Sidak).

and its metabolites were seven-fold higher in DCU than in RCU. (2) RCU and DCU did not differ significantly with regard to concentrations of amphetamines, methylphenidate, MDMA and opiates. (3) For both RCU and DCU, concentrations of amphetamines and opiates were below the recommended cut-off value of 200 pg/mg (Cooper *et al.* 2012), indicating no regular use of these drugs over the past 6 months. Although the MDMA concentrations for RCU and DCU were above the cut-off value for MDMA, it is noteworthy that these concentrations are rather low and substantially lower than cocaine concentrations. Therefore, the present cocaine user samples had little poly-toxic drug use and did not differ from one another with regard to drugs other than cocaine.

### Task correlations

Correlations revealed that SDM and NSDM tasks indeed measured different aspects of decision-making as neither the Distribution Game nor the Dictator Game was associated with the IGT ( $r = -0.01$  and  $-0.02$ ) and DD ( $r = -0.05$  and  $-0.11$ ). The Distribution Game and the Dictator Game correlated significantly ( $r = 0.61$ ,  $p < 0.001$ ), whereas the IGT and the DD did not correlate ( $r = 0.01$ ).

### SDM

In the Distribution Game, DCU ( $\beta = -0.20$ ,  $t = -2.40$ ,  $p < 0.05$ ) and by trend also RCU ( $\beta = -0.15$ ,  $t = -1.78$ ,  $p = 0.08$ ) chose point distributions that were more profitable for themselves and yielded lower payoffs for participant B (Table 4 and online Supplementary Table S2). Overall, participants from all three groups chose the fair distribution most often (Fig. 1, Table 1). However, DCU chose the unfair inefficient distributions more frequently compared with controls ( $\chi^2_1 = 10.74$ ,  $p < 0.01$ ) and RCU ( $\chi^2_1 = 8.3$ ,  $p < 0.01$ ) (online Supplementary Figs S1 and S2). Analyses regarding the Dictator Game showed trends that both RCU ( $\beta = -0.16$ ,  $t = -1.85$ ,  $p = 0.07$ ) and DCU ( $\beta = -0.15$ ,  $t = -1.73$ ,  $p = 0.09$ ) gave fewer points to their interaction partners compared with controls (Table 4 and online Supplementary Table S2). Analysis of the SDM composite score revealed that both RCU ( $\beta = -0.17$ ,  $t = -2.02$ ,  $p < 0.05$ ) and DCU ( $\beta = -0.20$ ,  $t = -2.35$ ,  $p < 0.05$ ) acted in a more self-serving manner than controls (Table 4 and online Supplementary Table S2). All groups reported to have had only a few doubts about the realness of the interaction partners. Interestingly, RCU ( $p < 0.05$ ) and by trend DCU ( $p < 0.1$ ) reported even fewer doubts than controls (Table 1). Introducing

**Table 2.** Self-reported drug use<sup>a</sup>

	Controls (n=68)	Recreational cocaine users (n=68)	Dependent cocaine users (n=30)	Value <sup>b</sup>	p <sup>b</sup>	df/df <sub>error</sub>
<b>Nicotine</b>						
Cigarettes per day	9.29 (9.73)	11.7 (8.77)	16.05 (13.77)**	4.59	0.01	2/163
Years of use	9.61 (9.54)	9.65 (6.37)	13.55 (8.54)	2.82	0.06	2/163
<b>Alcohol</b>						
g per week	110.49 (120.21)	167.8 (117.47)	199.7 (259.4)*	4.28	0.02	2/163
Years of use	13.62 (9.38)	11.23 (5.07)	12.89 (8.64)	1.66	0.19	2/163
<b>Cocaine</b>						
Times per week	0.00 (0.00)	1.07 (1.03)**	2.93 (2.53)**++	57.16	0.00	2/163
g per week	0.00 (0.00)	1.11 (1.41)	6.17 (8.70)**++	28.57	0.00	2/163
Years of use	0.00 (0.00)	6.47 (3.99)**	9.22 (6.43)**++	82.51	0.00	2/163
Maximum dose, g/day	–	3.46 (2.47)	8.75 (7.86)++	–3.61	0.00	96
Cumulative dose, g	0.00 (0.00)	519.69 (751.23)	4619.94 (8658.35)**++	17.55	0.00	2/163
Last consumption, days [n]	–	27.45 (37.6) [68]	20.43 (33.78) [30]	0.88	0.38	96
Urine toxicology, n (%)						
Positive	0 (0)	10 (15)	13 (43)	32.82 <sup>c</sup>	0.00	2
Negative	68 (100)	57 (85)	17 (57)++			
<b>Amphetamines</b>						
g per week	0.00 (0.00)	0.08 (0.21)**	0.05 (0.19)	4.65	0.01	2/163
Years of use	0.01 (0.00)	1.63 (2.97)**	1.54 (3.16)**	9.42	0.00	2/163
Cumulative dose, g	0.18 (1.42)	21.19 (56.77)*	22.26 (62.80)	4.52	0.01	2/163
Last consumption, days [n]	–	90.46 (145.48) [24]	78.38 (75.42) [6]	–0.09	0.93	60
<b>MDMA</b>						
Pills per week	0.00 (0.00)	0.08 (0.25)	0.41 (1.83)	2.88	0.06	2/163
Years of use	0.25 (1.64)	2.47 (3.76)**	3.06 (5.22)**	10.12	0.00	2/163
Cumulative dose, pills	0.73 (2.75)	35.86 (90.47)	157.38 (393.52)**++	8.37	0.00	2/163
Last consumption, days [n]	–	124.91 (167.18) [21]	82.13 (45.43) [9]	–1.32	0.19	68
<b>Cannabis</b>						
g per week	0.53 (1.50)	0.86 (2.05)	1.22 (3.74)	1.02	0.36	2/163
Years of use	4.68 (6.63)	7.74 (6.03)*	9.54 (8.94)**	6.26	0.00	2/163
Cumulative dose, g	479.16 (1083.03)	1042.85 (1780.04)	2626.67 (3857.12)**++	10.87	0.00	2/163
Last consumption, days [n]	39.02 (50.42) [29]	22.44 (32.57) [43]	72.75 (211.62) [18]	1.60	0.21	2/87
Urine toxicology, n (%)						
Positive	9 (13)	12 (18)	9 (30)	3.94 <sup>c</sup>	0.14	2
Negative	59 (87)	55 (82)	21 (70)			
<b>Serotonergic hallucinogens<sup>d</sup></b>						
Cumulative dose, times	0.80 (2.17)	6.03 (14.59)*	5.75 (10.47)	4.88	0.01	2/163
Last consumption, months [n]	97.57 (93.54) [14]	66.24 (61.18) [29]	181.99 (339.56) [18]	2.01	0.14	2/58
<b>GHB</b>						
Cumulative dose, times	0.00 (0.00)	1.76 (9.48)	1.28 (2.89)	1.43	0.24	2/163
Last consumption, months [n]	–	126.07 (31.37) [3]	30.00 [1]	–	–	–

Data are given as mean (standard deviation) unless otherwise indicated.

df, Degrees of freedom; MDMA, 3,4-methylenedioxy-N-methylamphetamine (methylenedioxymethamphetamine); GHB,  $\gamma$ -hydroxybutyric acid; LSD, lysergic acid diethylamide; DMT, *N,N*-dimethyltryptamine; 2-CB, 4-bromo-2,5-dimethoxyphenethylamine.

<sup>a</sup> Consumption per day or week captures the last 6 months; duration of use and cumulative dose are averaged within the total group. Last consumption is averaged only for subjects who used the drug in the last 6 months. In this case, sample size is shown.

<sup>b</sup> Analysis of variance or  $\chi^2$  test (all groups).

<sup>c</sup> *t* test (only cocaine user groups).

<sup>d</sup> Hallucinogens=psilocybin, LSD, DMT, 3,4,5-trimethoxyphenethylamine (mescaline), 2-CB.

Mean value was significantly different from that of the control group: \*  $p < 0.05$ , \*\*  $p < 0.01$  (*post-hoc* test, Sidak).

++ Value was significantly different from that of the recreational cocaine user group ( $p < 0.01$ , *post-hoc* test, Sidak).

**Table 3.** Hair toxicological analyses<sup>a</sup>

	Controls (n=68)	Recreational cocaine users (n=68)	Dependent cocaine users (n=30)	Value <sup>b</sup>	p <sup>b</sup>	df/df <sub>error</sub>
<b>Cocaine<sup>c</sup></b>						
Cocaine, pg/mg	0.00 (0.00)	2739.18 (4627.66)**	19135.67 (29168.78)**++	24.20	0.00	2/159
Benzoyllecgonine, pg/mg	0.00 (0.00)	545.82 (919.19)**	4002.67 (5733.19)**++	27.50	0.00	2/159
Ethylcocaine, pg/mg	0.00 (0.00)	275.89 (316.32)**	2034.33 (3644.51)**++	18.30	0.00	2/159
Norcocaine, pg/mg	0.00 (0.00)	62.44 (100.8)**	486.17 (586.29)**++	36.68	0.00	2/159
<b>Amphetamines<sup>d</sup></b>						
Amphetamine, pg/mg [n]	0.92 (7.44) [1]	76.34 (256.47)	59.67 (169.35)	3.04	0.05	2/159
Methamphetamine, pg/mg	0.00 (0.00)	1.19 (9.77)	1.33 (7.30)	0.61	0.55	2/159
Methylphenidate, pg/mg	0 (0)	10 (55)	5 (15)	1.00	0.37	2/159
<b>MDMA<sup>d</sup></b>						
MDMA, pg/mg [n]	1.81 (14.57) [1]	545.05 (1598.36)*	255.17 (652.54)	4.28	0.02	2/159
MDEA, pg/mg	0.00 (0.00)	2.16 (17.71)	0.00 (0.00)	0.71	0.50	2/159
MDA, pg/mg [n]	0.12 (0.93) [1]	18.66 (57.31)*	9.17 (28.29)	3.76	0.03	2/159
<b>Opiates<sup>d</sup></b>						
Morphine, pg/mg	0 (0)	3 (25)	70 (320)	3.06	0.05	2/159
Codeine, pg/mg	0 (2)	20 (115)	35 (115)	1.79	0.17	2/159
Methadone, pg/mg	0 (0)	1 (10)	40 (210)	2.18	0.12	2/159
EDDP, pg/mg	0 (0)	0 (0)	5 (25)	2.23	0.11	2/159
Tramadol, pg/mg	0 (0)	3 (17)	310 (1640)	2.39	0.10	2/159

Data are given as mean (standard deviation).

df, Degrees of freedom; MDMA, 3,4-methylenedioxy-N-methylamphetamine (methylenedioxyamphetamine); MDEA, methylenedioxyethylamphetamine; MDA, 3,4-methylenedioxyamphetamine; EDDP, primary methadone metabolite.

<sup>a</sup> The hair analysis was performed on two hair samples (each 3 cm in length) per participant capturing drug use over the last 6 months. Concentrations were averaged over the two samples. If the hair sample was not long enough, only one sample was analysed (3 cm, 3 months).

<sup>b</sup> Analysis of variance (all groups).

<sup>c</sup> Cut-off value for cocaine=500 pg/mg.

<sup>d</sup> Cut-off value for amphetamines, MDMA and opiates=200 pg/mg.

Mean value was significantly different from that of the control group: \*  $p < 0.05$ , \*\*  $p < 0.01$  (*post-hoc* test, Sidak).

++ Mean value was significantly different from that of the recreational cocaine user group ( $p < 0.01$ , *post-hoc* test, Sidak).

the factor 'doubts about the realness of the interaction' as an additional predictor into the analysis neither changed the results for the Distribution Game (controls *v.* RCU:  $\beta = -0.16$ ,  $t = -1.85$ ,  $p = 0.07$ ; controls *v.* DCU:  $\beta = -0.21$ ,  $t = -2.34$ ,  $p < 0.05$ ) nor for the Dictator Game (controls *v.* RCU:  $\beta = -0.18$ ,  $t = -2.04$ ,  $p < 0.05$ ; controls *v.* DCU:  $\beta = -0.16$ ,  $t = -1.77$ ,  $p = 0.08$ ) and was not a significant predictor variable for either of the tasks ( $p > 0.6$ ). Surprisingly, SCID-II ASPD symptoms were not correlated with SDM parameters. Importantly, introduction of ASPD symptoms as a further predictor in the regression analyses did not change the main results, indicating that SDM alterations in cocaine users could not be explained by the presence of increased ASPD symptoms in this group.

Because in the Distribution Game efficiency preferences cannot be assessed independently from fairness

preferences, adding a Dictator Game allowed us to isolate efficiency preferences in the Distribution Game. Thus, because efficiency preferences do not matter in the Dictator Game, it serves both as a clean measure for fairness preferences but also as a control for the fairness domain of the Distribution Game. We found that almost all participants remained 'fair' in the Dictator Game if they had already been 'fair' in the Distribution Game. In contrast, controls who chose one of the distributions classified as 'unfair efficient' in the Distribution Game often chose a fair point allocation in the Dictator Game, while DCU who chose unfair efficient distributions in the Distribution Game were more likely to allocate points in the Dictator Game in a self-serving manner ( $\chi^2 = 5.03$ ,  $p < 0.05$ ). Furthermore, the number of subjects choosing one of the unfair inefficient distributions in the Distribution

**Table 4.** Behavioural task parameters<sup>a</sup>

	Controls ( <i>n</i> =68)	Recreational cocaine users ( <i>n</i> =68)	Dependent cocaine users ( <i>n</i> =30)	Effect size: Cohen's <i>d</i> <sup>b</sup>	Effect size: Cohen's <i>d</i> <sup>c</sup>
Social decision-making tasks					
Distribution Game, payoff B <sup>d</sup>	20.33 (1.00)	17.82 (1.00)	15.92 (1.53)	0.30†	0.53*
Distribution Game, total amount	51.56 (0.57)	50.14 (0.57)	49.64 (0.87)	0.30	0.41
Distribution Game, <i>n</i>					
Fair	25	22	10		
Unfair efficient	23	23	6		
Unfair inefficient	20	23	14		
Dictator Game, payoff B <sup>d</sup>	18.45 (1.31)	15.02 (1.31)	14.28 (2.01)	0.31†	0.38†
Dictator Game, <i>n</i>					
Fair	44	36	14		
Unfair	24	32	16		
Social decision-making, <i>z</i> -standardized composite score	-0.01 (0.12)	-0.34 (0.12)	-0.52 (0.18)	0.34*	0.52*
Non-social decision-making tasks					
Iowa Gambling Task, total ratio – good:bad cards	18.63 (3.14)	13.64 (3.17)	7.91 (4.83)	0.19	0.41†
Iowa Gambling Task, total points	4301.21 (160.84)	4054.9 (162.44)	3785.32 (247.63)	0.19	0.39†
Delay Discounting					
<i>k</i> overall <sup>e</sup>	0.013 (0.004)	0.019 (0.004)	0.034 (0.006)	0.20	0.69**
<i>k</i> for large amounts <sup>e</sup>	0.009 (0.003)	0.016 (0.003)	0.031 (0.005)	0.24	0.75**
<i>k</i> for medium amounts <sup>e</sup>	0.014 (0.004)	0.020 (0.004)	0.037 (0.006)	0.19	0.72**
<i>k</i> for small amounts <sup>e</sup>	0.025 (0.005)	0.039 (0.006)	0.051 (0.008)	0.31†	0.57**

Data are given as mean (standard error) unless otherwise indicated.

<sup>a</sup> All parameters are corrected for age, sex and years of education.

<sup>b</sup> Controls *versus* recreational cocaine users.

<sup>c</sup> Controls *versus* dependent cocaine users.

<sup>d</sup> Used for the social decision-making composite score.

<sup>e</sup> *k*=free parameter that determines the discounting rate.

Significant *post-hoc* test: \* *p*<0.05, \*\**p*<0.01 (Sidak).

† Marginally significant *post-hoc* test (*p*<0.1, Sidak).

Game was substantially higher among RCU and DCU than among controls and both groups almost exclusively allocated points in an unfair manner in the Dictator Game ( $\chi^2=4.17-5.03$ , *p*<0.05).

### NSDM

A repeated-measures analysis of covariance revealed that in the IGT (Table 4 and online Supplementary Table S2), despite the fact that, overall, both RCU (net score: *d*=0.18) and particularly DCU (*d*=0.49) chose fewer favourable cards than controls, no statistically significant group effect emerged ( $F_{2,159}=1.81$ , *p*=0.17). As expected, the factor quartile was significant ( $F_{3,477}=2.83$ , *p*<0.05), reflecting a learning curve (online Supplementary Fig. S4A).


In the DD (Table 4 and online Supplementary Table S2), groups significantly differed in their preferences for smaller immediate and larger delayed

monetary rewards ( $F_{2,163}=6.52$ , *p*<0.01; online Supplementary Fig. S3B). Sidak-corrected *post-hoc* comparisons showed that DCU were more likely to choose immediate rewards compared with controls (*p*<0.01). As expected, discounting of delayed rewards varied with reward magnitude ( $F_{2,326}=34.79$ , *p*<0.001). Correlation analyses showed that *k* for medium amounts was strongly related to the cocaine metabolite ethylcocaine determined in the hair toxicology (*r*=0.37, *p*<0.0001), indicating that especially subjects who consumed cocaine in combination with alcohol showed increased levels of impulsivity with regard to reward (Pennings *et al.* 2002).

### Multiple regression and correlation analyses of substance use

Associations between drug use patterns and SDM and NSDM tasks were assessed by multiple regression





**Fig. 1.** Frequency (%) of how often the three groups chose the 'fair', 'unfair efficient' and 'unfair inefficient' distributions in the Distribution Game and whether they distributed the money in a fair or unfair manner in the Dictator Game. Almost all of the participants, independent of which group they belong to, remained 'fair' in the Dictator Game if they already had been 'fair' in the Distribution Game. In contrast, controls who chose one of the distributions classified as 'unfair efficient' in the Distribution Game often chose a fair point allocation in the Dictator Game, while dependent cocaine users (DCU) who also chose one of the distributions classified as 'unfair efficient' in the Distribution Game were more likely to allocate points in the Dictator Game in a selfish manner. Moreover, the number of subjects choosing one of the 'unfair inefficient' distributions in the Distribution Game was substantially higher among recreational cocaine users (RCU) and DCU than among controls, and these RCU and DCU almost exclusively allocated the money in an unfair manner in the Dictator Game.

models, with cumulative drug use, weekly consumption and duration of use as predictor variables. Only cocaine users were included for these analyses ( $n=98$ ). All three models had the common predictors of cocaine craving, positive cocaine urine toxicology, and age of cocaine use onset to control for recent drug effects, craving urges, and potentially more severe cocaine-related developmental effects due to early age of cocaine use onset. Only models explaining significant amounts of variance are reported in this section (see online Supplementary Tables S3 and S4).

None of the drug variables in the three models predicted behaviour in the Distribution and Dictator Games (online Supplementary Table S3) and the IGT (online Supplementary Table S4). In contrast, cumulative cocaine and cannabis use as well as years of cocaine and cannabis use were significant predictors for performance in the DD, reflecting that more intense and longer cocaine use was associated with stronger discounting of delayed rewards, whereas a higher and longer cannabis consumption was associated with lower discounting of delayed rewards (online Supplementary Table S4). Weekly consumption of cocaine, MDMA, cannabis, alcohol, and nicotine neither predicted performance in SDM nor NSDM tasks. Moreover, age of cocaine use onset was not a significant predictor in any of the SDM and NSDM tasks.

We conducted additional correlation analyses to examine potential associations between performance in SDM and NSDM tasks and cannabis use in controls ( $n=68$ ). However, in controls none of the cannabis use parameters correlated with any of the tasks ( $p>0.21$ ).

#### Potential co-factors

Additional analyses were conducted to investigate potential effects of psychiatric symptoms, cocaine craving, and recent drug use on performance in SDM and NSDM tasks (online Supplementary text, Supplementary Table S5, Supplementary Fig. S4). In the SDM tasks, cocaine users with and without clinically relevant ADHD and depression symptoms exhibited more self-serving money allocation behaviour. Moreover, mainly cocaine users with high cocaine craving scores allocated money in a more self-serving manner compared with controls. Recent cocaine and cannabis intake (positive urine toxicology) was not significantly associated with performance in SDM tasks, as both cocaine users with positive and negative cocaine and cannabis urine toxicology exhibited more self-serving behaviour. None of the IGT regression models explained a significant amount of variance. Lastly, cocaine users with elevated BDI symptoms discounted delayed reward significantly stronger than controls. All other DD regression models were not significant.

#### Discussion

In this study, we report on differences in individual and social decision-making in RCU and DCU in comparison with a control group. Careful psychiatric diagnostic procedures ensured that cocaine users had few psychiatric co-morbidities and detailed hair toxicology analyses showed relatively sparse poly-substance use. Our study yielded the following major findings. (I) during social interaction, both RCU

and DCU distributed money in a more self-serving manner than controls. More specifically, both groups took more money for themselves in the Distribution Game and gave less money to the second interaction partner in the Dictator Game. (II) DCU exhibited significantly elevated reward-related impulsivity in the DD ( $d=0.69$ ) and chose fewer advantageous cards in the IGT ( $d=0.41$ ), although the latter was not statistically significant. Higher cumulative doses of cocaine and longer duration of use were associated with a lower net score in the IGT and stronger discounting of delayed rewards in the DD. Taken together, our results indicate that both RCU and DCU show more self-serving behaviour in social interaction tasks, whereas only DCU show deficits in the processing of reward and punishment contingencies and exhibit increased reward-related impulsivity.

To our knowledge, no studies have assessed human social interaction using an experimental economic approach in cocaine addiction research so far. We observed that control subjects who chose one of the distributions classified as 'unfair efficient' in the Distribution Game often chose a fair point allocation in the Dictator Game, while DCU who also chose one of the distributions classified as 'unfair efficient' in the Distribution Game were more likely to allocate points in the Dictator Game in a selfish manner. Thus, it appears that the proportion of controls who seem to place a higher value in efficient distributions do this at the cost of fairness towards the other player. However, if efficiency preferences do not matter, as in the Dictator Game, the same subjects still care for fairness. This is not observed in those DCU who chose efficient distributions in the Distribution Game, as they seem to care only about efficiency and less about fairness. Furthermore, among RCU and DCU the number of subjects choosing one of the unfair inefficient distributions in the Distribution Game was substantially higher and, among these, almost everyone allocated points in an unfair manner in the Dictator Game. Consequently, these findings suggest that cocaine users are less concerned about fairness in dyadic interactions, compared with controls. Although the self-serving behaviour was more pronounced in DCU than RCU when compared with controls, the absence of a significant correlation between amount of cocaine use and self-serving SDM could signify that cocaine users may have a predisposition towards more self-serving behaviour. Additionally, also cocaine craving enhances the propensity to act selfishly, indicating that SDM preferences in cocaine users also have a state component.

Because of the cross-sectional design of our study, it is impossible to substantiate whether the differences in SDM among cocaine users and controls are due to a

certain predisposition, drug-induced cerebral alterations, or an interaction thereof. Nevertheless, the fact that both cocaine user groups exhibited more self-serving SDM than controls and the lacking relationship between cocaine-use patterns and SDM behaviour putatively support a stronger implication of pre-existent factors. Consistent with this notion, a prior study showed that chronic cocaine use is associated with selective deficits with regard to higher-level emotional reasoning such as understanding, managing and regulating emotions (Fox *et al.* 2011). Therefore, cocaine users might have a vulnerability that hinders them to adopt another person's perspective and to feel empathy.

Regarding performance in the IGT, the results of the present study were largely consistent with earlier data (Bechara *et al.* 2002; Verdejo-Garcia *et al.* 2007a; Kjome *et al.* 2010) in that they showed that particularly DCU chose fewer advantageous cards in the IGT than controls. However, although a medium effect size was present, the difference was not statistically significant, which is consistent with results from a prior study (Bolla *et al.* 2003) but not others (Bechara *et al.* 2002; Verdejo-Garcia *et al.* 2007a; Kjome *et al.* 2010). The lack of statistical significance in our study might be explained by two reasons. First, we applied stringent criteria to exclude subjects with severe psychiatric co-morbidities and toxicological hair analyses ensured that participants had little co-use of other illegal drugs. Therefore, even the DCU of the present study might have a higher level of general functioning compared with those of other study samples (Bechara *et al.* 2002; Verdejo-Garcia *et al.* 2007a,b; Kjome *et al.* 2010). Second, the IGT gains were paid, which is in line with observations of a previous study reporting impaired IGT performance in cocaine users only if monetary gains were hypothetical but not when the money they won was actually paid (Vadhan *et al.* 2009).

In the present study, we replicate previous results on intertemporal choice in DCU (Kirby & Petry, 2004; Heil *et al.* 2006; Bickel *et al.* 2011a). Importantly, although not statistically significant, also RCU exhibited slightly steeper discounting rates than controls with small effect sizes across reward magnitudes ( $d=0.19-0.31$ ; Table 4). The stronger effect found for DCU and the correlations between higher cumulative cocaine doses and longer duration of use with stronger discounting of delayed rewards might suggest that reward-related impulsivity is increased by the use of cocaine. This interpretation is concordant with animal studies showing that chronic administration of cocaine can cause sustained elevations in impulsive choice in rats and monkeys (Olausson *et al.* 2007; Mendez *et al.* 2010). Nevertheless, DD has also been shown to have

trait-like stability (Casey *et al.* 2011; Mischel *et al.* 2011; Odum, 2011) and a prospective study revealed that the ability to delay gratification in childhood predicted physical health, substance dependence, finances and criminal offences in adulthood (Moffitt *et al.* 2011). Thus, it is probable that predisposed tendencies of impulsive decision-making may render individuals more prone to initiate drug use, and, subsequently, neuroadaptations induced by repeated cocaine use may amplify pre-existing reward impulsivity resulting in the well-described compulsive drug-seeking behaviour—the inability to forego rewarding short-term effects of the drug in favour of the long-term benefits associated with abstinence (Bolla *et al.* 1998). A surprising finding in our study was that greater cumulative cannabis use and longer duration of use were associated with less pronounced discounting of delayed rewards in cocaine users. To our knowledge, the only study where the relationship between cannabis use and DD was systematically investigated showed no significant differences in the DD preferences between former dependent marijuana users and controls and only a trend for stronger DD in current dependent marijuana users (Johnson *et al.* 2010). Combined, these results may suggest that the association between cannabis use and DD is weaker than for cocaine and other drugs. The cocaine users in our sample who strongly co-use cannabis exhibited lower reward impulsivity as compared with users with less cannabis co-use (see online Supplementary Fig. S4). Interestingly, cannabis use was not correlated with DD in controls.

Because psychiatric co-morbidities such as ADHD are frequently present among addicted individuals (Ivanov *et al.* 2008; Perez de Los Cobos *et al.* 2011), we investigated how ADHD symptoms influence decision-making behaviour. However, ADHD symptoms were not a significant confounding factor regarding our results. Interestingly, cocaine users with high but not those with low levels of cocaine craving acted in a more self-serving manner compared with controls in the SDM while craving intensity was not related to performance in the NSDM tasks. One could speculate that strong craving urges may have fostered thoughts about obtaining cocaine as soon as possible, which could have led cocaine users to maximize their monetary profit. Furthermore, also recent drug use did not seem to influence behaviour in SDM and NSDM tasks. In line with the finding of altered intertemporal choice in depressive patients (Takahashi *et al.* 2008), particularly cocaine users with slightly elevated depression scores exhibited stronger DD than controls with low depression scores. However, it should be noted that specifically DCU reported more depressive symptoms so that our

DD results are not simply explained by depression. Finally, symptoms of depression did not seem to impact SDM.

The current findings should be interpreted bearing some limitations in mind. Given the cross-sectional design it is not possible to answer conclusively whether deficits in SDM and NSDM precede cocaine use or are due to cocaine-induced neuroadaptations. Therefore, data from longitudinal and prospective investigations are desirable to decompose further the effects of predisposition and sequelae of cocaine use. In the current study, we merely obtained behavioural results. Combining functional imaging with behavioural measures could be of great importance for future studies. Finally, in order to guarantee anonymity of the cocaine users, we had to use a cover story in the social interaction paradigms. However, we assessed whether participants had doubts about a real interaction and introduced this measure as a covariate into the statistical analyses, which did not change the results.

Identifying vulnerability markers and adverse drug-induced effects with regard to impaired decision-making in cocaine users is critical and may benefit the development of successful prevention and treatment strategies enhancing quality of life. For example, it was recently demonstrated that working memory training decreased the propensity to discount delayed rewards in stimulant addicts (Bickel *et al.* 2011b). Moreover, a large body of research has provided evidence that remediation efforts targeting neurocognitive and social cognitive skills in schizophrenia patients improve real-world psychosocial and disease outcomes (Lindenmayer *et al.* 2013; Medalia & Saperstein, 2013; Mueser *et al.* 2013). Likewise, knowledge from tasks measuring SDM could be integrated in therapeutic interventions for cocaine-addicted individuals, for example in the form of social skills trainings.

## Conclusion

In sum, these findings are the first to show that RCU and DCU both exhibit more self-serving behaviour regarding money allocation in social interaction paradigms. Interestingly, mainly the DCU performed worse in the IGT and showed elevated reward-related impulsivity compared with controls. The absence of significant correlations between SDM preferences and cocaine use implies that changes in SDM may have a trait component. In contrast, the intermediate performance of RCU compared with controls and DCU in NSDM tasks and the association of higher cumulative cocaine doses and longer duration of cocaine use with a lower net gain in the IGT and stronger DD suggest that NSDM may partially be influenced by cocaine

use. Our results might have implications for the conceptualization of treatment approaches that specifically target social interaction and decision-making deficits in cocaine users.

### Supplementary material

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S0033291713001839>.

### Acknowledgements

The study was supported by grants from the Swiss National Science Foundation (SNSF; grant no. PP00P1-123516/1) and the Olga Mayenfisch Foundation. C.E. received a personal grant from the SNSF (grant no. PA00P1\_134135). We are grateful to Alex Bücheli (Streetwork Zürich), Rudolf Stohler and Roland Kowalewski (Research Group Substance Use Disorders, Clinic for General and Social Psychiatry, University Hospital of Psychiatry), Lars Stark and Thilo Beck (ARUD, Zurich), Eric La Serra (Klinik St Pirminsberg, Psychiatrie-Dienste Süd, Kanton St. Gallen), and Michael Schaub (Research Institute for Public Health and Addiction, Zurich) for supporting the recruitment of participants. Moreover, we thank Joëlle Barthassat, Christina Gruber, Nina Ingold, Kathrin Küpeli, Franziska Minder, Valery Rohrbach, Claudia Schulz, and Stefanie Turin for the excellent technical support. Finally, we thank Urs Fischbacher for providing the opportunity to run pilot studies in his laboratory and James Hulka for carrying out programming adaptations in the IGT.

### Declaration of Interest

None.

### References

- Ainslie G (1975). Specious reward: a behavioral theory of impulsiveness and impulse control. *Psychological Bulletin* **82**, 463–496.
- Bechara A, Dolan S, Hindes A (2002). Decision-making and addiction (part II): myopia for the future or hypersensitivity to reward? *Neuropsychologia* **40**, 1690–1705.
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J (1961). An inventory for measuring depression. *Archives of General Psychiatry* **4**, 561–571.
- Bickel WK, Landes RD, Christensen DR, Jackson L, Jones BA, Kurth-Nelson Z, Redish AD (2011a). Single- and cross-commodity discounting among cocaine addicts: the commodity and its temporal location determine discounting rate. *Psychopharmacology* **217**, 177–187.
- Bickel WK, Yi R, Landes RD, Hill PF, Baxter C (2011b). Remember the future: working memory training decreases delay discounting among stimulant addicts. *Biological Psychiatry* **69**, 260–265.
- Bolla KI, Cadet JL, London ED (1998). The neuropsychiatry of chronic cocaine abuse. *Journal of Neuropsychiatry and Clinical Neurosciences* **10**, 280–289.
- Bolla KI, Eldreth DA, London ED, Kiehl KA, Mouratidis M, Contoreggi C, Matochik JA, Kurian V, Cadet JL, Kimes AS, Funderburk FR, Ernst M (2003). Orbitofrontal cortex dysfunction in abstinent cocaine abusers performing a decision-making task. *Neuroimage* **19**, 1085–1094.
- Bolla KI, Eldreth DA, Matochik JA, Cadet JL (2004). Sex-related differences in a gambling task and its neurological correlates. *Cerebral Cortex* **14**, 1226–1232.
- Buttner A (2012). Review: the neuropathology of drug abuse. *Neuropathology and Applied Neurobiology* **37**, 118–134.
- Casey BJ, Somerville LH, Gotlib IH, Ayduk O, Franklin NT, Askren MK, Jonides J, Berman MG, Wilson NL, Teslovich T, Glover G, Zayas V, Mischel W, Shoda Y (2011). Behavioral and neural correlates of delay of gratification 40 years later. *Proceedings of the National Academy of Sciences* **108**, 14998–15003.
- Charness G, Rabin M (2002). Understanding social preferences with simple tests. *Quarterly Journal of Economics*, 817–869.
- Colzato LS, Huizinga M, Hommel B (2009a). Recreational cocaine polydrug use impairs cognitive flexibility but not working memory. *Psychopharmacology* **207**, 225–234.
- Colzato LS, van den Wildenberg WP, Hommel B (2007). Impaired inhibitory control in recreational cocaine users. *PLoS One* **2**, 1143.
- Colzato LS, van den Wildenberg WP, Hommel B (2009b). Reduced attentional scope in cocaine polydrug users. *PLoS One* **4**, e6043.
- Cooper GA, Kronstrand R, Kintz P (2012). Society of Hair Testing guidelines for drug testing in hair. *Forensic Science International* **218**, 20–24.
- Couture SM, Penn DL, Roberts DL (2006). The functional significance of social cognition in schizophrenia: a review. *Schizophrenia Bulletin* **32**, 44–63.
- Cunha PJ, Bechara A, de Andrade AG, Nicastrì S (2011). Decision-making deficits linked to real-life social dysfunction in crack cocaine-dependent individuals. *American Journal on Addictions* **20**, 78–86.
- Engelmann D, Strobel M (2004). Inequality aversion, efficiency, and maximin preferences in simple distribution experiments. *American Economic Review* **94**, 857–869.
- European Monitoring Centre for Drugs and Drug Addiction (2012). *Annual report 2012: the state of the drugs problem in Europe*. Publications Office of the European Union: Luxembourg.
- Fehr E, Camerer CF (2007). Social neuroeconomics: the neural circuitry of social preferences. *Trends in Cognitive Sciences* **11**, 419–427.
- Fellows LK (2004). The cognitive neuroscience of human decision making: a review and conceptual framework. *Behavioral and Cognitive Neuroscience Reviews* **3**, 159–172.

- Fischbacher U (2007). z-Tree: Zurich toolbox for ready-made economic experiments. *Experimental Economics* **10**, 171–178.
- Fox HC, Bergquist KL, Casey J, Hong KA, Sinha R (2011). Selective cocaine-related difficulties in emotional intelligence: relationship to stress and impulse control. *American Journal on Addictions* **20**, 151–160.
- Haber S (2008). Parallel and integrative processing through the basal ganglia reward circuit: lessons from addiction. *Biological Psychiatry* **64**, 173–174.
- Heil SH, Johnson MW, Higgins ST, Bickel WK (2006). Delay discounting in currently using and currently abstinent cocaine-dependent outpatients and non-drug-using matched controls. *Addictive Behaviors* **31**, 1290–1294.
- Homer BD, Solomon TM, Moeller RW, Mascia A, DeRaleau L, Halkitis PN (2008). Methamphetamine abuse and impairment of social functioning: a review of the underlying neurophysiological causes and behavioral implications. *Psychological Bulletin* **134**, 301–310.
- Hser YI (1997). Self-reported drug use: results of selected empirical investigations of validity. *NIDA Research Monograph* **167**, 320–343.
- Hulka LM, Wagner M, Preller KH, Jenni D, Quednow BB (2013). Blue-yellow colour vision impairment and cognitive deficits in occasional and dependent stimulant users. *International Journal of Neuropsychopharmacology* **16**, 535–547.
- Ivanov I, Schulz KP, London ED, Newcorn JH (2008). Inhibitory control deficits in childhood and risk for substance use disorders: a review. *American Journal of Drug and Alcohol Abuse* **34**, 239–258.
- Johnson MW, Bickel WK, Baker F, Moore BA, Badger GJ, Budney AJ (2010). Delay discounting in current and former marijuana-dependent individuals. *Experimental and Clinical Psychopharmacology* **18**, 99–107.
- Jovanovski D, Erb S, Zakzanis KK (2005). Neurocognitive deficits in cocaine users: a quantitative review of the evidence. *Journal of Clinical and Experimental Neuropsychology* **27**, 189–204.
- Kirby KN, Petry NM (2004). Heroin and cocaine abusers have higher discount rates for delayed rewards than alcoholics or non-drug-using controls. *Addiction* **99**, 461–471.
- Kirby KN, Petry NM, Bickel WK (1999). Heroin addicts have higher discount rates for delayed rewards than non-drug-using controls. *Journal of Experimental Psychology: General* **128**, 78–87.
- Kishida KT, King-Casas B, Montague PR (2010). Neuroeconomic approaches to mental disorders. *Neuron* **67**, 543–554.
- Kjome KL, Lane SD, Schmitz JM, Green C, Ma L, Prasla I, Swann AC, Moeller FG (2010). Relationship between impulsivity and decision making in cocaine dependence. *Psychiatry Research* **178**, 299–304.
- Koob GF (2009). Dynamics of neuronal circuits in addiction: reward, antireward, and emotional memory. *Pharmacopsychiatry* **42**, 32–41.
- Lehrl S (1999). *Mehrfachwahl-Wortschatz-Intelligenztest (MWT-B) (Multiple Choice Vocabulary Intelligence Test)*. Hogrefe: Göttingen.
- Leland DS, Paulus MP (2005). Increased risk-taking decision-making but not altered response to punishment in stimulant-using young adults. *Drug and Alcohol Dependence* **78**, 83–90.
- Lindenmayer JP, McGurk SR, Khan A, Kaushik S, Thanju A, Hoffman L, Valdez G, Wance D, Herrmann E (2013). Improving social cognition in schizophrenia: a pilot intervention combining computerized social cognition training with cognitive remediation. *Schizophrenia Bulletin* **39**, 507–517.
- Lucantonio F, Stalnaker TA, Shaham Y, Niv Y, Schoenbaum G (2012). The impact of orbitofrontal dysfunction on cocaine addiction. *Nature Neuroscience* **15**, 358–366.
- Medalia A, Saperstein AM (2013). Does cognitive remediation for schizophrenia improve functional outcomes? *Current Opinion in Psychiatry* **26**, 151–157.
- Mendez IA, Simon NW, Hart N, Mitchell MR, Nation JR, Wellman PJ, Setlow B (2010). Self-administered cocaine causes long-lasting increases in impulsive choice in a delay discounting task. *Behavioral Neuroscience* **124**, 470–477.
- Mischel W, Ayduk O, Berman MG, Casey BJ, Gotlib IH, Jonides J, Kross E, Teslovich T, Wilson NL, Zayas V, Shoda Y (2011). ‘Willpower’ over the life span: decomposing self-regulation. *Social Cognitive and Affective Neuroscience* **6**, 252–256.
- Moffitt TE, Arseneault L, Belsky D, Dickson N, Hancox RJ, Harrington H, Houts R, Poulton R, Roberts BW, Ross S, Sears MR, Thomson WM, Caspi A (2011). A gradient of childhood self-control predicts health, wealth, and public safety. *Proceedings of the National Academy of Sciences* **108**, 2693–2698.
- Mueser KT, Deavers F, Penn DL, Cassisi J (2013). Psychosocial treatments for schizophrenia. *Annual Review of Clinical Psychology* **9**, 465–497.
- Nutt D, King LA, Saulsbury W, Blakemore C (2007). Development of a rational scale to assess the harm of drugs of potential misuse. *Lancet* **369**, 1047–1053.
- O’Brien CP (2005). Anticraving medications for relapse prevention: a possible new class of psychoactive medications. *American Journal of Psychiatry* **162**, 1423–1431.
- Odum AL (2011). Delay discounting: trait variable? *Behavioural Processes* **87**, 1–9.
- Olausson P, Jentsch JD, Krueger DD, Tronson NC, Nairn AC, Taylor JR (2007). Orbitofrontal cortex and cognitive-motivational impairments in psychostimulant addiction: evidence from experiments in the non-human primate. *Annals of the New York Academy of Sciences* **1121**, 610–638.
- Olesen J, Gustavsson A, Svensson M, Wittchen HU, Jonsson B (2012). The economic cost of brain disorders in Europe. *European Journal of Neurology* **19**, 155–162.
- Pennings EJ, Leccese AP, Wolff FA (2002). Effects of concurrent use of alcohol and cocaine. *Addiction* **97**, 773–783.
- Perez de Los Cobos J, Sinol N, Puerta C, Cantillano V, Lopez Zurita C, Trujols J (2011). Features and prevalence of patients with probable adult attention deficit

- hyperactivity disorder who request treatment for cocaine use disorders. *Psychiatry Research* **185**, 205–210.
- Preller KH, Ingold N, Hulka LM, Vonmoos M, Jenni D, Baumgartner MR, Vollenweider FX, Quednow BB** (2013). Increased sensorimotor gating in recreational and dependent cocaine users is modulated by craving and ADHD symptoms. *Biological Psychiatry* **73**, 225–234.
- Quednow BB, Kuhn KU, Hoenig K, Maier W, Wagner M** (2004). Prepulse inhibition and habituation of acoustic startle response in male MDMA ('ecstasy') users, cannabis users, and healthy controls. *Neuropsychopharmacology* **29**, 982–990.
- Quednow BB, Kuhn KU, Hoppe C, Westheide J, Maier W, Daum I, Wagner M** (2007). Elevated impulsivity and impaired decision-making cognition in heavy users of MDMA ('ecstasy'). *Psychopharmacology* **189**, 517–530.
- Rachlin H, Green L** (1972). Commitment, choice and self-control. *Journal of the Experimental Analysis of Behavior* **17**, 15–22.
- Rahman Q, Clarke CD** (2005). Sex differences in neurocognitive functioning among abstinent recreational cocaine users. *Psychopharmacology (Berlin)* **181**, 374–380.
- Reske M, Delis DC, Paulus MP** (2011). Evidence for subtle verbal fluency deficits in occasional stimulant users: quick to play loose with verbal rules. *Journal of Psychiatry Research* **45**, 361–368.
- Reske M, Eidt CA, Delis DC, Paulus MP** (2010). Nondependent stimulant users of cocaine and prescription amphetamines show verbal learning and memory deficits. *Biological Psychiatry* **68**, 762–769.
- Rilling JK, Sanfey AG** (2011). The neuroscience of social decision-making. *Annual Review of Psychology* **62**, 23–48.
- Rosler M, Retz W, Retz-Junginger P, Thome J, Supprian T, Nissen T, Stieglitz RD, Blocher D, Hengesch G, Trott GE** (2004). Tools for the diagnosis of attention-deficit/hyperactivity disorder in adults. Self-rating behaviour questionnaire and diagnostic checklist. *Nervenarzt* **75**, 888–895.
- Rounsaville BJ** (2004). Treatment of cocaine dependence and depression. *Biological Psychiatry* **56**, 803–809.
- Soar K, Mason C, Potton A, Dawkins L** (2012). Neuropsychological effects associated with recreational cocaine use. *Psychopharmacology (Berlin)* **222**, 633–643.
- Stewart JL, Flagan TM, May AC, Reske M, Simmons AN, Paulus MP** (2013). Young adults at risk for stimulant dependence show reward dysfunction during reinforcement-based decision making. *Biological Psychiatry* **73**, 235–241.
- Sussner BD, Smelson DA, Rodrigues S, Kline A, Losonczy M, Ziedonis D** (2006). The validity and reliability of a brief measure of cocaine craving. *Drug and Alcohol Dependence* **83**, 233–237.
- Takahashi T, Oono H, Inoue T, Boku S, Kako Y, Kitaichi Y, Kusumi I, Masui T, Nakagawa S, Suzuki K, Tanaka T, Koyama T, Radford MH** (2008). Depressive patients are more impulsive and inconsistent in intertemporal choice behavior for monetary gain and loss than healthy subjects – an analysis based on Tsallis' statistics. *Neuro Endocrinology Letters* **29**, 351–358.
- Tiffany ST, Singleton E, Haertzen CA, Henningfield JE** (1993). The development of a cocaine craving questionnaire. *Drug and Alcohol Dependence* **34**, 19–28.
- United Nations Office on Drugs and Crime** (2011). *World Drug Report 2011*. United Nations Office on Drugs and Crime: Vienna, Austria.
- Vadhan NP, Hart CL, Haney M, van Gorp WG, Foltin RW** (2009). Decision-making in long-term cocaine users: effects of a cash monetary contingency on Gambling task performance. *Drug and Alcohol Dependence* **102**, 95–101.
- Verdejo-Garcia A, Benbrook A, Funderburk F, David P, Cadet JL, Bolla KI** (2007a). The differential relationship between cocaine use and marijuana use on decision-making performance over repeat testing with the Iowa Gambling Task. *Drug and Alcohol Dependence* **90**, 2–11.
- Verdejo-Garcia AJ, Perales JC, Perez-Garcia M** (2007b). Cognitive impulsivity in cocaine and heroin polysubstance abusers. *Addictive Behaviors* **32**, 950–966.
- Vonmoos M, Hulka LM, Preller KH, Jenni D, Baumgartner MR, Stohler R, Bolla KI, Quednow BB** (2013). Cognitive dysfunctions in recreational and dependent cocaine users: the role of ADHD, craving, and early age of onset. *British Journal of Psychiatry*. **203**, 35–43.
- Wagner FA, Anthony JC** (2007). Male–female differences in the risk of progression from first use to dependence upon cannabis, cocaine, and alcohol. *Drug and Alcohol Dependence* **86**, 191–198.
- Washio Y, Higgins ST, Heil SH, McKerchar TL, Badger GJ, Skelly JM, Dantona RL** (2011). Delay discounting is associated with treatment response among cocaine-dependent outpatients. *Experimental and Clinical Psychopharmacology* **19**, 243–248.
- Wittchen HU, Wunderlich U, Gruschwitz S, Zaudig M** (1997a). *Strukturiertes Klinisches Interview für DSM-IV. Achse I: Psychische Störungen (Structured Clinical Interview for DSM-IV. Axis I: Mental Disorders)*. Hogrefe: Göttingen.
- Wittchen HU, Wunderlich U, Gruschwitz S, Zaudig M** (1997b). *Strukturiertes Klinisches Interview für DSM-IV. Achse II: Persönlichkeitsstörungen (Structured Clinical Interview for DSM-IV. Axis II: Personality Disorders)*. Hogrefe: Göttingen.