

**Disclosure:** The results of this study were from an educational programme that was developed through independent educational funding from Janssen Neuroscience.

**Keywords:** Suicide; major depressive disorder; Psychiatric emergencies; MDD

## Schizophrenia and other Psychotic Disorders 2

### O0131

#### Clinical Indicators of Symptom Dimensions and Cognitive Ability in Schizophrenia

L. Farakish<sup>1\*</sup>, S. Legge<sup>1</sup>, M. Owen<sup>1</sup>, M. O'Donovan<sup>1</sup>, J. Walters<sup>1</sup> and A. Cardno<sup>2</sup>

<sup>1</sup>Cardiff University, Mrc Centre For Neuropsychiatric Genetics And Genomics, Division Of Psychological Medicine And Clinical Neurosciences, School Of Medicine, Cardiff, United Kingdom and <sup>2</sup>University of Leeds, Division Of Psychological And Social Medicine, Leeds Institute Of Health Sciences, Faculty Of Medicine And Health, University Of Leeds, Leeds, Uk, Leeds, United Kingdom

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.317

**Introduction:** Schizophrenia is a heterogeneous disorder and it is unknown what causes individual variability in symptoms and cognitive ability.

**Objectives:** To examine the association between nine clinical predictors measurable at the onset of schizophrenia and five phenotype dimensions: positive, negative (diminished expressivity), negative (motivation and pleasure), disorganised symptoms and cognitive ability.

**Methods:** 852 participants (mean age 49 years old) with a diagnosis of schizophrenia or schizoaffective depression were included from the CardiffCOGS cross-sectional sample. Phenotype dimensions were created using confirmatory factor analysis and a 5-factor model. Associations were tested using linear regression, adjusting for age and sex. A Bonferroni correction was applied for ( $p < 1.1 \times 10^{-3}$ ) for multiple testing.

**Results:** Age of onset of psychosis was significantly associated with positive symptoms ( $\beta = -0.18$ ,  $p = 4.0 \times 10^{-6}$ ). Lower premorbid IQ was associated with diminished expressivity ( $\beta = -0.25$ ,  $p = 7.0 \times 10^{-13}$ ), reduced motivation and pleasure ( $\beta = -0.23$ ,  $p = 4.3 \times 10^{-11}$ ), disorganised symptoms ( $\beta = -0.14$ ,  $p = 7.6 \times 10^{-5}$ ) and reduced cognition ( $\beta = 0.54$ ,  $p = 4.8 \times 10^{-77}$ ). Poor premorbid social adjustment held associations with all except positive. Developmental delay was associated with reduced cognition ( $\beta = -0.35$ ,  $p = 4.3 \times 10^{-5}$ ). Cannabis use (year before onset), psychosocial stressors (within 6 months), childhood abuse and family history of schizophrenia held no associations.

**Conclusions:** Clinical indicators measurable at schizophrenia onset are associated with lifetime symptom variability. A younger psychosis onset is associated with more severe positive symptoms, suggesting possible age-targeted management. Pre-established links of lower premorbid IQ with poor premorbid social adjustment and negative symptom severity with cognition are strengthened. Further investigation could potentially improve diagnosis and guide treatment choice for aspects of schizophrenia with poor outcomes.

**Disclosure:** No significant relationships.

**Keywords:** Clinical indicators; Phenotype dimensions; Aetiology; schizophrenia

### O0132

#### Patient violence towards their family carers: A qualitative exploration of carers' experiences in psychosis

E. Wildman<sup>1\*</sup>, E. Kuipers<sup>1,2</sup>, D. Macmanus<sup>3</sup> and J. Onwumere<sup>1,2</sup>

<sup>1</sup>King's College London, Institute of Psychiatry, Psychology, and Neuroscience, Department Of Psychology, London, United Kingdom;

<sup>2</sup>Bethlem Royal Hospital, South London And Maudsley Nhs Foundation Trust, London, United Kingdom and <sup>3</sup>King's College London, Department Of Forensic And Neurodevelopmental Science, London, United Kingdom

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.318

**Introduction:** Compared to the general population, people living with schizophrenia spectrum disorders (SSD) are more likely to perpetrate acts of violence. When this happens, family members (informal carers) are most commonly the victims. However, family violence by people with SSD is often a taboo topic and largely neglected within public discourse, research, and clinical domains. Consequently, our understanding of families' experiences and support needs are limited.

**Objectives:** To develop a detailed understanding of the subjective experiences, and impact, of patient violence towards family carers.

**Methods:** Individual semi-structured interviews were held with family carers of adults with SSD and a history of violence perpetration towards their family carer. Interview data were subject to thematic analyses using NVivo software.

**Results:** Twenty-one UK based carers that were predominately White British (90%) and female (81%) were interviewed. Key themes highlight the range of physical and mental injuries endured by carers following patient violence, and speak to carers' experiences of suffering, living in a constant state of hypervigilance, as well as social isolation in the context of shame, stigma, and an absence of professional and informal support.

**Conclusions:** Family violence by people living with SSD can and does happen. Yet, too often, carers are left with no option but to continue supporting their relative in the absence of support, even in contexts where this compromises their own safety. The devastating impact of violence is far-reaching, across all levels of the family-system. The findings highlight the danger of neglecting family violence by people with SSD in research and clinical fields.

**Disclosure:** No significant relationships.

**Keywords:** Carers; violence; aggression; Psychosis

### O0133

#### A systematic review and meta-analysis of implicit Theory of Mind in schizophrenia

T. Csulak<sup>1\*</sup>, A. Hajnal<sup>1</sup>, S. Kiss<sup>2</sup>, F. Dembrowszky<sup>2</sup>, Z. Sipos<sup>2</sup>, M. Varjú-Solymár<sup>2</sup>, M. Kovács<sup>1</sup>, M. Herold<sup>1</sup>, E. Varga<sup>3</sup>, P. Hegyi<sup>2</sup>, T. Tényi<sup>1</sup> and R. Herold<sup>1</sup>

<sup>1</sup>University of Pécs, Department Of Psychiatry And Psychotherapy, Pécs, Hungary; <sup>2</sup>University of Pécs, Institute For Translational Medicine, Pécs, Hungary and <sup>3</sup>University of Pécs, Department Of Paediatrics, Pécs, Hungary

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.319