

# Biodiversity pattern of subtidal sponges (Porifera: Demospongiae) in the Penghu Archipelago (Pescadores), Taiwan

YUSHENG M. HUANG<sup>1,2,3</sup>, NICOLE J. DE VOOGD<sup>3</sup>, DANIEL F. R. CLEARY<sup>4</sup>, TSUNG-HSUAN LI<sup>5</sup>, HIN-KIU MOK<sup>5</sup> AND JINN-PYNG UENG<sup>2</sup>

<sup>1</sup>Department of Marine Sports and Recreation, National Penghu University of Science and Technology, Magong City, Taiwan,

<sup>2</sup>Department of Aquaculture, National Penghu University of Science and Technology, Magong City, Taiwan, <sup>3</sup>Naturalis Biodiversity Center, Leiden, the Netherlands, <sup>4</sup>Departamento de Biologia, University of Aveiro, Aveiro, Portugal, <sup>5</sup>Department of Oceanography, National Sun Yat-Sen University, Kaohsiung, Taiwan

*Sponge-related research in Taiwan has primarily focused on natural product exploration. This research has, however, been hampered by a lack of fundamental work on sponge taxonomy and ecology. In the present study, subtidal sponges were photo-recorded in situ and collected by scuba diving at a depth range of 2–20 m from 2009 to 2012 in 16 different sites surrounding the Penghu Archipelago, Taiwan. Sponge samples were identified to the lowest taxonomic level based on skeletal morphology and spicules. A total of 53 species belonging to 24 families and 10 orders were identified in this study. The number of sponge species per site ranged from 0 to 24. The most widely distributed sponge species was *Callyspongia (Euoplacella) cf. communis* (Carter, 1881) followed by *Haliclona (Gellius) cymaeformis* (Esper, 1794), and *Aaptos suberitoides* (Brøndsted, 1934). At one location, Chipeiyu, no sponges were observed. Non-metric multidimensional scaling (NMDS) ordination revealed relatively low similarity among most sampling sites. Large- and small-scale hydrological and habitat features are probably responsible for compositional variation of sponge assemblages among groups of sampling sites. Our richness analyses suggest that many more sponge species remain to be discovered in the Penghu Archipelago.*

**Keywords:** biodiversity, sponges, Penghu Archipelago, distribution, Demospongia

Submitted 3 November 2014; accepted 6 February 2015; first published online 5 March 2015

## INTRODUCTION

Sponges inhabit numerous marine ecosystems from polar seas to temperate and tropical waters and also thrive over a large depth range (van Soest, 1993; Fromont, 1999; Diaz & Rützler, 2001; Zea, 2002; Bell & Smith, 2004; de Voogd *et al.*, 2004; McClintock *et al.*, 2005; Wulff, 2006). Sponges have demonstrated an impressive ability to adapt and survive in marginal habitats (McClintock *et al.*, 2005; Roberts *et al.*, 2006; Yahel *et al.*, 2007; Lesser *et al.*, 2009; Olson & Kellogg, 2010). The ecological roles and biotechnological potential of marine sponges have been broadly reported in the past few decades. As benthic filter feeders, sponges play important and versatile roles including reef degraders and builders, links in benthic-pelagic nutrient coupling (Lesser, 2006), and chemical and physical media for sympatric organisms (Pawlik *et al.*, 1995; Huang *et al.*, 2008). A global assessment of sponge biodiversity was recently facilitated by the advent of the World Porifera Database (WPD) (van Soest *et al.*, 2015) which provides a single unified classification, the Systema Porifera (SP) (Hooper & van Soest, 2002). Specific areas with high sponge diversity (>201

species) such as the Mediterranean, north-eastern Atlantic, Caribbean, southern India, eastern Australia and subtropical Japan are at least partially due to the intensity of collection in these areas (van Soest *et al.*, 2012). The low numbers of recorded species from other areas such as Taiwan (61 species) can be attributed to the low level of sampling intensity. The number of taxonomic and biodiversity studies of marine sponges in the vicinity of Taiwan has substantially increased in the last few decades, including South Korea (Kang *et al.*, 2013; Shim & Sim, 2013), Singapore (Lim *et al.*, 2009; Lim *et al.*, 2012), Japan (Hoshino, 1980; Khodakovskaya, 2005) and Indonesia (Bell & Smith, 2004; Becking *et al.*, 2006; Cleary & de Voogd, 2007; de Voogd & Cleary, 2008; de Voogd, 2012). In Taiwan, however, much remains to be done to document the Taiwanese sponge fauna.

Taiwan is located at the centre of the Philippine–Japan island arc and lies within the Tropic of Cancer. Southern fringing coral reefs and northern rocky shore habitats dominate the coastline, except for western Taiwan. Marine climate in the winter (November to February) is significantly influenced by the combination of the north-eastern monsoon and Chinese Coastal Current (CCC), Kuroshio Current (KC) and South China Sea Current (SCSC). Tropical typhoons occasionally provoke regional disturbances in summer between July and October. Annual variation of mean sea surface temperature and salinity ranges from 21 to 28°C and 33.9 to 34.6 ‰, respectively (Jan *et al.*, 2010; and references

**Corresponding author:**

Y.M. Huang

Email: yusheng@gms.npu.edu.tw

therein). The first scientifically documented sponge specimen in Taiwan can be traced back to the late 19th century when the species, *Theonella swinhoei*, was first described; its holotype is presently preserved in the British Museum (Gray, 1868). Three newly described sponge species, *Chalinula amoyensis*, *Chondrocladia (Chondrocladia) arenifera* and *Echinodictyum axinelloides*, and one new record species, *Lissodendoryx (Lissodendoryx) ternatensis*, from the Taiwan Strait were first documented in the early 20th century (Brøndsted, 1929). There has, however, been a marked dearth of information on the ecology and distribution of sponges in Taiwan since then. The majority of sponge studies in Taiwan have focused on the natural products sequestered in marine sponges for medical and pharmaceutical applications (Lin, 1995; Lo, 1999; Chen, 2000; Huang, 2002; Liao, 2003; Jhou, 2004; Su *et al.*, 2011). However, little was done to identify the sponges. The majority of the studied sponge specimens were only identified to the genus level. Other studies have focused on the ecology and reproduction of the intertidal bath sponge, *Spongia ceylonensis* (Chung *et al.*, 2010), *Cinachyrella australiensis* (Chen, 1988; Huang, 1996; Chen *et al.*, 1997; Lu, 2003) and on the symbiotic microbial community associated with the Cyanobacteriosponge, *Terpios hoshinota*, which caused the 'black disease' of reef-building corals at Green Island, Taiwan (Liao *et al.*, 2007). In 2007, a preliminary survey on the sponge assemblages in southern Taiwan, including Green Island and Kenting, was conducted as part of a sponge taxonomy workshop and the results were published as a checklist with 65 sponge species (Shao *et al.*, 2008). The first morphological taxonomic study and preliminary survey on the biodiversity of sponges in a specific region in Taiwan included 28 sponge species belonging to seven orders, including 11 species that are newly recorded species for Taiwan (Li, 2013).

The objectives of the present study were to: (1) investigate the biodiversity and distribution of sponges in the Penghu Archipelago; (2) compare sponge assemblages in different locations of the Penghu Archipelago; (3) estimate the potential species richness of the subtidal sponge fauna in the Penghu Archipelago, Taiwan.

## MATERIALS AND METHODS

### Study area

The Pescadores, also known as the Penghu Islands, is an archipelago off the western coast of Taiwan in the Taiwan Strait. The archipelago consists of approximately 90 small islands and islets covering an area of 128 square kilometres. With the exception of the most western islet, HuaYu, which consists of andesite, the islands mainly consist of oceanic basalt that resulted from the slow eruption and discharge of magma from submarine volcanic crevasses 8–16 million years ago (Chen & Liu, 1996 and references therein). Tropical fringing reefs have formed extensively around most parts of the islands. The hydrology around the islands is unique. There are three major currents, the cold, southward CCC, a branch of warm, northward KC and the SCSC (Jan *et al.*, 2010; Tzeng *et al.*, 2012). The anfractuous coastline of Penghu, the largest island, and the convergence of three different currents (CCC, KC and SCSC) results in a complex

hydrological condition, which also influences seawater temperature, salinity, turbidity, dynamics and habitat stability.

### Sponge collection and taxonomy

Subtidal sponges were sampled during 60-min dives using the Roving Diver Technique (Munro, 2005) from 16 sites off the coast of the Penghu Archipelago (Fig. 1 and Table 1). Sponges (cryptic and boring specimens were excluded) were collected using scuba diving between July 2009 and July 2012, photographed *in situ*, and brought back to the laboratory for further identification based on the keys provided in Systema Porifera (Hooper & van Soest, 2002) and the World Porifera Database (van Soest *et al.*, 2015). Identifications were made to the lowest taxonomic level possible. Sponge vouchers are incorporated in the sponge collection of Naturalis Biodiversity Center, Leiden, the Netherlands.

### Statistical analysis

Species number ( $S$ ), richness ( $d$ ) and biodiversity index ( $H'$ ) of sponge assemblage data (presence/absence) were obtained from each sampling site using PRIMER-E (Plymouth Marine Laboratory, Plymouth, UK) (Clarke & Warwick, 2001a). Zero and outlier data from ChiPeiYu (CPY) and Hujing Bunker North (HBN) were excluded in the following analyses. We included diversity measures that take into account the relationships between species following a Linnaean classification, average taxonomic distinctness (AvTD,  $\Delta^+$ ) and variation in taxonomic distinctness (VarTD,  $\Lambda^+$ ) were calculated. These analyses are an indicative measure of the taxonomic distance of an assemblage and the relatedness of its constituent species based on presence/absence data (Clarke & Warwick, 2001b). Sponge similarity among sites was compared by first transforming the presence/absence matrix into a triangular similarity matrix based on the Bray–Curtis similarity index (Clarke & Warwick, 2001a). Variation in similarity was subsequently visually compared using non-metric multidimensional scaling (NMDS). The stress coefficient of Kruskal was used to test the ordination significance (McCune & Grace, 2002). Species richness in the Penghu Archipelago was estimated using EstimateS with different models (Standard, Uniques, Chao 2 and Jack 1) (Colwell, 2013).

## RESULTS

### Sponge distribution, biodiversity and taxonomic distinctness

Fifty-three subtidal sponge species belonging to 10 orders, 24 families and 34 genera were identified during this study (Table 2). The number of sponge species ( $S$ ) was the highest in Chingwan and Suogang (24) and lowest in Hujing bunker north (1) (Table 2). The most widely distributed sponge species was *Callyspongia (Euplaccella) cf. communis* (Carter, 1881) (11 sites) followed by *Haliclona (Gellius) cymaeformis* (Esper, 1794) (10 sites) and *Aaptos suberitoides* (Brøndsted, 1934) (8 sites). *Agelas nemoechinata* Hoshinota, 1985 was the most abundant species in both Dongyuping (DYP) and Siyuping (SYP). Only one sponge species, *Tetilla* sp.,




Fig. 1. Sampling locations of subtidal sponges off the Penghu Archipelago, Taiwan. Sixteen sampling sites across islands are indicated by site abbreviations given in Table 1.

growing in sandy sediment was observed in Hujing bunker north. Surprisingly, no sponges were observed in Chiipeiyu. Average taxonomic distinctness among sites varied markedly (Table 2, Fig. 2A). The average taxonomic distinctness value was highest in Wanan west (WAW, 98.21) and lowest in Sanshuie beach west (SSBW, 80.00). The AvTD values of seven sites were under the simulated average and those of most sites; only Shili beach west (SLBW) and SSBW fell below the 95% simulated range. Variation in taxonomic distinctness among sites also differed markedly (Table 2, Fig. 2B). VarTD was highest in Siyuping (SYP, 656.3) and lowest in Wanan west (WAW, 41.5). VarTD values of eight sites were higher than the simulated average. Two-dimensional plots of 95% probability ellipses from the simulated distribution (for sampling subsets  $M = 10-50$ ), with observed values of the AvTD and VarTD pairs superimposed within probability ellipses (Fig. 3), show that all faunas lie within the  $M = 10$  simulation envelope, except for SSBW, SYP, DYP and Hujing harbour west (HHW). The observed values of AvTD and VarTD for these 14 sites were quite strongly negatively linear-related with a product-moment correlation coefficient of  $r = -0.67$  ( $P = 0.009$ ) (Fig. 3).

### Comparison of sponge assemblages in different locations

Results of the Bray–Curtis cluster analysis revealed high similarities of sponge assemblages collected from islands and

locations close to one other (Fig. 4A). The sponge assemblages of the southernmost sampling sites, Dongyuping (DYP) and Siyuping (SYP), were similar (>60%), but dissimilar from those of the northern sites (<20%). Sponge assemblages of Hujing harbour east (HHE) and west (HHW) had the highest similarity (75%). Fongguie (FG), Cingwan (CW) and Suogang (SG), located along the southern coastline of the Penghu Island and within 10 km of one another, were also similar (>60%). Four groups with similarity >40% were distinguished based on the presence/absence data matrix of sponge assemblages from different sites: (Group 1) the northernmost site Kopoyu (KPY) and Wukan (WK); (Group 2) FG, CW, SG, Sanshuie cliff (SSC) and Wanan north (WAN); (Group 3) all sampling sites in Hujungyu (HHE and HHW), Shili beach west (SLBW) and Sanshuie beach west (SSBW) and (Group 4) the southernmost islands (DYP and SYP). Variation in similarity among different sites and groups were also revealed in MDS plot (Fig. 4B; stress = 0.11).

### Estimation of sponge biodiversity

The results of the observed and estimated accumulative number of subtidal sponge species as a function of sampling sites is shown in Fig. 5. The total number of sponge species recorded in the present study was 53 after surveying 16 different sites across a range of 60 km. The standard rarefaction estimate appeared to reach an asymptote at 105 species with a sampling effort of more than 100 sites. Estimations of Chao 2 and Jack 1 models reached 110 and 78 species, respectively.

Table 1. Localities visited during the course of the present study in the Penghu Archipelago, Taiwan from 2009–2012.

Location	Site coding	Coordinates	Depth (m)	Habitat
Dongyuping, south-west	DYP	23°14'53.14"N 119°30'40.08"E	11	Coral reefs, coarse coral sand
Siyuping, east	SYP	23°16'9.88"N 119°30'39.62"E	10	Coral reefs, coarse coral sand and basalt boulder
Wanan, west	WAW	23°23'21.56"N 119°29'27.02"E	14	Coral reefs and basalt boulder
Wanan, north	WAN	23°24'13.19"N 119°29'33.40"E	20	Coral reefs and basalt boulder
Hujingyu, west of the harbour	HHW	23°29'19.07"N 119°30'48.73"E	12	<i>Acropora</i> spp. coral reefs and sand
Hujingyu, east of the harbour	HHE	23°29'19.35"N 119°30'58.03"E	12	<i>Acropora</i> spp. coral reefs and sand
Hujingyu, north of the bunker	HBN	23°29'46.14"N 119°31'46.90"E	7	Sandy bottom
Fongguei	FG	23°32'15.34"N 119°32'39.23"E	6	Coral reefs and basalt boulder
Chingwan	CW	23°31'45.23"N 119°33'14.07"E	9	Basalt boulders and <i>Acropora</i> spp.
Shihli beach, west	SLBW	23°31'33.51"N 119°33'46.16"E	8	Basalt boulders and sandy bottom
Shanshuei beach, west	SSBW	23°30'39.75"N 119°35'17.27"E	7	Basalt boulders and sandy bottom
Shanshuei, cliff	SSC	23°30'31.05"N 119°36'30.73"E	13	Basalt boulders and coral reefs
Suogang	SG	23°31'08.68"N 119°36'41.79"E	15	Basalt boulders, grooves, and sand
Wukan	WK	23°32'40.41"N 119°37'57.76"E	16	Desalination plant underwater discharge pipe, basalt boulders and sandy bottom
Kopoyu, south	KPY	23°42'45.70"N 119°33'08.60"E	22	Huge basalt boulders on sandy bottom
Chipeiyu, south-west	CHI	23°44'12.78"N 119°35'47.29"E	11	Dead coral debris and, <i>Peysommelia</i> spp.

## DISCUSSION

## Sponge distribution, biodiversity and taxonomic distinctness

Studies of sponge taxonomy and biodiversity in Taiwan have been largely neglected over the past decades. This is the first study on the distribution and biodiversity of sponge in the Penghu Islands, Taiwan. In the present study 16 different sites were surveyed within an area of approximately 150 km<sup>2</sup>. Three sponges, *Callyspongia* (*Euplaccella*) cf. *communis* (Carter, 1881), *Haliclona* (*Gellius*) *cymaeformis* (Esper, 1794) and *Aptos suberitoides* (Brøndsted, 1934), were the most commonly found species (present at 11, 10 and 8 of 16 sites, respectively). The global distribution pattern of *C. communis* was mainly in the Indo-Pacific Ocean including Australia, New Zealand and Singapore (van Soest *et al.*, 2015); however, the studies about the natural history of *C. communis* are scarce. In contrast, the anti-fouling activity, symbiotic microorganisms and the function of secondary metabolites of *H. cymaeformis* and *A. suberitoides* have been studied (Pile *et al.*, 2003; Dobretsov *et al.*, 2005a, b; Tsukamoto *et al.*, 2010). The undescribed *Tetilla* sp. was only observed during the summer in Hujing Bunker North, where the substrate is a sandy bottom. The Indo-Pacific giant barrel sponges, *Xestospongia testudinaria* (Lamarck, 1815), were found in relatively high population densities in both Fongguei (7.6 sponges 100 m<sup>-2</sup>) and Chingwan (3.3 sponges 100 m<sup>-2</sup>) (Huang *et al.*, unpublished data). The coral-killing cyanobacteriosponge, *Terpios hoshinota*, was not observed in the Penghu Islands. This sponge species was very abundant at Green Island (eastern Taiwan), where there are occasional outbreaks (Wang *et al.*, 2012). Among the 53 sponge species recorded in the present study, 34 species (>60%) were newly recorded species, which is substantially higher than previous studies of sponge biodiversity (Bell & Barnes, 2000; van Soest *et al.*, 2007; de Voogd & Cleary, 2008). The AvTD of half of the analysed sampling sites fell below the mean estimated value, which is likely due to low sampling effort and habitat heterogeneity. The habitats of Shihli beach west (SLBW) and Shanshuei beach west (SSBW) were similar with sandy bottom and scattered cobbles and boulders eroded from the adjacent basalt columns. The highest AvTD value, but with the lowest VarTD, of WAW resulted from eight collected species representing eight different orders, respectively. The AvTD value of SSBW was the lowest because four of the five collected species were in the same order, Haplosclerida, and three of them were in the same family, Chalinidae. Most VarTDs of sampling sites were above the estimated mean; the VarTD of Siyuping (SYP) was even higher than the upper 95% limit. These results were likely due to the fact that there were several different orders represented only by a single species, but also with some species-rich genera (Clarke & Warwick, 2001b).

## Sponge assemblages in different sampling locations

Numerous sponge species have been shown to have a limited dispersal ability (Zea, 2002). Although few cosmopolitan sponge species have been studied, molecular evidence has tended to show that some of these 'cosmopolitan' species


**Table 2.** Records (presence/absence) of sponge species collected in the Penghu Archipelago, Taiwan at 15 localities from 2009–2012. Asterisks (\*) indicate new record species in the present study and Taiwan area.

Sponge taxa	Sampling sites														
	DYP	SYP	WAW	WAN	HHW	HHE	HBN	FG	CW	SLBW	SSBW	SSC	SG	WK	KPY
Spirophorida, Tetillidae															
<i>Cinachyrella australiensis</i> (Carter, 1886)	1												1		
<i>Tetilla</i> sp.*							1								
Astrophorida, Ancorinidae															
<i>Jaspis splendens</i> (de Laubenfels, 1954)			1						1				1		
<i>Rhabdastrella globostellata</i> (Carter, 1883)									1						
Astrophorida, Geodiidae															
<i>Erylus proximus</i> Dendy, 1916*												1			
Hadromerida, Clionaidae															
<i>Spheciospongia</i> sp.				1				1	1			1	1	1	
Hadromerida, Suberitidae															
<i>Aaptos suberitoides</i> (Brøndsted, 1934)	1	1		1				1	1			1	1	1	
Chondrosida, Chondrillidae															
<i>Chondrosia</i> aff. <i>corticata</i> Thiele, 1900*									1						
Poecilosclerida, Microcionidae															
<i>Clathria (Thalysias) reinwardti</i> Vosmaer, 1880			1					1	1			1	1		
<i>Clathria</i> sp. 1 'red, encrusting'									1			1			
<i>Clathria</i> sp. 2 'yellow encrusting'										1	1	1	1		
Poecilosclerida, Raspailiidae															
<i>Echinodictyum asperum</i> Ridley & Dendy, 1886*									1						
Poecilosclerida, Mycalina															1
Poecilosclerida, Mycalidae															
<i>Mycale (Zygomycale) parishii</i> Bowerbank, 1875*			1	1	1	1		1	1				1		
Halichondrida, Dictyonellidae															
<i>Acanthella cavernosa</i> Dendy, 1922		1	1					1	1				1	1	1
<i>Stylissa carteri</i> (Dendy, 1889)	1	1						1				1	1		
Halichondrida, Halichondriidae															
<i>Ciocalyptra</i> aff. <i>tyleri</i> Bowerbank, 1873*				1				1	1			1	1		1
<i>Axinyssa</i> sp.												1			
<i>Halichondria</i> sp.*													1		
Agelasida, Agelasiidae															
<i>Agelas cavernosa</i> Thiele, 1903*	1	1							1				1		
<i>Agelas nemoechinata</i> Hoshino, 1985	1	1											1		
Haplosclerida, Callyspongiidae															
<i>Callyspongia</i> cf. <i>communis</i> (Carter, 1881)*			1	1	1	1		1	1	1	1	1	1		1
<i>Callyspongia</i> sp. 'erected, fuzzy'								1	1				1		
<i>Callyspongia</i> sp. 'grey, repent'								1							
<i>Callyspongia</i> sp. 'light purple'									1						
<i>Callyspongia</i> sp.*				1											
<i>Callyspongia (Callyspongia)</i> sp.*								1	1						
Haplosclerida, Chalinidae															

Continued

Table 2. Continued


Sponge taxa	Sampling sites														
	DYP	SYP	WAW	WAN	HHW	HHE	HBN	FG	CW	SLBW	SSBW	SSC	SG	WK	KPY
<i>Cladocroce</i> sp.*								1		1	1		1		
<i>Haliclona cymaiformis</i> (Esper, 1794)*					1	1		1	1	1	1	1	1	1	1
<i>Haliclona</i> ( <i>Soestella</i> ) sp. 'purple'*				1											
<i>Haliclona</i> sp. 'light pink'*				1											
<i>Haliclona</i> sp. 'purple'*												1	1		
<i>Haliclona</i> sp. 'yellow, fine tube'*															1
<i>Haliclona</i> sp. 'orange, massive'*											1				
Haplosclerida, Niphatidae															
<i>Amphimedon</i> sp. 1 'yellow green'*										1					
<i>Amphimedon</i> sp. 2 'purplish'*										1					
<i>Dasychalina</i> aff. <i>fragilis</i> Ridley & Dendy, 1886				1				1	1			1	1		
<i>Niphates</i> sp.*				1											
Haplosclerida, Phloeodictyidae															
<i>Oceanapia</i> sp.*									1						
<i>Oceanapia</i> sp. 'brown tube'*					1			1							
Haplosclerida, Petrosiidae															
<i>Petrosia</i> ( <i>Petrosia</i> ) sp.												1	1		
<i>Xestospongia testudinaria</i> (Lamarck, 1815)			1	1				1	1				1	1	1
<i>Xestospongia vansoesti</i> Bakus & Nishiyama, 2000													1		
Dictyoceratida, Irciniidae															
<i>Ircinia ramosa</i> (Keller, 1889)*								1							
Dictyoceratida, Thorectidae															
<i>Hyrtios erectus</i> (Keller, 1889)	1														
<i>Hyrtios</i> sp.	1														
Dictyoceratida, Spongiidae															
<i>Spongia</i> sp.*													1		
Dictyoceratida, Dysideidae															
<i>Dysidea</i> aff. <i>frondosa</i> Berquist, 1995*						1		1	1					1	
<i>Dysidea</i> sp.									1			1		1	
<i>Euryspongia delicatula</i> Berquist, 1995*			1						1	1			1		
Verongida, Ianthellidae															
<i>Anomoianthella</i> sp.*															1
Verongida, Aplysinellidae															
<i>Aplysinella</i> sp.*			1												
<i>Suberea</i> sp.	1														
Species no. ( <i>S</i> )	8	5	8	11	4	4	1	18	24	7	5	16	24	7	8
Species richness ( <i>d</i> )	3.4	2.5	3.4	4.2	2.2	2.2	–	5.9	7.2	3.1	2.5	5.4	7.2	3.1	3.4
Diversity index ( <i>H'</i> )	2.1	1.6	2.1	2.4	1.4	1.4	–	2.9	3.2	1.9	1.6	2.8	3.2	1.9	2.1
Average Taxonomic Distinctness (AvTD, $\Delta^+$ )	94.6	87.5	98.2	87.7	87.5	95.8	–	90.9	93.8	83.3	80.0	94.4	94.3	94.1	92.0
Variation in Taxonomic Distinctness (VarTD, $\Lambda^+$ )	373	656	42	315	156	87	–	308	243	437	475	244	201	292	270




**Fig. 2.** Probability funnels (95% confidence interval) of (A) average taxonomic distinctness (AvTD, Δ<sup>+</sup>) and (B) variation in taxonomic distinctness (VarTD, Δ<sup>+</sup>) values. The central dotted line indicates the mean while solid lines constitute the 95% limits of the simulated values of 1000 random selections from the master list of 53 species for each sub-list ( $M = 10-50$ ). The solid dots represent the AvTD and VarTD values for each sampling site plotted against the number of species. (Site abbreviations given in Table 1.)

were in fact cryptic species complexes (Lazoski *et al.*, 2001; Xavier *et al.*, 2010; Swierts *et al.*, 2013). Our results showed that sponge assemblages recorded from different locations within a <60 km range were significantly dissimilar. Substrates, hydrological conditions, predation, nutrition

supplement, salinity and seawater temperature, and anthropological activities have been shown to influence sponge distributions (van Soest, 1993; Wulff, 2001; Bell & Smith, 2004; de Voogd *et al.*, 2004; Roberts *et al.*, 2006). The hydrological conditions near Penghu Islands are also significantly influenced by the cold China Coastal Current (CCC), a warm branch of the Kuroshio Current (KC) and the South China Sea Current (SCSC) (Jan *et al.*, 2002; Hu *et al.*, 2010; Tzeng *et al.*, 2012). Oceanic current patterns in the vicinity of Taiwan were shown to influence the geographic distribution of fiddler crabs (Shih, 2012), corals (Chen & Shashank, 2009) and intertidal barnacles (Chan & Lee, 2012). In addition to this, the combination of currents responsible for seawater surface temperature (SST) fluctuations to the north and south of the Penghu islands varies considerably across seasons (Jan *et al.*, 2002; Qiu *et al.*, 2011). Due to the combination of the north-eastern monsoon and CCC, SST around the northern islands of the Penghu Archipelago can drop to 11 °C and cause catastrophic mortality of coral reef fishes during the winter (Hsieh *et al.*, 2008). Many studies have focused on the effects of seawater warming on sponges (Webster *et al.*, 2008; Massaro, 2009; Duckworth & Peterson, 2012; Duckworth *et al.*, 2012; Kelmo *et al.*, 2013), but little is known about the effect of cold SST on the growth, reproduction and distribution of sponges (Maldonado & Young, 1998). Abnormally low SST could adversely affect sponge species survival,



**Fig. 3.** Scatter plot of average taxonomic distinctness (AvTD, Δ<sup>+</sup>) and variation in taxonomic distinctness (VarTD, Δ<sup>+</sup>) values for each sites (zero and outliers excluded). Solid ellipses indicate the 95% probabilities of simulated data from each sublist ( $M = 10-50$ ). Number of species in each site is shown between parentheses. (Site abbreviations given in Table 1.)



**Fig. 4.** (A) Bray–Curtis cluster analysis dendrogram generated from presence/absence data of subtidal sponge species collected from 14 sampling sites (zero and outliers excluded). The per cent similarity between sites is shown. (B) NMDS ordination analysis of 14 sampling sites based on sub-tidal sponge species collected. The NMDS analysis was conducted using presence/absence data and presented as a 2-dimensional scatter plot using *x* and *y* variables (Stress = 0.11). Area of shaded circles indicates species richness (*d*). (Site abbreviations given in Table 1.)

particularly for those species sensitive to temperature fluctuations and thus structure sponge assemblages. Current directions and flooding also change dramatically during the diurnal cycle. The neritic tidal current switches directions south-northward in the vicinity of the southern Penghu Islands (Wang, 2012) and east-westward along the coastline of the southern mainland Penghu Island (per. obs.). Although efficient swimming larvae with some cue response (phototaxis and/or chemotaxis) can effectively act against hydrodynamic forces in determining small-scale (<40 m) patterns of dispersal, the dispersal capability of sponge larvae is still restricted when compared with that of other invertebrate larvae. Small-scale random factors related to larval dispersal and settlement in the Caribbean have been shown to be crucial to unexplained heterogeneity among sponge communities (Zea, 1993, 2002). Among the four groups of sponge assemblage, Groups 1, 2 and 3 belong to the northern system in which the mean annual SST is slightly lower than the southern system (Group 4). Moreover, the period where SST is lower than 18°C is considerably longer in the north than in the south. The dissimilarity between sponge assemblages in the north and south is to be expected due to the differences in temperature. Sponge species that

are able to survive low SST survive in the north. The dissimilarity between Groups 2 and 3 is likely due to differences in the substrates (Table 1). The sponge assemblages of five locations in Group 2 with high similarity (>40%) is likely due to the homogeneity of hydrological and substrate features.

### Estimation of sponge biodiversity and conservation implications

Based on our results, the estimated number of sponge species in the Penghu Islands was 113; however, only 53 species were recorded in the present study. Sponges with an excavating or thin (<3 mm) encrusting morphology or cryptic features were not included. Hence, our results suggest that the potential biodiversity of sponges of the Penghu Archipelago is likely to be substantially higher than expected, which would make this area a potential 'hotspot' of sponge faunal diversity. The Penghu Archipelago is not as densely populated (<60 000 residents) as many islands in Indonesia. With the exception of the larger islands, the majority of islands/islets are unpopulated; however, intense tourism and marine recreational activities in the summer may adversely affect environmental






Fig. 5. Observed sample-based species richness (solid dots) of subtidal sponge species collected from 15 different sampling sites and estimated species richness based on the standard rarefaction curves (solid line), Chao 2 (Y), and Jack 1 (p) formulas. Dotted lines indicate the upper and lower 95% confidence interval of the standard rarefaction estimation.

conditions. In addition to this, there is widespread marine cage culture of commercially important fish species, e.g. amberjack (*Seriola dumerili*), cobia (*Rachycentron canadum*) and red seabream (*Pagrus major*), in the Penghu Archipelago, particularly in the inner bay of Magongwan (not included in the present study). Studies suggest that marine cage farming has led to chronic nutrient enrichment in the surrounding waters, which may have accounted for the deterioration of suitable habitats for coral reef organisms (Chen & Hsu, 2006; Huang *et al.*, 2011). Recently (October 2014), the Four Islands of Southern Penghu, including Dongyuping, Siyuping, Dongjiyu and Sijiyu (the latter two were not included in the present study), and proximate islets, have been proposed for implementation of the South Penghu Marine National Park. The establishment of a baseline of marine biodiversity including the sponge fauna is essential for future conservation policymaking and park management. For protecting the local sponge biodiversity from decline, the assessment of the capacity and environmental impact of various tourism activities, regulation of neritic fishing behaviour (both sport and commercial), and long-term monitoring of biotic and abiotic factors should be considered.

#### ACKNOWLEDGEMENTS

The authors would like to thank Y. Su, C.-C. Huang, P.-T. Wu, J.-C. Liang and K. Liao for their assistance with the fieldwork; Ocean Data Bank of the Ministry of Science and Technology, Taiwan for providing bathymetric maps; and K. Liao and F.-L. Kuo for refining maps.

#### FINANCIAL SUPPORT

This study was supported by a research grant awarded to Y.M.H. by the Ministry of Science and Technology, Taiwan (NSC 100-2621-B-346-001) and the Temminck Fellowship from Naturalis Biodiversity Center, the Netherlands.

#### REFERENCES

- Becking L.E., Cleary D.F.R., de Voogd N.J., Renema W., de Beer M., Van Soest R.W.M. and Hoeksema B.W. (2006) Beta diversity of tropical marine benthic assemblages in the Spermonde Archipelago, Indonesia. *Marine Ecology* 27, 76–88.
- Bell J.J. and Barnes D.K.A. (2000) A sponge diversity centre within a marine 'island'. *Hydrobiologia* 440, 55–64.
- Bell J.J. and Smith D. (2004) Ecology of sponges (Porifera) in the Wakatobi region, south-eastern Sulawesi, Indonesia: richness and abundance. *Journal of the Marine Biology Association of the United Kingdom* 84, 1–11.
- Brøndsted H.V. (1929) Neue Schwämme aus Amoy an der Formosa-Strasse. *Zoologischer Anzeiger* 81, 224–229.
- Chan B.K.K. and Lee P.-F. (2012) Biogeography of intertidal barnacles in different marine ecosystems of Taiwan – potential indicators of climate change? In Stevens L. (ed.) *Global Advances in Biogeography*. Rijeka: InTech. pp. 119–136. Available from <http://www.intechopen.com/books/global-advances-in-biogeography/biogeography-of-intertidal-barnacles-in-different-marine-systems-of-taiwan-potential-indicators-for->
- Chen C.A. and Shashank K. (2009) Taiwan as a connective stepping-stone in the Kuroshio Triangle and the conservation of coral ecosystems under the impacts of climate change. *Kuroshio Science* 3, 15–22.
- Chen C.-Y. (2000) *Studies on bioactive constituents from selected Taiwanese marine sponges*. PhD dissertation. National Sun Yat-Sen University, Kaohsiung, Taiwan, 174 pp.
- Chen Y.-G. and Liu T.-K. (1996) Sea level changes in the last several thousand years, Penghu Islands, Taiwan Strait. *Quaternary Research* 45, 254–262.
- Chen Y.-H. (1988) *Morphology of budding and adult Cinachya australiensis (Carter, 1886)*. M.S. Master's thesis. National Sun Yat-Sen University, Kaohsiung, Taiwan, 42 pp.
- Chen Y.-H., Chen C.-P. and Chang K.-H. (1997) Budding cycle and bud morphology of the globe-shaped sponge *Cinachya australiensis*. *Zoological Science* 36, 194–200.
- Chen Y.-S. and Hsu C.-Y. (2006) Ecological considerations of cage aquaculture in Taiwan. *Journal of the Fisheries Society of Taiwan* 33, 139–146.

- Chung I.-F., Huang Y.-M., Lee T.-H. and Liu L.-L. (2010) Reproduction of the bath sponge *Spongia ceylonensis* (Dictyoceratida: Spongiidae) from Penghu, Taiwan. *Zoological Studies* 49, 601–607.
- Clarke K.R. and Warwick R.M. (2001a) *Change in marine communities: an approach to statistical analysis and interpretation*. 2nd edition. Plymouth: PRIMER-E.
- Clarke K.R. and Warwick R.M. (2001b) A further biodiversity index applicable to species lists: variation in taxonomic distinctness. *Marine Ecology Progress Series* 216, 265–278.
- Cleary D.F.R. and De Voogd N.J. (2007) Environmental associations of sponges in the Spermonde Archipelago, Indonesia. *Journal of the Marine Biology Association of the United Kingdom* 87, 1669–1676.
- Colwell R.K. (2013) EstimateS: statistical estimation of species richness and shared species from samples. Ver. 9.1 for Mac OS. Available at <http://viceroy.eeb.uconn.edu/EstimateS>.
- de Voogd N.J. (2012) On sand-bearing myxillid sponges, with a description of *Psammochela tutiae* sp. nov. (Poecilosclerida, Myxillina) from the northern Moluccas, Indonesia. *Zootaxa* 3155, 21–28.
- de Voogd N.J., Becking L.E., Hoeksema B.W., Noor A. and Van Soest R.W.M. (2004) Sponge interactions with spatial competitors in the Spermonde Archipelago. *Bollettino dei Musei e Degli Istituti Biologici dell'Università di Genova* 68, 253–261.
- de Voogd N.J. and Cleary D.F.R. (2008) An analysis of sponge diversity and distribution at three taxonomic levels in the Thousand Islands/Jakarta Bay reef complex, West-Java, Indonesia. *Marine Ecology* 29, 205–215.
- Diaz M.C. and Rützler K. (2001) Sponges: an essential component of Caribbean coral reefs. *Bulletin of Marine Science* 69, 535–546.
- Dobretsov S., Dahms H.-U. and Qian P.-Y. (2005a) Antibacterial and anti-diatom activity of Hong Kong sponges. *Aquatic Microbial Ecology* 38, 191–201.
- Dobretsov S., Dahms H.-U., Tsoi M.Y. and Qian P.-Y. (2005b) Chemical control of epibiosis by Hong Kong sponges: the effect of sponge extracts on micro- and macrofouling communities. *Marine Ecology Progress Series* 297, 119–129.
- Duckworth A.R. and Peterson B.J. (2012) Effects of seawater temperature and pH on the boring rates of the sponge *Cliona celata* in scallop shells. *Marine Biology* 160, 27–35.
- Duckworth A.R., West L., Vansach T., Stubler A. and Hardt M. (2012) Effects of water temperature and pH on growth and metabolite biosynthesis of coral reef sponges. *Marine Ecology Progress Series* 462, 67–77.
- Fromont J. (1999) Demosponges of the Houtman Abrolhos. *Memoirs of the Queensland Museum* 44, 175–183.
- Gray J.E. (1868) Note on Theonella, a new genus of Coralloid sponges from Formosa. *Proceedings of the Zoological Society of London* 37, 565.
- Hooper J.N.A. and Van Soest R.W.M. (2002) *Systema Porifera: a guide to the classification of sponges*. 1st edition. New York, NY: Kluwer Academic/Plenum Publishers.
- Hoshino T. (1980) A guide to identification of principal fouling organism. (4) Sponges (Calcareous sponges and Demosponges). *Marine Fouling* 2, 53–58 [in Japanese].
- Hsieh H.J., Hsien Y.-L., Tsai W.-S., Su W.-C., Jeng M.-S. and Chen C.A. (2008) Tropical fishes killed by the cold. *Coral Reefs* 27, 599.
- Hu J., Kawamura H., Li C., Hong H. and Jiang Y. (2010) Review on current and seawater volume transport through the Taiwan Strait. *Journal of Oceanography* 66, 591–610.
- Huang H.Z. (1996) *The influence of environmental factors on the filtration rate of sponge*, *Cinachyra australiensis*. M.S. Master's thesis. National Sun Yat-Sen University, Kaohsiung, Taiwan.
- Huang J.P., McClintock J.B., Amsler C.D. and Huang Y.M. (2008) Mesofauna associated with the marine sponge *Amphimedon viridis*. Do its physical or chemical attributes provide a prospective refuge from fish predation? *Journal of Experimental Marine Biology and Ecology* 362, 95–100.
- Huang Y.-C.A., Hsieh H.J., Huang S.-C., Meng P.-J., Chen Y.-S., Keshavmurthy S., Nozawa Y. and Chen C.A. (2011) Nutrient enrichment caused by marine cage culture and its influence on subtropical coral communities in turbid waters. *Marine Ecology Progress Series* 423, 83–93.
- Huang Y.L. (2002) *Diversity and antibacterial activity of heterotrophic bacteria associated with sponges*. M.S. Master's thesis. National Taiwan University, Taipei, Taiwan, 80 pp.
- Jan S., Tseng Y.-H. and Dietrich D.E. (2010) Sources of water in the Taiwan Strait. *Journal of Oceanography* 66, 211–221.
- Jan S., Wang J., Chern C.-S. and Chao S.-Y. (2002) Seasonal variation of the circulation in the Taiwan Strait. *Journal of Marine Systems* 35, 249–268.
- Jhou G.-R. (2004) *Antitumor constituents from Formosan marine sponge Negombata corticata*. M.S. Master's thesis. National Sun Yat-Sen University, Kaohsiung, Taiwan, 146 pp.
- Kang D.W., Lee K.J. and Sim C.J. (2013) Two new marine sponges of the genus *Haliclona* (Haplosclerida: Chalinidae) from Korea. *Animal Systematics, Evolution and Diversity* 29, 51–55.
- Kelmo F., Bell J.J. and Attrill M.J. (2013) Tolerance of sponge assemblages to temperature anomalies: resilience and proliferation of sponges following the 1997–8 El Niño Southern Oscillation. *PLoS ONE* 8, e76441. doi: 76410.71371/journal.pone.0076441.
- Khodakovskaya A.V. (2005) Fauna of sponges (Porifera) of Peter the Great Bay, Sea of Japan. *Russian Journal of Marine Biology* 31, 209–214.
- Lazoski C., Solé-Cava A.M., Boury-Esnault N., Klautau M. and Russo C.A.M. (2001) Cryptic speciation in a high gene flow scenario in the oviparous marine sponge *Chondrosia reniformis*. *Marine Biology* 139, 421–429.
- Lesser M.P. (2006) Benthic-pelagic coupling on coral reefs: feeding and growth of Caribbean sponges. *Journal of Experimental Marine Biology and Ecology* 328, 277–288.
- Lesser M.P., Slattery M. and Leichter J.J. (2009) Ecology of mesophotic coral reefs. *Journal of Experimental Marine Biology and Ecology* 375, 1–8.
- Li T.-H. (2013) *Preliminary survey on the biodiversity of marine sponges (Porifera: Demospongiae) in the southern water off the Penghu Archipelago*. M.S. Master's thesis. National Sun Yat-sen University, Kaohsiung, Taiwan, 152 pp.
- Liao C.-C. (2003) *I. Studies on the bioactive constituents and acylated derivatives of Taiwanese Spongia sp. II. Studies on the lignans of Taiwanese Kadsura philippinensis Elmer*. M.S. Master's thesis, National Sun Yat-Sen University, Kaohsiung, Taiwan.
- Liao M.-H., Tang S.L., Hsu C.M., Wen K.C., Wu H., Chen W.M., Wang J.T., Meng P.-J., Twan W.H., Dai C.F., Soong K. and Chen C.A. (2007) The “black disease” of reef-building corals at Green Island, Taiwan – outbreak of a cyanobacteriosponge, *Terpios hoshinota* (Suberitidae; Hadromerida). *Zoological Studies* 46, 520.
- Lim S.-C., De Voogd N.J. and Tan K.-S. (2009) Fouling sponges (Porifera) on navigation buoys from Singapore waters. *Raffles Bulletin of Zoology* 22, 41–58.
- Lim S.-C., De Voogd N.J. and Tan K.-S. (2012) Biodiversity of shallow-water sponges (Porifera) in Singapore and description of a new species of *Forcepia* (Poecilosclerida: Coelosphaeridae). *Contributions to Zoology* 81, 55–71.

- Lin S.L. (1995) *Antitumor constituents from Formosan marine sponge Strongylophora durissima*. M.S. Master's thesis. National Sun Yat-Sen University, Kaohsiung, Taiwan, 84 pp.
- Lo K.-L. (1999) *Antitumor constituents from Formosan marine sponge Xestospongia sp.* M.S. Master's thesis. National Sun Yat-Sen University, Kaohsiung, Taiwan, 50 pp.
- Lu D.-K. (2003) *Fluorescence in situ hybridization of symbiotic chemotrophic sulfur-oxidizing bacteria of the sponge, Cinachyra australiensis*. M.S. Master's thesis. National Sun Yat-Sen University, Kaohsiung, Taiwan, 133 pp.
- Maldonado M. and Young C.M. (1998) Limits on the bathymetric distribution of keratose sponges: a field test in deep water. *Marine Ecology Progress Series* 174, 123–139.
- Massaro A. (2009) *Selective filtration in the tropical marine sponge Rhopaloeides odorabile: impacts of elevated seawater temperature on feeding behavior*. Independent Study Project (ISP) Collection. Paper 774. [http://digitalcollections.sit.edu/isp\\_collection/774](http://digitalcollections.sit.edu/isp_collection/774).
- McClintock J.B., Amsler C.D., Baker B.J. and Van Soest R.W.M. (2005) Ecology of Antarctic marine sponges: an overview. *Integrative and Comparative Biology* 45, 359–368.
- Mccune B. and Grace J.B. (2002) *Analysis of ecological communities*. Glenden Beach, Oregon, USA: MjM Software Design.
- Munro C. (2005) Diving systems. In Eleftheriou A. and McIntyre A. (eds) *Methods for the study of marine benthos*. 3rd edition. Oxford: Blackwell Science, pp. 112–159.
- Olson J.B. and Kellogg C.A. (2010) Microbial ecology of corals, sponges, and algae in mesophotic coral environments. *FEMS Microbiology Ecology* 73, 17–30.
- Pawlik J.R., Chanas B., Toonen T.J. and Fenical W. (1995) Defenses of Caribbean sponges against predatory reef fish. I. Chemical deterrence. *Marine Ecology Progress Series* 127, 183–194.
- Pile A.J., Grant A., Hinde R. and Borowitzka M.A. (2003) Heterotrophy on ultraplankton communities is an important source of nitrogen for a sponge-rhodophyte symbiosis. *Journal of Experimental Biology* 206, 4533–4538.
- Qiu Y., Chen C.-T.A., Guo X. and Jing C. (2011) Currents in the Taiwan Strait as observed by surface drifters. *Journal of Oceanography* 67, 395–404.
- Roberts D.E., Davis A.R. and Cummins S.P. (2006) Experimental manipulation of shade, silt, nutrients and salinity on the temperate reef sponge *Cymbastela concentrica*. *Marine Ecology Progress Series* 143, 143–154.
- Shao K.-T., Peng C.-I. and Wu W.J. (2008) *2008 Taiwan species diversity II. Species checklist*. Taipei: Forestry Bureau, Council of Agriculture, Executive Yuan, Taiwan.
- Shih H.-T. (2012) Distribution of fiddler crabs in East Asia, with a note on the effect of the Kuroshio Current. *Kuroshio Science* 6, 83–89.
- Shim E.J. and Sim C.J. (2013) Two marine sponges of the Family Ancorinidae (Demospongiae: Astrophorida) from Korea. *Animal Systematics, Evolution and Diversity* 29, 31–35.
- Su J.-H., Tseng S.-W., Lu M.-C., Liu L.-L., Chou Y. and Sung P.-J. (2011) Cytotoxic C21 and C22 terpenoid-derived metabolites from the sponge *Ircinia sp.* *Journal of Natural Products* 74, 2005–2009.
- Swierts T., Peijnenburg K.T.C.A., Christiaan de Leeuw C., Cleary D.F.R., Hörnlein C., Setiawan E., Wörheide G., Erpenbeck D. and de Voogd N.J. (2013) Lock, stock and two different barrels: comparing the genetic composition of morphotypes of the Indo-Pacific sponge *Xestospongia testudinaria*. *PLoS ONE* 8, e74396. doi:74310.71371/journal.pone.0074396.
- Tsakamoto S., Yamanokuchi R., Yoshitomi M., Sato K., Ikeda T., Rotinsulu H., Mangindaan R.E.P., de Voogd N.J., van Soest R.W.M. and Yokosawa H. (2010) Aaptamine, an alkaloid from the sponge *Aaptos suberitoides*, functions as a proteasome inhibitor. *Bioorganic and Medicinal Chemistry Letters* 20, 3341–3343.
- Tzeng M.-T., Lan K.-W. and Chan J.-W. (2012) Interannual variability of wintertime sea surface temperatures in the eastern Taiwan Strait. *Journal of Marine Science and Technology* 20, 707–712.
- Van Soest R.W.M. (1993) Distribution of sponges on the Mauritanian continental shelf. *Hydrobiologia* 258, 95–106.
- Van Soest R.W.M., Boury-Esnault N., Hooper J.N.A., Rützler K., de Voogd N.J., Alvarez de Glasby B., Hajdu E., Pisera A.B., Manconi R., Schoenberg C., Janussen D., Tabachnick K.R., Klautau M., Picton B., Kelly M., Vacelet J., Dohrmann M., Díaz M.-C. and Cárdena P. (eds) (2015) *World Porifera database*, available at <http://www.marinespecies.org/porifera>.
- Van Soest R.W.M., Boury-Esnault N., Vacelet J., Dohrmann M., Erpenbeck D., De Voogd N.J., Santodomingo N., Vanhoorne B., Kelly M. and Hooper J.N.A. (2012) global diversity of sponges (Porifera). *PLoS ONE* 7, e35105.
- Van Soest R.W.M., Cleary D.F.R., De Kluijver M.J., Lavaleye M.S.S., Maier C. and Van Duyl F.C. (2007) Sponge diversity and community composition in Irish bathyal coral reefs. *Contributions to Zoology* 76, 121–142.
- Wang J.-T., Hirose E., Hsu C.-M., Chen Y.-Y., Meng P.-J. and Chen C.A. (2012) A coral-killing sponge, *Terpios hoshinota*, releases larvae harboring cyanobacterial symbionts: an implication of dispersal. *Zoological Studies* 51, 314–320.
- Wang Y.-H. (2012) Habitats and environmental investigation of the Penghu 4-South Islands. *Marine National Park Headquarters*, pp. 264 [in Chinese with English abstract].
- Webster N.S., Cobb R.E. and Negri A.P. (2008) Temperature thresholds for bacterial symbiosis with a sponge. *ISME Journal* 2, 830–842.
- Wulff J.L. (2001) Assessing and monitoring coral reef sponges: why and how? *Bulletin of Marine Science* 69, 831–846.
- Wulff J.L. (2006) Rapid diversity and abundance decline in a Caribbean coral reef sponge community. *Biological Conservation* 127, 167–176.
- Xavier J.R., Rachello-Dolmen P.G., Parra-Velandia F.J., Schönberg C.H.L., Breeuwer J.A.J. and Van Soest R.W.M. (2010) Molecular evidence of cryptic speciation in the “cosmopolitan” excavating sponge *Cliona celata* (Porifera, Clionidae). *Molecular Phylogenetics and Evolution* 56, 13–20.
- Yahel G., Whitney F., Reising H.M., Eerkes-Medrano D.I. and Leys S.P. (2007) *In situ* feeding and metabolism of glass sponges (Hexactinellida, Porifera) studied in a deep temperate fjord with a remotely operated submersible. *Limnology and Oceanography* 52, 428–440.
- Zea S. (1993) Patterns of coral and sponge abundance in stressed coral reefs at Santa Marta, Colombian Caribbean. In Van Soest R.W.M., Van Kempen T.M.G. and Braekman J.-C. (eds) *Proceedings of the 4th International Porifera Congress, Amsterdam, the Netherlands, 1993*. Rotterdam: A. A. Balkema, pp. 257–264.
- and
- Zea S. (2002) Patterns of sponge (Porifera, Demospongiae) distribution in remote oceanic reef complexes of the southwestern Caribbean. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 25, 579–592.

#### Correspondence should be addressed to:

Y.M. Huang  
 Department of Marine Sports and Recreation, National  
 Penghu University of Science and Technology, Magong City,  
 Taiwan  
 email: [yusheng@gms.npu.edu.tw](mailto:yusheng@gms.npu.edu.tw)