

RISK AND MAIZE-BASED CROPPING SYSTEMS FOR SMALLHOLDER MALAWI FARMERS USING CONSERVATION AGRICULTURE TECHNOLOGIES

By A. R. NGWIRA†‡, C. THIERFELDER§, N. EASH¶ and D. M. LAMBERT††

†*Department of International Environment and Development Studies, Norwegian University of Life Sciences, P.O. Box 5003, NO-1432 Aas, Norway*, §*CIMMYT, P.O. Box MP 163, Mount Pleasant, Harare, Zimbabwe*, ¶*Department of Biosystems Engineering and Soil Science, University of Tennessee Institute of Agriculture, 2506 E. J. Chapman Drive, Knoxville, TN 37996-4518, USA* and ††*Department of Agricultural & Resource Economics, University of Tennessee Institute of Agriculture, 321 Morgan Hall, 2621 Morgan Circle, Knoxville, TN 37996-4518, USA*

(Accepted 8 April 2013; First published online 13 May 2013)

SUMMARY

Agricultural production in southern Africa is constrained by numerous factors, including low soil fertility, frequent droughts and flooding, limited access to fertilizers and the use of unsustainable management techniques that increase soil erosion rates. Conservation agriculture (CA) is based on the principles of minimum soil disturbance, crop residue retention and crop rotations. CA systems have been proposed to alleviate the negative externalities associated with conventional crop management systems. This study was conducted to examine the riskiness of economic returns of CA technologies based on maize grain yield evaluated in 12 target communities in Malawi from 2005–2011. On average, maize grain yields on both CA treatments exceeded the conventional control treatment by 22.1–23.6%, with differences more distinct in low altitude areas with low rainfall and frequent seasonal dry spells. Stochastic dominance analysis suggest that CA technologies would be preferred by risk-averse farmers, with corresponding differences in risk premiums (compared to conventional maize production systems) ranging between US\$40 and US\$105. However, these rankings are sensitive to the agroecological zones where the experiments were conducted. The risk premiums associated with the CA technologies in low elevation regions are unambiguous. Risk-averse farmers in higher elevations may need substantial incentives to adopt some CA technologies.

INTRODUCTION

Food security is a major concern for eastern and southern Africa where agriculture is predominantly rain-fed (Rockström *et al.*, 2002). Despite technological solutions and policy support, rainfall variability and degraded soils resulting from tillage and insufficient levels of organic matter recycling are significant factors contributing to chronically low crop yields (Kumwenda *et al.*, 1997; Rockström *et al.*, 2009; Wall, 2007). In many farming systems, soil organic matter has declined to unsustainably low levels due to continuous cropping; all of which are important causes of low soil moisture, poor nutrient use efficiency and lower productivity (Parr *et al.*, 1990). These problems reduce the benefits of improved genomic varieties and increase the inherent

‡Corresponding author. Email: robert.ngwira@umb.no

risks of farming (Thierfelder and Wall, 2009) and discourage adoption of improved practices. Conservation agriculture (CA) addresses these problems at several levels, and may provide a means whereby the yield risk that smallholder farmers face can be moderated.

Three actions define conservation agriculture: (1) Minimizing soil disturbance by direct sowing of crops into the soil, (2) protecting soil with cover crops or crop residues and (3) intercropping and/or crop rotation (FAO, 2002). Soil disturbance by mechanical tillage is reduced to an absolute minimum, and the use of agrochemicals and nutrients of mineral or organic origin are applied at optimal levels if known. The interactions between minimal soil disturbance, managing crop residues, applying optimal nutrient levels and controlling weed populations are often considered more consequential than the individual effects of these management activities alone. Instead of maximizing crop yields, the managerial objectives under conservation agriculture are to optimize long-term soil fertility through crop residue management and crop rotations, resulting in sustainably higher maize yields at reduced input costs.

Conservation agriculture systems improve water infiltration, decrease soil erosion and run-off, increase soil moisture retention and improve soil fertility and growth of beneficial microbial communities (Mando *et al.*, 1999; McGarry *et al.*, 2000; Roth *et al.*, 1988; Thierfelder and Wall, 2009; Verhulst *et al.*, 2011). CA systems resulted in yield increases for several crops, including maize, legumes, sorghum, cotton, sunflower, potatoes, finger millet, pigeonpea and cotton (Ngwira *et al.*, 2012; Rockström *et al.*, 2009; Thierfelder and Wall, 2010). A meta-analysis of the long-term effects of conservation agriculture on maize yield found increases in yield over time with practices that included crop rotations and high input use in low rainfall areas (Rusinamhodzi *et al.*, 2011), but most yield increases are not observed in the early years of adoption. Cases reporting lower yields from conservation agriculture compared to conventional technologies are often associated with learning curves producers face when adapting conservation agriculture to their production systems. One probable cause of the yield drag is likely due to the crop nutrients temporarily immobilized in the soil by increased number of microbes below the surface mulch, which need nitrogen from the soil for their population increase, thus leaving little nitrogen for early plant development. In addition, the nutrients in the residue break down more slowly and are consumed by microbes and become unavailable to the plant (Doran *et al.*, 1998). The short-term risk of lower production and therefore lower household consumption is an unattractive prospect for food insecure smallholder farmers (Shively, 2001), which makes it even more difficult to challenge farmer preconceptions of conservation agriculture. In addition to individual learning curves, newly introduced ideas or systems may appear as failures to would-be adopters if those technologies were implemented during periods of either extraordinary rainfall or extended dry spells on already-degraded, nutrient-poor soils. This short-term perspective challenges the widespread adoption of CA practices; highlighting the fact that farmer acceptance of the idea that agriculture is possible without tilling is critical (Wall, 2007).

This study examines conservation agriculture as a risk-mitigating technology for smallholder farmers in Malawi using standard risk analysis methods. Malawi is a

sub-tropical country situated between latitude 9° and 18°S and 33° and 36°E in South Eastern Africa with a land area of over 118,000 km², of which 23,000 km² is covered by Lake Malawi. The country is divided into three main regions: North, Central and South. Agriculture is the mainstay of the economy, contributing nearly 35% of the gross domestic product and employing more than 80% of the total labor force. It is estimated that 56.6% of the rural population live below the poverty line (NSO, 2012). The achievement of the Millennium Development Goals and growth and poverty reduction targets in the southern African region are especially challenging. This region is barely self-sufficient in food grains, with a net import of 10% if South Africa is excluded (FAOSTAT, 2010). With growth of population and incomes, the demand for maize is projected to increase approximately 3–4% annually over the next 10 years, leading to an increase in demand for maize grain by at least 40% (FAOSTAT, 2010). Typical cultivated land holding sizes range between 0.2 ha and 3 ha (Ellis *et al.*, 2003), with maize the principal crop occupying about 85% of arable land area under cultivation (MoAFS, 2011; Smale *et al.*, 1991).

Standard risk analysis methods are used to determine the risk premium associated with conservation agriculture and conventional technologies, given different assumptions about farmer aversion to risk. Risk premiums are the dollar amounts that producers would be willing to pay to eliminate the risk associated with a given technology. Therefore, the risk premium of CA technologies compared with those associated with conventional maize production practices provides insight into the potential monetary value of CA technologies to risk-averse smallholder producers.

Risk is quantified as the dispersion around an expected outcome; e.g. yields ('yield risk') or net returns ha⁻¹ ('economic' or 'price' risk) (Anderson and Dillon, 1990). In addition to uncertain rainfall and impending disease, access to equity and capital markets, household resources, social capital and liquidity constraints also influence technology adoption or abandonment through producers' perceptions of risk (Cavatassi *et al.*, 2011; Moser and Barrett, 2006; Wendland and Sills, 2008). Commodity prices and subsidies also shape risk perception, thereby motivating the adoption of novel production technologies (Serra, 2008).

In sub-Saharan Africa economic analyses of CA technologies as potentially risk-mitigating are relatively sparse. Technology adoption studies from Nigeria (Lal, 1986), Zimbabwe (Vogel, 1993) and Zambia (Thierfelder and Wall, 2010) found that crop losses were reduced in fields managed under conservation agriculture. Cavatassi *et al.* (2011) found that Ethiopian farmers appeared to adopt high yielding sorghum varieties to moderate production risk, yet in regions where rainfall was highly variable there were lower adoption rates of these same improved varieties. Recent partial budget analyses of conservation agriculture in sub-Saharan smallholder farming systems compared the marginal benefit of the technologies with conventional practices. For example, in their study of minimum tillage systems in Kenya, Guto *et al.* (2011) conclude that risk-averse farmers preferred CA technologies to conventional practices because the discounted marginal rates of return from conservation agriculture were higher than those from the conventional systems. Mazvimavi and Twomlow's (2009) profitability analysis used a partial budgeting approach to compare returns ha⁻¹ for conservation

agriculture and conventional practices assuming low, normal and high rainfall periods. Based on partial budgeting exercises, it is often concluded that CA technologies are 'risk-reducing' based on superior net returns ha^{-1} compared to conventional tillage systems (Guto *et al.*, 2011).

However, assertions about risk based on linear cost-benefit analysis are only relevant for risk-neutral producers (Mas-Colell *et al.*, 1995). Based on partial budgeting exercises it is often concluded that CA technologies are 'risk-reducing' based on superior net returns ha^{-1} compared with conventional tillage systems without actually measuring risk at all. Few studies have accounted for risk in net return comparisons of CA/non-CA production systems in Africa. For example, using stochastic dominance analysis of partial budget results Bekele (2005) found that risk-averse producers preferred soil and water conserving practices. Farmers associated a premium with the technologies that moderated the production risks accompanying drought. In general, if a producer is risk-averse, then risk impedes production by inducing lower input use than would otherwise prevail (Anderson and Dillon, 1990). This study examines the riskiness of net returns of conservation agriculture using non-parametric methods, stochastic dominance, target revenue sensitivity analysis and risk premium determination based on six-year (2005–2011) on-farm maize production data in Malawi. We hypothesize that the distribution of net returns generated from CA technologies will exhibit higher expected returns with lower variance compared with conventional agronomic practices. That is, the CA technologies will dominate conventional tillage practices in terms of higher risk premium at all levels of risk aversion. Thus, producers who are relatively more risk-averse will prefer CA technologies if these technologies reduce variability in net returns because of higher yields, or increase profit margins through lower input costs. We also hypothesize that these results will be generalizable across growing areas at different elevations.

MATERIALS AND METHODS

The study was conducted over six years (2005–2011) in 12 target communities of 10 extension planning areas (EPAs) in several districts in northern, central and southern Malawi (Table 1). The following districts were targeted for this research (from south to north): Zomba, Machinga, Balaka, Dowa Salima, Nkhotakota and Mzimba. All EPAs are characterized by unimodal rainfall patterns with a rainy season from November to April. The average annual rainfall was variable across the study communities (Table 1). All the study communities in Dowa, Mzimba and Zomba districts belong to the mid-altitude areas, while the rest of the communities in the remaining districts fall under low altitude areas (Brown and Young, 1966). The dominant soil types found in these EPAs are Chromic Luvisols, Haplic Lixisols, Eutric Fluvisols and Cambisols and some alluvial soils (WRB, 1998; Table 1).

Experimental design for on-farm field trials: Malawi 2005–2011

The study was conducted on farms with a total of six farmers hosting validation trials replicated across sites and spread across each target community (Table 1). However,

Table 1. Communities, geographical location and soil types of target villages.

District	Community	EPA	Agroecological zone	Altitude	Rainfall (mm)	Soil type
Balaka	Herbert	Bazale	Low altitude	635 m	684	Chromic Luvisols
Balaka	Lemu	Bazale	Low altitude	720 m	862	Chromic Luvisols
Balaka	Malula	Bazale	Low altitude	605 m	717	Eutric Fluvisols
Dowa	Chipeni	Mvera	Medium altitude	1166 m	883	Chromic Luvisols
Dowa	Chisepo	Chisepo	Medium altitude	1090 m	1013	Alfisols
Machinga	Matandika	Ntumbi	Low altitude	688 m	874	Cambic Arenosols
Mzimba	Enyezini	Emsizini	Medium altitude	1337 m	1194	Ferrallitic soils
Nkhotakota	Linga	Linga	Low altitude	629 m	1237	Alluvial soils
Nkhotakota	Mwansambo	Mwansambo	Low altitude	632 m	1371	Haplic Lixisols
Nkhotakota	Zidyana	Zidyana	Low altitude	535 m	1429	Haplic Luvisols
Salima	Chinguluwe	Chinguluwe	Low altitude	657 m	1241	Eutric Cambisols
Zomba	Songani	Malosa	Medium altitude	791 m	1371	Ferrallitic soils

the number of farmers per community varied according to the number of years of CA practice. In addition, the number of farmers changed due to farmer adherence to protocol: some years not as many, some years more than six. Following a good CA year more farmers generally followed the protocol. The main cropping systems studied in all EPAs were monocrop maize (*Zea mays* L.) and maize–legume intercropping. Farmer managed plots were 3000 m². Each plot was divided into three equally sized subplots (treatments) of 1000 m² as follows:

1. Treatment 1: Conventional control plot consisting of traditional ridge and furrow land preparations planted with continuous monocrop maize (CT or check). Residues were managed using methods commonly practiced in each EPA; i.e. residues were incorporated into the ridges in Lemu and removed in Zidyana. Continuous monocrop maize was planted on the ridges.
2. Treatment 2: CA plot with continuous monocrop maize (CA) planted into previous years' ridges (where they still existed) or directly into the plot without previous ridge formation. Crop residues from previous years' harvests were retained as surface mulch. Maize seeds were planted as sole crops by no till methods using a pointed stick (dibble stick).
3. Treatment 3: CA plot with maize intercropped with a legume (cowpea (*Vigna unguiculata* L.) in Zidyana or pigeonpea (*Cajanus cajan* L.) in Lemu) (CA–legumes). Both crops were planted with the dibble stick into previous years' ridges (where they still existed) or directly into the plot without further ridging. Crop residues were retained as surface mulch as in treatment 2.

Farmers with support from extension officers in all target communities managed the trials. Field staff and technicians provided plot management recommendations. Hybrid maize DKC8033 was used for the duration of the experiments. Sudan cowpea and ICEAP0040 pigeonpea varieties were the intercrop grain legumes. All plots were seeded after the first effective rains (e.g. rainfall greater than 30 mm after 15 November). Ridges in the conventional tillage practice were prepared using hand hoes around October, and planting was done again with hand hoes. Seed was planted in CA plots

with a pointed dibble stick, which aims at minimum soil disturbance by only creating a planting hole where seeds are placed. Both legumes and maize were seeded at the same time. Ridge spacing was kept constant in the CA treatments: 75 cm between maize rows, 25 cm between planting stations and one living plant per station. The CT plots followed seed population and spacing patterns normally used by the farmer. In most cases farmers have adopted similar 75×25 -cm plant spacing according to the recommendations from Sasakawa Global 2000 (Ito *et al.*, 2007). Intercropped legumes were seeded between maize rows planting two seeds spaced at 60 cm and 40 cm apart for pigeonpea and cowpea respectively.

All treatments received uniform fertilizer rates of 69 kg N ha^{-1} , which was supplied by applying two bags ha^{-1} of 23:21:0 + 4S at seeding and two bags ha^{-1} of urea (46% N) approximately three weeks after planting. Manual weeding as necessary did weed control in CT plots. In treatment 2, a tank mix of 2.5 L ha^{-1} glyphosate (N-(phosphono-methyl)glycine) and 6 L ha^{-1} bullet (which contains 25.4% Alachlor (2-chloro-N-(2,6-diethylphenyl)-N-(methoxymethyl)acetamide) and 14.5% Atrazine (2-chloro-4-ethylamino-6-isopropylamino-1,3,5-triazine)) was applied as post-planting and pre-emergence herbicide respectively. In treatment 3, only 2.5 L ha^{-1} post-planting herbicide glyphosate was used followed by manual weeding when weeds were 10-cm tall or 10 cm in circumference.

Maize grain yield

At physiological maturity, maize was harvested from 10 samples of two rows by 5 m from each treatment. A sub-sample of 20 cobs per treatment was shelled to calculate grain yield, to record the moisture percentage and to calculate final grain yield ha^{-1} basis at 12.5% moisture. The harvest area of the net plots was used to extrapolate yields to an areal basis. All maize stalks and leaves without cobs were weighed at harvest; 10 sub-samples per plot were air-dried for at least four weeks before final dry weights were taken and biomass was calculated to an areal basis.

Net returns and costs

Revenue ha^{-1} from maize grown with conservation agriculture and conventional tillage systems are based on yields recorded from 344 replicates (three treatment plots per farmer) from on-farm trials. In all, there are $3 \times 334 = 1032$ observations available for analysis. Net return (profit) ha^{-1} was estimated for each maize yield observation (kg ha^{-1}) produced by each technology based on the 2005–2011 domestic maize price series and the variable costs of each technology (conventional, CA, and CA + legume intercrop system; $n = 344$ for each technology). District maize prices were converted from Malawi Kwacha kg^{-1} to US dollars (US\$) kg^{-1} using the official exchange rates for this time period posted by the Reserve Bank of Malawi.¹ Maize prices were then converted into real 2010 US\$ (Table 2).

Extension agents working with farmers over the life of the project recorded variable costs. The most reliable cost estimates were based on farmer records in the Balaka and

¹http://www.rbm.mw/archive_dfbbr.aspx (Accessed 17 October 2011).

Table 2. Prices and costs used to generate net revenue conventional, conservation agriculture (CA) and CA with legume intercrops.

	Maize price (2010 US\$ kg ⁻¹)						
	2005	2006	2007	2008	2009	2010	2011
Balaka	0.15	0.21	0.16	0.29	0.27	0.21	0.12
Nkhotakota	0.17	0.22	0.15	0.43	0.21	0.21	0.15
Country average	0.15	0.21	0.23	0.29	0.35	0.26	0.19
	Costs (US\$ ha ⁻¹)						
	Check	CA	CA + legumes				
Balaka	378.10	412.25	430.47				
Nkhotakota and Dowa	345.80	382.45	440.67				

Dowa districts. The Dowa variable cost estimates were used to proxy variable costs in the Nkhotakota and Salima districts in the central region whereas the Balaka variable cost estimates were used to proxy variable costs in Machinga and Zomba districts in the southern region. Both regions have access to all-weather roads, which provide relatively efficient transportation of produce and farm inputs. In all, there were $7 \times (2 \times 3)$ possible price-by-cost combinations (Table 2). Net revenue ha⁻¹ from maize was then determined for each price-cost combination and each technology, given the maize yields recorded at each farmer plot over the duration of the trials. The risk analysis therefore assumes that recorded maize yields at each site, year and for each technology had an equal probability of being observed. Non-parametric procedures and descriptive statistics were used to compare the empirical distributions of the cultivation technology/net revenue ha⁻¹ combinations and the risk associated with each technology.

Risk analysis methods

The empirical distributions of net returns ha⁻¹ to conventional, CA and CA + legume intercrop systems were compared using three criteria: (1) stochastic dominance (Mas-Colell *et al.*, 1995), (2) a mean-variance approach (Lambert and Lowenberg-DeBoer, 2003) and (3) a relative risk criterion (Richardson, 2002). Two sensitivity analyses were conducted to measure the risk associated with each technology. The first method used a target revenue approach suggested by Richardson and Mapp (1976) to estimate the likelihood of achieving a net revenue target, given the returns from a risky technology. The second approach calculates the risk premium (based on certainty equivalents; Mas-Colell *et al.*, 1995), or the amount producers would be willing to pay to eliminate the risk associated with a technology at different levels of risk preference. Each method is discussed in sequel.

Stochastic dominance. Mean-variance and relative risk criterion provide information about the first and second moments of the empirical distributions of maize net returns ha⁻¹. However, skewness and extreme downside risk may be important determinants

of the perceived riskiness of an alternative and the readiness of producers to adopt new technologies. Stochastic dominance compares the cumulative distributions of outcomes (e.g. net returns ha^{-1}) based on two observations about humans: (i) people generally prefer more to less, and (ii) people prefer to avoid low-value outcomes. The second observation implies that people are generally risk-averse; however, this is not the same as saying that individuals avoid variability. Most people enjoy upside variability as long as they benefit from the outcome (e.g. higher yields, crop prices and profits) but are averse to downside variability. Those preferring more to less but are not averse to variability are risk-neutral.

These two observations about human behaviour are measurable in terms of the empirical distributions of outcomes using two decision rules which correspond with the assertions made above regarding human behaviour: first-degree stochastic dominance (FSD) and second-degree stochastic dominance (SDSD) rules. FSD assumes that decision makers prefer more to less. Under this criterion, a technology is preferred if it generates higher net returns ha^{-1} at every level of probability compared with those generated by competing technologies. Graphically, the preferred distribution is always to the right of other distributions.

Second-degree stochastic dominance assumes that producers are risk-averse. The propensity of a technology to produce low-value outcomes is measured as the area under its empirical distribution, with the alternative dominating others if the area under its empirical density is smaller at every outcome level. Non-parametric methods are typically used to detect SDSD. In the simple case of a cumulative distribution starting to the right of an alternative distribution and crossing over only once, the distribution to the right at the horizontal axis dominates if the area between the distributions below the crossover is greater than the area between the distributions above the intersection. Anderson (1974) provides discussion and examples of stochastic dominance use in interpreting agronomic research data.

We apply (Meyer, 1977) a generalized stochastic dominance approach (or stochastic dominance analysis with respect to utility function), using the negative exponential utility function to compare the net return ha^{-1} distributions of each alternative across the Risk Aversion Coefficient (RAC) interval; $0 \leq \text{RAC} < 0.03$, with the upper RAC limit determined using the method outlined in Lambert and Lowenberg-DeBoer (2003). The RAC is the Arrow–Pratt constant absolute risk aversion coefficient (Moschini and Hennessy, 2001). The lower bound was set to zero (risk neutrality) and the upper bound was equal to the level of risk that drove the certainty equivalent associated with a technology to zero (determined to be $\text{RAC} = 0.03$). Meyer's approach (1977) provides a method whereby inference can be made about alternative rankings, given heterogeneous risk preferences. Generalized stochastic dominance results were estimated using Simetar[©] software (Richardson *et al.*, 2006). This approach admits the possibility that risk preferences, as a function of risk-aversion, may change as the empirical distributions of returns cross one another.

Mean variance (MV) and relative risk criterion. The MV rule assumes that the dominant alternative must have either a higher mean for a given variance or a lower variance

for a given mean (Lambert and Lowenberg-DeBoer, 2003). Rankings generated by this criterion depend on the producer's preference for the trade-off between net returns ha^{-1} and the variance of these returns. A closely related procedure for ranking preferred alternatives is the relative risk approach, which uses the coefficient of variation (CV) calculated from each distribution of net returns ha^{-1} (Richardson *et al.*, 2006). Both methods provide a cursory assessment of risk, whereby summary statistics of the net returns from each technology can be used to rank the technologies in terms of riskiness.

Target net return probability analysis. This risk assessment method uses the empirical distributions of the net returns ha^{-1} to conduct sensitivity analyses regarding a target rate of return. For example, given a minimal rate of return, the likelihood of this occurring can be discerned from the empirical densities of net returns generated from each technology. The target probability methodology proposed by Richardson and Mapp (1976) was conducted using the Simetar[©] (Richardson *et al.*, 2006). This risk assessment tool ranks the profitability of the technologies according to the probability of achieving a given net return target. Lower target return levels were set at US\$0 (breakeven returns) comparing the conventional and CA technologies. Upper net return target levels ranged between US\$100 ha^{-1} and US\$1200 ha^{-1} in US\$100 intervals.

Certainty equivalent (CE) and risk premium analysis. Certainty equivalents for each alternative technology were estimated at different risk tolerance levels. A certainty equivalent value is the payoff (as net returns ha^{-1}) for which a producer is indifferent between an uncertain outcome and receiving a certain payment. Certainty equivalents for risk-averse individuals are always less than the expected monetary payoff of an uncertain project (i.e. the simple weighted average of net returns). When faced with several technology alternatives with different uncertain returns, a risk-averse individual would always choose the alternative with the highest certainty equivalent (Lambert and Lowenberg-DeBoer, 2003).

Beliefs about uncertainty maintained by risk-averse individuals are typically analysed using concave, twice-differentiable functions; for example, the negative exponential utility function $U(\pi; \lambda) = -\exp(-\text{RAC} \cdot \pi)$, where π is the expected net return ha^{-1} from a technology. Larger values of the RAC correspond with increased aversion to risk. The certainty equivalent is determined by solving the following expression with respect to the CE:

$$U(\text{CE}; \text{RAC}) = \sum_i p_i U(\pi_i; \text{RAC}),$$

where p_i is the probability weight associated with observing outcome π_i . When all observations are equally likely to occur, the probability weights are $1/n$. When $\text{RAC} = 0$, the producer is risk-neutral and the certainty equivalent is equal to the expected monetary value of the payoff (i.e. the simple average of the payoff distribution). As the RAC is increased, the certainty equivalent decreases relative to

Table 3. Conventional tillage (check), conservation agriculture (CA), and CA and legume intercrop treatment means (kg ha⁻¹) ± standard errors of means (SE), Malawi, 2005–2011.

District	Community	<i>n</i>	Check		CA		CA + Legumes	
			kg ha ⁻¹	SE	kg ha ⁻¹	SE	kg ha ⁻¹	SE
Balaka	Herbert	54	2319	±149 a	3,533	±209b	3,249	±214 b
Balaka	Lemu	84	2488	±186 a	4172	±254b	4115	±208 b
Balaka	Malula	105	3050	±247 a	4059	±226b	4006	±268 ab
Dowa	Chipeni	147	5733	±308 a	5690	±245a	5959	±275 a
Dowa	Chisepo	81	4063	±301 a	5028	±337b	4953	±386 a
Enyezini	Enyezini	27	3252	±243 a	2895	±584a	2642	±189 a
Machinga	Matandika	87	3442	±324 a	5029	±356b	5164	±364 b
Nkhotakota	Linga	69	4164	±334 a	5076	±377a	5191	±343 b
Nkhotakota	Mwansambo	141	3890	±229 a	4405	±252a	4542	±262 a
Nkhotakota	Zidyana	153	3988	±236 a	5247	±261b	4915	±242 b
Salima	Chinguluwe	54	4898	±340 a	6343	±313b	5478	±217 a
Zomba	Songani	30	4490	±488 a	5762	±688a	5659	±542 a
Weighted mean ± SE*			3934	±270	4862	±296	4803	±283

Notes: *The number of replicates in each community are determined by dividing the sample size in each community by 3 (the number of treatments; check CA and CA + legumes). Thus, in Herbert, there were 18 on-farm replicate trials. Weighted average across all sites (±SE of the weighted mean) based on sample size (*n*) of each site. Rows sharing the same letter are not different at the 5% level of significance (t-test).

the expected monetary value with the difference – the ‘risk premium’ – reflecting the amount of the expected payoff one would be willing to forgo to avoid risk. In terms of policies promoting the adoption of agronomic technologies among risk-averse producers, accurate estimation of risk premium may be useful for program budgeting, extension efforts and planning purposes.

RESULTS AND DISCUSSION

Agronomic results

Maize grain yields were highest under both CA systems across almost all target communities (Table 3). Explanations of yield differences between the three treatments across communities have been summarized by Thierfelder *et al.* (2013); here we only show overall yields across communities since maize grain yields are used as a basis of calculating economic returns. On average, maize yields on CA treatments out-yielded conventional control plots by 23.6% (CA) and 22.1% (CA + legume) over all years and communities. The highest yields for the CA system with monocrop maize were recorded in Chinguluwe (6343 kg ha⁻¹), Songani (5767 kg ha⁻¹) and Chipeni (5690 kg ha⁻¹) whereas the lowest yields were produced in Enyezini (2895 kg ha⁻¹) and Herbert (3533 kg ha⁻¹). For CA with maize–legume intercropping system, the highest yields were achieved in Chipeni (5959 kg ha⁻¹), Songani (5659 kg ha⁻¹) and Chinguluwe (5478 kg ha⁻¹) whereas the lowest yields were achieved again in Enyezini (2642 kg ha⁻¹) and Herbert (3249 kg ha⁻¹).

Table 4. Summary statistics comparing net returns ha^{-1} for conventional, conservation agriculture (CA) and CA + legume intercrop technologies, Malawi, 2005–2011.

	Check	CA	CA + legumes
Combined:		2010 US\$ ha^{-1} *	
Mean	562	732	662
CV [†]	98%	79%	81%
Standard deviation	551	577	537
Minimum	–296	–173	–193
Maximum	2848	3012	2476
<i>n</i>	344	344	344
≤750 m:			
Mean	446	659	579
CV [†]	116%	86%	87%
Standard deviation	518	569	506
Minimum	–329	–173	–193
Maximum	2905	3073	2525
<i>n</i>	249	249	249
>750 m:			
Mean	912	979	931
CV [†]	60%	59%	62%
Standard deviation	550	575	573
Minimum	–17	–136	–161
Maximum	2504	2407	2410
<i>n</i>	95	95	95
Non-parametric comparison of empirical distributions			
Combined:		D-statistic [‡]	Pr (D ≥ 0)
Check vs. CA		0.17	0.0001
Check vs. CA + legumes		0.14	0.0028
CA vs. CA + legumes		0.06	0.5934
≤750 m:			
Check vs. CA		0.21	0.0001
Check vs. CA + legumes		0.19	0.0002
CA vs. CA + legumes		0.07	0.5925
>750 m:			
Check vs. CA		0.15	0.2337
Check vs. CA + legumes		0.14	0.3133
CA vs. CA + legumes		0.06	0.9892

Notes: *2010 US\$.

[†]Coefficient of variation.[‡]Distance statistics, Kolmogorov–Smirnov non-parametric test of equality between empirical distributions.

Net returns

On average, highest maize net returns ha^{-1} were produced under the CA monocrop maize (732 ± 1.73 , mean \pm standard error of the mean, 2010 US\$), followed by the net returns ha^{-1} from the CA maize–legume intercrop system (662 ± 1.61) and then the net returns ha^{-1} from conventional tillage (562 ± 1.65 ; Table 4). Pairwise t-tests suggest that observed differences in expected net returns ha^{-1} were significant ($p < 0.01$ for all comparisons). The coefficient of variation of net returns

ha⁻¹ for the CA monocrop maize was lowest (79%), followed by the CA maize–legume intercrop (81%) and then the conventional tillage technology (98%; Table 4). There were no ties in the CVs, therefore the mean-variance rankings are the same as those categorized by the relative risk criteria. By these criteria it appears that risk-averse farmers would prefer both CA technologies.

Stochastic dominance

The stochastic dominance analysis corresponds to the relative ranking and mean-variance criteria. Based on the generalized stochastic dominance analysis, the CA monocrop maize would be preferred by risk neutral and risk-averse individuals over the CA maize–legume intercrop and continual tillage systems. The CA maize–legume system would be preferred over the conventional tillage system when these two technologies were compared side-by-side.

Net returns ha⁻¹ were lowest for the conventional tillage practice (–US\$296 ha⁻¹; Table 4 and Figure 1); therefore conventional tillage practices cannot dominate the CA systems by the first or second decision rules. Risk-averse producers would always rank the conventional technology last compared with the two CA systems. By similar reasoning the CA monocrop maize would be preferred to the CA maize–legume intercrop system by risk-averse individuals, even though the empirical distributions were not significantly different (Kolmogorov–Smirnov D-statistic = 0.06, $p = 0.59$; Table 4). The empirical distributions of the CA systems were significantly different from the empirical distribution of the net returns ha⁻¹ produced under the conventional tillage system ($p < 0.01$; Table 4). Thus, risk averse and risk neutral individuals would prefer both CA systems over the conventional system technology, with the CA maize–legume system ranking second.

Target return sensitivity analysis

The target return analysis uses empirical distributions (Figure 1) to estimate the probability of achieving a target value (Figure 2). The CA monocrop maize and the CA maize–legume intercrop systems yielded the most optimistic return profiles at all target net return levels. For example, the likelihood of a net revenue outcome occurring below breakeven levels was 14% on the conventional system, followed by 9% for the CA maize–legume system and 5% for the CA monocrop maize. Thus, the smaller the difference between the probability of achieving the breakeven and upper target values, the more likely returns ha⁻¹ exceed the breakeven profit. Consider, for example, the upper target of US\$600 ha⁻¹. In this scenario, the likelihood of observing net returns falling above the breakeven-upper target interval for the conventional tillage practice was 49% whereas for the CA monocrop maize and CA maize–legume intercrop technologies the probabilities of this occurring were 43% and 44% respectively. Thus, the likelihood of exceeding the US\$600 ha⁻¹ target would be $1.00 - 0.44 - 0.05 = 51\%$ and $1.00 - 0.43 - 0.09 = 48\%$ for the CA monocrop maize and CA maize–legume systems respectively. In this scenario, surpassing the upper level of US\$600 ha⁻¹ target, given the conventional tillage system, was lowest at 37%.




Figure 1. Empirical distribution of net returns (2010 US\$ ha⁻¹) from conventional ('Control'), conservation agriculture (CA) and CA + intercropped legume cultivation; Balaka, Dowa and Nkhotakota districts, Malawi, 2005–2011.




Figure 2. Target net returns ha⁻¹ and associated probabilities for each technology.

The aggregate areas generated by the target profiles also provide some indication of the degree of risk associated with the expected net returns ha^{-1} (Figure 2, inset values and corresponding percentages). For example, over the simulated outcomes, net returns ha^{-1} surpassed target returns 55% and 51% of the time for the CA monocrop maize and CA maize–legume intercrop systems whereas the return target surpassed only 44% of the time for the conventional tillage system.

Certainty equivalents and risk premiums

Certainty equivalents (the amount a risk-averse individual would accept instead of a risky outcome, in 2010 US\$ ha^{-1}) preserved the rankings found in the mean variance and stochastic dominance analyses (Figure 3). The certainty equivalent associated with the conventional tillage technology was driven to zero more quickly than those of the alternative technologies because the returns were relatively more risky ($\text{CE} = 0$ at $\text{RAC} = 0.0177$). Certainty equivalents were driven to zero for the CA monocrop maize and CA maize–legume intercrop at $\text{RAC} = 0.0237$ and 0.0296 respectively. Thus, the ‘risk-preferred’ technologies are those where the certainty equivalent remains positive as risk aversion increases. That the certainty equivalent curves do not overlap supports the technology rankings determined by the stochastic dominance analysis.

What dollar amount would producers be indifferent between CA monocrop maize, a CA maize–legume intercrop system and conventional tillage systems? Using the CEs calculated for each technology at the upper RAC bound of 0.0296 , if producers who already adopted CA monocrop maize were offered US\$40.42 (US\$105.45), then they would be indifferent between using the CA monocrop maize and the CA maize–legume intercrop (conventional tillage) systems (Figure 3). Focusing on the next dominant technology – the CA maize–legume intercropping – producers would be indifferent between using this technology and a conventional tillage system if they were offered US\$65.03 (Figure 3). In other words, from the perspective of risk-averse smallholder farmers in Malawi, these confidence premiums suggest a reasonable valuation of the CA monocrop maize and CA maize–legume intercrop systems compared with conventional tillage systems. As evidenced in Figure 3, the risk premiums associated with the CA technologies are positive when compared with the conventional technology at all levels of risk aversion. An important caveat is that these findings are based on pooled (combined) data and do not take into consideration potential differences in the performance of these technologies in different elevations in terms of production. We explore this caveat, keeping in mind that yield differences generated by the technologies were less evident in the high elevation sites (Figure 1).

Sensitivity analysis: net return distributions and elevation

We hypothesize that the degree to which technologies will be risk-preferred may differ according to the growing conditions typical of agroecological zones. We test this hypothesis by conducting the stochastic dominance and certainty equivalent analyses on the subsets of the data, categorizing the study sites into two agroecological zones according to elevation: communities situated at elevations less than 750 m (low altitude




Figure 3. Certainty equivalents (2010 US\$ ha⁻¹) and confidence premium associated with each technology.

sites), and those located above this level (mid-altitude sites; Table 1) (Brown and Young, 1966).

Net returns for maize produced using conventional tillage were 30% and 47% lower than net returns generated by the CA and CA + legume systems in lower elevation communities (Table 4). Pairwise t-tests suggest that these differences are significant at the 5% level for all comparisons. The CV of net returns ha^{-1} for the CA technologies were similar (about 87%), and considerably lower than the net returns ha^{-1} CV associated with the conventional maize production technology. By these criteria it appears that risk-averse farmers in low-elevation communities would prefer both CA technologies. The stochastic dominance ordering was similar to the pooled data results, with the conventional practice dominated by the CA technologies (Figure 1, Table 4), as were the results of the certainty equivalent analysis (Figure 3). At the RAC level of 0.029, producers in the lower altitude region would be indifferent between using the CA monocrop system and a conventional tillage system if they were offered US\$137.71; whereas for the CA + legume system the risk premium was US\$89.05. Consistent with the stochastic dominance results, the risk premium relative to the check plot was always positive across a range of RAC (Figure 3).

In the higher elevation communities, differences between the net return distributions were less evident (Figure 1, Table 4). The net returns from the conventional and CA monocrop plots were not different at the 5% level, and the CVs of the conventional and monocrop CA net returns were similar (about 60%). In contrast to the results based on the pooled data, the CA technologies do not exhibit stochastic dominance over the conventional technology as evidenced by the D-statistics of the Kolmogorov–Smirnov tests and the net return distributions, and also because net return minimum of the CA technologies was lower than the minimum net revenue of the conventional system (Figure 1). Net returns are clearly higher for the CA systems (relative to the conventional tillage system) between 25th and 50th percentiles; otherwise, the three distributions cross numerous times. At the higher elevations, risk neutral producers (RAC = 0) and producers with relatively low aversion to risk (RAC < 0.0012) would still prefer the CA technologies over the conventional system. The risk premium associated with the CA technologies becomes negative as risk aversion increases above an RAC of 0.0012. In other words, in the higher elevation, the more risk-averse a producer is, the higher the incentive would have to be for them to switch from a conventional tillage/maize production system to either of the CA systems. Clearly, these results are driven by the yield profiles of the technologies (Figure 1). At higher elevations, the distinction between the technologies disappears under excellent growing conditions.

CONCLUSIONS

This study examined the riskiness of net returns from maize production using monocrop maize and maize–legume intercropping CA technologies based on six-year on-farm production data in Malawi (2005–2011). The non-parametric methods used in this analysis, stochastic dominance and risk premium determination, account for upside variability in returns without resorting to strong distribution assumptions.

Maize grain yields in different treatments did not follow a similar trend across the target communities confirming the need to target and adapt conservation agriculture to local conditions (Rusinamhodzi *et al.*, 2011; Wall, 2007). In low rainfall, low altitude areas of Balaka and Machinga, conservation agriculture resulted in yield increases of 31.3% and 67.7%, respectively, over conventional tillage systems. In high rainfall, mid-altitude areas of Chipeni and Enyezini, conservation agriculture resulted in productivity decrease of 1.6% and 14.9%, respectively compared to conventional tillage systems. The lower differences in yields according to elevation had important implications with respect to analyzing the riskiness of the CA technologies compared to conventional tillage systems. Using the pooled data (low and high elevations combined), stochastic dominance analysis of the net returns ha^{-1} for maize production suggest that direct planting with solely maize and maize intercropped with legumes would be preferred by risk-averse producers compared to returns from maize produced using conventional methods. These findings were corroborated with a sensitivity analysis where the risk premiums associated with conservation agriculture and conventional maize production systems were estimated and compared. The differences between the conventional and conservation agriculture risk premium suggest that risk-averse producers would generally place more value on CA technologies (in terms of eliminating return risk). Given target return levels, the CA technologies analysed here could also be risk-preferred in terms of the likelihood target returns are obtainable. These findings were generally consistent when the yield data from lower altitude sites were analysed separately. However, the narrow margin in the percentage change in yield differences between technologies translated into profound differences in the risk analysis distribution comparisons analysing the higher elevation sites. At the higher elevation sites, under exceptional growing conditions the differences in net returns from the technologies appear less evident, and the case that CA technologies are 'risk-reducing' (in terms of expected net returns) becomes less clear. Indeed, the risk premiums determined by the ex-post analysis suggest that stronger incentives may be required to entice adoption of CA technologies at higher elevations. The reversal in preference ordering of the technologies is likely driven by the similarity in yields between the CA technologies studied relative to the conventional tillage system used as a reference practice.

Although based on six years of production data over a variety of growing conditions, the risk analysis only provides an ex-post profile assessment of the technologies. The results should be put into perspective in terms of the price data used in the analysis. Future studies could analyse the sensitivity of these findings with respect to stochastic prices using the Monte Carlo analysis. A more in-depth analysis would also entail correlating these findings with individual preferences. This would require detailed household socioeconomic and farm production obtained from household surveys. In addition, while some insight into the value of CA technologies was provided by the analysis, more structured interviews would be needed to generate the willingness-to-adopt estimates. There are other factors that are important in farmers' decisions to adopt conservation agriculture. For example Kassam *et al.* (2009) conclude that widespread sustained adoption of conservation agriculture

will require a *deeper understanding* by producers of the ecological systems which drive residue management and soil fertility dynamics. This deeper understanding suggests that farmers need to understand that the payoff from CA systems may not occur during the first year but in subsequent years once the improvement in soil properties are manifested. Comprehension of these biophysical interactions by producers underscores the relatively large investments (e.g. in fertilizers, equipment, herbicides, education and farmer training) needed to enhance farmer success with CA systems. Extensive adoption of conservation agriculture by small-scale farmers has also been constrained by institutional capacity for transferring new knowledge to agriculturalists (Gowing and Palmer, 2008) as well as lack to deliver new practical knowledge. Additional barriers to CA adoption include removal of crop residues for fodder, lack of appropriate seed drills that can sow in unploughed fields, high crop residues and the lack of availability or prohibitively high costs of herbicides or other inputs. On top of these problems are weak or nonexistent credit markets for purchasing inputs and markets for selling product (Giller *et al.*, 2009; Lal, 2009; Silici, 2010). Without reliable knowledge transfer systems and resilient social capital networks (e.g. Silici, 2010), partial, localized and sporadic adoption of CA technologies will likely to continue to be the norm (Gowing and Palmer, 2008; Mazvimavi *et al.*, 2008).

Finally, the impact of policies designed to extend the adoption of CA technologies may depend more on the ability of incentives to compensate producers for anticipated losses and the extent to which information is shared rather than on the 'band-wagon' effect produced by early adopters or targeted incentives (Baerenklau, 2005). Risk preferences and learning are key determinants in the adoption decision whereas peer-group influence could be less relevant (Baerenklau, 2005). The risk premium found here are first steps towards this end, acknowledging that a more in-depth microeconomic analysis of household and production data would be needed. Access to inputs may also be an important determinant of how would-be adopters welcome or dismiss CA technologies. Gowing and Palmer (2008) concluded that conservation agriculture would deliver the productivity gains required to achieve food security and poverty targets only if farmers have access to fertilizers and herbicides and knowledge how to use them. For labour-intensive production systems that require extensive weeding during critical growth periods and access to herbicides may provide impetus to integrate CA technologies into smallholder production systems. If net returns of farmers practicing conservation agriculture are not higher than from conventional practice but the ecosystem services (reduced soil erosion or carbon sequestration) are enhanced, then risk premium analyses may help formulate incentives provided by the government or non-government organizations. Such information bears important messages for technology promoters and policy makers.

Acknowledgements. We wish to thank the farmers, Total Land Care staff and Agricultural Extension Development Officers (AEDO) of Chinguluwe, Chipeni, Chisepo, Enyezini, Herbert, Lemu, Linga, Malula, Matandika, Mwansambo, Songani and Zidyana sections for their enthusiasm, collaboration and support during project implementation phase. Special thanks go to John Chisui, Richard Museka, Mphatso

Gama, Ivy Ligowe, Florence Kamwana and Pacsu Simwaka for their tireless efforts in coordination of project activities. We also wish to acknowledge the International Maize and Wheat Improvement Centre (CIMMYT) for logistical support, and the German Technical Cooperation (GTZ) and International Fund for Agriculture Development (IFAD) for funding project activities.

REFERENCES

- Anderson, J. (1974). Risk efficiency in the interpretation of agricultural production research. *Review of Marketing and Agricultural Economics* 42:131–184.
- Anderson, J. R. and Dillon, J. L. (1990). *The Analysis of Response in Crop and Livestock Production*. New York: Pergamon Press.
- Baerenklau, K. A. (2005). Toward an understanding of technology adoption: risk, learning, and neighborhood effects. *Land Economics* 81(1):1–19.
- Bekele, W. (2005). Stochastic dominance analysis of soil and water conservation in subsistence crop production in the eastern Ethiopian highlands: the case of the Hunde-Lafto area. *Environmental & Resource Economics* 32(4):533–550.
- Brown, P. and Young, A. (1966). *The Physical Environment of Central Malawi: With Special Reference to Soils and Agriculture*. Zomba, Malawi: Government Press.
- Cavatassi, R., Lipper, L. and Narloch, U. (2011). Modern variety adoption and risk management in drought prone areas: insights from the sorghum farmers of eastern Ethiopia. *Agricultural Economics* 42(3):279–292.
- Doran, J. W., Elliott, E. T. and Paustian, K. (1998). Soil microbial activity, nitrogen cycling, and long-term changes in organic carbon pools as related to fallow tillage management. *Soil & Tillage Research* 49(1–2):3–18.
- Ellis, F., Kutengule, M. and Nyasulu, A. (2003). Livelihoods and rural poverty reduction in Malawi. *World Development* 31(9):1495–1510.
- FAO. (2002). *Conservation Agriculture: Case Studies in Latin America and Africa*. FAO Soils Bulletin Nos. 77, 78. Rome, Italy: FAO.
- FAOSTAT. (2010). <http://faostat3.fao.org/home/index.html>; last accessed 20 July 2010.
- Giller, K. E., Witter, E., Corbeels, M. and Tittonell, P. (2009). Conservation agriculture and smallholder farming in Africa: the heretics' view. *Field Crops Research* 114(1):23–34.
- Gowing, J. W. and Palmer, M. (2008). Sustainable agricultural development in sub-Saharan Africa: the case for a paradigm shift in land husbandry. *Soil Use and Management* 24:92–99.
- Guto, S. N., Pypers, P., Vanlauwe, B., de Ridder, N. and Giller, K. E. (2011). Tillage and vegetative barrier effects on soil conservation and short-term economic benefits in the Central Kenya highlands. *Field Crops Research* 122(2):85–94.
- Ito, M., Matsumoto, T. and Quinones, M. A. (2007). Conservation tillage practice in sub-Saharan Africa: the experience of Sasakawa Global 2000. *Crop Protection* 26:417–423.
- Kassam, A., Friedrich, T., Shaxson, F. and Pretty, J. (2009). The spread of conservation agriculture: justification, sustainability and uptake. *International Journal of Agricultural Sustainability* 7(4):292–320.
- Kumwenda, J. D. T., Waddington, S. R., Snapp, S. S., Jones, R. B. and Blackie, M. J. (1997). Soil fertility management in Southern Africa. In *Africa's Emerging Maize Revolution*, 157–172 (Eds D. Byerlee and C. K. Eicher). Boulder, CO: Lynne Rienner.
- Lal, R. (1986). Soil surface management in the tropics for intensive land use and high and sustained production. *Advances in Soil Sciences* 5:1–109.
- Lal, R. (2009). The plow and agricultural sustainability. *Journal of Sustainable Agriculture* 33(1):66–84.
- Lambert, D. M. and Lowenberg-DeBoer, J. (2003). Economic analysis of row spacing for corn and soybean. *Agronomy Journal* 95:564–573.
- Mando, A., Brussaard, L. and Stroosnijder, L. (1999). Termite- and mulch-mediated rehabilitation of vegetation on crusted soil in West Africa. *Restoration Ecology* 7(1):33–41.
- Mas-Colell, A., Whinston, M. D. and Green, J. R. (1995). *Microeconomic Theory*. Oxford, UK: Oxford University Press.
- Mazvimavi, K. and Twomlow, S. (2009). Socioeconomic and institutional factors influencing adoption of conservation farming by vulnerable households in Zimbabwe. *Agricultural Systems* 101(1–2):20–29.
- Mazvimavi, K., Twomlow, S. J., Belder, P. and Howe, L. (2008). An assessment of the sustainable uptake of conservation farming in Zimbabwe. *Global Theme on Agroecosystems Report*, Vol. 39, 66pp. Bulawayo, Zimbabwe: International Crops Research Institute for the Semi-Arid Tropics.

- McGarry, D., Bridge, B. J. and Radford, B. J. (2000). Contrasting soil physical properties after zero and traditional tillage of an alluvial soil in the semi-arid subtropics. *Soil and Tillage Research* 53(2):105–115.
- Meyer, J. (1977). Choice among distributions. *Journal of Economic Theory* 14:326–336.
- MoAFS. (2011). *Ministry of Agriculture and Food Security Crop Production Estimates (MoAFS)*. Lilongwe, Malawi: MoAFS.
- Moschini, G. and Hennessy, D. A. (2001). Uncertainty, risk aversion, and risk management for agricultural producers. In *Handbook of Agricultural Economics*, 88–153 (Eds B. L. Gardner and G. C. Rausser). New York: Elsevier.
- Moser, C. M. and Barrett, C. B. (2006). The complex dynamics of smallholder technology adoption: the case of SRI in Madagascar. *Agricultural Economics* 35(3):373–388.
- Ngwira, A. R., Aune, J. B. and Mkwinda, S. (2012). On-farm evaluation of yield and economic benefit of short-term maize legume intercropping systems under conservation agriculture in Malawi. *Field Crops Research* 132:149–157.
- NSO. (2012). *Integrated Household Survey 2010–2011 – Household Socioeconomic Characteristics Report*. Zomba, Malawi: NSO.
- Parr, J. E., Stewart, B. A., Hornick, S. B. and Singh, R. P. (1990). Improving the sustainability of dryland farming systems: a global perspective. In *Dryland Agriculture Strategies for Sustainability*, Vol. 13, 1–8 (Eds R. P. Singh, J. F. Parr and B. A. Stewart). New York: Springer-Verlag New York Inc.
- Richardson, J. W. (2002). *SIMETAR, Simulation Software for Applied Risk Management*. College Station, TX: Department of Agricultural Economics, Texas A&M University.
- Richardson, J. W. and Mapp, H. P. (1976). Use of probabilistic cash flows in analyzing investment under conditions of risk and uncertainty. *Journal of Agricultural Economics* 8(2):19–24.
- Richardson, J. W., Schumann, K. D. and Feldman, P. A. (2006). *Simetar*® *Simulation for Applied Risk Management*. College Station, TX: Texas A&M University.
- Rockström, J., Barron, J. and Fox, P. (2002). Rainwater management for increased productivity among smallholder farmers in drought prone environments. *Physics and Chemistry of the Earth* 27(11–22):949–959.
- Rockström, J., Kaumbutho, P., Mwalley, J., Nzabi, A. W., Temesgen, M., Mawenya, L., Barron, J., Mutua, J. and Damgaard-Larsen, S. (2009). Conservation farming strategies in East and Southern Africa: yields and rain water productivity from on-farm action research. *Soil and Tillage Research* 103(1):23–32.
- Roth, C. H., Meyer, B., Frede, H. G. and Derpsch, R. (1988). Effect of mulch rates and tillage systems on infiltrability and other soil physical properties of an Oxisol in Paraná, Brazil. *Soil and Tillage Research* 11(1):81–91.
- Rusinamhodzi, L., Corbeels, M., Wijk, M., Rufino, M., Nyamangara, J. and Giller, K. (2011). A meta-analysis of long-term effects of conservation agriculture on maize grain yield under rain-fed conditions. *Agronomy for Sustainable Development* 31(4):657–673.
- Serra, T. (2008). Differential uncertainties and risk attitude between conventional and organic producers: the case of Spanish COP farmers. *Paper Presented at American Agricultural Economics Association Annual Meeting*, July 29–August 1, 2007, Portland, OR.
- Shively, G. E. (2001). Poverty, consumption risk, and soil conservation. *Journal of Development Economics* 65(2):267–290.
- Silici, L. (2010). *Conservation Agriculture and Sustainable Crop Intensification in Lesotho*, Integrated Crop Management Series, Vol. 10. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO).
- Smale, M., Kaunda, Z. H. W., Makina, H. L., Mkandawire, M. M. M. K., Msowoya, M. N. S., Mwale, D. J. E. K. and Heisey, P. W. (1991). Chimanga cha makolo, hybrid and composites: an analysis of farmer's adoption of technology of maize technology in Malawi, 1989–91. *CIMMYT Economic Working Paper 91/04*, Mexico, DF.
- Thierfelder, C., Chisui, J. L., Gama, M., Cheesman, S., Jere, Z. D., Trent Bunderson, W., Eash, N. S. and Rusinamhodzi, L. (2013). Maize-based conservation agriculture systems in Malawi: long-term trends in productivity. *Field Crops Research* 142(0):47–57.
- Thierfelder, C. and Wall, P. C. (2009). Effects of conservation agriculture techniques on infiltration and soil water content in Zambia and Zimbabwe. *Soil and Tillage Research* 105(2):217–227.
- Thierfelder, C. and Wall, P. C. (2010). Rotation in conservation agriculture systems in Zambia: effects on soil quality and water relations. *Experimental Agriculture* 46(3):309–325.
- Verhulst, N., Nelissen, V., Jespers, N., Haven, H., Sayre, K. D., Raes, D., Deckers, J. and Govaerts, B. (2011). Soil water content, maize yield and its stability as affected by tillage and crop residue management in rainfed semi-arid highlands. *Plant and Soil* 344(1–2):73–85.
- Vogel, H. (1993). An evaluation of five tillage systems for smallholder agriculture in Zimbabwe. *Der Tropenlandwirt* 94:21–36.
- Wall, P. (2007). Tailoring conservation agriculture to the needs of small farmers in developing countries: an analysis of issues. *Journal of Crop Improvement* 19:137–155.
- Wendland, K. J. and Sills, E. O. (2008). Dissemination of food crops with nutritional benefits: adoption and disadoption of soybeans in Togo and Benin. *Natural Resources Forum* 32(1):39–52.
- WRB. (1998). *World Reference Base for Soil Resources*, Vol. 88. Rome, Italy: FAO-ISRIC.