
BRIEF COMMUNICATION

Beyond Measures of Central Tendency: Novel Methods to Examine Sex Differences in Neuropsychological Performance Following Sports-Related Concussion in Collegiate Athletes

Victoria C. Merritt¹* , Liora S. Greenberg², Erin Guty², Megan L. Bradson², Amanda R. Rabinowitz³ and Peter A. Arnett²

¹VA San Diego Healthcare System, San Diego, CA 92161, USA

²The Pennsylvania State University, PA 16801, USA

³Moss Rehabilitation Research Institute, Elkins Park, PA 19027, USA

(RECEIVED February 5, 2019; FINAL REVISION June 28, 2019; ACCEPTED July 11, 2019; FIRST PUBLISHED ONLINE September 3, 2019)

Abstract

Objective: The purpose of this study was to examine sex differences in neuropsychological functioning after sports-related concussion using several approaches to assess cognition: mean performance, number of impaired scores, and intraindividual variability (IIV). **Method:** In the study, 152 concussed college athletes were administered a battery of neuropsychological tests, on average, 10 days post-concussion ($SD = 12.75$; $Mdn = 4$ days; $Range = 0–72$ days). Mean performance was evaluated across 18 individual neuropsychological variables, and the total number of impaired test scores ($>1.5 SD$ below the mean) was calculated for each athlete. Two measures of IIV were also computed: an intraindividual standard deviation (ISD) score and a maximum discrepancy (MD) score. **Results:** Analyses of covariance revealed that, compared with males, females had significantly more impaired scores and showed greater variability on both IIV indices (ISD and MD scores) after adjusting for time since injury and post-concussive symptoms. In contrast, no significant effects of sex were found when examining mean neuropsychological performance.

Conclusion: Although females and males demonstrated similar mean performance following concussion, females exhibited a greater level of cognitive impairment and larger inconsistencies in cognitive performance than males. These results suggest that evaluating cognitive indices beyond mean neuropsychological scores may provide valuable information when determining the extent of post-concussion cognitive dysfunction.

Keywords: Cognition, Cognitive impairment, Intraindividual variability, Cognitive dispersion, Sports concussion, Gender differences

INTRODUCTION

As the science regarding sports-concussion management continues to evolve, the need to consider individual differences as possible modifiers of outcome and recovery has remained salient (McCrory et al., 2013, 2017). A wide range of demographic, premorbid, and injury-specific variables have been evaluated in this context (Iverson et al., 2017), with biological sex emerging as a primary variable of interest. Despite numerous studies examining sex and concussion outcome and recovery, the literature remains equivocal as to its significance (Iverson et al., 2017). Further research is necessary to

establish the extent to which males and females experience differential outcomes following sports-related concussion (SRC).

Neuropsychological compromise is perhaps the most well-known sequela of SRC. The broader neuropsychological literature has reported premorbid sex differences in cognition (Lezak, Howieson, Bigler, & Tranel, 2012), and sex differences have also specifically been identified in athletes undergoing baseline neuropsychological testing (Merritt et al., 2017). Given these premorbid differences in cognition, it is imperative to consider the extent to which biological sex influences post-concussion cognitive functioning.

A number of studies have compared males and females on the Immediate Post-Concussion Assessment and Cognitive

*Correspondence and reprint requests to: Victoria C. Merritt, VA San Diego Healthcare System (151B), 3350 La Jolla Village Drive, San Diego, CA 92161, USA. Phone: 858-558-8585 x2670, Fax: 858-642-6340. E-mail: victoria.merritt@va.gov

Testing (ImPACT) cognitive composites, with two recent reviews providing evidence that females tend to demonstrate greater impairment than males on measures of visual memory and reaction time in the acute period following concussion (Covassin, Savage, Bretzin, & Fox, 2018; Merritt, Padgett, & Jak, 2019). Moreover, a meta-analysis by Dougan, Horswill, and Geffen (2014) found that female sex was associated with greater neuropsychological deficits acutely following injury, though specific domains of cognitive functioning were not described. While the exact cause of these differences is not known, neuroanatomical, physiological, and/or hormonal differences between the sexes have been hypothesized as reasons for the performance discrepancies in males and females on cognitive testing following concussion (Covassin et al., 2018; Dougan et al., 2014; Merritt et al., 2019).

Taken together, the above findings suggest that females may demonstrate greater neuropsychological compromise than males, at least acutely, following concussion. However, post-concussion sex differences in cognitive performance have not been consistently reported (Covassin et al., 2018; Merritt et al., 2019), as other factors such as age, education, history of developmental disorders, preexisting mental health and/or neurobehavioral (i.e., “post-concussion” like) symptoms, number of lifetime concussions, and time since injury may also influence cognition (Dougan et al., 2014; Karr, Areshenkoff, & Garcia-Barrera, 2014; Kontos, Sufrinko, Womble, & Kegel, 2016). For example, with respect to number of lifetime concussions, Covassin, Elbin, Kontos, and Larson (2010) showed that males with a history of *multiple* concussions performed more poorly than females with a similar concussion history on the ImPACT visual and verbal memory composites.

With this in mind, the purpose of the present study was to increase understanding of sex differences in neuropsychological functioning following SRC. To minimize the impact of possible confounding variables, males and females were matched on important demographic variables with adjustments for time since injury and post-concussive symptoms. In addition to evaluating mean cognitive performance, we examined neurocognitive impairment (total number of impaired scores across the test battery) and measures of intra-individual variability (IIV) across a comprehensive neuropsychological assessment that included both computerized and paper-and-pencil measures. We hypothesized that females would show poorer neurocognitive performance and greater dispersion across the neurocognitive test battery relative to males.

METHOD

Participants and Procedures

Participants included 152 concussed college athletes who were enrolled in a clinically based sports-concussion management program at a large university. Team physicians are responsible for identifying and diagnosing concussions at the time of injury and referring athletes for

neuropsychological evaluations post-concussion. Athletes who are referred to the sports-concussion management program undergo a post-concussion clinical interview and a comprehensive neuropsychological assessment for clinical purposes, and athletes simultaneously consent to having their clinical data be used for research purposes. (To date, no athletes have declined using their clinical data for research). During the clinical interview, athletes are asked about the current concussion (i.e., presence and duration of loss of consciousness [LOC] and retrograde/anterograde amnesia, mechanism of injury, date of injury, etc.) and any previous concussions. The clinical interview is conducted by a PhD-level clinical neuropsychologist or a trained doctoral student, and the neuropsychological assessment is administered by a trained doctoral student or an undergraduate research assistant under the supervision of a PhD-level clinical neuropsychologist. All athletes referred for post-concussion evaluations were seen individually, and assessment procedures (including completion of the clinical interview and neuropsychological tests) took approximately 2 hr.

For the purpose of this study, a concussion was defined using criteria set forth by Ruff et al. (2009) (i.e., LOC less than or equal to 30 min, retrograde or anterograde amnesia lasting less than 24 hr, *or* an alteration in mental status at the time of injury). In order to be included in the present study, athletes must have (1) sustained a concussion according to the above criteria; (2) completed the post-concussion testing as soon as clinically indicated following their injury, but no greater than 3 months post-injury; and (3) passed performance validity testing (see “Measures” section for validity criteria). All participants included in the study provided informed consent, and the study was approved by the university’s institutional review board.

Measures

A comprehensive neuropsychological assessment, including traditional paper-and-pencil measures and computerized testing, was administered to all athletes. The Wechsler Test of Adult Reading (WTAR) was administered to assess premorbid functioning. Additional measures broadly assessing learning and memory, attention, processing speed, and executive functioning included the Brief Visuospatial Memory Test-Revised (BVMT-R), Comprehensive Trail-Making Test (CTMT), a modified version of the Digit Span subtest from the Wechsler Adult Intelligence Scale—Third Edition (WAIS-III), Hopkins Verbal Learning Test-Revised (HVLTR), Penn State University (PSU) Cancellation Test, Stroop Color-Word Test (SCWT), Symbol-Digit Modalities Test (SDMT), and Vigil/W Continuous Performance Test. Finally, the ImPACT computerized program was administered, which generates five composite scores, including four core cognitive composites—Verbal Memory, Visual Memory, Visual Motor Speed, and Reaction Time—and a fifth composite, the Impulse Control Composite (ICC), which serves as a measure of effort

or performance validity. For the purpose of the present study, invalid test performance was defined as scoring above 30 on the ImPACT ICC (Lovell, 2016). The ImPACT also includes the 22-item Post-Concussion Symptom Scale (PCSS), from which a total symptom score was computed.

Neuropsychological Data Transformations

Data transformations of the neurocognitive variables were carried out using procedures described previously (Merritt et al., 2017; Rabinowitz & Arnett, 2013). Briefly, 18 neurocognitive variables were selected from the above measures (see Table 2) and converted from raw scores to standard scores ($M = 100$, $SD = 15$) using sex-specific means and standard deviations from a large normative sample of college athletes at baseline (Merritt et al., 2017). All standard scores were calculated so that higher values reflect better performance.

Neurocognitive performance was evaluated using three techniques: (1) mean performance, (2) number of impaired scores, and (3) IIV.

Mean neurocognitive performance was evaluated across the 18 individual neuropsychological variables. To determine the number of impaired scores across the neurocognitive test battery, the total number of scores falling below a designated impairment level ($>1.5 SD$ below the mean) was counted for each participant (possible range: 0–18). Although there is no universally accepted definition of “impairment,” many prior studies have utilized $>1.5 SD$ below the mean to reflect impaired performance (Iverson & Brooks, 2011). Moreover, prior research examining base rates of impairment in a sample similar to the present study (i.e., collegiate athletes) also utilized the $>1.5 SD$ below the mean threshold to indicate impaired performance (Arnett et al., 2014). Finally, two measures of IIV were calculated: (1) an intraindividual standard deviation (ISD) score and (2) a maximum discrepancy (MD), or range, score. As per Merritt, Rabinowitz, and Arnett (2018), the ISD score was calculated for each athlete by taking the standard deviation of the standard scores across all 18 neuropsychological variables. The MD score was calculated by subtracting each athlete’s lowest cognitive test score from their highest test score. The MD score thus reflects the range of an athlete’s overall cognitive performance, accounting for their absolute highest score and lowest score in the test battery. For both IIV indices, higher scores represent greater IIV, or inconsistency in performance.

Data Analyses

Descriptive statistics were conducted on the overall sample, and males and females were compared on demographic and injury characteristics (independent samples t tests for continuous variables and chi-square or Fisher’s exact tests for categorical variables). Analyses of covariance (ANCOVAs) adjusting for time since injury and post-concussive symptoms were used to evaluate whether males

and females differed across the 18 individual neurocognitive variables. Given the number of analyses proposed, we determined *a priori* that we would use $p < .01$ as our criterion for significance for all analyses pertaining to the individual cognitive variables to reduce the likelihood of Type 1 error. We chose this approach over the Bonferroni correction, which is recognized as overly conservative in many applications, and increases Type II error rates (Rothman, 1990). Finally, ANCOVAs adjusting for time since injury and post-concussive symptoms were conducted to compare males and females on the neurocognitive summary indices (i.e., number of impaired scores and the ISD and MD scores). Given the limited number of planned *a priori* analyses (three in total), we used $p < .05$ as our criterion for significance to interpret these latter findings. Effect sizes are listed as partial eta-squared values (η_p^2), interpreted as small = .01, medium = .06, and large = .14. All analyses were conducted using the Statistical Package for the Social Sciences (SPSS; Version 25).

RESULTS

Participant Demographic and Injury Characteristics

The overall sample ($N = 152$) was 75.7% male ($n = 115$) and 24.3% female ($n = 37$). On average, participants were 20.13 years of age ($SD = 1.38$) and had completed 13.60 years of education ($SD = 1.24$). Athletes included in the present study participated in the following sports: football ($n = 46$), lacrosse ($n = 23$), basketball ($n = 21$), ice hockey ($n = 14$), rugby ($n = 14$), soccer ($n = 12$), wrestling ($n = 10$), and other ($n = 12$). Participants were tested, on average, 9.59 days ($SD = 12.75$; $Mdn = 4$ days; $Range = 0–72$ days) following their concussion (66.4% of athletes were tested within 1 week post-injury; 80.9% were tested within 2 weeks post-injury). At the time of the post-concussion assessment, no athletes had returned to play.

Table 1 displays participant demographic and injury characteristics by sex. There were no significant differences between males and females with regard to age, years of education, premorbid IQ, and post-concussive symptoms. However, males and females differed with regard to the timing of their neuropsychological assessment, such that females were tested, on average, over 2 weeks post-injury whereas males were tested, on average, 1 week post-injury.

Neuropsychological Assessment: Individual Scores

There were no significant differences between males and females in mean performance on any of the individual neurocognitive variables after adjusting for time since injury and post-concussive symptoms ($p = .012–.816$; $\eta_p^2 = .00–.04$, small effect sizes). See Table 2.

Table 1. Participant demographic and injury characteristics

Variables	Males (<i>n</i> = 115)		Females (<i>n</i> = 37)		<i>p</i> ^a
	M	SD	M	SD	
Age	20.17	1.43	20.00	1.22	.474
Education (years)	13.56	1.31	13.73	0.99	.398
WTAR FSIQ	104.39	7.19	104.58	7.43	.890
Time since injury (days)	7.12	10.65	17.24	15.59	.001
PCSS total score	8.67	12.41	13.62	17.12	.110
	<i>N</i>	%	<i>N</i>	%	<i>p</i> ^b
Race					
Caucasian	76	66.1	30	81.1	
Non-Caucasian	37	32.2	7	18.9	.109
Not reported/missing	2	1.7	0	0	
History of ADHD/LD					
Yes	9	7.8	0	0	
No	103	89.6	35	94.6	.162
Unknown	3	2.6	2	5.4	
History of concussion					
Yes	64	55.7	20	54.1	
No	49	42.6	16	43.2	.909
Not reported/missing	2	1.7	1	2.7	
Loss of consciousness					
Yes	19	16.5	6	16.2	
No	91	79.1	30	81.1	.902
Unknown	5	4.3	1	2.7	
Retrograde amnesia					
Yes	22	19.1	5	13.5	
No	88	76.5	30	81.1	.726
Unknown	5	4.3	2	5.4	
Anterograde amnesia					
Yes	38	33.0	10	27.0	
No	69	60.0	25	67.6	.711
Unknown	8	7.0	2	5.4	

Note. WTAR = Wechsler Test of Adult Reading; FSIQ = Full-Scale IQ; PCSS = Post-Concussion Symptom Scale; ADHD = attention-deficit/hyperactivity disorder; LD = learning disability.

^aIndependent samples *t* tests were used to evaluate whether males and females differed with regard to age, years of education, premorbid intellectual functioning (as assessed by WTAR FSIQ), time since injury, and PCSS total score.

^bChi-square or Fisher's exact tests were used to evaluate whether males and females differed with regard to race, presence of ADHD/LD, history of concussion, presence of loss of consciousness, retrograde amnesia, and anterograde amnesia.

Neuropsychological Assessment: Summary Scores

A significant effect of sex was found for the number of impaired scores after adjusting for time since injury and post-concussive symptoms, such that females exhibited greater neurocognitive impairment than males ($p = .045$; $\eta_p^2 = .03$, small effect size). Additionally, females demonstrated significantly greater IIV than males, as reflected by the ISD ($p = .001$; $\eta_p^2 = .07$, medium effect size) and MD ($p = .004$; $\eta_p^2 = .06$, medium effect size) scores, after adjusting for the same variables; see Table 2.

DISCUSSION

To improve the understanding of sex differences following SRC, the present study evaluated a wide range of neurocognitive outcomes, matching groups on important demographic variables and accounting for time since injury and post-concussive symptoms. This is the first study, to our knowledge, to examine sex differences in cognition using indices beyond mean performance, including neurocognitive impairment and measures of IIV, across a comprehensive neuropsychological battery. Results showed that, compared with males, females had significantly more impaired scores and showed greater cognitive variability across the neuropsychological test battery. However, no significant effects of sex were found when examining mean neuropsychological performance.

Existing literature regarding the influence of biological sex on neurocognitive outcomes following SRC is somewhat equivocal. While previous empirical research has demonstrated that females perform worse than males in the domains of visual memory and reaction time, not all studies have identified sex differences in cognitive functioning post-concussion (Covassin et al., 2018; Merritt et al., 2019). Notably, past studies have focused exclusively on evaluating *mean* neuropsychological performance, and previous work has largely reported on findings pertaining to the ImPACT cognitive battery. Furthermore, potential confounding variables have been inconsistently reported and controlled for in past research.

In the present study, we administered a comprehensive neuropsychological assessment, spanning the domains of learning and memory, attention, processing speed, and executive functioning, and found that males and females demonstrated comparable mean performance across all tests, which is somewhat of a departure from previous studies (Covassin et al., 2018; Merritt et al., 2019). This may be because groups were matched on variables known to influence cognitive functioning, such as age, education, preexisting developmental disorders, and history of concussion (Dougan et al., 2014; Karr et al., 2014; Kontos et al., 2016). Furthermore, we adjusted for time since injury and post-concussive symptoms in our analyses, and excluded for poor effort, which has not always been the case in previous research. Finally, although some of the neurocognitive variables evaluated in the present study (i.e., BVMT-R immediate recall, SDMT incidental memory, SCWT word time) demonstrated a trend in the direction of females performing more poorly than males, the effect size comparisons were small, and not necessarily clinically meaningful.

In contrast to the above null findings, there were significant sex differences when evaluating the neurocognitive summary scores, such that females demonstrated greater impairment and larger inconsistencies in performance across the neuropsychological test battery. Although the effect size for greater impaired scores in females was small, it may be especially meaningful given that past research using a comparable sample of athletes on a similar test battery showed

Table 2. Neurocognitive variables: means and standard deviations by group, ANCOVA results, and effect sizes

Neurocognitive Variables	Males			Females			<i>F</i>	<i>p</i>	η_p^2
	N	M	SD	N	M	SD			
<i>Mean Performance (SS)</i>									
BVMT-R Immediate Recall	115	101.94	13.99	37	94.02	14.30	4.35	.039	.029
BVMT-R Delayed Recall	115	99.08	16.64	37	91.42	22.37	3.38	.068	.022
CTMT 1 Time	115	101.78	15.58	37	97.03	19.93	1.59	.210	.011
CTMT 2 Time	115	99.43	19.80	37	96.72	14.57	.50	.480	.003
HVLT-R Immediate Recall	115	98.80	16.31	37	101.66	18.68	1.45	.231	.010
HVLT-R Delayed Recall	115	95.44	18.88	37	94.77	23.34	.05	.816	.000
ImPACT Verbal Memory Composite	115	98.82	14.80	37	93.78	16.01	2.21	.139	.015
ImPACT Visual Memory Composite	112	99.87	15.84	37	93.46	16.41	3.60	.060	.024
ImPACT Visuomotor Speed Composite	115	100.73	16.35	37	99.71	11.38	.07	.797	.000
ImPACT Reaction Time Composite	115	97.84	18.36	37	101.52	20.42	1.60	.208	.011
PSU Cancellation Test Total Correct	112	98.64	14.32	37	100.29	16.87	.34	.561	.002
SDMT Total Correct	114	99.87	15.82	37	97.94	23.14	.48	.489	.003
SDMT Incidental Memory	114	98.63	17.16	37	92.07	20.56	6.40	.012	.042
SCWT Word Time	114	98.22	16.85	36	89.93	30.59	5.35	.022	.035
SCWT Color-word Time	111	101.60	18.94	37	96.37	19.56	.97	.326	.007
Vigil Average Delay	111	95.50	15.72	37	91.03	21.14	1.19	.277	.008
WAIS-III Digit Span Forward	114	104.47	13.17	37	101.63	19.15	1.94	.166	.013
WAIS-III Digit Span Backward	115	104.26	17.56	37	102.34	16.94	1.02	.314	.007
<i>Neurocognitive Impairment</i>									
Number of Impaired Scores	115	1.58	2.44	37	2.73	3.46	4.10	.045	.027
<i>IIV (Dispersion) Indices</i>									
Intraindividual SD	115	13.45	4.28	37	16.25	4.07	11.20	.001	.070
Maximum Discrepancy	115	51.57	20.24	37	62.52	20.99	8.64	.004	.055

Notes: SS = standard score; BVMT-R = Brief Visuospatial Memory Test-Revised; CTMT = Comprehensive Trail-Making Test; HVLT-R = Hopkins Verbal Learning Test-Revised; ImPACT = Immediate Post-Concussion Assessment and Cognitive Testing; SDMT = Symbol-Digit Modalities Test; SCWT = Stroop Color-Word Test; WAIS-III = Wechsler Adult Intelligence Scale—Third Edition; IIV = intraindividual variability. ANCOVA results are presented above, adjusting for time since injury and post-concussive symptoms (PCSS total score). η_p^2 = effect size interpretation: small = .01, medium = .06, large = .14.

that males had a higher base rate of impairment at baseline than females (Arnett, Meyer, Merritt, & Guty, 2016). With regard to the variability findings, few studies are available for comparison. Among these, reaction time variability as opposed to cognitive dispersion has been examined, and results show greater IIV in adult females relative to males (Dykiert, Der, Starr, & Deary, 2012; Ghisletta, Renaud, Fagot, Lecerf, & De Ribaupierre, 2018). Thus, our finding that females demonstrated greater cognitive dispersion relative to males is consistent with this prior work. In clinical populations outside of concussion, greater IIV has repeatedly been associated with neurobiological integrity (i.e., increased white matter hyperintensities), and could reflect underlying central nervous system dysfunction (Bunce et al., 2013; MacDonald, Nyberg, & Bäckman, 2006). Moreover, there is accumulating evidence to suggest that, relative to mean cognitive performance, IIV may serve as a better predictor of cognitive outcome in a variety of clinical samples (Cole, Weinberger, & Dickinson, 2011; Haynes, Bauermeister, & Bunce, 2017). Although more research is necessary to determine the significance of IIV in the context of SRC, our results highlight the importance of attending to these nuanced differences in males' and females' cognitive profiles

following concussion. Though speculative, it is possible that increased IIV in the acute post-injury phase may serve as an early indicator of a more complicated recovery or greater CNS vulnerability following SRC.

When considering the clinical implications of this research, our findings support previous recommendations regarding the need to consider individual differences when making determinations about concussion management, and add to the growing body of evidence emphasizing the development of individually tailored treatments for concussed athletes. There are, however, limitations of the study that should be considered. First, the generalizability of our findings may be limited, as we focused exclusively on collegiate athletes with relatively acute concussions. Future research will need to examine cognitive sex differences in older and younger samples and in individuals in the chronic phase of injury to determine whether the findings hold in other populations. Second, it is important to recognize that females underwent post-concussion testing, on average, over 2 weeks following their injury whereas males were tested, on average, 1 week post-injury. It is thus possible that there may be sampling bias with regard to when athletes are referred for

post-concussion evaluations, and that females who are referred for testing may be those with more complicated recoveries. Importantly though, time since injury was accounted for in our analyses. Another related limitation is the smaller proportion of females in the sample relative to males, which appears to be a common issue when examining sex differences in the context of SRC. Given this discrepancy, it is possible that we were underpowered to detect group differences; nonetheless, despite potential power limitations, the present study offers preliminary support for the importance of evaluating cognitive scores beyond just mean performance. A final limitation of this study is that pre-injury, or baseline, data were not available for all athletes and so we were unable to make direct pre/post-injury comparisons. Nevertheless, we accounted for premorbid sex differences in cognitive functioning by utilizing sex-specific normative data.

In conclusion, our results suggest that measures of IIV and impairment scores may provide valuable information about neurological health and potentially aid in the identification of post-concussion cognitive dysfunction. Findings should be replicated in larger samples with more equal representation from both sexes, and ongoing research will need to continue evaluating the clinical significance of IIV and impairment scores and determine the extent to which they may influence, or guide, return-to-play decisions.

ACKNOWLEDGMENTS

The authors would like to thank Wayne Sebastianelli at Penn State Sports Medicine for his generous support of our research. We would also like to thank the athletes who participated in the sports-concussion management program.

CONFLICTS OF INTEREST

The authors have nothing to disclose.

REFERENCES

- Arnett, P., Meyer, J., Merritt, V., & Gutty, E. (2016). Neuropsychological testing in mild traumatic brain injury: What to do when baseline testing is not available. *Sports Medicine and Arthroscopy Review*, 24(3), 116–122.
- Arnett, P., Rabinowitz, A., Vargas, G., Ukueberuwa, D.M., Merritt, V.C., & Meyer, J. (2014). Neuropsychological testing in sports concussion management: An evidence-based model when baseline is unavailable. In S. Slobounov & W. Sebastianelli (Eds.), *Concussion in athletics: From brain to behavior*. New York: Springer.
- Bunce, D., Bielak, A.A., Cherbuin, N., Batterham, P.J., Wen, W., Sachdev, P., & Anstey, K.J. (2013). Utility of intraindividual reaction time variability to predict white matter hyperintensities: A potential assessment tool for clinical contexts? *Journal of the International Neuropsychological Society*, 19(9), 971–976.
- Cole, V.T., Weinberger, D.R., & Dickinson, D. (2011). Intraindividual variability across neuropsychological tasks in schizophrenia: A comparison of patients, their siblings, and healthy controls. *Schizophrenia Research*, 129(1), 91–93.
- Covassin, T., Elbin, R., Kontos, A., & Larson, E. (2010). Investigating baseline neurocognitive performance between male and female athletes with a history of multiple concussion. *Journal of Neurology, Neurosurgery & Psychiatry*, 81(6), 597–601.
- Covassin, T., Savage, J.L., Bretzin, A.C., & Fox, M.E. (2018). Sex differences in sport-related concussion long-term outcomes. *International Journal of Psychophysiology*, 132, 9–13.
- Dougan, B.K., Horswill, M.S., & Geffen, G.M. (2014). Athletes' age, sex, and years of education moderate the acute neuropsychological impact of sports-related concussion: A meta-analysis. *Journal of the International Neuropsychological Society*, 20(1), 64–80.
- Dykiert, D., Der, G., Starr, J.M., & Deary, I.J. (2012). Sex differences in reaction time mean and intraindividual variability across the life span. *Developmental Psychology*, 48(5), 1262–1276.
- Ghisletta, P., Renaud, O., Fagot, D., Lecerf, T., & De Ribaupierre, A. (2018). Age and sex differences in intra-individual variability in a simple reaction time task. *International Journal of Behavioral Development*, 42(2), 294–299.
- Haynes, B.I., Bauermeister, S., & Bunce, D. (2017). A systematic review of longitudinal associations between reaction time intraindividual variability and age-related cognitive decline or impairment, dementia, and mortality. *Journal of the International Neuropsychological Society*, 23(5), 431–445.
- Iverson, G.L. & Brooks, B.L. (2011). Improving accuracy for identifying cognitive impairment. In M.R. Schoenberg & J.G. Scott (Eds.), *The little black book of neuropsychology* (pp. 923–950). New York: Springer.
- Iverson, G.L., Gardner, A.J., Terry, D.P., Ponsford, J.L., Sills, A.K., Broshek, D.K., & Solomon, G.S. (2017). Predictors of clinical recovery from concussion: A systematic review. *British Journal of Sports Medicine*, 51(12), 941–948.
- Karr, J.E., Areshenkoff, C.N., & Garcia-Barrera, M.A. (2014). The neuropsychological outcomes of concussion: A systematic review of meta-analyses on the cognitive sequelae of mild traumatic brain injury. *Neuropsychology*, 28(3), 321–336.
- Kontos, A.P., Sufrinko, A., Womble, M., & Kegel, N. (2016). Neuropsychological assessment following concussion: An evidence-based review of the role of neuropsychological assessment pre-and post-concussion. *Current Pain and Headache Reports*, 20(6), 38.
- Lezak, M.D., Howieson, D.B., Bigler, E.D., & Tranel, D. (2012). *Neuropsychological assessment* (5th ed.). New York: Oxford University Press.
- Lovell, M.R. (2016). *IMPACT: Administration and technical manual*. IMPACT Applications, Inc.
- MacDonald, S.W., Nyberg, L., & Bäckman, L. (2006). Intraindividual variability in behavior: Links to brain structure, neurotransmission and neuronal activity. *Trends in Neurosciences*, 29(8), 474–480.
- McCrorry, P., Meeuwisse, W., Dvorak, J., Aubry, M., Bailes, J., Broglio, S., Cantu, R.C., Cassidy, D., Echemendia, R.J., Castellani, R.J., Davis, G.A., Ellenbogen, R., Emery, C., Engebretsen, L., Feddermann-Demont, N., Giza, C.C., Guskiewicz, K.M., Herring, S., Iverson, G.L., Johnston, K.M., Kissick, J., Kutcher, J., Leddy, J.J., Maddocks, D., Makkissi, M., Manley, G.T., McCrea, M., Meehan, W.P., Nagahiro, S., Patricios, J., Putukian, M., Schneider, K.J., Sills, A., Tator, C.H., Turner, M., & Vos, P.E. (2017). Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. *British Journal of*

- Sports Medicine*, 51, 838–847. doi:[10.1136/bjsports-2017-097699](https://doi.org/10.1136/bjsports-2017-097699)
- McCrory, P., Meeuwisse, W.H., Aubry, M., Cantu, R.C., Dvorak, J., Echemendia, R.J., Engebretsen, L., Johnston, K., Kutcher, J.S., Raftery, M., Sills, A., Benson, B.W., Davis, G.A., Ellenbogen, R.G., Guskiewicz, K., Herring, S.A., Iverson, G.L., Jordan, B.D., Kissick, J., McCrea, M., McIntosh, A.S., Maddocks, D., Makkdissi, M., Purcell, L., Putukian, M., Schneider, K., Tator, C.H., & Turner, M. (2013). Consensus statement on concussion in sport—the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Journal of Athletic Training*, 5(4), 255–279.
- Merritt, V.C., Meyer, J.E., Cadden, M.H., Roman, C.A., Ukueberuwa, D.M., Shapiro, M.D., & Arnett, P.A. (2017). Normative data for a comprehensive neuropsychological test battery used in the assessment of sports-related concussion. *Archives of Clinical Neuropsychology*, 32(2), 168–183.
- Merritt, V.C., Padgett, C.R., & Jak, A.J. (2019). A systematic review of sex differences in concussion outcome: What do we know? *The Clinical Neuropsychologist*, 33(6), 1016–1043. doi:[10.1080/13854046.2018.1508616](https://doi.org/10.1080/13854046.2018.1508616)
- Merritt, V.C., Rabinowitz, A.R., & Arnett, P.A. (2018). The influence of the apolipoprotein E (APOE) gene on subacute post-concussion neurocognitive performance in college athletes. *Archives of Clinical Neuropsychology*, 33(1), 36–46.
- Rabinowitz, A.R. & Arnett, P.A. (2013). Intraindividual cognitive variability before and after sports-related concussion. *Neuropsychology*, 27(4), 481–490.
- Rothman, K.J. (1990). No adjustments are needed for multiple comparisons. *Epidemiology*, 1(1), 43–46.
- Ruff, R.M., Iverson, G.L., Barth, J.T., Bush, S.S., Broshek, D.K., & NAN Policy Planning Committee. (2009). Recommendations for diagnosing a mild traumatic brain injury: A national academy of neuropsychology education paper. *Archives of Clinical Neuropsychology*, 24(1), 3–10.