ERRATUM

The following is a correction for an error that occurred in the *Journal of the International Neuropsychological Society*, Vol. 8, No. 6, pp. 781–793. The article titled "Neuropsychological functioning in a cohort of HIV infected women: Importance of antiretroviral therapy," by J.L. Richardson et al., had errors in Table 5, which appeared on page 788. Please refer to the numbers in boldface that shows the incorrect and correct values, respectively.

ORIGINAL

Table 5. Predictors of abnormal NP protocols 1,2

Variable	Normal No. (%)	Abnormal No. (%)	Univariate risk of NP abnormal protocol <i>OR</i> (95% <i>CI</i>)	Model I: Adjusted risk of NP abnormal protocol by disease status <i>OR</i> (95% <i>CI</i>)	Model II: Adjusted risk of NP abnormal protocol by treatment status OR (95% CI)
Ethnicity					
White	65 (54.6)	54 (45.4)	1.0	1.0	1.0
Africam American	32 (72.7)	12 (27.3)	2.22 (1.04, 4.72)*	1.81 (0.74, 4.45)	1.96 (0.79, 4.89)
Hispanic	45 (66.2)	23 (33.8)	1.36 (0.59, 3.13)	0.96 (0.31, 2.60)	1.01 (0.37, 2.78)

 $p \le .05; p \le .01; p \le .001; p \le .0001.$

CORRECTED

Table 5. Predictors of abnormal NP protocols ^{1,2}

Variable	Normal No. (%)	Abnormal No.# (%)	Univariate risk of NP abnormal protocol <i>OR</i> (95% <i>CI</i>)	Model I: Adjusted risk of NP abnormal protocol by disease status OR (95% CI)	Model II: Adjusted risk of NP abnormal protocol by treatment status OR (95% CI)
Ethnicity					
White	32 (72.7)	12 (27.3)	1.0	1.0	1.0
African American	65 (54.6)	54 (45.4)	2.22 (1.04, 4.72)*	1.81 (0.74, 4.45)	1.96 (0.79, 4.89)
Hispanic	45 (66.2)	23 (33.8)	1.36 (0.59, 3.13)	0.96 (0.31, 2.60)	1.01 (0.37, 2.78)

 $[*]p \le .05; **p \le .01; ***p \le .001; †p \le .0001.$

Cambridge University Press and the authors regret the inconvenience that this inadvertent error may have caused.

¹Multivariate analysis adjusted for all other variables listed in the model.

²Tertile cuts of summed number of yrs used marijuana, cocaine, heroin, or alcohol (3 or more glasses per day).

¹Multivariate analysis adjusted for all other variables listed in the model.

²Tertile cuts of summed number of yrs used marijuana, cocaine, heroin, or alcohol (3 or more glasses per day).