

# Central coherence in eating disorders: a systematic review

C. Lopez<sup>1\*</sup>, K. Tchanturia<sup>1</sup>, D. Stahl<sup>2</sup> and J. Treasure<sup>3</sup>

<sup>1</sup> Division of Psychological Medicine, Eating Disorders Research Unit, Institute of Psychiatry, King's College London, London, UK

<sup>2</sup> Department of Biostatistics and Computing, Institute of Psychiatry, King's College London, London, UK

<sup>3</sup> Eating Disorders Research Unit, Department of Academic Psychiatry, King's College London, London, UK

**Background.** This review systematically appraised the research evidence for local *versus* global information processing to test the hypothesis that people with eating disorders (ED) had weak central coherence.

**Method.** Searches on Medline, EMBASE, PsycINFO and ISI Web of Science databases were conducted in November 2006 and subsequently updated in September 2007. Each search was conducted in two steps: (1) neuropsychological tasks measuring central coherence and (2) words related to cognitive functioning in eating disorders. Data were summarized in a meta-analysis if the number of studies for a given test was >5.

**Results.** Data were extracted from 16 studies. Meta-analyses were conducted for four tasks obtaining moderate effect sizes. The majority of studies found global processing difficulties across the ED spectrum. The results are less clear regarding local processing.

**Conclusions.** People with ED have difficulties in global processing. It is less certain as to whether they have superior local processing. Currently, there is insufficient evidence to refute the weak central coherence hypothesis.

Received 13 December 2007; Revised 17 March 2008; Accepted 31 March 2008; First published online 30 April 2008

**Key words:** Anorexia nervosa, bulimia nervosa, cognitive function, global processing, local processing, weak coherence.

## Introduction

Pre-existing cognitive abnormalities may be relevant to both the development (Lena *et al.* 2004) and maintenance of eating disorders (ED) (Schmidt & Treasure, 2006). For example, poor performance in set-shifting tasks (a component of executive functioning) is a possible candidate endophenotype for ED (Bulik *et al.* 2007; Treasure *et al.* 2007) as it has been found present in the acute illness, after recovery and in healthy relatives (Tchanturia *et al.* 2004; Holliday *et al.* 2005; Roberts *et al.* 2007).

It has been hypothesized that a cognitive style characterized by weak central coherence may also be linked to ED (Treasure *et al.* 2006; Gillberg *et al.* 2007; Southgate *et al.* 2007; Treasure, 2007; Lopez *et al.* 2008*a,b*). Weak coherence refers to the cognitive style where there is a bias towards detail accompanied by difficulties in the integrative processing of information (Happé & Booth, 2008) and is characteristically associated with autistic spectrum disorders (ASD) (Frith, 1989). This trait may explain the commonality

between some individuals with anorexia nervosa (AN) and those with ASD (Gillberg *et al.* 1996; Wentz *et al.* 1999).

The purpose of this review was to systematically summarize the evidence that would support the hypothesis of weak central coherence in ED. Our hypothesis was that people with ED would show superior performance on tasks in which a detail processing style was beneficial and difficulties in those tasks in which a global processing strategy was required.

## Method

### Selection of studies

A two-step search of the following electronic databases was used to identify relevant papers for inclusion in the review: Medline, EMBASE, PsycINFO and ISI Web of Science. Searches were conducted in November 2006 and subsequently updated in September 2007. In step 1, a literature search was conducted using the term 'central coherence' to generate all the neuropsychological tests used to explore this concept. Then the search was narrowed to articles in which these tasks were undertaken in people

\* Address for correspondence: C. Lopez, Eating Disorders Research Unit, 5th Floor, Academic Psychiatry Department, Bermondsey Wing, Guy's Hospital, London SE1 9RT, UK.  
(Email: carolina.lopez@iop.kcl.ac.uk)

with ED. In step 2, databases were searched for articles including words related to cognitive assessment in ED. The search included combinations of key words regarding diagnostic categories (anorexia nervosa, bulimia nervosa, eating disorders, binge eating disorder, bulimic disorders) and key words related to cognitive functioning (cognition, information processing, neuropsychology, cognitive functioning, cognitive styles, local/global processing, field dependency/independence, holistic/analytic style, abstract thinking) to ensure that papers not explicitly advertising the use of the mentioned tasks would be included.

Relevant published citations for all articles obtained were also pursued. Key investigators were contacted asking for unpublished material. A manual search was also carried out. Thirty-five tests were generated from the search of the central coherence concept mainly in relation to ASD. Only four articles used the concept in ED (Gillberg *et al.* 2007; Southgate *et al.* 2007; Lopez *et al.* 2008a,b).

After the second search, only the following relevant tests were found to have been used in ED: the Block Design and Object Assembly subscales of the Wechsler Intelligence Scales, the Group/Embedded Figures Test (EFT), Rey-Osterrieth Complex Figure Test (ROFT), Matching Familiar Figures Test (MFFT), Sentence Completion Task (SCT), Homograph Reading Task (HRT) and California Verbal Learning Test (CVLT).

#### *Paper retrieval*

The search was conducted by two researchers on the basis of the following inclusion criteria:

- (1) Population: participants suffering or recovered from an ED and a healthy control (HC) group.
- (2) Instruments: studies specifying the use of at least one of the aforementioned tasks.
- (3) Study designs: cross-sectional studies.
- (4) The outcome variable was functioning on the information processing aspect of the tasks.

#### *Selection results*

Thirty-seven relevant abstracts were found and the full papers were retrieved. One paper could not be obtained (Talarczyk & Rajewski, 2001). Fifteen of the remaining studies were excluded as they did not include an HC group (Sours, 1969; Hamsher *et al.* 1981; Small *et al.* 1982, 1983; Gordon *et al.* 1984; Touyz *et al.* 1986; Dura & Bornstein, 1989; Ranseen & Humphries, 1992; Bowers, 1994; Kaye *et al.* 1995; Nakasuji, 1999; Bayless *et al.* 2002; McDowell *et al.* 2003; Kitabayashi *et al.* 2004; Key *et al.* 2006). Some papers omitted the raw data and the corresponding

author was contacted for these if the study was published after 1996. Four studies were eliminated because the relevant outcomes were not available, including the only study using the CVLT (Fox, 1981; Horne *et al.* 1991; Bradley *et al.* 1997; Steinglass *et al.* 2006). Finally, one study reported data published previously (Murphy *et al.* 2004). Data from two studies included repeated measures in a longitudinal study (Szmukler *et al.* 1992; Gillberg *et al.* 2007). These data were included in the review table but were not included (because of lack of independence) in the meta-analysis. Therefore, the review included a total of 17 studies.

#### *Description of the tasks*

All of the following tasks have been used to explore central coherence with the caveat that most also involve other cognitive processes.

According to the ASD literature (Happé & Frith, 2006), tasks in which a superior detail focused processing would benefit performance are:

*Block Design Test (BD; Wechsler, 1949, 1974).* This construction test requires putting sets of blocks together to replicate complex designs depicted on a diagram. Time and errors are measured. It has been argued that a possible mechanism involved in faster performance is a superior detail processing (Shah & Frith, 1993). This hypothesis has been tested using a modified version of the task in which performance in two conditions of the designs, unsegmented (whole designs) and segmented (designs are broken into their constituents parts), are compared. The central coherence account predicts that those with weak coherence would benefit less from the segmentation of the designs (Shah & Frith, 1993; Happé *et al.* 2001).

*Group/Embedded Figures Test (EFT; Witkin *et al.* 1971, 2002).* This perceptual test measures the time taken to locate and trace 12 (or 18, in the group version) simple shapes embedded in complex designs. The main outcome is the time taken to locate the hidden shapes. A shorter time has been described as the result of a strong local processing or a bias towards detail (Baron-Cohen & Hammer, 1997; Jolliffe & Baron-Cohen, 1997).

*Matching Familiar Figures Test (MFFT; Kagan *et al.* 1964; Kagan, 1966).* This visual perceptual test was designed to measure cognitive impulsivity. The participant is asked to identify the exact replica of a familiar object among eight very similar alternatives. The latency to identify the 'matching' figure and the number of errors are scored. Shorter latencies may

reflect an impulsive cognitive style. However, combining latency and errors provides a measure of efficiency. People with an efficient detail focused processing are both quick and error free.

Tasks that would be favoured by global strategies are:

*Object Assembly (OA; Wechsler, 1974, 1981).* This task involves solving five jigsaw-type puzzles depicting familiar objects and the outcome is the time taken to complete the puzzles. A shorter time suggests a better ability to create an integrated global representation from its parts (Tokley & Kemps, 2007).

*Rey–Osterrieth Complex Figure Test (ROFT; Osterrieth, 1944).* This visual perceptual task requires copying a diagram of a complex figure and then recalling it from memory without previous warning. The interval between copy and recall may vary from 3 to 60 min. In general, the accuracy of the drawing at copy and recall phases has been scored by a system defined by Taylor (adapted from Osterrieth, 1944; Spreen & Strauss, 1998). Process measures to assess the quality of copy strategies have been developed (Savage *et al.* 1999; Booth, 2006). It has been found that the organizational strategies used to copy the figure (detail *versus* global) mediate the percentage of recall (e.g. Savage *et al.* 1999; Lopez *et al.* 2008a). A global approach favours recall whereas a detail approach affects it negatively. The assumption from this mediation hypothesis is that poor recall might reflect weak visual spatial coherence.

*Sentence Completion Task (SCT; Happé et al. 2001).* This verbal test consists of a series of sentences constructed to provoke conflict between local and global processing. The outcome is the number of 'local' associations and hesitation time, which is thought to be indicative of a bias to local processing or difficulties in global processing (Booth, 2006).

*Homograph Reading Task (HRT; Happé, 1997; Jolliffe & Baron-Cohen, 1999).* This verbal task tests the ability to process a sentence as a whole rather than on a local level. The participant is asked to read out a set of 16 sentences where the context of the sentence determines how a homograph within it should be pronounced. The number of sentences pronounced correctly indicates stronger verbal coherence.

### Data synthesis

Outcomes clustered by tasks were summarized by a meta-analysis if the number of studies available was >5; this applied to BD, EFT (accuracy) and OA. For

the rest of the measures Cohen's *d* individual effect sizes were calculated.

Meta-analyses were carried out in Stata 9.1 (Stata Corporation, College Station, TX, USA) using the user-contributed commands for meta-analyses METAN (Bradburn *et al.* 1998) and METABIAS (Steichen, 1998). Forrest plots are used to show the meta-analysis with all the independent data available for each measure.

Cohen's *d*, the difference between ED and HC divided by the pooled standard deviation, was calculated for each study (Wilson, 2001). Cohen's *d* is understood as negligible ( $\geq 0$  and  $< 0.15$ ), small ( $\geq 0.15$  and  $< 0.40$ ), medium ( $\geq 0.40$  and  $< 0.75$ ), large ( $\geq 0.75$  and  $< 1.10$ ), very large ( $\geq 1.10$  and  $< 1.45$ ) and huge ( $> 1.45$ ).

The standardized effect sizes were subsequently analysed using METAN. The standard error of each study's standardized effect size was calculated using the method of Cooper & Hedges (1994) and pooled using a random-effect model that allowed for between-study variation of effect sizes (Everitt, 2003).

Homogeneity between the trials was analysed using Cochran's *Q* test. Because of the small sample sizes an additional measure of heterogeneity or inconsistency  $I^2 [(Q - df) / Q]$  was calculated (Higgins *et al.* 2003).  $I^2$  ranges between 0% (no inconsistency) and 100% (high heterogeneity).

The presence of publication bias was assessed informally by visual inspection of funnel plots that were corroborated by its corresponding statistical analogue, Begg's adjusted rank test (Begg & Mazumdar, 1994), and additionally by Egger's test (Egger *et al.* 1997) as implemented in *metabias*.

## Results

Table 1 shows a summary of the main characteristics of the studies included in the review. Most of the studies ( $n = 13$ ) compared people with AN with an HC group. Only six studies included people with bulimia nervosa (BN). One study included a group with weight-recovered AN (WRAN). Two studies included mixed AN groups, that is patients at various stages of illness (acute and partial recovery).

### Summary by task

#### Block Design Test (BD)

BD was used in nine studies. In general, BD was used as a part of the standard measure of intelligence quotient (IQ). Lopez *et al.* (2008a, b) reported the only studies using segmented *versus* unsegmented versions of the BD. They found no statistical differences between people with acute BN and HC across the two conditions of the tasks although people with BN



Gillberg <i>et al.</i> (1996)	Mixed AN	51	21	21.2 (3.5)	102.9		10 ASD	Mixed AN v. HC	-0.30				-0.65
	HC	51	20.8	21.2 (2.3)	106.5								
Mathias & Kent (1998)	AN	34	22.7 (7.4)	15.3 (1.7)	96.1 (8.8)*	33%	↑anxiety; ↑depression	AN v. HC	-0.43	-0.52		-0.67*	-0.50
	HC	31	20.8 (2.4)	22.8 (2.4)	101.1 (6.9)*	No							
Murphy <i>et al.</i> (2002, 2004)	AN	16	22.3 (4.4)	14.8 (1.2)	119.3 (9.8)	13%	↑OCD	AN v. HC		-0.18		-0.45	
	BN	16	22.0 (4.5)	20.1 (2.3)	122.7 (4.3)	25%		BN v. HC		-0.18		0.61*	
	HC	16	25.3 (2.6)	22.0 (2.6)	117.1 (5.6)	No							
Galderisi <i>et al.</i> (2003)	AN	14	23.7 (4.54)**	15.3 (2.23)	100 (10.8)**	No		AN v. HC	0				0.31
	BN	31	23.7 (4.54)	21.5 (2.30)	100 (10.8)	No	4 depression; 3 PD	BN v. HC	-0.29				-0.45
	HC	45	24.2 (4.8)	N.R.	101.1 (13.6)	No							
Sherman <i>et al.</i> (2006)	AN	18	25.6 (5.8)	16.7 (1.1)	N.R.	61%	6 depression; 3 dysthymia; 5 GAD	AN v. HC		-0.64*	-0.87*	-1.42*	
	HC	19	25.7 (5.3)	22.2 (1.8)	N.R.	No							
Gillberg <i>et al.</i> (2007)	Mixed AN	47	24.5	22.2 (4.1)	105.2*			Mixed AN v. HC	-0.3				-0.46*
	HC	51	24.2	22.2 (3.4)	109.4								
Southgate <i>et al.</i> (in press)	AN	20	26.8 (8.5)	16.3 (2.6)	116.8 (4.8)	No	↑OCD; ↑anxiety; ↑depression	AN v. HC		-0.38	-0.3		
	BN	14	25.7 (4.9)	21.1 (6.7)	114.4 (7.3)	No	↑anxiety; ↑depression	BN v. HC		-0.34	-0.01		
	HC	26	27.3 (11.5)	22.0 (3.4)	114.8 (5.8)	No							
Tockley & Kemps (2007)	AN	24	21.9 (5.3)	16.8 (1.5)	matched	N.R.	↑OCD; ↑anxiety; ↑depression	AN v. HC	-0.75*				-0.31*
	HC	24	22.0 (5.0)	22.7 (4.3)									
Lopez <i>et al.</i> (2008a)	AN	42	28.4 (9.6)	15.8 (1.7)	112.8 (6.8)	40.5%	↑OCD; ↑anxiety; ↑depression	AN v. HC	-0.51*	0	-0.70*	-0.51*	SCT = 0.65* HRT = 0.0
Lopez <i>et al.</i> (2008b)	BN	42	27.0 (7.2)	21.7 (2.4)	111.8 (6.9)	16.7%	↑OCD; ↑anxiety; ↑depression	BN v. HC	-0.69*	-1.3*	-0.37	-0.97*	SCT = 1.1* HRT = -0.45
	HC	42	26.3 (6.4)	21.9 (2.7)	112.2 (5.4)								

BMI, Body mass index; BD, Block Design Test; EFT, Embedded Figures Test; MFFT, Matching Familiar Figures Test; ROFT, Rey–Osterrieth Complex Figure Test; OA, Object Assembly; AN, anorexia nervosa; WRAN, weight-recovered AN; N.R., not reported; HC, healthy control group; ANR, restricting anorexia nervosa; SA, substance abuse; OCD, obsessive–compulsive disorder; ASD, autistic spectrum disorder; PD, personality disorder; GAD, generalized anxiety disorder; SCT, Sentence Completion Task; HRT, Homograph Reading Task.

\* Significant differences between groups.

\*\* The authors reported Age and IQ grouping AN and BN.




Fig. 1. Forrest plot for the Block Design (BD) meta-analysis excluding repeated measures: ■, anorexia nervosa (AN); □, bulimia nervosa (BN); ▨, mixed AN; ■, weight-recovered AN (WRAN).

performed slightly better in the unsegmented trial ( $d=0.2$ ). The AN group performed worse than HC in the segmented designs ( $d=0.46$ ). The BN but not the AN group benefited less from segmentation (BN  $d=0.37$ ; AN  $d=0.02$ ).

All of the studies using the original form of the task were included in a meta-analysis (Fig. 1). People with ED performed slower with a moderate standardized effect size ( $d=0.43$ ,  $z=5.24$ ,  $p<0.001$ ). There was no evidence of heterogeneity across studies [ $\chi^2(9)=6.38$ ,  $p=0.70$ ,  $I^2=0.0\%$ ]. Two of the studies were outliers with a larger effect size (Pendleton Jones *et al.* 1991; Szmukler *et al.* 1992). No reason could be found from scrutiny of the studies as to why their results were atypical. The larger effect size in the study by Pendleton Jones *et al.* (1991) may have been confounded by the lower IQ level in the ED groups. Participants with BN also showed poor performance on this task (Pendleton Jones *et al.* 1991; Galderisi *et al.* 2003). No evidence of publication bias was found.

#### Embedded Figures Test (EFT)

The EFT was used in six studies (Fig. 2) with an ED population. One study (Basseches & Karp, 1984) also administered the test to an obese group. The latter was not included in the meta-analysis but the study revealed that obese people performed worse than an HC group ( $d=-1.3$ ).

The meta-analysis revealed a high degree of heterogeneity across the studies [ $\chi^2(7)=58.8$ ,  $p<0.001$ ] with an index of inconsistency reaching 86.3%. Beggs' test revealed the existence of publication bias ( $z=1.98$ ,  $p=0.048$ ).

Visual inspection of the data suggested that there was a distinction between the studies carried out before 1991 and the three more recent studies. There may be technical and administration differences that account for the highly divergent results. All studies except that of Tokley & Kemps (2007), which used the Group EFT, used the original EFT. The procedure for the Group EFT is clearly defined: participants have unlimited access to the 'hidden' shape. Lopez *et al.* (2008a,b) used the modification of the task made by Happé & Booth (2008), in which both the 'hidden' shape and the complex figure were placed simultaneously in front of the participant. This modification in the administration could explain why these more recent studies found superior function (moderate effect sizes) in the ED groups. Studies using the original version of the task, which was confounded by a memory component, found that people with ED performed poorly on the task (moderate effect sizes). In addition, in two of the studies (Basseches & Karp, 1984; Pendleton Jones *et al.* 1991) the IQ in the ED group was lower than in the HC and this may have confounded the results as there is some evidence that EFT is related to intellectual ability (Flexer & Roberge, 1980; Riding & Pearson, 1994).




Fig. 2. Forrest plot for the Embedded Figures Test (EFT) meta-analysis: ■, anorexia nervosa (AN); □, bulimia nervosa (BN); ■, weight-recovered AN (WRAN).

#### Matching Familiar Figures Test (MFFT)

Only two studies using the MFFT fulfilled the criteria to be included in this review (Toner *et al.* 1987; Southgate *et al.* in press). One study (Toner *et al.* 1987) found that the binge-purge AN group was less accurate than both the HC ( $d=0.81$ ) and the restricting AN ( $d=0.74$ ) groups. They found no differences in terms of response latency. Those partially recovered from a binge-purging AN were the fastest group ( $d=0.41$ ) but the least accurate ( $d=1.23$ ), indicative of cognitive impulsivity. Southgate *et al.* (in press) reported no statistical differences between the ED and HC groups in number of errors, response latency and impulsivity. However, the ED groups were in general faster and more accurate than HC. The AN group was more efficient (combining speed and accuracy) than the BN and HC groups ( $d=0.89$ ).

Differences in the status of the illness as well as the diagnostic subgrouping may explain some differences across studies. More studies using the efficiency index are needed.

#### Rey–Osterrieth Complex Figure Test (ROFT)

**Copy accuracy.** Six studies included an AN sample (Thompson, 1993; Kingston *et al.* 1996; Mathias & Kent, 1998; Murphy *et al.* 2002; Sherman *et al.* 2006; Lopez *et al.* 2008a) and two studies also included a BN group (Murphy *et al.* 2002; Lopez *et al.* 2008b). All were integrated in a meta-analysis. People with ED

were less accurate than HC ( $d=0.49$ , 95% CI  $-0.88$  to  $-0.1$ ). There was high heterogeneity [ $\chi^2(7)=26.23$ ,  $p<0.001$ ,  $I^2=73.3\%$ ], mainly due to two studies with outlying results (Thompson, 1993; Lopez *et al.* 2008b). Inspection of these two studies revealed elements such as a failure to match groups (Mathias & Kent, 1998) or a BN group highly affected by co-morbid conditions (Lopez *et al.* 2008b), which could account for these differences. Both studies considered the effect of confounding variables such as depression. No publication bias was found.

**Recall accuracy.** Six studies incorporated a group with AN, three a BN group, and one a recovered AN group (Fig. 3). The time intervals between copy and delayed recall varied from 20 min to 30 min. People with ED performed poorly on this task, with an overall effect size of  $d=0.41$  and high indices of heterogeneity [ $\chi^2(9)=26.9$ ,  $p<0.001$ ;  $I^2=66\%$ ]. The heterogeneity could be due to the inclusion of populations with different diagnoses and in different stages of the illness. No evidence of publication bias was found.

A meta-analysis of the six studies comparing AN and HC groups only was conducted and found no evidence of heterogeneity [ $\chi^2(5)=7.4$ ,  $p=0.19$ ] and a low level of inconsistency ( $I^2=32.8\%$ ). This suggests that poor function on this task is more related to AN ( $d=0.55$ ) There was no evidence of publication bias. Sherman *et al.* (2006) found the largest effect size. It is




Fig. 3. Forrest plot for accuracy in recall Rey–Osterrieth Complex Figure Test (ROFT) meta-analysis: ■, anorexia nervosa (AN); □, bulimia nervosa (BN); ▣, weight-recovered AN (WRAN).

possible that co-morbidity with anxiety and depression and the fact that over 60% of the sample in his study were on medication might account for this effect.

The data from BN group are less clear. The effect size of the three studies available varied from  $d = -0.96$  to  $d = 0.60$ . Murphy *et al.* (2002) reported a superior performance of the BN group whereas Pendleton *et al.* (1991) and Lopez *et al.* (2008*b*) obtained opposite results. IQ differences and age of the participants may confound these results.

*Organizational strategies (copy).* Three studies (Sherman *et al.* 2006; Lopez *et al.* 2008*a,b*) explored this component and confirmed that the organizational approach mediates the effect of grouping (e.g. AN versus HC) on the deficit in visual memory in the recall, that is a detailed rather than a global approach impaired recall accuracy. The effect sizes are shown in Table 1. Lopez *et al.* (2008*a,b*) also found that a process measure specifically developed for central coherence (Booth, 2006) mediated the percentage of recall.

#### Object Assembly (OA)

Six of the studies used this task to compare groups with AN or mixed AN and HC. Only one included a BN group (Fig. 4) and one of the studies reported repeated measures 3 years apart (Gillberg *et al.* 2007).

People with ED take longer on this task with a moderate effect size ( $d = 0.41$ ,  $z = 2.74$ ,  $p = 0.006$ ), with no evidence of heterogeneity [ $\chi^2(5) = 8.65$ ,  $p = 0.12$ ] and moderate inconsistency ( $I^2 = 42.2\%$ ).

Although statistically there was no publication bias (all  $p > 0.70$ ), the trim-and-fill procedure on the funnel plot indicated that there is a small publication bias towards larger studies. After correction, the overall effect remained at a moderate level ( $d = 0.35$ ,  $p = 0.02$ ).

Two studies found that people with AN completed this task faster than controls and the studies were scrutinized to find any explanations for this (Thompson, 1993; Galderisi *et al.* 2003). It is possible that IQ level could be a confounding variable in the study of Thompson (1993) but the atypical results from Galderisi *et al.* (2003) are unexplained.

Gillberg *et al.* (2007) readministered the task to a mixed group of AN (most of them recovered) and an HC group after 3 years of first assessment and found persisting difficulties in this task in the mixed AN group (first assessment:  $d = -0.65$ , second assessment:  $d = -0.46$ ).

#### Verbal tasks

*Sentence Completion Task (SCT).* Two studies have used this task (Lopez *et al.* 2008*a,b*). Difficulties in global processing expressed in longer hesitations to provide appropriate completions were found in both AN ( $d = 0.65$ ) and BN ( $d = 1.1$ ) groups. No differences were found in the number of local completions.






Fig. 4. Forrest plot for Object Assembly meta-analysis: ■, anorexia nervosa (AN); □, bulimia nervosa (BN); ▨, mixed AN.

*Homograph Reading Task (HRT)*. Two studies used this task (Lopez *et al.* 2008*a,b*). People with BN made more errors on this task ( $d=0.45$ ) whereas people with AN performed on this task like the controls.

#### Co-morbidity

Co-morbid conditions were reported in most of the studies (see Table 1) and in some studies this was considered as a potential confounder. The majority of studies found no confounding effects of anxiety or depression (Kingston *et al.* 1996; Mathias & Kent, 1998; Sherman *et al.* 2006; Steinglass *et al.* 2006; Southgate *et al.* 2007; Lopez *et al.* 2008*a*). Three studies found that levels of anxiety were associated with poor cognitive performance (Pendleton Jones *et al.* 1991; Szmukler *et al.* 1992; Lopez *et al.* 2008*b*). One study found that the level of depression decreased copy accuracy on the ROFT (Thompson, 1993). Obsessive-compulsive disorder symptoms were associated with poor recall in the ROFT in two studies (Thompson, 1993; Lopez *et al.* 2008*a*).

One study in which 10% of the AN sample met criteria for ASD (Gillberg *et al.* 1996) observed that this subgroup had poorer performance in the OA test.

Possible confounding by medicines was reported in some of the studies. Lopez *et al.* (2008*a,b*) and Sherman *et al.* (2006) reported no difference between those taking medicines and those who did not, whereas Kingston *et al.* (1996) found that medicines were associated with poor performance in the BD test.

#### Discussion

The aim of this review was to summarize the evidence that would support the hypothesis that people with ED have weak coherence and to compile, wherever possible, a meta-analysis of the results on tasks that are purported to involve global or detail strategies.

Overall, poorer performance (moderate effect sizes) was found in tests that may benefit from a global strategy, for example the ROFT (recall accuracy and organizational strategies), OA, SCT and HRT. Recent studies that have removed the memory component from the EFT show superior performance of the ED groups ( $d$  varied from  $-0.51$  to  $-0.70$ ), which may be attributed to a detailed strategy. This is in contrast to the older studies that found inferior performance on this task ( $d$  ranged from  $0.47$  to  $1.02$ ) in which recall would have been a confounding factor. People with AN and BN performed more efficiently on the MFFT, which may reflect a bias for detail. Thus, overall, these findings support the hypothesis that people with ED have weak coherence and superior detail function.

However, performance on some of the tasks, for example BD, do not fit with the hypothesis. Faster performance in this task has been thought to be the result of a superior analytical ability in people with autism (Shah & Frith, 1993). However, as noted by Happé & Frith (2006), errors in the task could result from poor global integration; that is, performance on this task involves a balance between local and global approaches (Strauss *et al.* 2006). In this review, people with ED showed slower performance relative to HC

with moderate effect size across the studies. The crucial element that is used to highlight superiority in detail functioning is purported to be the lack of advantage from segmentation. This modification of the task was used in only two studies and people with BN did have less benefit from segmentation ( $d=0.37$ ). Such an effect was not found in AN.

In general, the performance on these tasks across the ED diagnostic spectrum showed little variation. Exceptions were found in the ROFT (recall) and in the verbal coherence tasks. Difficulties in the former were consistent among AN groups, whereas people with BN and those in recovery showed more variation. Thus, it is possible that poor nutrition could account for poor performance in this task. In the verbal tasks, people with BN performed poorer than those with AN. It has been hypothesized that impulsivity in BN accounts for differences in the HRT.

This review has limitations. First, only three studies aimed to test the central coherence hypothesis directly. Second, none of the neuropsychological tasks used can be said to examine purely local or global processing, some examine the trade-off between the two and some tap into other aspects of executive function and attention. Third, many studies do not provide sufficient information about how the measures were administered or whether they have been modified, and the assumption that there is enough similarity to justify summarizing the data may be flawed. Likewise, differences in the case mix, such as use of medicines, and diagnostic issues may make it difficult to merge the data.

In conclusion, the hypothesis that people with ED have weak coherence remains unproved. There is consistency in the evidence suggesting weak global processing but less suggesting superiority in local processing. This contrasts with the conclusion drawn in the systematic review of studies examining this hypothesis in ASD, which showed strong evidence for superiority in local processing but weaker evidence for a deficit in global processing (Happé & Frith, 2006). Therefore, further research to examine specifically whether the difficulties in global processing are accompanied by superiority in detail processing is needed.

### Acknowledgements

This work was made possible by a grant from the Nina Jackson Foundation for Research into Eating Disorders. C.L. was in receipt of a Ph.D. scholarship from Mideplan-Chile.

### Declaration of Interest

None.

### References

- Baron-Cohen S, Hammer J** (1997). Parents of children with Asperger syndrome: what is the cognitive phenotype? *Journal of Cognitive Neuroscience* **9**, 548–554.
- Basseches HI, Karp SA** (1984). Field dependence in young anorectic and obese women. *Psychotherapy and Psychosomatics* **41**, 33–37.
- Bayless J, Kanz J, Moser D, McDowell B, Bowers W, Andersen A, Paulsen J** (2002). Neuropsychological characteristics of patients in a hospital-based eating disorder program. *Annals of Clinical Psychiatry* **14**, 203–207.
- Begg CB, Mazumdar M** (1994). Operating characteristics of a rank correlation test for publication bias. *Biometrics* **50**, 1088–1101.
- Booth R** (2006). *Local–Global Processing and Cognitive Style in Autism Spectrum Disorders and Typical Development*. Institute of Psychiatry, King's College London: London.
- Bowers W** (1994). Neuropsychological impairment among anorexia nervosa and bulimia patients. *Eating Disorders: The Journal of Treatment and Prevention* **2**, 42–46.
- Bradburn MJ, Deeks JJ, Altman DG** (1998). sbe24: metan – an alternative meta-analysis command. *Stata Technical Bulletin* **44**, 1–15.
- Bradley SJ, Taylor MJ, Rovet JF, Goldberg E, Hood J, Wachsmuth R, Azcue MP, Pencharz PB** (1997). Assessment of brain function in adolescent anorexia nervosa before and after weight gain. *Journal of Clinical and Experimental Neuropsychology* **19**, 20–33.
- Bulik CM, Hebebrand J, Keski-Rahkonen A, Klump KL, Reichborn-Kjennerud T, Mazzeo SE, Wade TD** (2007). Genetic epidemiology, endophenotypes, and eating disorder classification. *International Journal of Eating Disorders* **40**, S52–S60.
- Cooper H, Hedges L** (1994). *The Handbook of Research Synthesis*. The Russell Sage Foundation: New York.
- Dura JR, Bornstein RA** (1989). Differences between IQ and school achievement in anorexia nervosa. *Journal of Clinical Psychology* **45**, 433–435.
- Egger M, Smith GD, Schneider M, Minder C** (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal* **315**, 629–634.
- Everitt BS** (2003). *Modern Medical Statistics*. Arnold Publishers: London.
- Flexer BK, Roberge JJ** (1980). IQ, field dependence–independence, and the development of formal operational thought. *Journal of General Psychology* **103**, 191–201.
- Fox C** (1981). Neuropsychological correlates of anorexia nervosa. *International Journal of Psychiatry in Medicine* **11**, 285–290.
- Frith U** (1989). *Autism: Explaining the Enigma*. Basil Blackwell: Oxford.
- Galderisi S, Mucci A, Monteleone P, Sorrentino D, Piegari G, Maj M** (2003). Neurocognitive functioning in subjects with eating disorders: the influence of neuroactive steroids. *Biological Psychiatry* **53**, 921–927.
- Gillberg I, Gillberg C, Rastam M, Johansson M** (1996). The cognitive profile of anorexia nervosa: a comparative

- study including a community-based sample. *Comprehensive Psychiatry* 37, 23–30.
- Gillberg I, Rastam M, Wentz E, Gillberg C** (2007). Cognitive and executive functions in anorexia nervosa ten years after onset of eating disorder. *Journal of Clinical and Experimental Neuropsychology* 29, 170–178.
- Gordon DP, Halmi KA, Ippolito PM** (1984). A comparison of the neuropsychological evaluation of adolescents with anorexia nervosa and of adolescents with conduct disorders. *Journal of Adolescence* 7, 245–266.
- Hamsher K, Halmi K, Benton A** (1981). Prediction of outcome in anorexia nervosa from neuropsychological status. *Psychiatry Research* 4, 79–88.
- Happé F** (1997). Central coherence and theory of mind in autism: reading homographs in context. *British Journal of Developmental Psychology* 15, 1–12.
- Happé F, Booth R** (2008). The power of the positive: revisiting weak coherence in autism spectrum disorders. *Quarterly Journal of Experimental Psychology* 61, 50–63.
- Happé F, Briskman J, Frith U** (2001). Exploring the cognitive phenotype of autism: weak ‘central coherence’ in parents and siblings of children with autism. I. Experimental tests. *Journal of Child Psychology and Psychiatry* 42, 299–307.
- Happé F, Frith U** (2006). The weak coherence account: detail-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders* 36, 5–25.
- Higgins JPT, Thompson SG, Deeks JJ, Altman DG** (2003). Measuring inconsistency in meta-analyses. *British Medical Journal* 327, 557–560.
- Holliday J, Tchanturia K, Landau S, Collier DA, Treasure J** (2005). Is impaired set-shifting an endophenotype of anorexia nervosa? *American Journal of Psychiatry* 162, 2269–2275.
- Horne RL, Van Vactor JC, Emerson S** (1991). Disturbed body image in patients with eating disorders. *American Journal of Psychiatry* 148, 211–215.
- Jolliffe T, Baron-Cohen S** (1997). Are people with autism and Asperger syndrome faster than normal on the Embedded Figures Test? *Journal of the American Academy of Child and Adolescent Psychiatry* 38, 527–534.
- Jolliffe T, Baron-Cohen S** (1999). A test of central coherence theory: linguistic processing in high-functioning adults with autism or Asperger syndrome: is local coherence impaired? *Cognition* 22, 149–185.
- Kagan J** (1966). Reflection-impulsivity: the generality and dynamics of conceptual tempo. *Journal of Abnormal Psychology* 1, 17–24.
- Kagan J, Rosman B, Day D, Albert J, Phillips W** (1964). Information processing and the child: significance of analytic and reflective attitudes. *Psychological Monographs* 78, No. 578.
- Kaye WH, Bastiani AM, Moss H** (1995). Cognitive style of patients with anorexia nervosa and bulimia nervosa. *International Journal of Eating Disorders* 18, 287–290.
- Key A, O’Brien A, Gordon I, Christie D, Lask B** (2006). Assessment of neurobiology in adults with anorexia nervosa. *European Eating Disorders Review* 14, 308–314.
- Kingston K, Szmukler G, Andrewes D, Tress B, Desmond P** (1996). Neuropsychological and structural brain changes in anorexia nervosa before and after refeeding. *Psychological Medicine* 26, 15–28.
- Kitabayashi Y, Ueda H, Kashima A, Okamoto A, Kooguchi Y, Narumoto J, Wada Y, Yamashita T, Fukui K** (2004). Change of cognitive function following body weight recovery in anorexia nervosa. *Seishin Igaku* 46, 243–248.
- Lena SM, Fiocco AJ, Leyenaar JK** (2004). The role of cognitive deficits in the development of eating disorders. *Neuropsychological Review* 14, 99–113.
- Lopez C, Tchanturia K, Stahl D, Booth R, Holliday J, Treasure J** (2008a). An examination of the concept of central coherence in women with anorexia nervosa. *International Journal of Eating Disorders* 41, 143–152.
- Lopez C, Tchanturia K, Stahl D, Treasure J** (2008b). Central coherence in women with bulimia nervosa. *International Journal of Eating Disorders* 41, 340–347.
- Mathias J, Kent P** (1998). Neuropsychological consequences of extreme weight loss and dietary restriction in patients with anorexia nervosa. *Journal of Clinical and Experimental Neuropsychology* 20, 548–564.
- McDowell BD, Moser DJ, Fernyhough K, Bowers WA, Andersen AE, Paulsen JS** (2003). Cognitive impairment in anorexia is not due to depressed mood. *International Journal of Eating Disorders* 33, 351–355.
- McLaughlin EF, Karp SA, Herzog DB** (1985). Sense of ineffectiveness in woman with eating disorders: a clinical study of anorexia nervosa and bulimia. *International Journal of Eating Disorders* 4, 511–523.
- Murphy R, Nutzinger DO, Paul T, Leplow B** (2002). Dissociated conditional-associative learning in anorexia nervosa. *Journal of Clinical and Experimental Neuropsychology* 24, 176–186.
- Murphy R, Nutzinger DO, Paul T, Leplow B** (2004). Conditional-associative learning in eating disorders: a comparison with OCD. *Journal of Clinical and Experimental Neuropsychology* 26, 190–199.
- Nakasuji T** (1999). Clinical and psychological characteristics in anorexia nervosa and bulimia nervosa. *Journal of the Osaka City Medical Center* 48, 153–170.
- Osterrieth P** (1944). The test of copying a complex figure: a contribution to the study of perception and memory. *Archives de Psychologie* 30, 206–356.
- Pendleton Jones B, Duncan C, Brouwers P, Mirsky A** (1991). Cognition in eating disorders. *Journal of Clinical and Experimental Neuropsychology* 13, 711–728.
- Ranseen JD, Humphries LL** (1992). The intellectual functioning of eating disorder patients. *Journal of the American Academy of Child and Adolescent Psychiatry* 31, 844–846.
- Riding RJ, Pearson F** (1994). The relationship between cognitive style and intelligence. *Educational Psychology* 14, 413–425.
- Roberts ME, Tchanturia K, Stahl D, Southgate L, Treasure J** (2007). A systematic review and meta-analysis of set shifting ability in eating disorders. *Psychological Medicine* 37, 1075–1084.

- Savage CR, Baer L, Keuthen N, Brown HD, Rauch SL, Jenike MA** (1999). Organizational strategies mediate nonverbal memory impairment in obsessive-compulsive disorder. *Biological Psychiatry* **45**, 905–916.
- Schmidt U, Treasure J** (2006). Anorexia nervosa: valued and visible. A cognitive-interpersonal maintenance model and its implications for research and practice. *British Journal of Clinical Psychology* **45**, 343–366.
- Shah A, Frith U** (1993). Why do autistic individuals show superior performance on the block design task? *Journal of Child Psychology and Psychiatry* **34**, 1351–1364.
- Sherman BJ, Savage CR, Eddy KT, Blais MA, Deckersbach T, Jackson SC, Franko DL, Rauch SL, Herzog DB** (2006). Strategic memory in adults with anorexia nervosa: are there similarities to obsessive compulsive spectrum disorders? *International Journal of Eating Disorders* **39**, 468–476.
- Small A, Madero J, Teagno L, Ebert M** (1983). Intellect, perceptual characteristics, and weight gain in anorexia nervosa. *Journal of Clinical Psychology* **39**, 780–782.
- Small A, Teagno L, Madero J, Gross H, Ebert M** (1982). A comparison of anorexics and schizophrenics on psychodiagnostic measures. *International Journal of Eating Disorders* **1**, 49–56.
- Sours JA** (1969). The anorexia nervosa syndrome: phenomenologic and psychodynamic components. Clinical heterogeneity in four cases. *Psychiatric Quarterly* **43**, 240–256.
- Southgate L, Tchanturia K, Collier D, Treasure J** (in press). The development of the Maudsley Childhood Obsessive–Compulsive Personality Trait Questionnaire in an eating disorder sample. *European Eating Disorders Review*.
- Southgate L, Tchanturia K, Treasure J** (2007). Information processing bias in anorexia nervosa. *Psychiatry Research*. doi: 10.1016/j.psychres.2007.07.017.
- Spreen O, Strauss E** (1998). *A Compendium of Neuropsychological Tests: Administration, Norms and Commentary*. Oxford University Press: New York.
- Steichen T** (1998). sbe19: tests for publication bias in meta-analysis. *Stata Technical Bulletin* **41**, 9–15.
- Steinglass JE, Walsh T, Stern Y** (2006). Set shifting deficit in anorexia nervosa. *Journal of the International Neuropsychological Society* **12**, 431–435.
- Strauss E, Sherman E, Spreen O** (2006). *A Compendium of Neuropsychological Tests: Administration, Norms, and Commentary*. Oxford University Press: Oxford.
- Szmukler GI, Andrewes D, Kingston K, Chen L, Stargatt R, Stanley R** (1992). Neuropsychological impairment in anorexia nervosa: before and after refeeding. *Journal of Clinical and Experimental Neuropsychology* **14**, 347–352.
- Talarczyk M, Rajewski A** (2001). Self-acceptance level in patients with anorexia nervosa. *Psychiatria Polska* **35**, 389–398.
- Tchanturia K, Morris R, Anderluh M, Collier D, Nikolaou V, Treasure J** (2004). Set shifting in anorexia nervosa: an examination before and after weight gain, in full recovery and relationship to childhood and adult OCPD traits. *Journal of Psychiatric Research* **38**, 545–552.
- Thompson S** (1993). Implications of neuropsychological test results of women in a new phase of anorexia nervosa. *European Eating Disorders Review* **1**, 152–165.
- Tokley M, Kemps E** (2007). Preoccupation with detail contributes to poor abstraction in anorexia nervosa. *Journal of Clinical and Experimental Neuropsychology* **29**, 734–741.
- Toner BB, Garfinkel PE, Garner DM** (1987). Cognitive style of patients with bulimic and diet-restricting anorexia nervosa. *American Journal of Psychiatry* **144**, 510–512.
- Touyz SW, Beumont PJV, Johnstone LC** (1986). Neuropsychological correlates of dieting disorders. *International Journal of Eating Disorders* **5**, 1025–1034.
- Treasure J** (2007). Getting beneath the phenotype of anorexia nervosa: the search for viable endophenotypes and genotypes. *Canadian Journal of Psychiatry* **52**, 212–209.
- Treasure J, Lopez C, Roberts M** (2007). Endophenotypes in eating disorders: moving toward etiologically based diagnosis and treatment focused on pathophysiology. *Pediatric Health* **1**, 171–181.
- Treasure J, Southgate L, Tchanturia K, Lopez C, Collier D** (2006). Gene environment and developmental issues in eating disorders. *Association for Child and Adolescent Mental Health (ACAMH) Occasional Paper* **25**, 51–62.
- Wechsler D** (1949). *Wechsler Intelligence Scale for Children (WISC)*. The Psychological Corporation: San Antonio, TX.
- Wechsler D** (1974). *Wechsler Intelligence Scale for Children – Revised (WISC-R)*. The Psychological Corporation: San Antonio, TX.
- Wechsler D** (1981). *Wechsler Adult Intelligence Scale – Revised (WAIS-R)*. The Psychological Corporation: San Antonio, TX.
- Wentz E, Gillberg C, Gillberg IC, Rastam M** (1999). Ten-year follow-up of adolescent-onset anorexia nervosa: personality disorders. *Journal of the American Academy of Child and Adolescent Psychiatry* **38**, 1389–1395.
- Wilson DB** (2001). *Effect Size Determination Program*. University of Maryland: College Park, MD.
- Witkin H, Oltman P, Raskin E, Karp S** (1971). *A Manual for the Embedded Figures Test*. Consulting Psychologists Press: Palo Alto, CA.
- Witkin H, Oltman P, Raskin E, Karp S** (2002). *Group Embedded Figures Test Manual*. Mind Garden, Inc.: Redwood City, CA.