



COMPOSITIO MATHEMATICA

Degeneracy loci, virtual cycles and nested Hilbert schemes II

Amin Gholampour and Richard P. Thomas

Compositio Math. **156** (2020), 1623–1663.

[doi:10.1112/S0010437X20007290](https://doi.org/10.1112/S0010437X20007290)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Degeneracy loci, virtual cycles and nested Hilbert schemes II

Amin Gholampour and Richard P. Thomas

ABSTRACT

We express nested Hilbert schemes of points and curves on a smooth projective surface as ‘virtual resolutions’ of degeneracy loci of maps of vector bundles on smooth ambient spaces. We show how to modify the resulting obstruction theories to produce the virtual cycles of Vafa–Witten theory and other sheaf-counting problems. The result is an effective way of calculating invariants (VW, SW, local PT and local DT) via Thom–Porteous-like Chern class formulae.

Contents

1	Introduction and summary of results	1624
1.1	Overview	1624
1.2	Degeneracy loci	1624
1.3	Resolutions	1625
1.4	Nested Hilbert schemes	1626
1.5	Nested Hilbert schemes of points	1626
1.6	Nested Hilbert schemes of curves and points	1627
1.7	Reduced virtual cycle	1627
1.8	Nonreduced virtual cycle	1628
1.9	Computing virtual cycles by comparison	1629
1.10	ℓ -step Hilbert schemes	1629
1.11	Vafa–Witten invariants	1629
1.12	Laarakker	1631
1.13	Notation	1632
1.14	Splitting trick	1632
2	Virtual resolutions of degeneracy loci	1632
2.1	Perfect obstruction theory	1634
2.2	Digression: description as a deepest degeneracy locus	1638
2.3	Generalised Carlsson–Okounkov vanishing	1639
3	Comparison	1640
4	Nested Hilbert schemes	1643
4.1	Embedding and reduced virtual cycle	1645
4.2	Comparison of reduced cycles	1647
4.3	Comparison of virtual cycles	1648

Received 17 May 2019, accepted in final form 21 April 2020, published online 1 October 2020.

2010 Mathematics Subject Classification. 14D20, 14J60, 14N35, 14C05 (primary).

Keywords: Hilbert scheme, degeneracy locus, Thom–Porteous formula, local Donaldson–Thomas theory, Vafa–Witten invariants.

This journal is © Foundation Compositio Mathematica 2020.

4.4	Extension to ambient space	1649
4.5	Summary	1651
5	ℓ-step nested Hilbert schemes	1653
6	Computations	1655
6.1	Vafa–Witten invariants	1655
	Acknowledgements	1662
	References	1662

1. Introduction and summary of results

1.1 Overview

It is usually hard to calculate with the virtual fundamental class $[M]^{\text{vir}}$ of a moduli space M . In the rare situation that M (and its perfect obstruction theory) is cut out of a smooth ambient space $\iota: M \hookrightarrow X$ by a section of a vector bundle E , the pushforward of the virtual cycle $\iota_*[M]^{\text{vir}} = e(E)$ is the Euler class of E , making calculation possible.

In [GT20] we described a generalisation to the case where M is instead the *deepest* degeneracy locus of a map of vector bundles on a smooth ambient space. This we applied to nested Hilbert schemes of *points* on smooth surfaces, reproducing the virtual cycles of [GSY20a], which agree with those arising in Vafa–Witten theory [TT20] or reduced local DT theory [GSY20b].

That technique does not extend to other degeneracy loci, but it does apply to their *resolutions* (1.2). We apply this to nested Hilbert schemes of *points and curves* on smooth surfaces with their virtual cycles [GSY20a] coming from Vafa–Witten theory [TT20] or reduced DT theory [GSY20b]. The result is an effective new tool for computing sheaf-theoretic enumerative invariants of projective surfaces, such as those arising in Seiberg–Witten, Vafa–Witten, local DT and local PT theories.

1.2 Degeneracy loci

Let X be a smooth complex quasi-projective variety and

$$\sigma : E_0 \longrightarrow E_1 \tag{1.1}$$

a map of vector bundles of ranks e_0 and e_1 over X . For a positive integer $r \leq e_0$, consider the r th degeneracy locus of σ ,

$$D_r(\sigma) := \{x \in X : \dim \ker(\sigma_x) \geq r\}$$

with scheme structure given by viewing it as the zero locus of

$$\wedge^{e_0-r+1} \sigma : \wedge^{e_0-r+1} E_0 \longrightarrow \wedge^{e_0-r+1} E_1.$$

That is, $D_r(\sigma)$ is the $(e_1 - e_0 + r - 1)$ th Fitting scheme of the first cohomology sheaf $h^1(E_\bullet) = \text{coker } \sigma$ of the complex E_\bullet (1.1). In particular, it depends on E_\bullet only up to quasi-isomorphism and we often denote it by $D_r(E_\bullet)$.

For the nested Hilbert scheme applications in this paper we will take only $r = 1$. On a first read-through of the abstract construction the reader may find it simpler to do the same, substituting projective bundles $\mathbb{P}(-)$ for the Grassmann bundles $\text{Gr}(r, -)$ that follow.

1.3 Resolutions

Over points $x \in D_r$ with $\dim \ker(\sigma_x) = r$ there is a *unique* r -dimensional subspace of $E_0|_x$ on which σ_x vanishes. Taking the set of all such r -dimensional subspaces for *every* point of D_r defines a natural space \tilde{D}_r dominating D_r ,

$$\begin{array}{c} \tilde{D}_r(\sigma) := \{(x, U) \in \text{Gr}(r, E_0) : U \subseteq \ker(\sigma_x) \subseteq E_0|_x\} \\ p \downarrow \\ D_r(\sigma), \end{array} \tag{1.2}$$

whose fibre over x is $\text{Gr}(r, \ker \sigma_x)$. Its scheme structure¹ comes from viewing $\tilde{D}_r(\sigma) \subset \text{Gr}(r, E_0) \xrightarrow{p} X$ as the vanishing locus of the composition

$$\mathcal{U} \hookrightarrow p^*E_0 \xrightarrow{p^*\sigma} p^*E_1. \tag{1.3}$$

Here $\mathcal{U} \subset p^*E_0$ denotes the universal rank r subbundle on $\text{Gr}(r, E_0)$ (or its restriction to \tilde{D}_r) whose fibre at (x, U) (1.2) is U . If σ is appropriately transverse, then each D_r has codimension $r(e_1 - e_0 + r)$, singular locus D_{r+1} and $\tilde{D}_r \rightarrow D_r$ is a *resolution of singularities*. For arbitrary σ we call \tilde{D}_r the *virtual resolution* of D_r since it is *virtually smooth*.

PROPOSITION 1. *Expressing $\tilde{D}_r(E_\bullet)$ as the zero locus of the section (1.3) of the bundle $\mathcal{U}^* \otimes p^*E_1$ endows it with a perfect obstruction theory*

$$\{p^*T_X \longrightarrow \mathcal{U}^* \otimes \text{Cone}(\mathcal{U} \rightarrow p^*E_\bullet)[1]\}^\vee \longrightarrow \mathbb{L}_{\tilde{D}_r}$$

of virtual dimension $\text{vd} := \dim X - r(e_1 - e_0 + r)$. It depends only on the quasi-isomorphism class of the complex E_\bullet (1.1). The resulting virtual cycle

$$[\tilde{D}_r(E_\bullet)]^{\text{vir}} \in A_{\text{vd}}(\tilde{D}_r(E_\bullet)),$$

when pushed forward to X , has class given by the Thom–Porteous formula²

$$\Delta_{r-\text{rk}(E_\bullet)}^r(c(-E_\bullet)) \in A_{\text{vd}}(X).$$

This is a rewriting of parts of [Ful98, Chapter 14] in the language of [BF97]. It is useful if we recognise a moduli space M as having the same perfect obstruction theory as a \tilde{D}_r . Then invariants defined by integration against $[M]^{\text{vir}}$ can be calculated more easily on the smooth space X if the integrand can be expressed as a pullback from X .

The drawback is that the map $\tilde{D}_r \rightarrow X$ contracts³ the exceptional locus of $\tilde{D}_r \rightarrow D_r$, so most integrands (such as the one in (1.15) below) will not be pullbacks from X . We could instead work in $\text{Gr}(r, E_0)$, to which $[\tilde{D}_r]^{\text{vir}}$ pushes forward to give $c_{re_1}(\mathcal{U}^* \otimes p^*E_1)$, but this description is not an invariant of the quasi-isomorphism class of E_\bullet .

¹ A more efficient description of $\tilde{D}_r(\sigma)$ is as the relative Grassmannian $\text{Gr}(\text{coker } \sigma^*, r)$ of r -dimensional quotients of the sheaf $\text{coker}(\sigma^*)$.

² Here $\Delta_b^a(c) := \det(c_{b+j-i})_{1 \leq i, j \leq a}$, as in [Ful98, Chapter 14].

³ Except when D_r is the *deepest* degeneracy locus, i.e. when $D_{r+1} = \emptyset$. Then $\tilde{D}_r \cong D_r$ and $\tilde{D}_r \rightarrow X$ is an embedding. This is the case studied in [GT20].

However, in examples like ((1.10) and (1.12)) below each kernel $\ker \sigma_x = h^0(E_\bullet|_x)$ naturally embeds in (the fibre over x of) a certain vector bundle $B \rightarrow X$,

$$h^0(E_\bullet|_x) \subseteq B_x \quad \forall x \in X. \tag{1.4}$$

The global condition is that B^* surjects onto $\text{coker}(\sigma^*) = h^0(E_\bullet^\vee)$ over D_r (2.18). Then \tilde{D}_r naturally embeds in $\text{Gr}(r, B)/X$ and we can push forward the virtual class.

THEOREM 1. *Under the embedding $\iota: \tilde{D}_r(E_\bullet) \hookrightarrow \text{Gr}(r, B)$, the pushforward of the virtual cycle to $A_{\text{vd}}(\text{Gr}(r, B))$ is given by*

$$\iota_* [\tilde{D}_r(E_\bullet)]^{\text{vir}} = \Delta_{\text{rk}(B) - \text{rk}(E_\bullet)}^r (c(\mathcal{Q} - q^*E_\bullet)), \tag{1.5}$$

where \mathcal{Q} is the universal quotient bundle over $q: \text{Gr}(r, B) \rightarrow X$.

1.4 Nested Hilbert schemes

For simplicity consider first *two-step* Hilbert schemes of nested subschemes (of dimensions 0 and 1) of a fixed smooth projective surface S . Given $\beta \in H^2(S, \mathbb{Z})$ and integers $n_1, n_2 \geq 0$, we set

$$S_\beta^{[n_1, n_2]} := \{I_1(-D) \subseteq I_2 \subseteq \mathcal{O}_S : [D] = \beta, \text{length}(\mathcal{O}_S/I_i) = n_i\}. \tag{1.6}$$

When $n_1 = 0 = n_2$ we use $S_\beta = S_\beta^{[0, 0]}$ to denote the Hilbert scheme of divisors in class β . Conversely when $\beta = 0$ we get the nested Hilbert scheme of points $S^{[n_1, n_2]}$ studied in [GT20]; we review this briefly first.

1.5 Nested Hilbert schemes of points

For simplicity we set $H^{\geq 1}(\mathcal{O}_S) = 0$ for now. For points $I_1, I_2 \subset \mathcal{O}_S$ of $S^{[n_1]}, S^{[n_2]}$, we have

$$\text{Hom}(I_1, I_2) = \begin{cases} \mathbb{C}, & I_1 \subseteq I_2, \\ 0, & I_1 \not\subseteq I_2. \end{cases} \tag{1.7}$$

Therefore, $\iota: S^{[n_1, n_2]} \hookrightarrow S^{[n_1]} \times S^{[n_2]}$ is the degeneracy locus $D_1(E_\bullet)$ of the two-term complex⁴ of vector bundles

$$E_\bullet = R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2) \quad \text{over } S^{[n_1]} \times S^{[n_2]},$$

which, when restricted to the point (I_1, I_2) , has 0th cohomology (1.7). Moreover, (1.7) shows that all higher degeneracy loci are empty, so we are in the situation of Footnote 3 with $r = 1$. Thus, the Thom–Porteous formula gives

$$\iota_* [S^{[n_1, n_2]}]^{\text{vir}} = c_{n_1+n_2}(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2)[1]) \cap [S^{[n_1]} \times S^{[n_2]}].$$

In [GT20] we also showed that the relevant perfect obstruction theory agrees (in K-theory at least) with that coming from Vafa–Witten theory. Therefore,⁵ Vafa–Witten invariants can

⁴ This uses our simplifying assumption $H^{\geq 1}(\mathcal{O}_S) = 0$. (For general S we modify E_\bullet in [GT20], removing $H^{\geq 1}(\mathcal{O}_S)$ terms to make it two-term.) The notation will be defined in §4; the \mathcal{I}_i are the universal ideal sheaves on $S \times S^{[n_1]} \times S^{[n_2]}$ and π denotes the projection down S to $S^{[n_1]} \times S^{[n_2]}$.

⁵ Siebert’s formula [Sie04, Theorem 4.6] for the virtual class shows that it depends on the perfect obstruction theory only through the K-theory class of its virtual tangent bundle.

be written as integrals over the smooth ambient space $S^{[n_1]} \times S^{[n_2]}$, as exploited to great effect in [Laa18, Laa20] for instance.

1.6 Nested Hilbert schemes of curves and points

When $\beta \neq 0$ we consider

$$\text{Hom}(I_1(-D), I_2) \quad \text{for } D \text{ in class } \beta. \tag{1.8}$$

In contrast to (1.7), this can become arbitrarily big, with different elements corresponding (up to scale) to different nested subschemes of S . Therefore, the corresponding nested Hilbert scheme $S_\beta^{[n_1, n_2]}$ dominates the degeneracy locus $D_1(E_\bullet)$ of the complex of vector bundles

$$E_\bullet = R\mathcal{H}om_\pi(\mathcal{I}_1(-D), \mathcal{I}_2) \quad \text{over } S^{[n_1]} \times S^{[n_2]}, \tag{1.9}$$

which, when restricted to the point $(I_1, I_2, \mathcal{O}(D))$, has 0th cohomology (1.8). Since a point of the nested Hilbert scheme $S_\beta^{[n_1, n_2]}$ is a one-dimensional subspace of the kernel (1.8) of the complex (1.9), we see that

$$S_\beta^{[n_1, n_2]} = \tilde{D}_1(E_\bullet)$$

is the virtual resolution of the degeneracy locus $D_1(E_\bullet)$ with $r = 1$. Since

$$\text{Hom}(I_1(-D), I_2) \subseteq H^0(\mathcal{O}(D)), \tag{1.10}$$

the 0th cohomology of E_\bullet embeds in $H^0(\mathcal{O}(D))$. So, we get an example of (1.4) with $r = 1$ and B the trivial vector bundle with fibre $H^0(\mathcal{O}(D))$. The embedding $\tilde{D}_1(E_\bullet) \hookrightarrow \mathbb{P}(B)$ of Theorem 1 then becomes the following.

THEOREM 2. *Suppose that $H^{\geq 1}(\mathcal{O}_S) = 0$ for now. Then $S_\beta^{[n_1, n_2]}$ has a virtual cycle of dimension $n_1 + n_2 + h^0(\mathcal{O}(D)) - 1$, which, under the embedding*

$$\iota : S_\beta^{[n_1, n_2]} \hookrightarrow S^{[n_1]} \times S^{[n_2]} \times \mathbb{P}(H^0(\mathcal{O}(D))),$$

pushes forward to

$$\iota_* [S_\beta^{[n_1, n_2]}]^{\text{vir}} = c_{n_1+n_2}(-\mathcal{O}_{\mathbb{P}(H^0(\mathcal{O}(D)))}(-1) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2(D))).$$

1.7 Reduced virtual cycle

For general S with possibly nonzero $H^{\geq 1}(\mathcal{O}_S)$, we work instead with

$$E_\bullet = R\mathcal{H}om_\pi(\mathcal{I}_1(-\mathcal{L}_\beta), \mathcal{I}_2) \quad \text{over } X := S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S), \tag{1.11}$$

which, when restricted to the point $(I_1, I_2, \mathcal{O}(D))$, has 0th cohomology (1.8). Here \mathcal{L}_β is a Poincaré line bundle over $S \times \text{Pic}_\beta(S)$, normalised (by tensoring by the pullback of $\mathcal{L}_\beta^{-1}|_{\{x\} \times \text{Pic}_\beta(S)}$ if necessary) so that $\mathcal{L}_\beta|_{\{x\} \times \text{Pic}_\beta(S)}$ is trivial on some fixed basepoint $x \in S$.

Again, a point of $S_\beta^{[n_1, n_2]}$ is a one-dimensional subspace of the kernel (1.8) of the complex (1.11), so

$$S_\beta^{[n_1, n_2]} = \tilde{D}_1(E_\bullet)$$

is the virtual resolution of the degeneracy locus $D_1(E_\bullet)$ with $r = 1$. This gives a description of $S_\beta^{[n_1, n_2]}$ as a projective cone over $S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$; see Proposition 4.7, generalising [DKO07, Lemma 2.15].

We can no longer take B to have fibre $H^0(L)$ over $L \in \text{Pic}_\beta(S)$ since this may jump in dimension. Instead we fix a sufficiently ample divisor $A \subset S$ and let B be the bundle with fibre $H^0(L(A))$:

$$B := \pi_*(\mathcal{L}_\beta(A)) \quad \text{over } X.$$

Therefore, the inclusions

$$\text{Hom}(I_1(-L), I_2) \subseteq H^0(L) \subseteq H^0(L(A)) \tag{1.12}$$

give the required embeddings $h^0(E_\bullet|_x) \subset B_x$ of (1.4). Thus, $\tilde{D}_1 \subset \mathbb{P}(B)$ such that $\mathcal{O}_{\mathbb{P}(B)}(-1)$ restricts to $\mathcal{U} \rightarrow \tilde{D}_1$ and Theorem 1 gives the following.

THEOREM 3. *Fix a surface S and $\beta \in H^2(S, \mathbb{Z})$.*

- *If $H^2(L) = 0$ for all effective $L \in \text{Pic}_\beta(S)$, then $S_\beta^{[n_1, n_2]} \cong \tilde{D}_1(E_\bullet)$ inherits a reduced virtual cycle of dimension $\chi(L) + n_1 + n_2 + h^1(\mathcal{O}_S) - 1$.*
- *If $H^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$, then via $\iota: S_\beta^{[n_1, n_2]} = \tilde{D}_1(E_\bullet) \hookrightarrow \mathbb{P}(B)$ it pushes forward to*

$$\iota_*[S_\beta^{[n_1, n_2]}]^{\text{red}} = c_{n_1+n_2+d}(q^*B(1) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))),$$

where $q: \mathbb{P}(B) \rightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$ is the projection, $\mathcal{O}(1) := \mathcal{O}_{\mathbb{P}(B)}(1)$ and $d = \frac{1}{2}A(2\beta + A - K_S) = \chi(L(A)) - \chi(L)$ for any $L \in \text{Pic}_\beta(S)$.

The $H^2(L) = 0$ condition ensures that E_\bullet is two-term. (We handle $H^2(L) \neq 0$ in the next section.) When $n_1 = 0$, this theorem was proven in [KT14a, Theorem A.7]. Here, as in [KT14a, GSY20a], we call the virtual cycle the *reduced* cycle $[S_\beta^{[n_1, n_2]}]^{\text{red}}$ since it differs from the one appearing in Vafa–Witten theory⁶ by an $H^2(\mathcal{O}_S)$ term in the obstruction theory.

The apparent difference between the formulae in Theorems 2 and 3 will be explained by the formula (2.25) in Theorem 2.24.

1.8 Nonreduced virtual cycle

In general, when $H^2(L)$ need not vanish, E_\bullet (1.11) is a three-term complex with nonzero $h^2(E_\bullet)$ cohomology sheaf. So, in order to apply our theory we use the *splitting trick* from [GT20, §6.1] to remove $R^2\pi_*\mathcal{O}[-2]$ from $E_\bullet(1)$ after pulling back to an affine bundle over X . (Since an affine bundle has the same Chow groups as its base, no information is lost from the virtual cycle.)

We only manage this on a neighbourhood of $S_\beta^{[n_1, n_2]} \subset \mathbb{P}(B)$, making $E_\bullet(1)$ a two-term complex there. By Proposition 1, this is enough to prove the following.

THEOREM 4. *This construction defines a virtual fundamental class*

$$[S_\beta^{[n_1, n_2]}]^{\text{vir}} \in A_{n_1+n_2+\text{vd}(S_\beta)}(S_\beta^{[n_1, n_2]}), \tag{1.13}$$

where $\text{vd}(S_\beta) = \frac{1}{2}\beta(\beta - K_S)$. It agrees with the virtual cycle of [GSY20a].

⁶ Or indeed reduced DT theory, confusingly. Removing *one* copy of $H^2(\mathcal{O}_S)$ from the DT obstruction space of $S_\beta^{[n_1, n_2]}$ gives the Vafa–Witten/reduced DT obstruction theory. Removing *two* copies gives the degeneracy locus obstruction theory of Proposition 1.

1.9 Computing virtual cycles by comparison

Since the above neighbourhood need not be all of $\mathbb{P}(B)$, we get no useful pushforward formula for (1.13) in general.⁷ This problem is down to the curve class β alone; if $\beta = 0$ it does not arise, whereas if $n_1 = 0 = n_2$ the problem already arises for the Hilbert scheme S_β of pure curves in class β . However,

$$[S_\beta]^{\text{vir}} \in A_{\text{vd}_\beta}(S_\beta), \quad \text{vd}_\beta := \text{vd}(S_\beta) = \frac{1}{2}\beta(\beta - K_S),$$

is a well-studied class that has been understood by other methods [DKO07, Koo16] due to its importance in Seiberg–Witten theory. Separating out the parts of the obstruction theory governing the curve and the points, and applying the degeneracy locus technique to the latter only, we are able to prove the following comparison result.

Let $\mathcal{D}_\beta \subset S \times S_\beta$ denote the universal curve. We call

$$\text{CO}_\beta^{[n_1, n_2]} := R\pi_* \mathcal{O}(\mathcal{D}_\beta) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2(\mathcal{D}_\beta)) \tag{1.14}$$

the Carlsson–Okounkov K-theory class [CO12] on $S^{[n_1]} \times S^{[n_2]} \times S_\beta$.

THEOREM 5. *Pushing the virtual and reduced classes forwards along*

$$\iota: S_\beta^{[n_1, n_2]} \hookrightarrow S^{[n_1]} \times S^{[n_2]} \times S_\beta$$

gives

$$\iota_* [S_\beta^{[n_1, n_2]}]^{\text{vir}} = c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) \cap [S^{[n_1]} \times S^{[n_2]}] \times [S_\beta]^{\text{vir}}$$

and, if $H^2(L) = 0$ for all effective $L \in \text{Pic}_\beta(S)$,

$$\iota_* [S_\beta^{[n_1, n_2]}]^{\text{red}} = c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) \cap [S^{[n_1]} \times S^{[n_2]}] \times [S_\beta]^{\text{red}}.$$

When $n_1 = 0$ the first of these results was proved in [Koo16] and the second follows from [KT14a, Appendix A].

1.10 ℓ -step Hilbert schemes

The generalisation of Theorems 3 and 5 to ℓ -step nested Hilbert schemes

$$S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}$$

is straightforward. See § 5 for details.

1.11 Vafa–Witten invariants

Vafa–Witten invariants [TT20] are made up of ‘instanton contributions’ and ‘monopole contributions’. The former are virtual Euler characteristics of moduli spaces of rank r sheaves on S , as studied in [GK20] for instance. The latter are integrals over moduli spaces of chains of sheaves on S of total rank r with nonzero maps between them.

When $p_g(S) > 0$ and r is prime, a vanishing result [Tho20, Theorem 5.23] implies that the only nonzero monopole contributions come from moduli spaces of chains of rank 1 sheaves. After

⁷ The exceptions are when $p_g(S) = 0$; then (1.13) is the reduced virtual class of Theorem 3, or when $H^2(L) = 0$ for all effective $L \in \text{Pic}_\beta(S)$, so (1.13) is zero, as the perfect obstruction theory which produced it can be reduced to give the class $[S_\beta^{[n_1, n_2]}]^{\text{red}}$.

tensoring with a line bundle, these are nested Hilbert schemes. For instance, in rank 2 the relevant integrals take the form

$$\int_{[S_\beta^{[n_1, n_2]}]^{vir}} \sum_i \rho^* \alpha_i \cup h^i = \sum_i \int_{\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir})} \alpha_i, \tag{1.15}$$

where $\alpha_i \in H^*(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$, $h := c_1(\mathcal{O}_{\mathbb{P}(B)}(1))$ and

$$\rho: S_\beta^{[n_1, n_2]} \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S).$$

Using Theorem 5, we can express this in terms of (a) integrals over the smooth space $S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$ and (b) integrals over S_β . The latter give the Seiberg–Witten invariants [DKO07, CK13],

$$\text{SW}_\beta := \begin{cases} \text{deg } [S_\beta]^{vir}, & \text{vd}_\beta = \frac{1}{2}\beta \cdot (K_S - \beta) = 0, \\ 0, & \text{otherwise.} \end{cases}$$

THEOREM 6. Suppose that $p_g(S) > 0$. Then, in $H_*(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$,

$$\begin{aligned} \rho_* [S_\beta^{[n_1, n_2]}]^{vir} &= \text{SW}_\beta \cdot c_{n_1+n_2}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)) \times [L], \\ \rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir}) &= 0 \quad \text{for } i > 0, \end{aligned}$$

where $L \in \text{Pic}_\beta(S)$, so $[L]$ is the generator of $H_0(\text{Pic}_\beta(S), \mathbb{Z})$.

When $p_g(S) = 0$, then S may not be of ‘Seiberg–Witten simple type’, which means that we have to consider higher dimensional Seiberg–Witten moduli spaces. Letting $\text{AJ}: S_\beta \rightarrow \text{Pic}_\beta(S)$ denote the Abel–Jacobi map, the higher Seiberg–Witten invariants [DKO07] are

$$\text{SW}_\beta^j := \text{AJ}_*(h^j \cap [S_\beta]^{vir}) \in H_{2(\text{vd}_\beta - j)}(\text{Pic}_\beta(S)) = \wedge^{2(\text{vd}_\beta - j)} H^1(S).$$

THEOREM 7. If $p_g(S) = 0$, then, in $H_{2(n_1+n_2+\text{vd}_\beta-i)}(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$:

•

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir}) = \sum_{j=0}^{n_1+n_2} c_{n_1+n_2-j}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cup \text{SW}_\beta^{i+j};$$

• if $H^2(L) = 0 \forall L \in \text{Pic}_\beta(S)$, then, setting $d = n_1 + n_2 + h^1(\mathcal{O}_S) - \text{vd}_\beta$,

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir}) = c_{d+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta));$$

• if not,

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir}) = \text{SW}_\beta \cdot c_{n_1+n_2+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)) \times [L]$$

and both sides vanish when $i > 0$ (cf. Theorem 6).

Now set $\beta^\vee := K_S - \beta$, so that the condition that $H^2(L) \neq 0$ for some $L \in \text{Pic}_\beta(S)$ is the condition that β^\vee is effective. Seiberg–Witten theory has a duality under $\beta \leftrightarrow \beta^\vee$. This gives interesting dualities between invariants of nested Hilbert schemes under $\beta \leftrightarrow \beta^\vee$, $n_1 \leftrightarrow n_2$. Define

the map

$$\rho^\vee : S_{\beta^\vee}^{[n_2, n_1]} \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$$

by replacing $\beta \leftrightarrow \beta^\vee$, $n_1 \leftrightarrow n_2$ in $\rho : S_\beta^{[n_1, n_2]} \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$ and then composing with $L \mapsto K_S \otimes L^{-1} : \text{Pic}_{\beta^\vee}(S) \rightarrow \text{Pic}_\beta(S)$.

THEOREM 8 (Duality). *In $H_{2(n_1+n_2+\text{vd}_\beta-i)}(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$:*

- if $p_g(S) > 0$,

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) = (-1)^{s+i} \rho_*^\vee(h^i \cap [S_{\beta^\vee}^{[n_2, n_1]}]^{\text{vir}});$$

- if $p_g(S) = 0$,

$$\begin{aligned} \rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) &= (-1)^{s+i} \rho_*^\vee(h^i \cap [S_{\beta^\vee}^{[n_2, n_1]}]^{\text{vir}}) \\ &\quad + c_{d+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)), \end{aligned}$$

where $s = n_1 + n_2 - \chi(\mathcal{O}_S) - \text{vd}_\beta = d - 1$.

THEOREM 9. *If $p_g(S) > 0$, the contributions of the nested Hilbert schemes $S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}$ to the Vafa–Witten invariants of S are invariants of its oriented diffeomorphism type.*

When $p_g(S) = 0$, this can fail because only the unordered pair $(\text{SW}_\beta, \text{SW}_{\beta^\vee})$ is invariant under oriented diffeomorphisms, rather than the individual invariant SW_β .

1.12 Laarakker

In [Laa18] (stable case) and [Laa20] (general case), Laarakker used these results to compute Vafa–Witten invariants [TT17, TT20] and refined Vafa–Witten invariants [Tho20] on surfaces with $p_g > 0$ (and $h^{0,1} = 0$ for now).

By the comparison result for virtual cycles of Theorem 5, the contributions from points and curves split, with the curves contributing Seiberg–Witten invariants, i.e. well-understood integrals over linear systems of curves. Laarakker evaluated the contributions of Hilbert schemes of points via the method of [EGL01]. The result depends only on the curve class $\beta \in H_2(S, \mathbb{Z})$ and the cobordism class of the surface, and thus only on $c_1(S)^2$, $c_2(S)$ and β^2 . Therefore, these contributions can be calculated on K3 surfaces and toric surfaces (despite these not having $p_g > 0$!).

The results are very general, but the absolute simplest to state is for S minimal of general type with $p_g(S) > 0$ and $H_1(S, \mathbb{Z}) = 0$ such that K_S is not divisible by 2.

THEOREM 10 [Laa18]. *Let S be as above. The monopole branch contributions⁸ of rank 2 Higgs pairs with $\det = K_S$ to the refined Vafa–Witten generating series $\sum_n \text{VW}_{2, K_S, n}(t) q^n$ can be written*

$$A(t, q)^{\chi(\mathcal{O}_S)} B(t, q)^{c_1(S)^2},$$

where

$$A(t, q), B(t, q) \in \mathbb{Q}(t^{1/2})(q^{1/2})$$

are universal functions, independent of S .

⁸ See [Tho20] for definitions. The monopole branch contributions come from \mathbb{C}^* -fixed Higgs pairs (E, ϕ) on S with $\det E = K_S$, $\text{tr } \phi = 0$ and $\phi \neq 0$.

Furthermore, K3 calculations [GK20, Tho20] determine $A(t, q)$ completely in terms of modular forms. And, $B(t, q)$ can be predicted by modularity and the results of Göttsche–Kool [GK20] on the instanton contributions; they checked this prediction in low degree by toric computations.

1.13 Notation

Given a map $f: X \rightarrow Y$, we often use the same letter f to denote its base change by any map $Z \rightarrow Y$, i.e. $f: X \times_Y Z \rightarrow Z$. In particular, projections $S \times M \rightarrow M$ down the surface S are all denoted by π . We often suppress pullback maps f^* on sheaves when it should not cause confusion.

For a vector bundle B over X , we denote by $\mathbb{P}(B)$ the projective bundle of lines in B . For a coherent sheaf \mathcal{F} on X , we denote by $\mathbb{P}^*(\mathcal{F}) = \text{Proj Sym}^\bullet \mathcal{F}$ the projective cone of quotient lines of \mathcal{F} . When \mathcal{F} is locally free, then of course $\mathbb{P}^*(\mathcal{F}) \cong \mathbb{P}(\mathcal{F}^*)$.

1.14 Splitting trick

Since we are ultimately interested in Chern class formulae, we often only care about the topological type of a given algebraic vector bundle. In particular, we will often wish to split nontrivial extensions of vector bundles, which does not change their topological type. One way to do this is by working with C^∞ bundles; we prefer to stay within algebraic geometry and use the *Jouanolou trick* [Jou73] instead.

Namely, if Y is quasi-projective, there exists an *affine bundle* $\rho: \tilde{Y} \rightarrow Y$ such that \tilde{Y} is an affine variety. Therefore, on pulling locally free sheaves back by ρ^* , they become *projective* $\mathcal{O}_{\tilde{Y}}$ -modules and any extensions become split. But ρ is a homotopy equivalence, inducing an isomorphism of Chow groups [Kre99, Corollary 2.5.7]

$$\rho^*: A_*(Y) \xrightarrow{\sim} A_{*+\dim \rho}(\tilde{Y}), \tag{1.16}$$

so formulae for cycles upstairs on \tilde{Y} induce similar formulae on Y .

A very explicit such affine bundle was used in [GT20, § 6.1] and all the necessary pullbacks were laboriously documented. In this paper, for the sake of clarity and economy of notation, we will often just say ‘*by the Jouanolou trick*’ and give no further details. So, we will pretend we are working on Y , suppressing mention of the fact that we have replaced Y by its homotopic affine model \tilde{Y} and then used the isomorphism (1.16) to recover results on Y at the end.

2. Virtual resolutions of degeneracy loci

Fix a smooth complex quasi-projective variety X and

$$\sigma: E_0 \longrightarrow E_1 \tag{2.1}$$

a map of vector bundles of ranks e_0 and e_1 over X . For each positive integer $r \leq e_0$, we let

$$D_r(\sigma) := \{x \in X: \dim \ker(\sigma_x) \geq r\}$$

denote the r th degeneracy locus of σ with scheme structure defined by the vanishing of

$$\wedge^{e_0-r+1} \sigma: \wedge^{e_0-r+1} E_0 \longrightarrow \wedge^{e_0-r+1} E_1. \tag{2.2}$$

This is the $(e_1 - e_0 + r - 1)$ th Fitting scheme [Eis94, § 20.2] of the first cohomology sheaf $h^1(E_\bullet) = \text{coker } \sigma$ of the complex E_\bullet (2.1). In particular, it depends on E_\bullet only up to

quasi-isomorphism and we often denote it by $D_r(E_\bullet)$. Then define

$$\tilde{D}_r(E_\bullet) = \tilde{D}_r(\sigma) := \text{Gr}(\text{coker } \sigma^*, r) \tag{2.3}$$

to be the relative Grassmannian of r -dimensional quotients of the cokernel sheaf $h^0(E_\bullet^\vee)$ of $\sigma^*: E_1^* \rightarrow E_0^*$. This is the moduli space of r -dimensional subspaces of fibres $(E_0)_x$ which are annihilated by σ , in the following sense.

Consider the functor

$$\begin{aligned} \text{Schemes}/X &\longrightarrow \text{Sets}, \\ (f: S \rightarrow X) &\longmapsto \{ \text{rank } r \text{ subbundles } U \subseteq f^*E_0 \text{ which are} \\ &\text{subsheaves of } h^0(f^*E_\bullet) \subseteq f^*E_0 \}. \end{aligned} \tag{2.4}$$

Let $p: \text{Gr}(r, E_0) \rightarrow X$ denote the relative Grassmannian with universal subbundle $\mathcal{U} \subset p^*E_0$.

PROPOSITION 2.5. *The functor (2.4) is represented by $\tilde{D}_r(\sigma)/X$. It embeds*

$$\iota: \tilde{D}_r \hookrightarrow \text{Gr}(r, E_0) \tag{2.6}$$

as the zero locus of $\tilde{\sigma} \in \Gamma(\mathcal{U}^* \otimes p^*E_1)$ defined by the composition

$$\mathcal{U} \hookrightarrow p^*E_0 \xrightarrow{p^*\sigma} p^*E_1. \tag{2.7}$$

Finally, the projection $p: \tilde{D}_r \rightarrow X$ has scheme-theoretic image inside D_r .

Remarks. Clearly, $p: \tilde{D}_r(\sigma) \rightarrow D_r(\sigma)$ is a set-theoretic surjection: the fibre over $x \in D_r(\sigma)$ is $\text{Gr}(r, \ker \sigma_x)$, but it need not be onto as a map of schemes. For instance, $x.\text{id}: \mathcal{O}^{\oplus 2} \rightarrow \mathcal{O}^{\oplus 2}$ on \mathbb{C}_x has $D_1 = (x^2 = 0) \subset \mathbb{C}_x$, the thickened origin, whereas \tilde{D}_1 is a reduced \mathbb{P}^1 over the origin.

By the definition of the functor (2.4), $\tilde{D}_r(\sigma)$ comes equipped with a universal subbundle $\mathcal{U} \subseteq p^*E_0$. It is the restriction of the universal subbundle on $\text{Gr}(r, E_0)$ and the dual of the universal quotient bundle on $\text{Gr}(\text{coker}(\sigma^*), r)$. Its fibre over the point $(x, U \subseteq \ker(\sigma_x))$ is U .

Proof. Since $\text{Gr}(\text{coker } \sigma^*, r)/X$ represents the functor

$$\begin{aligned} \text{Schemes}/X &\longrightarrow \text{Sets}, \\ (f: S \rightarrow X) &\longmapsto \{ \text{rank } r \text{ locally free quotients of } \text{coker } f^*\sigma^* \}, \end{aligned} \tag{2.8}$$

it is sufficient to prove that this is isomorphic to (2.4). A locally free quotient Q of $\text{coker}(f^*\sigma^*)$ gives a complex

$$f^*E_1^* \xrightarrow{f^*\sigma^*} f^*E_0^* \longrightarrow Q \longrightarrow 0, \tag{2.9}$$

which is exact at Q . Since Q is locally free, the dual complex

$$0 \longrightarrow U \longrightarrow f^*E_0 \xrightarrow{f^*\sigma} f^*E_1 \tag{2.10}$$

is exact at $U := Q^*$ after any base change: thus, $U \hookrightarrow f^*E_0$ is a subbundle, giving an element of the set (2.4). The converse is easier: any complex (2.10) with U a subbundle dualises to a complex (2.9) exact at $Q := U^*$.

The functor (2.4) is a subfunctor of the functor taking $f: S \rightarrow X$ to the set $\{\text{rank } r \text{ subbundles } U \subset f^*E_0\}$ represented by $\text{Gr}(r, E_0)/X$. This induces an embedding of X -schemes

$\tilde{D}_r(\sigma) \hookrightarrow \text{Gr}(r, E_0)$ under which the universal subbundle \mathcal{U} pulls back to the dual \mathcal{Q}^* of the universal quotient bundle. Its image is contained in the zero locus Z of (2.7) by applying (2.4) to $S = \tilde{D}_r$. So, $\tilde{D}_r(\sigma) \subseteq Z$.

Conversely, setting $f: Z \rightarrow X$ to be the projection, the restriction of (2.7) to Z gives a rank r subbundle $\mathcal{U}|_Z \subset f^*E_0$, which factors through $\ker(f^*\sigma)$ by the definition of Z as the zero locus. This gives an element of the set (2.4), classified by a map $Z \rightarrow \tilde{D}_r(\sigma)$. The two maps are mutual inverses by construction.

Finally, $p^*\sigma$ factors through p^*E_0/\mathcal{U} over $Z = \tilde{D}_r(\sigma)$ by (2.7). Therefore, $p^* \wedge^{e_0-r+1} \sigma$ (2.2) factors through $\wedge^{e_0-r+1} p^*E_0/\mathcal{U}$. But this is zero because $\text{rank}(p^*E_0/\mathcal{U}) = e_0 - r$, so $\tilde{D}_r(\sigma)$ projects into the zero locus D_r of (2.2). \square

2.1 Perfect obstruction theory

Let $I \subset \mathcal{O}_{\text{Gr}(r, E_0)}$ denote the ideal of \tilde{D}_r generated by $\tilde{\sigma}$ (2.7). Then \tilde{D}_r inherits the standard perfect obstruction theory as in the following diagram.

$$\begin{array}{ccc}
 \mathcal{U} \otimes p^*E_1^*|_{\tilde{D}_r} & \xrightarrow{d\tilde{\sigma}|_{\tilde{D}_r}} & \Omega_{\text{Gr}}|_{\tilde{D}_r} & & \mathbb{L}_{\tilde{D}_r}^{\text{vir}} \\
 \tilde{\sigma} \downarrow & & \parallel & = & \downarrow \\
 I/I^2 & \xrightarrow{d} & \Omega_{\text{Gr}}|_{\tilde{D}_r} & & \mathbb{L}_{\tilde{D}_r}.
 \end{array} \tag{2.11}$$

That is, the complex on the bottom row is a representative of the truncated cotangent complex $\mathbb{L}_{\tilde{D}_r}$ and the top row is the virtual cotangent bundle of the obstruction theory.

PROPOSITION 2.12. *The virtual cotangent bundle $\mathbb{L}_{\tilde{D}_r}^{\text{vir}}$ is a cone*

$$\text{Cone} \{ \mathcal{U} \otimes p^*E_\bullet^\vee|_{\tilde{D}_r}[-1] \longrightarrow (\mathcal{U}^* \otimes \mathcal{U}[-1] \oplus p^*\Omega_X)|_{\tilde{D}_r} \}$$

of virtual dimension $\text{vd} := \dim X - r(e_1 - e_0 + r)$. It depends only on the quasi-isomorphism class of the complex E_\bullet (1.1). The pushforward of the resulting virtual cycle

$$[\tilde{D}_r(E_\bullet)]^{\text{vir}} \in A_{\text{vd}}(\tilde{D}_r(E_\bullet))$$

to X is given by the Thom–Porteous formula

$$\Delta_{r-\text{rk}(E_\bullet)}^r(c(-E_\bullet)) \in A_{\text{vd}}(X).$$

Proof. Modifying $\mathbb{L}_{\tilde{D}_r}^{\text{vir}}$ (2.11) by an acyclic complex, it is quasi-isomorphic to the total complex

$$\begin{array}{ccc}
 \mathcal{U} \otimes p^*E_1^*|_{\tilde{D}_r} & \xrightarrow{d\tilde{\sigma}|_{\tilde{D}_r}} & \Omega_{\text{Gr}}|_{\tilde{D}_r} \\
 \text{id}_{\mathcal{U}} \otimes \downarrow p^*\sigma^* & & \downarrow \phi \\
 \mathcal{U} \otimes p^*E_0^*|_{\tilde{D}_r} & \xlongequal{\quad} & \mathcal{U} \otimes p^*E_0^*|_{\tilde{D}_r},
 \end{array} \tag{2.13}$$

where ϕ is the composition $\Omega_{\text{Gr}} \cong \mathcal{U} \otimes (p^*E_0/\mathcal{U})^* \rightarrow \mathcal{U} \otimes p^*E_0^*$. This diagram was shown to commute in [GT20, Claim 2, proof of Theorem 3.6].

Therefore, $\mathbb{L}_{\tilde{D}_r}^{\text{vir}} \cong \text{Cone}(\mathcal{U} \otimes p^*E_\bullet^\vee|_{\tilde{D}_r} \rightarrow \text{Cone}(\phi))[-1]$. To determine $\text{Cone}(\phi)$, we fit ϕ into the following commutative diagram, all of whose rows and columns are exact triangles.

$$\begin{array}{ccccc}
 p^*\Omega_X|_{\tilde{D}_r} & \longrightarrow & \Omega_{\text{Gr}}|_{\tilde{D}_r} & \longrightarrow & \Omega_{\text{Gr}/X}|_{\tilde{D}_r} \\
 & & \phi \downarrow & & \downarrow \\
 & & \mathcal{U} \otimes p^*E_0^*|_{\tilde{D}_r} & \cong & \mathcal{U} \otimes p^*E_0^*|_{\tilde{D}_r} \\
 & & \downarrow & & \downarrow \\
 & & \text{Cone}(\phi) & \longrightarrow & \mathcal{U} \otimes \mathcal{U}^*|_{\tilde{D}_r} \longrightarrow p^*\Omega_X|_{\tilde{D}_r}[2]
 \end{array} \tag{2.14}$$

We calculate the lower right hand arrow. By definition, it is the composition from the top left to the bottom right of the following diagram.

$$\begin{array}{ccccc}
 \mathcal{U} \otimes \mathcal{U}^*|_{\tilde{D}_r} & \longrightarrow & \Omega_{\text{Gr}/X}|_{\tilde{D}_r}[1] & \longrightarrow & \mathcal{U} \otimes p^*E_0^*|_{\tilde{D}_r}[1] \\
 & & \parallel & & \\
 \Omega_{\text{Gr}}|_{\tilde{D}_r}[1] & \longrightarrow & \Omega_{\text{Gr}/X}|_{\tilde{D}_r}[1] & \longrightarrow & p^*\Omega_X|_{\tilde{D}_r}[2]
 \end{array} \tag{2.15}$$

Here the upper row is the rightmost vertical exact triangle in (2.14), while the lower row is the exact triangle along the top row of (2.14). The first arrow in (2.15) is well known to be the relative (to X) Atiyah class of the bundle \mathcal{U} . So, using its absolute Atiyah class, we get the commutative diagram

$$\begin{array}{ccccc}
 \mathcal{U} \otimes \mathcal{U}^*|_{\tilde{D}_r} & \xrightarrow{\text{At}_{\text{Gr}/X}(\mathcal{U})} & \Omega_{\text{Gr}/X}|_{\tilde{D}_r}[1] & \longrightarrow & \mathcal{U} \otimes p^*E_0^*|_{\tilde{D}_r}[1] \\
 \text{At}_{\text{Gr}}(\mathcal{U}) \downarrow & & \parallel & & \\
 \Omega_{\text{Gr}}|_{\tilde{D}_r}[1] & \longrightarrow & \Omega_{\text{Gr}/X}|_{\tilde{D}_r}[1] & \longrightarrow & p^*\Omega_X|_{\tilde{D}_r}[2]
 \end{array}$$

which shows that the composition $\mathcal{U} \otimes \mathcal{U}^* \rightarrow p^*\Omega_X[2]$ is zero. Therefore, the bottom row of (2.14) gives $\text{Cone}(\phi) \cong \mathcal{U} \otimes \mathcal{U}^* \oplus p^*\Omega_X|_{\tilde{D}_r}[1]$, implying the claimed result.

Finally, since the perfect obstruction theory arises from the \tilde{D}_r description as the zero locus of a section of the bundle $\mathcal{U}^* \otimes p^*E_1 \rightarrow \text{Gr}$, the resulting virtual cycle is the localised top Chern class [Ful98, § 14.1] of that bundle. Its pushforward to Gr is

$$c_{re_1}(\mathcal{U}^* \otimes p^*E_1); \tag{2.16}$$

further pushing down $p: \text{Gr} \rightarrow X$ shows that the pushforward of $[\tilde{D}_r]^{\text{vir}}$ to X is given by

$$p_* c_{re_1}(\mathcal{U}^* \otimes p^*E_1) = \Delta_{e_1 - e_0 + r}^r(c(-E_\bullet))$$

by [Ful98, Theorem 14.4]. □

This proves Proposition 1 in the Introduction. But, in passing from $\text{Gr}(r, E_0)$ down to X , we have contracted \tilde{D}_r back to D_r and so lost information in general. (The exception is the case studied in [GT20], where r is the largest integer for which D_r is nonempty; then $p|_{\tilde{D}_r}$ is an embedding.)

We could work in $\text{Gr}(r, E_0)$, in which the class of the virtual cycle is $c_{re_1}(\mathcal{U}^* \otimes p^*E_1)$, but this is not a quasi-isomorphism invariant of the complex E_\bullet . In examples such as (1.10) and (1.12), we can replace E_0 by a more canonical bundle B , so now we assume that we have chosen B such that

$$h^0(E_\bullet|_x) \hookrightarrow B_x \quad \text{for each } x \in D_r. \tag{2.17}$$

For base-change reasons the precise condition is most easily stated via duals.

Choice. We choose a vector bundle B on X and a surjection

$$B^*|_{D_r} \longrightarrow h^0(E_\bullet^\vee)|_{D_r} \longrightarrow 0. \tag{2.18}$$

For some purposes we will require that the surjection (2.18) extends to all of X ,

$$B^* \longrightarrow h^0(E_\bullet^\vee) \longrightarrow 0. \tag{2.19}$$

For instance, if $r = 1$, this is immediate from (2.18).

Note that by restricting both exact sequences $E_1^* \rightarrow E_0^* \rightarrow h^0(E_\bullet^\vee) \rightarrow 0$ and (2.18) to $x \in D_r$ and dualising, we recover (2.17).

The surjection $B^*|_{D_r} \rightarrow (\text{coker } \sigma^*)|_{D_r}$ of (2.18) induces an embedding $\text{Gr}(\text{coker } \sigma^*, r) \subset \text{Gr}(B^*, r)$ since $\text{Gr}(\text{coker } \sigma^*, r)$ lies over $D_r \subset X$. By Proposition 2.5, this is an embedding

$$\iota_B : \tilde{D}_r \hookrightarrow \text{Gr}(r, B) \xrightarrow{q} X \tag{2.20}$$

such that $\iota_B^* \mathcal{U}_B \cong \iota^* \mathcal{U}$. Here \mathcal{U}_B and $\mathcal{Q}_B = q^*B/\mathcal{U}_B$ denote the universal sub- and quotient bundles on $\text{Gr}(r, B)$, and $\iota : \tilde{D}_r \hookrightarrow \text{Gr}(r, E_0)$ is the embedding (2.6). We will compare these via a third embedding,

$$I := \iota \times_X \iota_B : \tilde{D}_r \hookrightarrow \text{Gr}(r, E_0) \times_X \text{Gr}(r, B),$$

fitting into the following Cartesian diagram.

$$\begin{array}{ccc} \tilde{D}_r \times_X \text{Gr}(r, B) & \xhookrightarrow{\iota \times 1} & \text{Gr}(r, E_0) \times_X \text{Gr}(r, B) \\ q \downarrow \uparrow s & \nearrow I & \downarrow q_0 \\ \tilde{D}_r & \xhookrightarrow{\iota} & \text{Gr}(r, E_0) \end{array} \tag{2.21}$$

Here the vertical maps are flat and the section $s := \text{id}_{\tilde{D}_r} \times_X \iota_B$. As usual we suppress various pullback or base-change maps.

LEMMA 2.22. *The section s in (2.21) is a regular embedding, with image cut out by the regular section of $\mathcal{U}^* \otimes \mathcal{Q}_B$ defined by the composition*

$$\mathcal{U} \hookrightarrow B \twoheadrightarrow \mathcal{Q}_B \quad \text{on } \tilde{D}_r \times_X \text{Gr}(r, B). \tag{2.23}$$

Proof. Since $s^*q^*\iota^*\mathcal{U} = \iota^*\mathcal{U} = s^*\mathcal{U}_B$ on $\tilde{D}_r \times_X \text{Gr}(r, B)$, we see that s^* applied to (2.23) gives the same as s^* applied instead to the composition

$$\mathcal{U}_B \hookrightarrow B \twoheadrightarrow \mathcal{Q}_B \quad \text{on } \tilde{D}_r \times_X \text{Gr}(r, B).$$

This is zero by the definition of \mathcal{Q}_B , so $s(\tilde{D}_r)$ lies in the zero locus Z of (2.23). We are left with showing that $Z \subseteq s(\tilde{D}_r)$.

By the definition of Z , pulling back the composition (2.23) by $\iota_Z: Z \hookrightarrow \tilde{D}_r \times_X \text{Gr}(r, B)$ shows that the composition along the top row of the following diagram is zero. This uniquely fills in the dotted arrow ϕ to the lower short exact sequence in the following diagram.

$$\begin{CD} 0 @>>> \iota_Z^* \mathcal{U} @>>> \iota_Z^* B @>>> \iota_Z^* \mathcal{Q}_B @>>> 0 \\ @. @V \phi VV @| @. @. \\ 0 @>>> \iota_Z^* \mathcal{U}_B @>>> \iota_Z^* B @>>> \iota_Z^* \mathcal{Q}_B @>>> 0 \end{CD}$$

Since ϕ is a vector bundle injection, it is an isomorphism. And, since $\mathcal{U} = \iota_B^* \mathcal{U}_B = q^* s^* \mathcal{U}_B$ is pulled back from \tilde{D}_r , this gives the following diagram.

$$\begin{CD} \iota_Z^* q^* s^* \mathcal{U}_B @>>> \iota_Z^* B \\ @| @| \\ \iota_Z^* \mathcal{U}_B @>>> \iota_Z^* B \end{CD}$$

Thus, this subbundle is represented by both of the maps ι_Z and $s \circ q \circ \iota_Z$, but any subbundle is classified by a unique map $Z \rightarrow \tilde{D}_r \times_X \text{Gr}(r, B)$. So, $\iota_Z = s \circ q \circ \iota_Z$, which gives $Z \subseteq s(\tilde{D}_r)$.

Since $\text{codim}(s(\tilde{D}_r)) = \dim \text{Gr}(r, B) - \dim X = \text{rank}(\mathcal{U}^* \otimes \mathcal{Q}_B)$, it follows that s is a regular embedding. □

THEOREM 2.24. *Let $b := \text{rank } B$. The pushforward of $[\tilde{D}_r]^{\text{vir}}$ to $\text{Gr}(r, B)$ is given by*

$$\iota_{B*} [\tilde{D}_r]^{\text{vir}} = \Delta_{b+e_1-e_0}^r (c(\mathcal{Q}_B - E_\bullet)).$$

If (2.19), holds this also equals

$$c_{r(b-e_0+e_1)}(\mathcal{U}_B^* \otimes (B - E_\bullet)). \tag{2.25}$$

Proof. By Proposition 2.5 and flat base change around the diagram (2.21),

$$(\iota \times 1)_* q^* [\tilde{D}_r]^{\text{vir}} = q_0^* \iota_* [\tilde{D}_r]^{\text{vir}} = q_0^* c_{re_1}(\mathcal{U}^* \otimes E_1).$$

Therefore, Lemma 2.22 gives

$$\begin{aligned} I_* [\tilde{D}_r]^{\text{vir}} &= (\iota \times 1)_* s_* [\tilde{D}_r]^{\text{vir}} \\ &= (\iota \times 1)_* s_* s^* q^* [\tilde{D}_r]^{\text{vir}} \\ &= (\iota \times 1)_* (c_{r(b-r)}(q^* \mathcal{U}^* \otimes \mathcal{Q}_B) \cap q^* [\tilde{D}_r]^{\text{vir}}) \\ &= c_{r(b-r)}(q_0^* \mathcal{U}^* \otimes \mathcal{Q}_B) \cap (\iota \times 1)_* q^* [\tilde{D}_r]^{\text{vir}} \\ &= c_{r(b-r)}(q_0^* \mathcal{U}^* \otimes \mathcal{Q}_B) \cap c_{re_1}(q_0^* \mathcal{U}^* \otimes E_1) \\ &= c_{r(b-r+e_1)}(q_0^* \mathcal{U}^* \otimes (\mathcal{Q}_B \oplus E_1)). \end{aligned}$$

Pushing down $p: \text{Gr}(r, E_0) \times_X \text{Gr}(r, B) \rightarrow \text{Gr}(r, B)$ gives

$$\iota_{B*} [\tilde{D}_r]^{\text{vir}} = \Delta_{b+e_1-e_0}^r (c(\mathcal{Q}_B + E_1 - E_0)) \tag{2.26}$$

by [Ful98, Proposition 14.2.2].

To prove (2.25), we need to lift the surjection $B^* \rightarrow \text{coker}(\sigma^*)$ of (2.19) to a map $\psi: B^* \rightarrow E_0^*$; see (2.27) below for how to do this after using the Jouanolou trick of § 1.14. This then gives $h^0(E_\bullet|_x) \hookrightarrow B_x$ for all $x \in X$, which means that

$$E_0 \xrightarrow{(\sigma, \psi^*)} E_1 \oplus B$$

is a subbundle. Therefore, its cokernel C is a rank $(b + e_1 - e_0)$ vector bundle over X , equivalent in K-theory to $B - E_\bullet$. Thus, we can apply [Ful98, Example 14.4.12] to rewrite (2.26) as

$$\begin{aligned} \iota_{B^*}[\tilde{D}_r]^{\text{vir}} &= \Delta_{b+e_1-e_0}^r(c(C - \mathcal{U}_B)) \\ &= c_{r(b+e_1-e_0)}(\mathcal{U}_B^* \otimes C) \\ &= c_{r(b+e_1-e_0)}(\mathcal{U}_B^* \otimes (B - E_\bullet)). \end{aligned}$$

This formula should be compared with (2.16). □

2.2 Digression: description as a deepest degeneracy locus

As usual fix a smooth quasi-projective variety X carrying a two-term complex of bundles

$$\sigma: E_0 \longrightarrow E_1.$$

Almost by definition the virtual resolution \tilde{D}_r of the r th degeneracy locus can be seen as the deepest degeneracy locus of the complex $\mathcal{U} \rightarrow p^*E_1$ (1.3) up on $\text{Gr}(r, E_0)$. Such deepest degeneracy loci (those for which r is maximal, i.e. D_{r+1} is empty) were studied in [GT20].

Since E_0 is not a quasi-isomorphism invariant, we would again prefer to replace $\text{Gr}(r, E_0)$ with (an affine bundle over) $\text{Gr}(r, B)$. For this we require (2.19) to hold, giving the following diagram.

$$\begin{array}{ccccc} & & B^* & & \\ & & \downarrow & & \\ & \psi & & & \\ E_1^* & \xrightarrow{\sigma^*} & E_0^* & \longrightarrow & \text{coker}(\sigma^*) \longrightarrow 0 \end{array} \tag{2.27}$$

Using the Jouanolou trick described in § 1.14 we may, on replacing X by an affine bundle over it (and pulling everything back to it), assume that B^* is a projective \mathcal{O} -module. Then we may pick a lift $\psi: B^* \rightarrow E_0^*$.

Over $\text{Gr}(r, B)$, with its universal quotient bundle $\pi: B \rightarrow \mathcal{Q}_B$ (suppressing some pullback maps as usual), we consider the composition

$$E_0 \xrightarrow{(\sigma, \psi^*)} E_1 \oplus B \xrightarrow{(\text{id}, \pi)} E_1 \oplus \mathcal{Q}_B. \tag{2.28}$$

τ

At a closed point $(x, V) \in \text{Gr}(r, B)$ (i.e. a point $x \in X$ and an r -dimensional subspace $V \leq B_x$), we have

$$\ker \tau_x = \ker \sigma_x \cap \ker(\pi_x \circ \psi_x^*) = \ker \sigma_x \cap V,$$

where in the last expression we have used ψ_x^* to identify $\ker \sigma_x$ as a subspace of B_x by (2.17). In particular, $\ker \tau_x$ is at most r dimensional, so $D_r(\tau)$ is a deepest degeneracy locus, and the locus

of points where it is precisely r dimensional is

$$D_r(\tau) = \{(x, U) \in \text{Gr}(r, B) : U \subseteq \ker(\sigma_x)\} = \tilde{D}_r(\sigma). \tag{2.29}$$

This bijection of sets generalises to S -points of X : given $f: S \rightarrow X$, we get a bijection between the sets

$$\{\text{rank } r \text{ subbundles } U \subset f^*E_0 : U \hookrightarrow f^*E_0 \xrightarrow{f^*\sigma} f^*E_1 \text{ is zero}\}$$

and

$$\{\text{rank } r \text{ subbundles } (U \subset f^*E_0, V \subset f^*B) : \text{the composition } U \hookrightarrow f^*E_0 \xrightarrow{f^*\tau} f^*E_1 \oplus f^*B/V \text{ is zero}\}.$$

The functors taking $f: S \rightarrow X$ to either of these two sets are therefore isomorphic. The first is (2.4) and is represented by $\tilde{D}_r(\sigma)$ by Proposition 2.5. The second is represented by $D_r(\tau) \subset \text{Gr}(r, B)/X$. Therefore, (2.29) is an isomorphism of schemes.

Furthermore, we can compare the perfect obstruction theory of Proposition 2.12 for $\tilde{D}_r(\sigma)$ to the deepest degeneracy locus perfect obstruction theory of [GT20, Theorem 3.6] for $D_r(\tau)$. In both cases the K-theory class of the virtual cotangent bundle is the restriction of

$$\Omega_X + \mathcal{U} \otimes (E_\bullet^\vee - \mathcal{U}^*),$$

so their virtual cycles agree. Finally, the Thom–Porteous formula for $\tilde{D}_r(\sigma)$ of Theorem 2.24 and the Thom–Porteous formula for $D_r(\tau)$ of [GT20, Theorem 3.6] both give

$$\Delta_{b+e_1-e_0}^r(c(\mathcal{Q}_B - E_\bullet))$$

as the pushforward of the virtual cycle to $\text{Gr}(r, B)$.

THEOREM 2.30. *Suppose that (2.19) holds. Then, after pulling back to an affine bundle, the embedding $\iota_B: \tilde{D}_r(\sigma) \hookrightarrow \text{Gr}(r, B)$ of (2.20) is the deepest degeneracy locus of the map of vector bundles (2.28),*

$$\tilde{D}_r(\sigma) \cong D_r(\tau).$$

The two resulting virtual cycles and Thom–Porteous formulae agree.

2.3 Generalised Carlsson–Okounkov vanishing

PROPOSITION 2.31. *Under the assumptions of Theorem 2.30,*

$$c_{b+e_1-e_0+i}(B + E_1 - E_0) = 0 \quad \forall i > 0.$$

Proof. On an affine bundle over X , recall the map (2.28)

$$E_0 \xrightarrow{(\sigma, \psi^*)} E_1 \oplus B \tag{2.32}$$

of the last section. By (2.19), it is an injective map of bundles and so is quasi-isomorphic to a rank $(b - e_0 + e_1)$ vector bundle (its cokernel). Therefore, its higher Chern classes vanish on the affine bundle and so on X by (1.16). □

Taking $B = \mathcal{O}_X$ and $E_\bullet = R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2)$ (in the notation of § 4) over the product of Hilbert schemes of points $X = S^{[n_1]} \times S^{[n_2]}$, this recovers the original Carlsson–Okounkov vanishing of [CO12] and [GT20, Corollary 8.1], at least when $H^{\geq 1}(\mathcal{O}_S) = 0$ to ensure that E_\bullet is two-term. The general surface S can be handled by another splitting trick to remove $H^{\geq 1}(\mathcal{O}_S)$ from E_\bullet ; see [GT20, § 8].

3. Comparison

Given a map of two-term complexes $F_\bullet \rightarrow E_\bullet$, we can try to compare the virtual cycles of § 2. We begin with maps of a specific shape.

PROPOSITION 3.1. *Suppose that we are given a commutative diagram of vector bundles*

$$\begin{array}{ccc} F_0 & \xrightarrow{\tau} & F_1 \\ \downarrow u_0 & & \downarrow u_1 \\ E_0 & \xrightarrow{\sigma} & E_1 \end{array} \tag{3.2}$$

over a smooth quasi-projective variety X , with u_0 a vector bundle injection and u_1 a surjection. Then there is an embedding

$$\iota: \tilde{D}_r(F_\bullet) \hookrightarrow \tilde{D}_r(E_\bullet)$$

such that the tautological bundle \mathcal{U} on $\tilde{D}_r(E_\bullet)$ pulls back to that on $\tilde{D}_r(F_\bullet)$ and

$$\iota_* [\tilde{D}_r(F_\bullet)]^{\text{vir}} = c_{r, \text{rank}(E_\bullet - F_\bullet)}(\mathcal{U}^* \otimes (E_\bullet - F_\bullet)) \cap [\tilde{D}_r(E_\bullet)]^{\text{vir}}.$$

Proof. Recall that $\tilde{D}_r(E_\bullet)$ (and its perfect obstruction theory) is cut out of $\text{Gr}(r, E_0)$ by the section $\tilde{\sigma}$ of $\mathcal{U}^* \otimes E_1$ (2.7). Therefore, its virtual cycle is given by the localised top Chern class of $\mathcal{U}^* \otimes E_1$ [Ful98, § 14.1],

$$[\tilde{D}_r(E_\bullet)]^{\text{vir}} = \tilde{\sigma}^! [0_{\mathcal{U}^* \otimes E_1}] \in A_{\dim X + r(e_0 - e_1 + r)}(\tilde{D}_r(E_\bullet)),$$

where $0_{\mathcal{U}^* \otimes E_1} \cong \text{Gr}(r, E_0)$ is the zero section of $\mathcal{U}^* \otimes E_1$ and $e_i := \text{rank}(E_i)$.

Now base change by the embedding $u: \text{Gr}(r, F_0) \hookrightarrow \text{Gr}(r, E_0)$ induced by the vector bundle injection $u_0: F_0 \hookrightarrow E_0$. This is a regular embedding cut out by the canonical section of $\mathcal{U}^* \otimes (E_0/F_0)$ on $\text{Gr}(r, E_0)$.⁹ Therefore, [Ful98, Proposition 14.1(d)(ii)] gives

$$u^! [\tilde{D}_r(E_\bullet)]^{\text{vir}} = (u^* \tilde{\sigma})^! [0_{\mathcal{U}^* \otimes E_1}]. \tag{3.3}$$

Set $K_1 := \ker(u_1)$. Suppressing pullback maps, on $\text{Gr}(r, F_0)/X$ we have the exact sequence

$$0 \longrightarrow \mathcal{U}^* \otimes K_1 \longrightarrow \mathcal{U}^* \otimes F_1 \xrightarrow{\text{id} \otimes u_1} \mathcal{U}^* \otimes E_1 \longrightarrow 0$$

in which the section $\tilde{\tau}$ of the second bundle projects to $u^* \tilde{\sigma}$ in the third. Therefore, (3.3) and Lemma 3.4 below give the identity

$$c_{r(f_1 - e_1)}(\mathcal{U}^* \otimes K_1) \cap u^! [\tilde{D}_r(E_\bullet)]^{\text{vir}} = j_* \tilde{\tau}^! [0_{\mathcal{U}^* \otimes E_1}] = j_* [\tilde{D}_r(F_\bullet)]^{\text{vir}},$$

⁹ This is a special case of Proposition 2.5 applied to the two-term complex of bundles $\ker(u_0^*) \rightarrow E_0^*$ in place of E_\bullet .

where $j: \tilde{D}_r(F_\bullet) \hookrightarrow u^{-1}(\tilde{D}_r(E_\bullet))$ and $f_i := \text{rank}(F_i)$. Pushing forward by $i: u^{-1}(\tilde{D}_r(E_\bullet)) \hookrightarrow \tilde{D}_r(E_\bullet)$ and using $\iota = i \circ j$ gives

$$\iota_*[\tilde{D}_r(F_\bullet)]^{\text{vir}} = i_*u^!(c_{r(f_1-e_1)}(\mathcal{U}^* \otimes K_1) \cap [\tilde{D}_r(E_\bullet)]^{\text{vir}}).$$

We then apply [Ful98, Example 6.3.4] to the fibre square

$$\begin{array}{ccc} u^{-1}(\tilde{D}_r(E_\bullet)) & \xhookrightarrow{i} & \tilde{D}_r(E_\bullet) \\ \downarrow & & \downarrow \\ \text{Gr}(r, F_0) & \xhookrightarrow{u} & \text{Gr}(r, E_0) \end{array}$$

in which the bottom row is the regular embedding cut out by the canonical section of $\mathcal{U}^* \otimes (E_0/F_0)$. The result is

$$\begin{aligned} \iota_*[\tilde{D}_r(F_\bullet)]^{\text{vir}} &= c_{r(e_0-f_0)}(\mathcal{U}^* \otimes E_0/F_0) \cap c_{r(f_1-e_1)}(\mathcal{U}^* \otimes K_1) \cap [\tilde{D}_r(E_\bullet)]^{\text{vir}} \\ &= c_{r(e_0-f_0)+r(f_1-e_1)}(\mathcal{U}^* \otimes (E_0/F_0 \oplus K_1)) \cap [\tilde{D}_r(E_\bullet)]^{\text{vir}}. \end{aligned}$$

Substituting $K_1 = F_1 - E_1$ in K-theory gives the result. □

LEMMA 3.4. *Let $0 \rightarrow A \rightarrow B \xrightarrow{\pi} C \rightarrow 0$ be an exact sequence of vector bundles over a quasi-projective variety X . Given a section s of B , we let $t := \pi(s)$ be the induced section of C . Then the resulting localised top Chern classes of B and C are related by the identity*

$$c_a(A) \cap t^! [0_C] = \iota_* s^! [0_B],$$

where $\iota: Z(s) \hookrightarrow Z(t)$ is the embedding of zero schemes and $a := \text{rank}(A)$.

Proof. The total spaces of the bundles fit in the fibre square

$$\begin{array}{ccc} A & \xhookrightarrow{j} & B \\ \pi \downarrow & \nearrow s & \downarrow \pi \\ X & \xhookrightarrow{t} & C \end{array}$$

with both maps π being *flat*. Letting 0_C denote the zero section of C , the top Chern class of C localised by t is

$$t^! [0_C] = [0_C] \cdot [t(X)] = [0_C] \cdot \pi_*[s(X)],$$

where \cdot denotes the refined intersection product of [Ful98, Chapter 8], lying in the Chow group of the scheme-theoretic intersection $0_C \cap t(X) = Z(t)$.

So, by the projection formula [Ful98, Example 8.1.7],

$$t^! [0_C] = \pi_*(\pi^*[0_C] \cdot [s(X)]) = \pi_*(j_*[A] \cdot [s(X)]).$$

Now cap with the top Chern class of A to get its zero section:

$$\begin{aligned} c_a(A) \cap t^! [0_C] &= \pi_*(j_*[0_A] \cdot [s(X)]) \\ &= \pi_*([0_B] \cdot [s(X)]) \\ &= \pi_*(s^! [0_B]). \end{aligned}$$

But $\pi|_{0_B \cap s(X)} : 0_B \cap s(X) \rightarrow 0_C \cap t(X)$ is the inclusion $\iota : Z(s) \hookrightarrow Z(t)$. □

Proposition 3.1 now implies the following comparison results.

THEOREM 3.5. *Let X be a smooth quasi-projective variety. Suppose that*

$$F_\bullet \longrightarrow E_\bullet \longrightarrow G \tag{3.6}$$

is an exact triangle in $D(\text{Coh}(X))$ with F_\bullet, E_\bullet being two-term complexes of vector bundles supported in degrees 0, 1. If G is quasi-isomorphic to a rank g vector bundle, then there is an embedding $\iota : \tilde{D}_r(F_\bullet) \hookrightarrow \tilde{D}_r(E_\bullet)$ such that

$$\begin{aligned} \iota_* [\tilde{D}_r(F_\bullet)]^{\text{vir}} &= c_{r, \text{rank}(E_\bullet - F_\bullet)}(\mathcal{U}^* \otimes (E_\bullet - F_\bullet)) \cap [\tilde{D}_r(E_\bullet)]^{\text{vir}} \\ &= c_{rg}(\mathcal{U}^* \otimes G) \cap [\tilde{D}_r(E_\bullet)]^{\text{vir}}. \end{aligned}$$

Similarly, if $G[1]$ is a rank g vector bundle, then $\tilde{D}_r(E_\bullet) \cong \tilde{D}_r(F_\bullet)$ and

$$\begin{aligned} [\tilde{D}_r(E_\bullet)]^{\text{vir}} &= c_{r, \text{rank}(F_\bullet - E_\bullet)}(\mathcal{U}^* \otimes (F_\bullet - E_\bullet)) \cap [\tilde{D}_r(F_\bullet)]^{\text{vir}} \\ &= c_{rg}(\mathcal{U}^* \otimes G[1]) \cap [\tilde{D}_r(F_\bullet)]^{\text{vir}}. \end{aligned}$$

Proof. In the first case fix a representative vector bundle G and a locally free resolution of E_\bullet , sufficiently negative that the morphism $E_\bullet \rightarrow G$ in $D(\text{Coh}(X))$ is represented by a genuine map of complexes. Then trim E_\bullet by removing $E_{<0}$ and replacing E_0 by $E_0/\text{im}(E_{-1})$. Similarly, remove $E_{>1}$ and replace E_1 by $\ker(E_2 \rightarrow E_3)$. The upshot is a quasi-isomorphic two-term complex of bundles $\sigma : E_0 \rightarrow E_1$ with a map $f : E_0 \rightarrow G$.

Then, since $F_\bullet \cong \text{Cone}(E_\bullet \rightarrow G)[-1]$, we can realise the exact triangle (3.6) by the following short exact sequence of vertical complexes.

$$G[-1] \longrightarrow F_\bullet \longrightarrow E_\bullet \quad = \quad \begin{array}{ccc} E_0 & \xlongequal{\quad} & E_0 \\ \sigma \oplus \downarrow f & & \downarrow \sigma \\ G \xrightarrow{(0, \text{id})} E_1 \oplus G & \xrightarrow{(\text{id}, 0)} & E_1 \end{array} \tag{3.7}$$

Therefore, the map of two-term complexes $F_\bullet \rightarrow E_\bullet$ takes the form of (3.2) with $u_0 = \text{id}$ and $u_1 = (\text{id}, 0)$, and Proposition 3.1 now gives the result.

When $G[1]$ is a vector bundle H , a similar argument represents the exact triangle as follows.

$$F_\bullet \longrightarrow E_\bullet \longrightarrow G \quad = \quad \begin{array}{ccc} E_0 & \xlongequal{\quad} & E_0 \\ \tau \downarrow & & \downarrow \sigma \\ F_1 & \xrightarrow{u_1} & E_1 \twoheadrightarrow H \end{array}$$

So, now $\tilde{D}_r(F_\bullet)$ and $\tilde{D}_r(E_\bullet)$ are both cut out of $\text{Gr}(r, E_0)$ by sections $\tilde{\tau}$ and $\tilde{\sigma} = (\text{id} \otimes u_1) \circ \tilde{\tau}$ of $\mathcal{U}^* \otimes F_1$ and $\mathcal{U}^* \otimes E_1$, respectively, as in (2.7). Since the former is a subbundle of the latter, we find that $\tilde{D}_r(F_\bullet) \cong \tilde{D}_r(E_\bullet)$ and the claimed formula is the excess intersection formula¹⁰ of [Ful98, Theorem 6.3] applied to the fibre diagram

$$\begin{array}{ccc}
 \tilde{D}_r & \longrightarrow & \text{Gr}(r, E_0) \\
 \downarrow & & \downarrow \tilde{\tau} \\
 \text{Gr}(r, E_0) & \xrightarrow{0} & \mathcal{U}^* \otimes F_1 \\
 \parallel & & \text{id} \downarrow \otimes u_1 \\
 \text{Gr}(r, E_0) & \xrightarrow{0} & \mathcal{U}^* \otimes E_1
 \end{array}$$

with excess normal bundle $\mathcal{U}^* \otimes E_1 / \mathcal{U}^* \otimes F_1 \cong \mathcal{U}^* \otimes H$. □

4. Nested Hilbert schemes

A nested Hilbert scheme represents the functor which takes a base scheme B to the set of families of ideals

$$J_1 \subseteq J_2 \subseteq \dots \subseteq J_n \subseteq \mathcal{O}_{S \times B} \tag{4.1}$$

with each $\mathcal{O}_{S \times B} / J_i$ flat over B and of fixed topological class on each S fibre. By [Kol90, Proof of Lemma 6.13], the double duals $J_i^{**} \subseteq \mathcal{O}_{S \times B}$ are locally free of the form $\mathcal{O}(-D_i) \subseteq \mathcal{O}_{S \times B}$ for some Cartier divisor $D_i \subset S \times B$ flat over B . Therefore, $J_i \subseteq J_i^{**}$ can be written as $I_{Z_i}(-D_i) \subseteq \mathcal{O}(-D_i)$ for some subscheme $Z_i \subset S \times B$, flat over B of relative dimension 0. Writing $I_i := I_{Z_i}$, then (4.1) takes the form

$$I_1(-D_1) \subseteq I_2(-D_2) \subseteq \dots \subseteq I_n(-D_n) \subseteq \mathcal{O}_{S \times B}. \tag{4.2}$$

As we will see in §6, the Hilbert schemes S_β of pure divisors $\mathcal{O}(-D) \subseteq \mathcal{O}_S$ have been heavily studied for their relationship to Seiberg–Witten theory; see [DKO07] for instance. Taking the product $S_{\beta_i} \times S^{[n_i]}$ with a (smooth) Hilbert scheme of points gives the Hilbert scheme $S_{\beta_i}^{[n_i]}$ of ideals $I_i(-D_i) \subseteq \mathcal{O}_S$, where $n_i = \text{length}(Z_i)$. So, to understand *nested* Hilbert schemes what remains is to find a way to impose, one by one, the inclusions

$$I_i(-D_i) \subseteq I_{i+1}(-D_{i+1}) \tag{4.3}$$

on these products of Hilbert schemes. Taking double duals implies that $E_i := D_i - D_{i+1}$ is effective, so (4.3) is equivalent to the two inclusions

$$I_i(-E_i) \subseteq I_{i+1} \quad \text{and} \quad \mathcal{O}(-D_{i+1}) \subseteq \mathcal{O}_{S \times B}.$$

It is sufficient to study the first, since the second is a special case of it. So, we are reduced to studying the two-step nested Hilbert scheme

$$S_\beta^{[n_1, n_2]}, \quad \beta \in H^2(S, \mathbb{Z}), \quad n_1, n_2 \geq 0,$$

¹⁰ Equivalently, the obstruction sheaf of $\tilde{D}_r(E_\bullet)$ is an extension of $\mathcal{U}^* \otimes G$ by the obstruction sheaf of $\tilde{D}_r(F_\bullet)$, from which the formula also follows.

which represents the functor mapping B to the set of families of ideals

$$I_1(-D) \subseteq I_2 \subseteq \mathcal{O}_{S \times B}, \tag{4.4}$$

with D, I_1, I_2 flat over B and such that on closed fibres S_b we have $[D_b] = \beta$ and colength $(I_i|_{S_b}) = n_i$. The map to double duals recovers Z_1, Z_2 and D , giving a classifying map to a smooth space

$$S_\beta^{[n_1, n_2]} \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S) =: X. \tag{4.5}$$

Since (4.4) is the same data as a one-dimensional subspace of $\text{Hom}(I_1(-D), I_2)$, we see that, at the level of points, $S_\beta^{[n_1, n_2]}$ (4.5) is the virtual resolution \tilde{D}_1 of the $r = 1$ degeneracy locus of the complex over X which restricts to $R\text{Hom}(I_1, I_2 \otimes L)$ at $(I_1, I_2, L) \in X$. To make a scheme-theoretic statement, we use

$$\pi : S \times S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S) \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$$

with its universal ideal sheaves $\mathcal{I}_1, \mathcal{I}_2$ and a fixed choice of Poincaré line bundle¹¹ \mathcal{L}_β . Using base change, the Nakayama lemma and the fact that $\text{Ext}^i(I_1, I_2 \otimes L) = 0$ for $i \notin [0, 2]$ on any S fibre of π , we can trim

$$R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta) \cong \{E_0 \xrightarrow{\sigma} E_1 \longrightarrow E_2\} =: E_\bullet \tag{4.6}$$

to be a three-term complex of locally free sheaves. Using its stupid truncation $E_0 \rightarrow E_1$ gives the following result, which will be modified later to give versions invariant under quasi-isomorphisms.

PROPOSITION 4.7. *The nested Hilbert scheme (4.5) is the virtual resolution of the $r = 1$ degeneracy locus of $\sigma : E_0 \rightarrow E_1$ (4.6), giving isomorphisms*

$$S_\beta^{[n_1, n_2]} \cong \tilde{D}_1(\sigma) \cong \mathbb{P}^*(\mathcal{E}xt_\pi^2(\mathcal{I}_2, \mathcal{I}_1 \otimes K_S \otimes \mathcal{L}_\beta^*))$$

under which the tautological bundle $\mathcal{O}(-1)$ on $\tilde{D}_1(\sigma)$ corresponds to the dual of the tautological quotient bundle on \mathbb{P}^* .

Proof. By the $r = 1$ case of Proposition 2.5, $\tilde{D}_1(\sigma)$ represents the functor which maps an X -scheme $f : B \rightarrow X$ to the set of line subbundles $\mathcal{L} \hookrightarrow f^*E_0$ over B factoring through

$$\mathcal{L} \hookrightarrow h^0(f^*E_\bullet) = \pi_{B*} \mathcal{H}om(f^*\mathcal{I}_1, f^*(\mathcal{I}_2 \otimes \mathcal{L}_\beta)),$$

where $\pi_B : S \times B \rightarrow B$ is the base change of π . By adjunction, this is equivalent to the set of line bundles $\mathcal{L} \rightarrow B$ and maps

$$f^*\mathcal{I}_1 \hookrightarrow f^*(\mathcal{I}_2 \otimes \mathcal{L}_\beta) \otimes \pi_B^* \mathcal{L}^{-1} \quad \text{over } S \times B, \tag{4.8}$$

which are *nonzero on restriction to any fibre S_b* . We need to show this is the same as the functor (4.4) represented by $S_\beta^{[n_1, n_2]}$, i.e. that (4.8) is equivalent to the set of families

$$f^*\mathcal{I}_1(-D) \hookrightarrow f^*\mathcal{I}_2 \quad \text{over } S \times B \tag{4.9}$$

with $\mathcal{I}_1, \mathcal{I}_2$ and D flat over B of the correct topological type on each fibre.

¹¹ We always normalise \mathcal{L}_β (by tensoring it by the pullback of $\mathcal{L}_\beta^{-1}|_{\{x\} \times \text{Pic}_\beta(S)}$ if necessary) so that $\mathcal{L}_\beta|_{\{x\} \times \text{Pic}_\beta(S)}$ is trivial on some fixed basepoint $x \in S$.

Firstly, taking double duals in (4.8) gives a map $\mathcal{O} \rightarrow \mathcal{O}(D)$ for some divisor D which does not contain any fibres and so is flat over B . This gives (4.9), as required. Conversely, given a family (4.9), note that $\mathcal{O}_{S \times B}(D)$ is isomorphic to $f^* \mathcal{L}_\beta$ on each S fibre, so may be written globally as $f^* \mathcal{L}_\beta \otimes \pi_B^* \mathcal{L}^{-1}$, where \mathcal{L} is the line bundle $\pi_{B*}(f^* \mathcal{L}_\beta(D))$. This gives (4.8), with the map nonzero on every fibre (since D is flat over B and so does not contain any fibre).

The second claimed isomorphism follows from the definition (2.3) of $\tilde{D}_1(\sigma)$: it is the space of one-dimensional quotients of the cokernel of $\sigma^*: E_1^* \rightarrow E_0^*$,

$$\tilde{D}_1(\sigma) = \mathbb{P}^*(\text{coker } \sigma^*) = \mathbb{P}^*(h^0(E_\bullet^\vee)) = \mathbb{P}^*(\mathcal{E}xt_\pi^2(\mathcal{I}_2, \mathcal{I}_1 \otimes K_S \otimes \mathcal{L}_\beta^*))$$

by relative Serre duality down π . □

4.1 Embedding and reduced virtual cycle

Throughout this section we require β to be of Hodge type $(1, 1)$. Until now everything has been true (but vacuous) when this assumption fails, and the results deformation invariant. The constructions of this section are only invariant under deformations of S inside the Noether–Lefschetz locus in which β is of type $(1, 1)$.

As discussed in the Introduction, each fibre of $\tilde{D}_1(\sigma) \rightarrow D_1(\sigma)$ admits a natural embedding

$$\mathbb{P}(\text{Hom}(I_1, I_2 \otimes L)) \subseteq \mathbb{P}(H^0(L)) \xrightarrow{s_A} \mathbb{P}(H^0(L(A))). \tag{4.10}$$

Here we have fixed, once and for all, $s_A \in H^0(\mathcal{O}_S(A))$ cutting out a divisor $A \subset S$ sufficiently positive that

$$H^{\geq 1}(L(A)) = 0 \quad \forall L \in \text{Pic}_\beta(S).$$

(When $H^{\geq 1}(\mathcal{O}_S) = 0$ and L is effective, we may take $A = \emptyset$ and the proof of Theorem 3 below reduces to a proof of Theorem 2 instead.) So, we let B be the vector bundle

$$B := \pi_*(\mathcal{L}_\beta(A)) \quad \text{over } X = S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S). \tag{4.11}$$

Then (4.10) gives the embedding $h^0(E_\bullet) \subset B_x$ of (1.4) at each point $x = (I_1, I_2, L) \in X$. The global version (2.19) is the composition of surjections

$$\begin{aligned} B^* \cong R^2 \pi_*(\mathcal{L}_\beta^*(K_S - A)) &\xrightarrow{s_A} R^2 \pi_*(\mathcal{L}_\beta^* \otimes K_S) \cong R^2 \pi_*(\mathcal{I}_1 \otimes \mathcal{L}_\beta^* \otimes K_S) \\ &\longrightarrow \mathcal{E}xt_\pi^2(\mathcal{I}_2, \mathcal{I}_1 \otimes \mathcal{L}_\beta^* \otimes K_S). \end{aligned} \tag{4.12}$$

By (2.20), we get an embedding

$$\iota_B: S_\beta^{[n_1, n_2]} \cong \tilde{D}_1(E_\bullet) \hookrightarrow \mathbb{P}(B) \xrightarrow{q} X. \tag{4.13}$$

Suppose now that, as in the statements of Theorems 2 and 3,

$$H^2(\mathcal{O}(D)) = 0 \quad \text{for any effective divisor } D \text{ in class } \beta. \tag{4.14}$$

The resulting surjection

$$0 = H^2(L) \cong \text{Ext}^2(\mathcal{O}, I_2(L)) \longrightarrow \text{Ext}^2(I_1, I_2(L)) \cong h^2(E_\bullet|_x)$$

and base change show that E_\bullet has $h^{\geq 2} = 0$ over the image of $S_\beta^{[n_1, n_2]}$ in X and therefore also over a neighbourhood U thereof. Therefore, in (4.6), we can take E_\bullet to be a two-term complex of vector bundles

$$R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta) \cong \{E_0 \xrightarrow{\sigma} E_1\} =: E_\bullet \quad \text{over } U \subseteq X. \tag{4.15}$$

Proposition 4.7 then gives

$$S_\beta^{[n_1, n_2]} \cong \tilde{D}_1(E_\bullet)$$

and Proposition 2.12 endows it with a perfect obstruction theory, which we call the *reduced* obstruction theory.¹²

THEOREM 4.16. *Suppose that $H^2(\mathcal{O}(D)) = 0$ for every effective divisor D in class β . The above construction gives a reduced obstruction theory with virtual tangent bundle*

$$\begin{aligned} T_{S_\beta^{[n_1, n_2]}}^{\text{vir}} &= -R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_1)_0 - R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_2)_0 \\ &\quad + R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1)) + R^1\pi_* \mathcal{O} - \mathcal{O} \end{aligned} \tag{4.17}$$

in K -theory and reduced virtual cycle

$$[S_\beta^{[n_1, n_2]}]^{\text{red}} \in A_{n_1+n_2+\text{vd}_\beta+p_g}(S_\beta^{[n_1, n_2]}), \tag{4.18}$$

where $\mathcal{O}(1) := \mathcal{O}_{\mathbb{P}(B)}(1)$ and $\text{vd}_\beta := \beta(\beta - K_S)/2$.

If $H^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$, then the pushforward of (4.18) to $\mathbb{P}(B)$ is

$$\iota_{B*} [S_\beta^{[n_1, n_2]}]^{\text{red}} = c_{n_1+n_2+d}(B(1) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))),$$

where $d = \frac{1}{2}A(2\beta + A - K_S) = \chi(L(A)) - \chi(L)$ for any $L \in \text{Pic}_\beta(S)$.

Proof. The virtual tangent bundle is described by the $r = 1$ case of Proposition 2.12 with $\mathcal{U} = \mathcal{O}(-1)$, giving the restriction of

$$T_X - \mathcal{O}_X - \mathcal{O}(-1)^* \otimes R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)$$

to $S_\beta^{[n_1, n_2]}$. Since $X = S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$, we derive (4.17) from

$$T_{\text{Pic}_\beta(S)} \cong R^1\pi_* \mathcal{O} \quad \text{and} \quad T_{S^{[n_i]}} \cong \mathcal{E}xt_\pi^1(\mathcal{I}_i, \mathcal{I}_i)_0 \cong R\mathcal{H}om_\pi(\mathcal{I}_i, \mathcal{I}_i)_0[1].$$

When $H^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$, then E_\bullet is two-term over the whole of X , so we may apply Theorem 2.24 with $r = 1$ to give the formula for the pushforward. \square

Remark. This proves Theorems 2 and 3 from the Introduction. Since the virtual class depends only on the K -theory class of the virtual tangent bundle, Theorem 4.16 shows that $[S_\beta^{[n_1, n_2]}]^{\text{red}}$ coincides with the reduced virtual cycles of [KT14a] (when $n_1 = 0$) and [GSY20a, Proposition 2.10] (when the latter is defined).

¹² This is because (4.17) shows that at the level of virtual tangent bundles it is obtained from the Vafa–Witten perfect obstruction theory by removing a copy of $H^2(\mathcal{O}_S)$.

4.2 Comparison of reduced cycles

Next we will compare the reduced cycles on $S_\beta^{[n_1, n_2]}$ and S_β . Note that $S_\beta^{[n_1, n_2]}$ and $S_\beta \times S^{[n_1]} \times S^{[n_2]}$ are the virtual resolutions of the degeneracy loci of

$$R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta) \quad \text{and} \quad R\pi_* \mathcal{L}_\beta \quad \text{over } X,$$

respectively. These are related by the obvious diagram

$$\begin{array}{ccc}
 & & R\pi_* \mathcal{L}_\beta \\
 & \nearrow u & \downarrow \dashrightarrow \\
 R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta) & \longrightarrow & R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{L}_\beta) \\
 & & \downarrow \dashrightarrow \\
 & & H[-1]
 \end{array} \tag{4.19}$$

with the vertical column an exact triangle. Here $H := \mathcal{E}xt_\pi^2(\mathcal{O}/\mathcal{I}_1, \mathcal{L}_\beta)$, so

$$H^* \cong (\mathcal{L}_\beta^* \otimes K_S)^{[n_1]}$$

by relative Serre duality down π . Thus, H is a vector bundle.

So, we now use the Jouanolou trick of § 1.14, pulling back to an affine bundle over U , so that H becomes a projective \mathcal{O} -module. In particular, its connecting homomorphism to $R\pi_* \mathcal{L}_\beta[2]$ in the vertical exact triangle (4.19) is zero over the open neighbourhood U (4.15) over which it is supported in degrees -2 and -1 . (Recall our assumption (4.14) that $H^2(\mathcal{O}_S(D)) = 0$ for every effective D of class β .) So, over U , we may choose the splittings marked with dotted arrows in (4.19).

LEMMA 4.20. *Cone(u) is quasi-isomorphic to a rank $(n_1 + n_2)$ vector bundle.*

Proof. Taking H^0 of (4.19) at a closed point $x = (I_1, I_2, L)$ of U gives the following diagram.

$$\begin{array}{ccc}
 & & H^0(L) \\
 & \nearrow h^0(u) & \parallel \\
 \text{Hom}(I_1, I_2 \otimes L) & \hookrightarrow & \text{Hom}(I_1, L)
 \end{array}$$

Since this diagram commutes, the diagonal arrow is an injection. Thus, by the Nakayama lemma, $\text{Cone}(u)$, and any base change of it, has $h^{<0} = 0$.

Taking H^1 instead gives part of the following diagram.

$$\begin{array}{ccccc}
 H^1(I_2 \otimes L) & \longrightarrow & H^1(L) & \longrightarrow & 0 \\
 \downarrow & \nearrow h^1(u) & \downarrow \dashrightarrow & & \downarrow \\
 \text{Ext}^1(I_1, I_2 \otimes L) & \longrightarrow & \text{Ext}^1(I_1, L) & \xrightarrow{a} & \text{Ext}^1(I_1, L/(L \otimes I_2))
 \end{array}$$

Again, this commutes and the rows are exact. Since $\text{Ext}^1(I_1, I_2 \otimes L)$ surjects onto $\ker(a)$, in which $H^1(L)$ lies, the diagonal arrow is surjective. Thus, $\text{Cone}(u)$ has $h^{\geq 1} = 0$ and is a vector bundle, as claimed. \square

So, it follows immediately from Theorem 3.5 and the Jouanolou trick that

$$[S_\beta^{[n_1, n_2]}]^{\text{red}} = c_{n_1+n_2}(\text{Cone}(u)(1)) \cap [S_\beta]^{\text{red}} \times [S^{[n_1]} \times S^{[n_2]}].$$

On $S \times S_\beta$, the tautological composition $\mathcal{O}(-1) \rightarrow \pi^* \pi_* \mathcal{L}_\beta \rightarrow \mathcal{L}_\beta$ gives the universal section

$$\mathcal{O} \xrightarrow{s_\beta} \mathcal{L}_\beta(1) \quad \text{cutting out the universal divisor } \mathcal{D}_\beta \subset S \times S_\beta. \tag{4.21}$$

In particular, $\mathcal{L}_\beta(1) \cong \mathcal{O}(\mathcal{D}_\beta)$ on $S \times S_\beta$. Thus, $\text{Cone}(u)(1)$ represents the Carlsson–Okounkov K-theory class of (1.14):

$$\text{Cone}(u)(1) = \text{CO}_\beta^{[n_1, n_2]} = R\pi_* \mathcal{O}(\mathcal{D}_\beta) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2(\mathcal{D}_\beta)).$$

Hence, we have proved the second formula in Theorem 5:

COROLLARY 4.22. *If $H^2(\mathcal{O}_S(D)) = 0$ for all effective divisors in class β , then*

$$[S_\beta^{[n_1, n_2]}]^{\text{red}} = c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) \cap [S^{[n_1]} \times S^{[n_2]}] \times [S_\beta]^{\text{red}}.$$

4.3 Comparison of virtual cycles

To recover the full perfect obstruction theory of [GSY20a] or Vafa–Witten theory on $S_\beta^{[n_1, n_2]}$ from (the virtual resolution of) a degeneracy locus, we would have to modify the three-term complexes $R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)$ and $R\pi_* \mathcal{L}_\beta$ by $H^2(\mathcal{O}_S)$ terms to make them two-term. This cannot be done over $X = S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$.¹³

However, up on (the product of S with) $S^{[n_1]} \times S^{[n_2]} \times S_\beta$, we can use the universal section s_β (4.21) of $\mathcal{L}_\beta(1)$, notice the twist!, together with a choice of splitting $R\pi_* \mathcal{O} \cong \bigoplus_{i=0}^2 R^i \pi_* \mathcal{O}[-i]$, to induce maps

$$\begin{array}{ccc} & R^2 \pi_* \mathcal{O}[-2] & \\ & \swarrow \quad \downarrow s_\beta \quad \searrow & \\ R\pi_*(\mathcal{I}_2 \otimes \mathcal{L}_\beta(1)) & \longrightarrow & R\pi_* \mathcal{L}_\beta(1) \longrightarrow \pi_*((\mathcal{O}/\mathcal{I}_2) \otimes \mathcal{L}_\beta(1)) \quad \text{on } S_\beta^{[n_1, n_2]}. \end{array}$$

The lower row is an exact triangle. We apply the Jouanolou trick of § 1.14, pulling back from $S^{[n_1]} \times S^{[n_2]} \times S_\beta$ to an affine bundle over it. Here $R^2 \pi_* \mathcal{O}$ becomes a projective \mathcal{O} -module, so the right-hand diagonal arrow becomes zero. Thus, we can fill in the dotted arrow, giving the following commutative diagram.

$$\begin{array}{ccc} & R^2 \pi_* \mathcal{O}[-2] & H[-2] \\ & \swarrow \quad \searrow & \downarrow \\ R\pi_*(\mathcal{I}_2 \otimes \mathcal{L}_\beta(1)) & \longrightarrow & R\pi_* \mathcal{L}_\beta(1) \\ \downarrow & & \downarrow \\ R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1)) & \longrightarrow & R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{L}_\beta(1)) \end{array} \tag{4.23}$$

¹³ A referee suggested the following illustrative example. Let β be the fibre class in an elliptic K3 surface S and take $n_1 = 0 = n_2$. Then $S_\beta = \mathbb{P}^1$ and its Vafa–Witten virtual tangent bundle is easily worked out to be $T_{\mathbb{P}^1} - \mathcal{O}_{\mathbb{P}^1}$ (using (4.35), for example). To see \mathbb{P}^1 as \tilde{D}_1 of a complex over $S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S) = \text{point}$, we must take $\mathbb{C}^2 \xrightarrow{0} \mathbb{C}^q$, but this has virtual tangent bundle $T_{\mathbb{P}^1} - \mathcal{O}_{\mathbb{P}^1}^{\oplus q}(1)$.

Here the right-hand column is an exact triangle, with $H := \mathcal{E}xt_{\pi}^2(\mathcal{O}/\mathcal{I}_1, \mathcal{L}_{\beta}(1))$ a vector bundle Serre dual to

$$H^* \cong (K_S \otimes \mathcal{L}_{\beta}^{-1}(-1))^{[n_1]}.$$

On second cohomology sheaves h^2 we claim that the following diagram gives the surjections.

$$\begin{array}{ccc}
 & R^2\pi_* \mathcal{O} & \\
 a \swarrow & & \searrow b \\
 R^2\pi_*(\mathcal{I}_2 \otimes \mathcal{L}_{\beta}(1)) & \xlongequal{c} & R^2\pi_* \mathcal{L}_{\beta}(1) \\
 d \downarrow & & \downarrow e \\
 \mathcal{E}xt_{\pi}^2(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_{\beta}(1)) & \xrightarrow{f} & \mathcal{E}xt_{\pi}^2(\mathcal{I}_1, \mathcal{L}_{\beta}(1))
 \end{array} \tag{4.24}$$

The claim for the maps b, c, d, f follows from zeros in the obvious exact sequences in which each sits. It then follows from the diagram that a and e are also onto. So, if we use the notation

$$\tau(\mathcal{F}) := \text{Cone}(R^2\pi_* \mathcal{O}[-2] \rightarrow \mathcal{F})$$

for any of the maps $a, b, d \circ a$ or $e \circ b$ in (4.23), then each $\tau(\mathcal{F})$ has $h^{\geq 2} = 0$ and can be represented by a two-term complex of vector bundles. The bottom row and right-hand column of (4.23) now give the diagram

$$\begin{array}{ccc}
 & & \tau(R\pi_* \mathcal{L}_{\beta}(1)) \\
 & \xrightarrow{u} & \downarrow \dashrightarrow \\
 \tau(R\mathcal{H}om_{\pi}(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_{\beta}(1))) & \longrightarrow & \tau(R\mathcal{H}om_{\pi}(\mathcal{I}_1, \mathcal{L}_{\beta}(1))) \\
 & & \downarrow \dashrightarrow \\
 & & H[-1]
 \end{array} \tag{4.25}$$

which should be compared to (4.19). Again, the column is an exact triangle, and again H is a projective \mathcal{O} -module on the affine bundle, so has vanishing connecting homomorphism to the complex $\tau(R\pi_* \mathcal{L}_{\beta}(1))[2]$ supported in degrees -2 and -1 . Thus, we may split the vertical exact triangle and fill in the dotted arrows on $S^{[n_1]} \times S^{[n_2]} \times S_{\beta}$.

4.4 Extension to ambient space

Let $B = \pi_*(\mathcal{L}(A))$ be the bundle (4.11). Applying (4.13) when $n_1 = 0 = n_2$ gives an embedding $S_{\beta} \subseteq \mathbb{P}(B)/\text{Pic}_{\beta}(S)$ and so

$$S^{[n_1]} \times S^{[n_2]} \times S_{\beta} \subseteq \mathbb{P}(B)/X,$$

where $X = S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_{\beta}(S)$, as in (4.5).

We apply the Jouanolou trick to $\mathbb{P}(B)$, giving an affine bundle over it whose total space is an affine variety. The restriction of this affine bundle to $S^{[n_1]} \times S^{[n_2]} \times S_{\beta}$ is therefore also an affine variety, to which the constructions of the last § 4.3 apply. As usual we suppress pullbacks to these affine bundles.

All of the terms in (4.23) extend to (the affine bundle over) $\mathbb{P}(B)$, as do the horizontal and vertical arrows (canonically). As for the diagonal arrows, their targets are quasi-isomorphic to complexes of vector bundles $G_0 \rightarrow G_1 \rightarrow G_2$, and $R^2\pi_* \mathcal{O}$ is a projective \mathcal{O} -module on an

affine variety, so the arrows can be represented by maps $R^2\pi_* \mathcal{O} \rightarrow G_2$. Again, since $R^2\pi_* \mathcal{O}$ is a projective \mathcal{O} -module, these maps can be extended to (the affine bundle over) $\mathbb{P}(B) \supset S^{[n_1]} \times S^{[n_2]} \times S_\beta$. So, we get an extension of the diagram (4.25) to (the affine bundle over) $\mathbb{P}(B)$.

Since the surjectivity (4.24) is an open condition, all the terms of this extended diagram can be represented by two-term complexes of vector bundles over a neighbourhood U of (the affine bundle over) $S^{[n_1]} \times S^{[n_2]} \times S_\beta$ inside (the affine bundle over) $\mathbb{P}(B)$. By shrinking U if necessary, we may assume that it is affine.¹⁴

So, finally, the splitting map $H \rightarrow \tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{L}_\beta(1)))$ of the vertical exact sequence in (4.25) extends from (the affine bundle over) $S^{[n_1]} \times S^{[n_2]} \times S_\beta$ to its affine neighbourhood U since $H|_U$ is a projective \mathcal{O} -module (and hence has vanishing connecting homomorphism to the complex $\tau(R\pi_* \mathcal{L}_\beta(1))[2]$ supported in degrees -2 and -1). So, we can fill in the dotted arrows, giving

$$\begin{array}{ccc} & & \tau(R\pi_* \mathcal{L}_\beta(1)) \\ & \nearrow u & \downarrow \uparrow \\ \tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))) & \longrightarrow & \tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{L}_\beta(1))) \end{array} \quad \text{over } U. \tag{4.26}$$

Lemma 4.20 now applies verbatim to the map u of (4.25) to show that

$$\text{Cone}(u) \text{ is a vector bundle of rank } n_1 + n_2 \text{ over } U. \tag{4.27}$$

Next¹⁵ we describe a nice form of these complexes $\tau(\mathcal{F})$ on the open set U .

LEMMA 4.28. *Over U each of the terms $\tau(\mathcal{F})$ in (4.26) has a two-term resolution by vector bundles*

$$\tau(\mathcal{F})(-1) \simeq \{B \longrightarrow F\}, \tag{4.29}$$

which, on dualising, induces the surjection $B^* \twoheadrightarrow h^0((\tau(\mathcal{F})(-1))^\vee)$ of (4.12). Moreover, the arrows in (4.26) may be taken to be genuine maps of complexes between these resolutions.

Proof. The final claim follows from the fact that on any affine variety like U , maps in the derived category between complexes of locally free sheaves are quasi-isomorphic to genuine maps of complexes.

We start by working downstairs on X . The composition

$$R\pi_* \mathcal{L}_\beta \xrightarrow{s_A} R\pi_* \mathcal{L}_\beta(A) = \pi_* \mathcal{L}_\beta(A) = B \tag{4.30}$$

induces the inclusion $H^0(L) \hookrightarrow H^0(L(A))$ on restricting to any point $x = (I_1, I_2, L) \in X$ and taking h^0 . Therefore, its cone has cohomology in degrees $1, 2$ after base change to any point, so it can be represented by a two-term complex of vector bundles $G_1 \rightarrow G_2$, meaning that we can

¹⁴ Let $U = P \setminus Z$ be an open neighbourhood of a closed subvariety S of an affine variety P . Then Z is disjoint from S , so we get a surjection $\mathcal{O}[P] \twoheadrightarrow \mathcal{O}[S \sqcup Z] \cong \mathcal{O}[S] \oplus \mathcal{O}[Z]$. Let f map to $(1, 0)$. Then $P \setminus f^{-1}(0) \subseteq U$ is an affine neighbourhood of S .

¹⁵ Hence, the K-theory class $\text{CO}_\beta^{[n_1, n_2]}$ (1.14) is represented by a bundle after pullback to an affine bundle. By (1.16), this gives a more conceptual proof of the vanishing $c_{n_1+n_2+i}(\text{CO}_\beta^{[n_1, n_2]}) = 0 \forall i > 0$ of [GT20, Corollary 8.11]. As Aravind Asok kindly explained to us, it is possible that any bundle on the affine bundle is automatically a pullback from the base; this would prove that $\text{CO}_\beta^{[n_1, n_2]}$ is represented by a bundle on $S^{[n_1]} \times S^{[n_2]} \times S_\beta$.

write

$$R\pi_* \mathcal{L}_\beta \cong \{B \longrightarrow G_1 \longrightarrow G_2\}.$$

By construction, this resolution dualises to induce the surjection $B^* \rightarrow h^0((R\pi_* \mathcal{L}_\beta)^\vee)$ of (4.12).

Now pull back to (the affine bundle over) $\mathbb{P}(B)/X$, twist by $\mathcal{O}_{\mathbb{P}(B)}(1)$ and use the map $R^2\pi_* \mathcal{O}[-2] \rightarrow R\pi_* \mathcal{L}_\beta(1)$ of (4.23) (which as we already noted extends to the affine bundle over $\mathbb{P}(B)$). It can be represented by a genuine map of complexes, so we can represent its cone as

$$\tau(R\pi_* \mathcal{L}_\beta(1)) \cong \{B(1) \rightarrow G(1)\}, \quad G := \ker [G_1 \oplus (R^2\pi_* \mathcal{O})(-1) \rightarrow G_2]$$

over the affine open U on which the second arrow is a surjection. This gives the result for $\mathcal{F} = R\pi_* \mathcal{L}_\beta(1)$.

For $\mathcal{F} = \tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1)))$, we can now write the exact triangle defined by u (4.26) as

$$\tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))) \xrightarrow{u} \{B(1) \rightarrow G(1)\} \longrightarrow \text{Cone}(u)$$

with the last arrow a genuine map of complexes of vector bundles on U . Thus, the first term can be written as the two-term complex of vector bundles

$$\tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))) \cong \{B(1) \longrightarrow \text{Cone}(u) \oplus G(1)\}.$$

A similar argument applies to $\tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{L}_\beta(1)))$. □

So, we can now see (the affine bundle over) $S_\beta^{[n_1, n_2]}$ as the zeros in U of the section of $F(1)$ defined by the composition

$$\mathcal{O}_{\mathbb{P}(B)}(-1) \hookrightarrow B \longrightarrow F, \tag{4.31}$$

where $\{B \rightarrow F\}$ is the representative (4.29) of $\tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta))$. For the confused reader, we summarise the construction.

4.5 Summary

By Proposition 4.7, $S_\beta^{[n_1, n_2]}$ is the virtual resolution \tilde{D}_1 of the $r = 1$ degeneracy locus of the complex $R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)$ over X . Since this is three-term, $\{B \rightarrow G_1 \rightarrow G_2\}$, in general this does not give it a natural virtual cycle, but it does express it as the zero locus in $q: \mathbb{P}(B) \rightarrow X$ of the composition¹⁶

$$\mathcal{O}_{\mathbb{P}(B)}(-1) \hookrightarrow q^*B \longrightarrow q^*G_1. \tag{4.32}$$

We cannot modify $R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)$ by an $R^2\pi_* \mathcal{O}$ term on X to get a two-term degeneracy locus construction directly. But, we can after pulling back to $\mathbb{P}(B)$ and twisting by $\mathcal{O}(1)$, replacing (4.32) by

$$\mathcal{O}_{\mathbb{P}(B)}(-1) \hookrightarrow q^*B \longrightarrow \ker [q^*G_1 \oplus (R^2\pi_* \mathcal{O})(-1) \rightarrow q^*G_2] \tag{4.33}$$

on the open set U , where the last arrow is a surjection. This is (4.31).

THEOREM 4.34. $S_\beta^{[n_1, n_2]}$ is the zero locus of (4.31). The virtual cycle of the resulting perfect obstruction theory agrees with the nonreduced virtual cycles of [GSY20a, TT20] whenever they

¹⁶ Changing G_1 , for instance to $\ker(G_1 \rightarrow G_2)$ when this is locally free, gives the same zero locus but a different perfect obstruction theory.

are defined.¹⁷ It satisfies

$$\iota_* [S_\beta^{[n_1, n_2]}]^{\text{vir}} = c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) \cap [S^{[n_1]} \times S^{[n_2]}] \times [S_\beta]^{\text{vir}}$$

under pushforward by $\iota: S_\beta^{[n_1, n_2]} \hookrightarrow S^{[n_1]} \times S^{[n_2]} \times S_\beta$.

Proof. Since $\tau(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta))$ differs from $R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)$ only in $h^{\geq 1}$ (even after any base change), h^0 is unaffected. In other words, the zero loci of (4.31) and (4.32) are the same and the latter is $S_\beta^{[n_1, n_2]}$ by Proposition 4.7.

The virtual tangent bundle differs from the one in Theorem 4.16 by only $R^2\pi_* \mathcal{O}[-1]$, so its K-theory class is

$$-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_1)_0 - R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_2)_0 + R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2(\mathcal{D}_\beta)) - R\pi_* \mathcal{O}, \tag{4.35}$$

agreeing with the ones from [GSY20a, TT20]. Since virtual classes depend only on the K-theory class of the virtual tangent bundle, this makes the virtual classes of [GSY20a, TT20] equal to the localised top Chern class

$$[S_\beta^{[n_1, n_2]}]^{\text{vir}} = s^! [0_{F(1)}] \in A_{n_1+n_2+\text{vd}(S_\beta)}(S_\beta^{[n_1, n_2]}) \tag{4.36}$$

of the bundle $F(1) \rightarrow U$ and its section s defined by (4.31).

Setting $n_1 = 0 = n_2$, we get

$$[S_\beta]^{\text{vir}} = t^! [0_{G(1)}] \in A_{\text{vd}(S_\beta)}(S_\beta),$$

where $B \rightarrow G$ is the resolution of $\tau(R\pi_* \mathcal{L}_\beta)$ on the open subset of (the affine bundle over) $\mathbb{P}(B \rightarrow \text{Pic}_\beta(S))$, which is the image of U (in the affine bundle over $\mathbb{P}(B \rightarrow X)$), and t is the induced section of $G(1)$. Pulling back to U gives

$$[S_\beta]^{\text{vir}} \times S^{[n_1]} \times S^{[n_2]} = t^! [0_{G(1)}], \tag{4.37}$$

where as usual we have suppressed the pullback map on $G(1)$ and its section t . Now the map u of (4.26) induces, via Lemma 4.28, a map

$$\tilde{u}: F(1) \longrightarrow G(1)$$

such that $\tilde{u} \circ s = t$. By (4.27), \tilde{u} is a surjection with kernel the vector bundle $\text{Cone}(u)$, so applying Lemma 3.4 and using (4.37) gives

$$\begin{aligned} [S_\beta^{[n_1, n_2]}]^{\text{vir}} &= s^! [0_{F(1)}] \\ &= c_{n_1+n_2}(\text{Cone}(u)) \cap t^! [0_{G(1)}] \\ &= c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) \cap [S^{[n_1]} \times S^{[n_2]}] \times [S_\beta]^{\text{vir}}. \end{aligned} \quad \square$$

Remark. We have now proved Theorem 5 from the Introduction. When $p_g(S) > 0$ and $H^2(\mathcal{O}_S(D)) = 0$ for all effective D in class β , so that the reduced class is defined, the virtual

¹⁷ The Vafa–Witten moduli space of [TT20] depends on a stability condition. But, even in the unstable case, where we get a stack with possibly nontrivial stabilisers, we can pass to the \mathbb{C}^* -fixed locus and cut down the automorphisms to those which commute with the \mathbb{C}^* action. The nested Hilbert scheme then appears as a connected component, with the \mathbb{C}^* -fixed part of the Vafa–Witten obstruction theory [TT20] defining a perfect obstruction theory on it which recovers those of [GSY20a] and Theorem 4.34.

cycle is automatically zero by the identity

$$[S_{\beta}^{[n_1, n_2]}]^{vir} = c_{p_g}(\pi_* \mathcal{O}) \cap [S_{\beta}^{[n_1, n_2]}]^{red} = 0 \tag{4.38}$$

of Theorem 3.5.

5. ℓ -step nested Hilbert schemes

Fix curve classes $\beta_1, \dots, \beta_{\ell-1} \in H^2(S, \mathbb{Z})$ and integers n_1, \dots, n_{ℓ} . As a set, the ℓ -step nested Hilbert scheme is

$$S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_{\ell}]} := \{I_i(-D_i) \subseteq I_{i+1} \subseteq \mathcal{O}_S, i = 1, \dots, \ell - 1: \\ [D_i] = \beta_i, \text{ length}(\mathcal{O}_S/I_i) = n_i\}.$$

As a scheme, it represents the functor described in (4.2) (with $D_{\ell} = 0$). That description immediately implies that it can be written as the fibre product

$$S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_{\ell}]} = S_{\beta_1}^{[n_1, n_2]} \times_{S^{[n_2]}} S_{\beta_2}^{[n_2, n_3]} \times_{S^{[n_3]}} \cdots \times_{S^{[n_{\ell-1}]}} S_{\beta_{\ell-1}}^{[n_{\ell-1}, n_{\ell}]}.$$

It follows that $S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_{\ell}]}$ is the intersection of the $\ell - 1$ virtual resolutions $\tilde{D}_1(E^i)$ of the degeneracy loci $D_1(E^i)$ of the complexes

$$E^i_{\bullet} := R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_{i+1} \otimes \mathcal{L}_{\beta_i}) \quad \text{over } X := \prod_{j=1}^{\ell} S^{[n_j]} \times \prod_{j=1}^{\ell-1} \text{Pic}_{\beta_j}(S). \tag{5.1}$$

Here $i = 1, \dots, \ell - 1$ while $\pi: S \times X \rightarrow X$ is the projection and $\mathcal{L}_{\beta_i}, \mathcal{I}_i$ are (the pullbacks of) Poincaré line bundles and the universal ideal sheaves, respectively.

Taking $A \subset S$ to be a divisor which is sufficiently positive that

$$H^{\geq 1}(L_i(A)) = 0 \quad \forall L_i \in \text{Pic}_{\beta_i}(S), \quad i = 1, \dots, \ell - 1,$$

we get vector bundles

$$B_i := \pi_*(\mathcal{L}_{\beta_i}(A)), \quad i = 1, \dots, \ell - 1, \quad \text{over } X$$

and an embedding

$$\iota_B: S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_{\ell}]} \hookrightarrow \mathbb{P}(B_1) \times_X \cdots \times_X \mathbb{P}(B_{\ell-1}). \tag{5.2}$$

Suppose that $H^2(\mathcal{O}(D_i)) = 0$ for every effective D_i of class β_i , $i = 1, \dots, \ell - 1$. Then each of the complexes (5.1) can be represented by a two-term complex of vector bundles $E_0^i \rightarrow E_1^i$. So, we can mimic the constructions of §§ 2 and 4. We replace the projective bundles $\mathbb{P}(E_0)/X$ and $\mathbb{P}(B)/X$ by the fibre products $\mathbb{P}(E_0^1) \times_X \cdots \times_X \mathbb{P}(E_0^{\ell-1})$ and $\mathbb{P}(B_1) \times_X \cdots \times_X \mathbb{P}(B_{\ell-1})$, and cut out the ℓ -step nested Hilbert scheme by the obvious section of $V_1 \boxplus \cdots \boxplus V_{\ell-1}$, where $V_i = \mathcal{O}_{\mathbb{P}(E_0^i)}(-1)^* \otimes p_i^*(E_1^i)$. The upshot is a (multiply reduced) perfect obstruction theory on

$S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}$ with virtual tangent bundle

$$-\sum_{i=1}^{\ell} R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_i)_0 + \sum_{i=1}^{\ell-1} R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_{i+1}(\mathcal{D}_{\beta_i})) + (R^1\pi_*\mathcal{O})^{\oplus(\ell-1)} - \mathcal{O}^{\oplus(\ell-1)}$$

in K-theory, and reduced virtual class

$$[S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}]^{\text{red}} \in A_{n_1+n_\ell+\text{vd}_{\beta_1}+\dots+\text{vd}_{\beta_{\ell-1}}+(\ell-1)p_g(S)}(S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}). \tag{5.3}$$

THEOREM 5.4. *If $H^2(L_i) = 0$ for all $L_i \in \text{Pic}_{\beta_i}(S)$ and all $i = 1, \dots, \ell - 1$, then the pushforward of the reduced cycle (5.3) by the embedding (5.2) is*

$$\iota_{B*} [S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}]^{\text{red}} = \prod_{i=1}^{\ell-1} c_{b_i+n_i+n_{i+1}-\chi(\mathcal{O}_S)-\text{vd}_{\beta_i}}(B_i(1_i) - R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_{i+1}(\mathcal{L}_{\beta_i}(1_i))),$$

where $b_i := \text{rank}(B_i)$ and $\mathcal{O}(1_i)$ denotes the pullback of $\mathcal{O}_{\mathbb{P}(B_i)}(1)$.

For general classes β_i we follow the procedure of § 4.3 to modify each of the complexes (5.1) by $R^2\pi_*\mathcal{O}[-2]$ after pulling back to (the fibre product of the) $\mathbb{P}(B_i)$ and tensoring by $\mathcal{O}(1_i)$. The result is a perfect obstruction theory with virtual tangent bundle

$$-\sum_{i=1}^{\ell} R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_i)_0 + \sum_{i=1}^{\ell-1} R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_{i+1}(\mathcal{D}_{\beta_i})) + (R\pi_*\mathcal{O})^{\oplus(\ell-1)} \tag{5.5}$$

in K-theory, and virtual cycle

$$[S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}]^{\text{vir}} \in A_{n_1+n_\ell+\text{vd}_{\beta_1}+\dots+\text{vd}_{\beta_{\ell-1}}}(S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}).$$

Just as in (4.38), this vanishes if $p_g(S) > 0$ and $H^2(L) = 0$ for all $L \in \text{Pic}_{\beta_i}(S)$ for some $i = 1, \dots, \ell - 1$.

Iterating the construction of § 4.3, we get the following comparison result, a direct generalisation of Theorem 5.

THEOREM 5.6. *The pushforward of $[S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}]^{\text{vir}}$ by the embedding*

$$S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]} \hookrightarrow \prod_{j=1}^{\ell} S^{[n_j]} \times \prod_{j=1}^{\ell-1} S_{\beta_j}$$

is given by the formula

$$\prod_{i=1}^{\ell-1} c_{n_i+n_{i+1}}(R\pi_*\mathcal{O}(\mathcal{D}_{\beta_i}) - R\mathcal{H}om_{\pi}(\mathcal{I}_i, \mathcal{I}_{i+1}(\mathcal{D}_{\beta_i}))) \cap [S^{[n_1]} \times \dots \times S^{[n_\ell]}] \times [S_{\beta_1}]^{\text{vir}} \times \dots \times [S_{\beta_{\ell-1}}]^{\text{vir}}.$$

Moreover, if $H^2(L_i) = 0$ for all effective $L_i \in \text{Pic}_{\beta_i}(S)$, $i = 1, \dots, \ell - 1$, then the same formula holds with all virtual classes $[\cdot]^{\text{vir}}$ replaced by reduced classes $[\cdot]^{\text{red}}$.

Remark. Since (5.5) matches the virtual tangent bundle in [GSY20a], the virtual class $[S_{\beta_1, \dots, \beta_{\ell-1}}^{[n_1, \dots, n_\ell]}]^{\text{vir}}$ is the same class that arises in higher rank Vafa–Witten theory [TT20] or the reduced local DT theory of S [GSY20b].

6. Computations

6.1 Vafa–Witten invariants

On a polarised surface $(S, \mathcal{O}_S(1))$, Vafa–Witten theory is an enumerative sheaf-counting theory [TT20]. The ‘instanton contributions’ compute virtual Euler numbers of moduli spaces of Gieseker stable sheaves of fixed rank r and determinant L on S , as studied by Göttsche–Kool [GK20].

This leaves the ‘monopole contributions’, which are integrals over moduli spaces of chains of sheaves (of total rank r) with nonzero maps between them. When the individual sheaves have rank 1, we get nested ideal sheaves (all tensored by a line bundle). And, when r is prime and $p_g(S) > 0$, a vanishing theorem [Tho20, § 5.2] implies that all nonzero contributions come from nested Hilbert schemes. A similar result applies in the semistable case [TT17] and the refined version of both theories [Tho20].

The simplest case (for simplicity) is the monopole contribution to the Vafa–Witten invariants in rank 2 and fixed determinant when slope stability is the same as slope semistability. Here we get integrals over the virtual cycles of nested Hilbert schemes of pairs $I_1(-D) \subseteq I_2 \subseteq \mathcal{O}_S$ for any $[D] = \beta$ satisfying the slope stability¹⁸ condition $\text{deg}(K_S(-D)) > 0$. We then sum over all β, n_1, n_2 giving the same total Chern classes and then multiply by a factor¹⁹ of $2^{2h^1(\mathcal{O}_S)}$.

The integrand is $1/e(N^{\text{vir}})$, where N^{vir} is the moving part of the Vafa–Witten obstruction theory and e is the \mathbb{C}^* -equivariant Euler class. So, by [TT20, § 8], we want to compute the integral of

$$\frac{e(R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1(-\mathcal{D}_\beta)K_S \mathfrak{t})) \ e(R\mathcal{H}om_\pi(\mathcal{I}_1(-\mathcal{D}_\beta), \mathcal{I}_2K_S^{-1}\mathfrak{t}^{-1}))}{e(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_1K_S \mathfrak{t})_0) \ e(R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_2K_S \mathfrak{t})) \ e(R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1(-\mathcal{D}_\beta)K_S^2 \mathfrak{t}^2))}$$

over $[S_\beta^{[n_1, n_2]}]^{\text{vir}}$. Write $\mathcal{O}(\mathcal{D}_\beta)$ as $\mathcal{L}_\beta(1)$ as in (4.21). Let $h := c_1(\mathcal{O}(1))$ on $\mathbb{P}(B) = \mathbb{P}(\pi_* \mathcal{L}_\beta(A))$ and its subschemes $S_\beta^{[n_1, n_2]}$. Then each term $e(\cdot)$ takes the form

$$e((E_0 - E_1)(1)) = \frac{c_{r_0}(E_0(1))}{c_{r_1}(E_1(1))} = \frac{\sum_{i=0}^{r_0} c_{r_0-i}(E_0) \cup h^i}{\sum_{i=0}^{r_1} c_{r_1-i}(E_1) \cup h^i},$$

where the E_i have only nonzero weight spaces (so the quotient makes sense in localised equivariant cohomology) and ranks r_i . Therefore, expanding in powers of h^i with coefficients pulled back

¹⁸ Gieseker stability is slightly more subtle, with a further inequality on the $n_i = \text{length}(\mathcal{O}/I_i)$ in the strictly slope semistable case. Laarakker [Laa18, Proposition 3.5] has proved that Gieseker unstable Higgs pairs have $\iota_*[S_\beta^{[n_1, n_2]}]^{\text{vir}} = 0$, so we can sum over all β , independent of stability.

¹⁹ Thanks are due to Ties Laarakker for pointing out the $r^{2h^1(\mathcal{O}_S)}$ factor in total rank r . This counts the r -torsion line bundles we can twist by to get another Higgs pair with the same fixed determinant.

from $S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$, we get an expression

$$\sum_i \int_{[S_\beta^{[n_1, n_2]}]^{vir}} \rho^* \alpha_i \cup h^i = \sum_i \int_{\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir})} \alpha_i,$$

where ρ is part of the following commutative diagram.

$$\begin{CD} S_\beta^{[n_1, n_2]} @>\iota>> S^{[n_1]} \times S^{[n_2]} \times S_\beta \\ @V\rho VV @. @VV\text{id} \times \text{id} \times \text{AJ} V \\ @. @. S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S) \end{CD} \tag{6.1}$$

Here $\text{AJ}: S_\beta \rightarrow \text{Pic}_\beta(S)$ is the Abel–Jacobi map. So, we should calculate the pushdown $\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir})$. This will prove Theorems 6–8 from the Introduction, and compute Vafa–Witten invariants in terms of integrals over the smooth spaces $X = S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$.

Let

$$\text{SW}_\beta := \begin{cases} \text{deg } [S_\beta]^{vir}, & \text{vd}_\beta = \frac{1}{2}\beta(K_S - \beta) = 0, \\ 0, & \text{otherwise} \end{cases}$$

denote the Seiberg–Witten invariant [DKO07, CK13] of S in class $\beta \in H^2(S, \mathbb{Z})$. Things are easiest when $b^+(S) > 1$, i.e. $p_g(S) > 0$.

THEOREM 6.2. *Suppose that $p_g(S) > 0$. Then, in $H_*(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$,*

$$\begin{aligned} \rho_* [S_\beta^{[n_1, n_2]}]^{vir} &= \text{SW}_\beta \cdot c_{n_1+n_2}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)) \times [L], \\ \rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{vir}) &= 0 \quad \text{for } i > 0, \end{aligned}$$

where $L \in \text{Pic}_\beta(S)$, so $[L]$ is the generator of $H_0(\text{Pic}_\beta(S), \mathbb{Z})$.

Proof. If $\text{AJ}_*[S_\beta]^{vir} = 0$, then $\text{SW}_\beta = 0$, so by our comparison result of Theorem 4.34 (or Theorem 5 in the Introduction) both sides of the claimed identity are zero. So, we assume that $\text{AJ}_*[S_\beta]^{vir} \neq 0$.

By [DKO07, Definition 3.19], this means that β is a *basic class*. And, by [DKO07, Proposition 3.20] S is of *simple type*, which implies that $\text{vd}_\beta = 0$.

Thus, $[S_\beta]^{vir}$ is a zero-dimensional class. Firstly, this implies that $h^i|_{[S_\beta]^{vir}} = 0$ for $i > 0$, and so the second claimed result. And, secondly, on restriction to

$$S \times S^{[n_1]} \times S^{[n_2]} \times [S_\beta]^{vir},$$

the line bundle $\mathcal{L}_\beta(1) = \mathcal{O}(\mathcal{D}_\beta)$ is topologically equivalent to (the pullback of) L for any $L \in \text{Pic}_\beta(S)$. In particular, in the formula of Theorem 4.34,

$$\iota_* [S_\beta^{[n_1, n_2]}]^{vir} = c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) \cap [S^{[n_1]} \times S^{[n_2]}] \times [S_\beta]^{vir} \tag{6.3}$$

we may replace $\text{CO}_\beta^{[n_1, n_2]}$ (1.14) by $R\pi_* L - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)$. Applying $(\text{id} \times \text{id} \times \text{AJ})_*$ then gives

$$\rho_* [S_\beta^{[n_1, n_2]}]^{vir} = c_{n_1+n_2}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)) \times \text{deg } [S_\beta]^{vir} [\text{point}],$$

where $[\text{point}] = [L]$ is the generator of $H_0(\text{Pic}_\beta(S), \mathbb{Z})$. □

Now we suppose that $b^+(S) = 1$, i.e. $p_g(S) = 0$. Then S may not be of simple type, so we have to consider higher dimensional Seiberg–Witten moduli spaces and the higher Seiberg–Witten invariants [DKO07]

$$SW_\beta^j := \text{AJ}_*(h^j \cap [S_\beta]^{\text{vir}}) \in H_{2(\text{vd}_\beta - j)}(\text{Pic}_\beta(S)) = \wedge^{2(\text{vd}_\beta - j)} H^1(S).$$

THEOREM 6.4. *If $p_g(S) = 0$, then, in $H_{2(n_1+n_2+\text{vd}_\beta - i)}(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$,*

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) = \sum_{j=0}^{n_1+n_2} c_{n_1+n_2-j}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cup SW_\beta^{i+j}.$$

Proof. We again use our comparison formula (6.3). By [Man16, Proposition 1],²⁰

$$\begin{aligned} c_{n_1+n_2}(\text{CO}_\beta^{[n_1, n_2]}) &= c_{n_1+n_2}(R\pi_* \mathcal{L}_\beta(1) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))) \\ &= \sum_{j=0}^{n_1+n_2} c_{n_1+n_2-j}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cup h^j. \end{aligned}$$

So, $(\text{id} \times \text{id} \times \text{AJ})_*(h^i \cap (6.3)) = \rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}})$ is

$$\sum_{j=0}^{n_1+n_2} c_{n_1+n_2-j}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cup \text{AJ}_*(h^{i+j} \cap [S_\beta]^{\text{vir}}). \quad \square$$

We can get simpler formulae by splitting into the cases that $H^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$ or not. By Serre duality, this is the condition that $\beta^\vee := K_S - \beta$ is not effective or effective, respectively.

THEOREM 6.5. *Suppose that $p_g(S) = 0$. Fix $L \in \text{Pic}_\beta(S)$ and $i \geq 0$. Then the following results hold in $H_{2(n_1+n_2-i)}(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$.*

- If β^\vee is effective,

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) = SW_\beta \cdot c_{n_1+n_2+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)) \times [L]$$

and both sides vanish when $i > 0$ (cf. Theorem 6.2).

- If β^\vee is not effective,

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) = c_{d+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta))$$

with $d = n_1 + n_2 + h^1(\mathcal{O}_S) - \text{vd}_\beta$.

Proof. By the comparison result (6.3), both sides of the first identity vanish if $\text{vd}_\beta < 0$ or β is not effective. So, we may assume that both β, β^\vee are effective and $\text{vd}_\beta \geq 0$. By [DKO07, Corollary 3.15], this implies that $\text{vd}_\beta = 0$ and $h^1(\mathcal{O}_S) = 1$.

²⁰ Manivel extended the formula $c_r(E(1)) = \sum_{j=0}^r c_{r-j}(E) \cap c_1(\mathcal{O}(1))^j$ from rank r bundles to rank r perfect complexes. In fact, we showed in (4.27) that $\text{CO}_\beta^{[n_1, n_2]}$ has the K-theory class of a vector bundle on (an affine bundle over) $S^{[n_1]} \times S^{[n_2]} \times S_\beta$ anyway.

So, just as in (6.3), $\mathcal{O}(1)$ is trivial on the zero-dimensional $[S_\beta]^{\text{vir}}$ and we may replace $\text{CO}_\beta^{[n_1, n_2]}$ in (6.3) by $R\pi_* L - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)$ for one $L \in \text{Pic}_\beta(S)$. So, applying $(\text{id} \times \text{id} \times \text{AJ})_*$ to (6.3) gives

$$\rho_* [S_\beta^{[n_1, n_2]}]^{\text{vir}} = c_{n_1+n_2}(R\pi_* L - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes L)) \times \text{deg} [S_\beta]^{\text{vir}} \times [\text{point}].$$

This deals with $i = 0$. When $i > 0$, the left-hand side of the required identity vanishes by $\text{vd}_\beta = 0$ and (6.3), while the right-hand side vanishes by the generalised Carlsson–Okounkov vanishing of [GT20, Theorem 3].

For the second result, we may assume that β is of type $(1, 1)$, otherwise $\text{Pic}_\beta(S)$ is empty and both sides of the identity vanish. Since we also have $H^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$, the reduced cycle of § 4.1 is defined and, since $p_g(S) = 0$, it equals the virtual cycle. Therefore, by Theorem 4.16 (or Theorem 3 of the Introduction), the pushforward of the virtual cycle to $\mathbb{P}(B)$ is given by

$$\begin{aligned} & c_{n_1+n_2+k}(B(1) - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta(1))) \\ &= \sum_{j=0}^{n_1+n_2+k} c_{n_1+n_2+k-j}(B - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cup c_1(\mathcal{O}(1))^j, \end{aligned}$$

again by [Man16, Proposition 1]. Here $k = \text{rank}(B) - \text{vd}_\beta - \chi(\mathcal{O}_S)$.

We now cup with $h^i = c_1(\mathcal{O}(1))^i$ and push down $q: \mathbb{P}(B) \rightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$ (which restricts to the map ρ of (6.1)). Using the fact that $q_* h^{i+j}$ is the Segre class $s_{i+j-\chi(L(A))+1}(B)$ for $i + j \geq \chi(L(A)) - 1$, and zero otherwise, we find that $\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}})$ is

$$\begin{aligned} & \sum_{j=\chi(L(A))-1-i}^{n_1+n_2+k} c_{n_1+n_2+k-j}(B - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cup s_{i+j-\chi(L(A))+1}(B) \\ &= c_{n_1+n_2+i+k-\text{rank}(B)+1}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)). \quad \square \end{aligned}$$

Finally, we use the standard duality in Seiberg–Witten theory under $\beta \leftrightarrow \beta^\vee$ to give interesting dualities between invariants of nested Hilbert schemes under $\beta \leftrightarrow \beta^\vee$ and $n_1 \leftrightarrow n_2$. Define the map

$$\rho^\vee : S_{\beta^\vee}^{[n_2, n_1]} \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$$

by replacing $\beta \leftrightarrow \beta^\vee$, $n_1 \leftrightarrow n_2$ in $\rho: S_\beta^{[n_1, n_2]} \longrightarrow S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S)$ and then composing with $L \mapsto K_S \otimes L^{-1}: \text{Pic}_{\beta^\vee}(S) \rightarrow \text{Pic}_\beta(S)$.

THEOREM 6.6 (Duality). *In $H_{2(n_1+n_2+\text{vd}_\beta-i)}(S^{[n_1]} \times S^{[n_2]} \times \text{Pic}_\beta(S))$:*

- if $p_g(S) > 0$,

$$\rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) = (-1)^{s+i} \rho_*^\vee(h^i \cap [S_{\beta^\vee}^{[n_2, n_1]}]^{\text{vir}});$$

- if $p_g(S) = 0$,

$$\begin{aligned} \rho_*(h^i \cap [S_\beta^{[n_1, n_2]}]^{\text{vir}}) &= (-1)^{s+i} \rho_*^\vee(h^i \cap [S_{\beta^\vee}^{[n_2, n_1]}]^{\text{vir}}) \\ &\quad + c_{d+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)), \end{aligned}$$

where $s = n_1 + n_2 - \chi(\mathcal{O}_S) - \text{vd}_\beta = d - 1$.

Proof. By Theorem 6.2 with $\beta \leftrightarrow \beta^\vee$, $n_1 \leftrightarrow n_2$ and $L \leftrightarrow K_S \otimes L^{-1}$,

$$\rho_*^\vee [S_\beta^{[n_2, n_1]}]^\text{vir} = \text{SW}_{\beta^\vee} \cdot c_{n_1+n_2}(-R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1 \cdot K_S \cdot L^{-1})) \times [\text{point}]$$

for a fixed $L \in \text{Pic}_\beta(S)$. For $p_g(S) > 0$, Seiberg–Witten invariants have the standard duality

$$\text{SW}_\beta = (-1)^{\chi(\mathcal{O}_S)} \text{SW}_{\beta^\vee}.$$

(For instance, in [DKO07], this is part of Conjecture 0.1, and it is then shown that this would follow from proving that $\text{deg}[S_{K_S}]^\text{vir} = (-1)^{\chi(\mathcal{O}_S)}$ on all minimal general type surfaces. This latter identity was proved in [CK13].) And,

$$c_{n_1+n_2}(-R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1 \cdot K_S \cdot L^{-1})) = (-1)^{n_1+n_2} c_{n_1+n_2}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \cdot L))$$

by Serre duality down the fibres of π . This gives the first identity for $i = 0$. For $i > 0$, it is trivial by Theorem 6.2.

To prove the second identity, we distinguish three cases. If β^\vee is not effective, i.e. $H^2(L) = 0$ for all $L \in \text{Pic}_\beta(S)$, then the result is the last part of Theorem 6.5. Similarly, if β is not effective, then we apply the last part of Theorem 6.5 with $\beta \leftrightarrow \beta^\vee$, $n_1 \leftrightarrow n_2$. By $K_S \otimes \mathcal{L}_{\beta^\vee}^{-1} = \mathcal{L}_\beta$, $\text{vd}_\beta = \text{vd}_{\beta^\vee}$ and Serre duality down the fibres of π ,

$$c_{d+i}(-R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1 \otimes \mathcal{L}_{\beta^\vee})) = (-1)^{d+i} c_{d+i}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)),$$

which gives the required result.

Finally, we consider β and β^\vee both effective. If $\text{vd}_\beta < 0$, then the left-hand side of the identity is zero by the comparison result of Theorem 5. On the right-hand side, we have

$$\begin{aligned} & c_{n_1+n_2+h^1(\mathcal{O}_S)-\text{vd}_\beta}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \\ &= \sum_{i \geq 0} c_i(-R\pi_* \mathcal{L}_\beta) \cdot c_{n_1+n_2+h^1(\mathcal{O}_S)-\text{vd}_\beta-i}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)). \end{aligned}$$

The first term of the sum vanishes for $i > \dim \text{Pic}_\beta(S) = h^1(\mathcal{O}_S)$, while the second term vanishes for $i < h^1(\mathcal{O}_S) - \text{vd}_\beta$ by the generalised Carlsson–Okounkov vanishing of [GT20, Theorem 3]. When $\text{vd}_\beta < 0$, this makes the whole sum vanish.

So, we may assume that $\text{vd}_\beta \geq 0$ and β, β^\vee are both effective. This implies that $\text{vd}_\beta = 0$ and $h^1(\mathcal{O}_S) = 1$ by [DKO07, Corollary 3.15].

So, just as in (6.3), $\mathcal{O}(1)$ is trivial on the zero-dimensional $[S_\beta]^\text{vir}$ and we may replace $\text{CO}_\beta^{[n_1, n_2]}$ in (6.3) by the pullback of $R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)$. Thus, pushing (6.3) down by $(\text{id} \times \text{id} \times \text{AJ}_\beta)_*$ gives $\rho_*[S_\beta^{[n_1, n_2]}]^\text{vir}$ as

$$c_{n_1+n_2}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cap [S^{[n_1]} \times S^{[n_2]}] \times \text{AJ}_{\beta*} [S_\beta]^\text{vir}. \tag{6.7}$$

Swapping $\beta \leftrightarrow \beta^\vee$ and $n_1 \leftrightarrow n_2$ shows that $\rho_*^\vee [S_{\beta^\vee}^{[n_2, n_1]}]^\text{vir}$ is

$$c_{n_1+n_2}(R\pi_* \mathcal{L}_{\beta^\vee} - R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1 \otimes \mathcal{L}_{\beta^\vee})) \cap [S^{[n_1]} \times S^{[n_2]}] \times \text{AJ}_{\beta^\vee*} [S_{\beta^\vee}]^\text{vir}.$$

By Serre duality down π and the identity $c_i(E) = (-1)^i c_i(E^\vee)$, this is

$$(-1)^{n_1+n_2} c_{n_1+n_2}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cap [S^{[n_1]} \times S^{[n_2]}] \times \text{AJ}_{\beta^\vee*} [S_{\beta^\vee}]^\text{vir}. \tag{6.8}$$

We have noted that $h^1(\mathcal{O}_S) = 1$, so $\chi(\mathcal{O}_S) = 0$. Thus, identifying $\text{Pic}_\beta(S) \cong \text{Pic}_{\beta^\vee}(S)$ via $L \mapsto K_S \otimes L^{-1}$, [DKO07, Theorem 3.16] gives

$$\text{AJ}_{\beta^*} [S_\beta]^{\text{vir}} = \text{AJ}_{\beta^\vee} [S_{\beta^\vee}]^{\text{vir}} + c_1(-R\pi_* \mathcal{L}_\beta).$$

Since $\text{Pic}_\beta(S)$ is one dimensional, $c_{\geq 2}(-R\pi_* \mathcal{L}_\beta) = 0$. And, $c_{\geq n_1+n_2+1}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) = 0$ by the generalised Carlsson–Okounkov vanishing of [GT20, Theorem 3]. Therefore,

$$\begin{aligned} c_1(-R\pi_* \mathcal{L}_\beta) c_{n_1+n_2}(R\pi_* \mathcal{L}_\beta - R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \\ = c_{n_1+n_2+1}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)). \end{aligned}$$

Substituting this into (6.8) and comparing with (6.7) yields

$$\begin{aligned} \rho_*^\vee [S_{\beta^\vee}^{[n_2, n_1]}]^{\text{vir}} &= (-1)^{n_1+n_2} \rho_* [S_\beta^{[n_1, n_2]}]^{\text{vir}} \\ &\quad - (-1)^{n_1+n_2} c_{n_1+n_2+1}(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 \otimes \mathcal{L}_\beta)) \cap [S^{[n_1]} \times S^{[n_2]}]. \end{aligned}$$

Multiplying by $(-1)^{n_1+n_2}$ and noting that $d = n_1 + n_2 + 1$ then gives the required result for $i = 0$.

For $i > 0$, the restrictions of h^i to both $[S_\beta]^{\text{vir}}$ and $[S_\beta^{[n_1, n_2]}]^{\text{vir}}$ are trivial (the former by $\text{vd}_\beta = 0$, the latter by the comparison result Theorem 5). \square

THEOREM 6.9. *If $p_g(S) > 0$, the nested Hilbert scheme contributions to the Vafa–Witten invariants are invariants of the oriented diffeomorphism type of S .*

Proof. We sketch the argument for two-step nested Hilbert schemes; the general case is no more complicated. Handling stability as in Footnote 18, we express the invariants as a sum over β with $\text{deg } \beta < \text{deg } K_S$ of terms

$$\text{VW}_{2,\beta,n} = \sum_{n=n_1+n_2} \int_{[S_\beta^{[n_1, n_2]}]^{\text{vir}}} \frac{1}{e(N^{\text{vir}})}.$$

Let L be any line bundle with $c_1(L) = \beta$. Applying Theorem 6.2 gives

$$\text{VW}_{2,\beta,n} = \text{SW}_\beta \sum_{n=n_1+n_2} \int_{S^{[n_1]} \times S^{[n_2]}} A(\mathcal{I}_1, \mathcal{I}_2, L), \tag{6.10}$$

where the integrand $A(\mathcal{I}_1, \mathcal{I}_2, L)$ is

$$\frac{c_n(-R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 L)) e(R\mathcal{H}om_\pi(\mathcal{I}_2 L, \mathcal{I}_1 K_S \mathfrak{t})) e(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_2 L K_S^{-1} \mathfrak{t}^{-1}))}{e(R\mathcal{H}om_\pi(\mathcal{I}_1, \mathcal{I}_1 K_S \mathfrak{t}_0)) e(R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_2 K_S \mathfrak{t})) e(R\mathcal{H}om_\pi(\mathcal{I}_2, \mathcal{I}_1 L^{-1} K_S^2 \mathfrak{t}^2))}.$$

The coefficient SW_β is proved in [DKO07, CK13] to be a Seiberg–Witten invariant, which is an oriented diffeomorphism invariant when $p_g(S) > 0$.²¹

We would now like to apply the inductive method of [EGL01], which deals with integrals over single Hilbert schemes $S^{[n_1]}$. This is adapted in [GNY08, §5]²² to deal with products of Hilbert schemes $S^{[n_1]} \times S^{[n_2]}$, essentially by applying [EGL01] to the disjoint union $T := S \sqcup S$

²¹ When $p_g(S) = 0$, this need not be quite true; only the unordered pair $(\text{SW}_\beta, \text{SW}_{\beta^\vee})$ is invariant under oriented diffeomorphisms.

²² We thank a referee for suggesting this method and pointing out the reference [GNY08].

and its Hilbert schemes

$$T^{[n]} = \bigsqcup_{n=n_1+n_2} S^{[n_1]} \times S^{[n_2]}.$$

Let \mathcal{I}_1 and \mathcal{I}_2 denote the ideal sheaves on $S \times T^{[n]}$ whose restrictions to $S \times S^{[n_1]} \times S^{[n_2]}$ are \mathcal{I}_1 and \mathcal{I}_2 , respectively. Then (6.10) can be rewritten

$$\text{VW}_{2,\beta,n} = \text{SW}_\beta \int_{T^{[n]}} A(\mathcal{I}_1, \mathcal{I}_2, L).$$

The induction writes this as an integral over $T^{[n-1]} \times S$, then $T^{[n-2]} \times S^2$ and so on, finally giving an integral over S^n . The first step pulls back along the generically finite map ψ , divides by $\deg \psi$ and then pushes down σ , in the following diagram.

$$\begin{array}{ccc} T^{[n-1,n]} & \xrightarrow{\psi} & T^{[n]} \\ \sigma \downarrow & & \\ T^{[n-1]} \times S & & \end{array} \tag{6.11}$$

We split $T^{[n-1,n]}$ into two (unions of) connected components

$$\left(\bigsqcup_{n_1+n_2=n, n_1 \geq 1} S^{[n_1-1,n_1]} \times S^{[n_2]} \right) \sqcup \left(\bigsqcup_{n_1+n_2=n, n_2 \geq 1} S^{[n_1]} \times S^{[n_2-1,n_2]} \right)$$

and restrict the diagram to each in turn; we describe the first. On each of its components σ is the projection down the cone $\mathbb{P}^*(I_Z) \rightarrow S^{[n_1-1]} \times S$, where $Z = Z_{n_1-1} \subset S^{[n_1-1]} \times S$ is the universal subscheme, multiplied by the identity on $S^{[n_2]}$. As such it carries a line bundle $\mathcal{O}(1)$; it is the kernel of $\mathcal{O}^{[n_1]} \twoheadrightarrow \mathcal{O}^{[n_1-1]}$ in the obvious notation.

By [GNY08, (5.4)] and [EGL01, (10)], the integrand $A(\mathcal{I}_1, \mathcal{I}_2, L)$ pulls back to a rational function in \mathfrak{t} and the Chern classes of:

- (1) $R\mathcal{H}om_\pi(\mathcal{I}_i, \mathcal{I}_j \cdot \xi)$ pulled back from $T^{[n-1]}$, where ξ is a tensor product of powers of L and K_S ;
- (2) $\mathcal{I}_1, \mathcal{I}_2$ pulled back from $T^{[n-1]} \times S$;
- (3) L and T_S pulled back from S ; and
- (4) the line bundle $\mathcal{O}(1)$.

Taking the coefficient of \mathfrak{t}^0 gives a polynomial in these terms. Since (1)–(3) are pulled back from $T^{[n-1]} \times S$, we can push down by σ_* using the projection formula and [EGL01, Lemma 1.1] to replace powers of $c_1(\mathcal{O}(1))$ by polynomials in the Chern classes of \mathcal{I}_1 and \mathcal{I}_2 . This completes the first step.

The second (and later) steps are similar, with (6.11) replaced by the following diagram.

$$\begin{array}{ccc} T^{[n-2,n-1]} \times S & \xrightarrow{\psi \times \text{id}_S} & T^{[n-1]} \times S \\ \sigma \times \text{id}_S \downarrow & & \\ T^{[n-2]} \times S \times S & & \end{array}$$

The pullback and pushdown handle classes pulled back from either factor of $T^{[n-1]} \times S$, just as before. The new issue is to deal with Chern classes of \mathcal{J}_1 and \mathcal{J}_2 on $T^{[n-1]} \times S$. Here we use [GN Y08, (5.2)] to express them as pullbacks from $T^{[n-2]} \times S \times S$. So, the induction continues, just as in [EGL01, Proposition 3.1].

The final result is an integral over S^n , which can be written as a polynomial in the numbers $c_1(S)^2$, $c_2(S)$, β^2 and $c_1(S) \cdot \beta$. See also [GN Y08, Lemma 5.5] and [Laa18, Proposition 7.2], or [KT14b, § 3] when $n_1 = 0$.

Since $e(S)$, $p_1(S)$ and the intersection form are oriented diffeomorphism invariants of S , so are $c_1(S)^2$, $c_2(S)$, β^2 . This leaves $c_1(S) \cdot \beta$, but, as an observant referee pointed out, we may assume that this equals $-\beta^2$; otherwise $\text{vd}_\beta \neq 0$, so $\text{SW}_\beta = 0$ and the integral (6.10) vanishes. \square

ACKNOWLEDGEMENTS

We thank Martijn Kool and Ties Laarakker for help and useful conversations, Bhargav Bhatt for pointing out the Jouanolou trick and two thorough referees for useful suggestions. A.G. acknowledges partial support from NSF grant DMS-1406788. R.P.T. is partially supported by EPSRC grant EP/R013349/1.

REFERENCES

- BF97 K. Behrend and B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128** (1997), 45–88.
- CO12 E. Carlsson and A. Okounkov, *Exts and vertex operators*, Duke Math. J. **161** (2012), 1797–1815.
- CK13 H. L. Chang and Y. H. Kiem, *Poincaré invariants are Seiberg–Witten invariants*, Geom. Topol. **17** (2013), 1149–1163.
- DKO07 M. Dürr, M. A. Kabanov and C. Okonek, *Poincaré invariants*, Topology **46** (2007), 225–294.
- Eis94 D. Eisenbud, *Commutative algebra, with a view toward algebraic geometry*, Graduate Texts in Mathematics, vol. **150** (Springer, 1994).
- EGL01 G. Ellingsrud, L. Göttsche and M. Lehn, *On the cobordism class of the Hilbert scheme of a surface*, J. Algebraic Geom. **10** (2001), 81–100.
- Ful98 W. Fulton, *Intersection theory* (Springer, 1998).
- GSY20a A. Gholampour, A. Sheshmani and S.-T. Yau, *Nested Hilbert schemes on surfaces: virtual fundamental class*, Adv. Math. **365** (2020), doi:10.1016/j.aim.2020.107046.
- GSY20b A. Gholampour, A. Sheshmani and S.-T. Yau, *Localized Donaldson–Thomas theory of surfaces*, Amer. J. Math. **142** (2020), 405–442.
- GT20 A. Gholampour and R. P. Thomas, *Degeneracy loci, virtual cycles and nested Hilbert schemes I*, Tunis. J. Math. **2** (2020), 633–665.
- GK20 L. Göttsche and M. Kool, *Virtual refinements of the Vafa–Witten formula*, Comm. Math. Phys. **376** (2020), 1–49.
- GN Y08 L. Göttsche, H. Nakajima and K. Yoshioka, *Instanton counting and Donaldson invariants*, J. Differential Geom. **80** (2008), 343–390.
- Jou73 J. P. Jouanolou, *Une suite exacte de Mayer–Vietoris en K-théorie algébrique*, in *Higher K-theories* (Springer, 1973), 293–316.
- Kol90 J. Kollár, *Projectivity of complete moduli*, J. Differential Geom. **32** (1990), 235–268.
- Koo16 M. Kool, *Stable pair invariants of surfaces and Seiberg–Witten invariants*, Q. J. Math. **67** (2016), 365–386.
- KT14a M. Kool and R. P. Thomas, *Reduced classes and curve counting on surfaces I: theory*, Algebr. Geom. **1** (2014), 334–383.

- KT14b M. Kool and R. P. Thomas, *Reduced classes and curve counting on surfaces II: calculations*, *Algebr. Geom.* **1** (2014), 384–399.
- Kre99 A. Kresch, *Cycle groups for Artin stacks*, *Invent. Math.* **138** (1999), 495–536.
- Laa18 T. Laarakker, *Monopole contributions to refined Vafa–Witten invariants*, *Geom. Topol.*, to appear. Preprint (2018), [arXiv:1810.00385](https://arxiv.org/abs/1810.00385).
- Laa20 T. Laarakker, *Vertical Vafa–Witten invariants*, Preprint (2020), [arXiv:1906.01264](https://arxiv.org/abs/1906.01264).
- Man16 L. Manivel, *Chern classes of tensor products*, *Internat. J. Math.* **27** (2016), 1650079.
- Sie04 B. Siebert, *Virtual fundamental classes, global normal cones and Fulton’s canonical classes*, in *Frobenius manifolds* (Vieweg, 2004), 341–358.
- TT17 Y. Tanaka and R. P. Thomas, *Vafa–Witten invariants for projective surfaces II: semistable case*, *Pure Appl. Math. Q.* **13** (2017), 517–562, volume in honour of the 60th birthday of Simon Donaldson.
- TT20 Y. Tanaka and R. P. Thomas, *Vafa–Witten invariants for projective surfaces I: stable case*, *J. Algebr. Geom.* **29** (2020), 603–668.
- Tho20 R. P. Thomas, *Equivariant K-theory and refined Vafa–Witten invariants*, *Comm. Math. Phys.* **378** (2020), 1451–1500.

Amin Gholampour amigh@math.umd.edu

Department of Mathematics, University of Maryland, College Park, MD 20742, USA

Richard P. Thomas richard.thomas@imperial.ac.uk

Department of Mathematics, Imperial College London, London SW7 2AZ, UK