

Epidemiologic significance of *Toxoplasma gondii* infections in turkeys, ducks, ratites and other wild birds: 2009–2020

J. P. Dubey¹ , F. H. A. Murata¹, C. K. Cerqueira-Cézar¹, O. C. H. Kwok¹ and C. Su²

¹United States Department of Agriculture, Agricultural Research Service, Animal Parasitic Diseases Laboratory, Beltsville Agricultural Research Center, Building 1001, Beltsville, MD 20705-2350, USA and ²Department of Microbiology, University of Tennessee, Knoxville, TN 37996-0845, USA

Review

Cite this article: Dubey JP, Murata FHA, Cerqueira-Cézar CK, Kwok OCH, Su C (2021). Epidemiologic significance of *Toxoplasma gondii* infections in turkeys, ducks, ratites and other wild birds: 2009–2020. *Parasitology* **148**, 1–30. <https://doi.org/10.1017/S0031182020001961>

Received: 1 September 2020
Revised: 10 October 2020
Accepted: 12 October 2020
First published online: 19 October 2020

Key words:

Birds; clinical disease; epidemiology; genotype; oocyst; prevalence; *Toxoplasma gondii*

Author for correspondence:

J. P. Dubey,
E-mail: jitender.dubey@ars.usda.gov

Abstract

Toxoplasma gondii infections are common in humans and animals worldwide. Wild and domestic avian species are important in the epidemiology of *T. gondii* infections because felids prey on them and excrete millions of oocysts in the environment, disseminating the infection. Herbivorous birds are also excellent sentinels of environmental contamination with *T. gondii* oocysts because they feed on the ground. *Toxoplasma gondii* infections in birds of prey reflect infections in intermediate hosts. Humans can become infected by consuming undercooked avian tissues. Here, the authors reviewed prevalence, persistence of infection, clinical disease, epidemiology and genetic diversity of *T. gondii* strains isolated from turkeys, geese, ducks, ratites and avian species (excluding chickens) worldwide 2009–2020. Genetic diversity of 102 *T. gondii* DNA samples isolated worldwide is discussed. The role of migratory birds in dissemination of *T. gondii* infection is discussed.

Introduction

Toxoplasma gondii infections are prevalent in humans and animals worldwide. The ingestion of undercooked infected meat or consumption of food and water contaminated with oocysts excreted in cat feces are the main sources of infection. Cats are everywhere and a single cat can excrete millions of oocysts that can remain viable in the environment for months under natural conditions (Dubey, 2010). Estimation of oocyst contamination of the environment is difficult because of low numbers present in soil or water (Lélu *et al.*, 2012). Wild and domestic birds are excellent sentinels of environmental contamination with *T. gondii* oocysts because herbivorous birds feed on the ground, and birds of prey consume hundreds of rodents and other small mammals yearly that are important intermediate hosts of *T. gondii* (Dubey *et al.*, 2020; Lemmi *et al.*, 2020). Migratory birds (penguins, geese and others) can transport the parasite across seas (Sandström *et al.*, 2013). Some species (turkeys, geese, ducks and ostriches) are part of food supply for humans. A study estimated that billions of birds are consumed by cats yearly (Loss *et al.*, 2013). Thus, there is great potential for the spread of *T. gondii* oocysts in the environment.

We recently reviewed the biology of *T. gondii* infections in chickens (*Gallus domesticus*) (Dubey *et al.*, 2020). Here, *T. gondii* infections in other avian species, including domestic turkeys, ducks, geese, ratites and other avian species, are reviewed.

Turkeys (*Meleagris gallopavo*)

Antibodies to *T. gondii* were detected from 11.0 to 89.8% of turkeys surveyed (Table 1). Using a kinetic recombinant antigen (GRA7 and GRA8) ELISA, *T. gondii* antibodies were detected in 387 (20.2%) of 1913 sera from 14 turkey farms in different areas of Germany (Koethe *et al.*, 2011). Seroprevalence varied greatly among farms and within the individual farms, depending on fattening cycles and season which turkeys were slaughtered. Seroprevalences were higher in turkeys slaughtered in summer vs in fall or winter (Koethe *et al.*, 2011).

A very high rate of infection was reported in a study from Iran (Sarkari *et al.*, 2014). Antibodies to *T. gondii* were found in 89.8% of turkeys (Table 1) and *T. gondii* DNA was detected in 61.6% of turkey tissues (Table 2). Viable *T. gondii* was isolated from the muscle of 5 wild hunted turkeys in the USA (Table 3).

Turkeys are considered resistant to clinical toxoplasmosis and there were no reports of clinical toxoplasmosis since 2009. Experimentally, turkeys inoculated intravenously with *T. gondii* tachyzoites or oocysts orally remained healthy, irrespective of the dose (Bangoura *et al.*, 2013; Zöller *et al.*, 2013; Hotop *et al.*, 2014; Maksimov *et al.*, 2018). A kinetic ELISA was developed based on a mixture of recombinant dense granule antigens GRA7 and GRA8 using sera from turkeys inoculated intravenously with Me49 tachyzoites; SAG1 antigen was not suitable in this ELISA using recombinant SAG1 (Koethe *et al.*, 2011). In a subsequent study, information on a large panel of 101 synthetic peptides was obtained on sera from 18 turkeys intravenously inoculated with tachyzoites of 3 strains of *T. gondii* (RH-Type I, Me49-Type II and NED-Type III). The authors concluded that by using selected peptides, it was possible to

Table 1. Seroprevalence of *Toxoplasma gondii* in turkeys (*Meleagris gallopavo*)

Country	Area	Source	No. tested	No. positive	% Positive	Test	Cut-off	Remarks	Reference
Brazil	Pernambuco	28 Farms	204	21	11.0	MAT	1:25		Sá <i>et al.</i> (2016)
Egypt	Kafr El-Sheikh	Farms	17	5	29.4	IHA ^a	1:80		Harfoush and Tahoon (2010)
Germany	5 States	14 Farms, abattoirs	1913	387	20.2	ELISA-in house	1:50	Sex, season association	Koethe <i>et al.</i> (2011)
Iran	Fars	FR, abattoir	54	49	89.8	MAT	1:40	<i>T. gondii</i> DNA and isolation	Sarkari <i>et al.</i> (2014)
Iran	Shiraz	Farms, abattoir	9	1	11.1	MAT	1:20		Asgari <i>et al.</i> (2013)
Iraq	Ninevah	Farms	107	82	76.6	LAT ^b	1:20		Butty (2009)
Nigeria	Ondo, Osun, Oyo	Farms	320	13	4.1	MAT	1:20	Sex association	Ayinmode <i>et al.</i> (2017)
USA	Pennsylvania	Hunted	20	6	30.0	MAT	1:5	<i>T. gondii</i> isolated	Cerqueira-Cézar <i>et al.</i> (2019)

ELISA, enzyme-linked immunosorbent assay; FR, free-range; IHA, indirect haemagglutination assay; LAT, latex agglutination test; MAT, modified agglutination test (Dubey and Desmonts, 1987).

^aIHA (Toxo-IHA Fumouze, Diagnostics, France).

^bLAT (Toxo latex kit from Bio-kit-SA, Barcelona, Spain).

serotype strains up to 9 weeks post-inoculation (p.i.) (Maksimov *et al.*, 2018). Antibodies peaked at 5–7 weeks p.i., using the SAG1-ELISA, and the results varied with the *T. gondii* isolate. Similar results were obtained by the indirect fluorescent antibody assay (IFA) (Maksimov *et al.*, 2018).

In turkeys orally inoculated with oocysts, the parasite was widely disseminated in turkey tissues (Bangoura *et al.*, 2013). Inoculated turkeys were euthanized 6 or 12 weeks p.i. and parasite distribution was assessed by polymerase chain reaction (PCR); no difference was found with respect to 6 or 12 weeks p.i. Brain, heart and drumstick were most frequently infected tissues. The route of inoculation could affect the distribution of parasite DNA. In turkeys inoculated intravenously with tachyzoites, liver, pectoral muscle, heart and brain were affected in decreasing order (Zöller *et al.*, 2013). Judging from the histopathological results, the number of *T. gondii* in turkey tissues was low. Tissue cysts were found in imprints of 1 liver, and 2 pectoral muscles of turkeys parenterally inoculated with tachyzoites (Zöller *et al.*, 2013). By using magnetic-capture PCR and 100 g samples, most *T. gondii* were found in the brain and heart (Koethe *et al.*, 2015). Parasite burden was higher in the drumstick vs pectoral muscles.

Ducks (*Anas spp.*) and geese

Ducks are important for the economy of some countries, especially China. Ducks are a good source of meat and eggs for human consumption. To our knowledge, there are no reports of clinical toxoplasmosis in domestic ducks or geese, but antibodies are common (Tables 4 and 5).

Toxoplasma gondii DNA was found in 9 (7.8%) of 115 muscle samples from 115 ducks and 2 of 42 geese (4.7%) (Zou *et al.*, 2017). Viable *T. gondii* was isolated from tissues of ducks in China, France and Malaysia (Table 3).

Ostriches (*Struthio camelus*) and other ratites

Ratite meat is lean and its human consumption is increasing. Serologic data are summarized in Table 6. Antibodies to *T. gondii* were detected up to 80.0% of sera (Table 6).

Viable *T. gondii* was demonstrated in ostrich tissues. In a Brazilian report, *T. gondii* was detected in brains of ostriches and in soil samples from paddocks (da Silva and Langoni, 2016). Brain (25 g) samples of 38 seropositive and 20 seronegative ostriches were bioassayed in mice. *Toxoplasma gondii* was isolated

from 14 seropositive but not from seronegative ostriches. All strains were apparently pathogenic for mice. *Toxoplasma gondii* DNA was found in peritoneal exudates of mice inoculated from tissues of 8 ostriches, and in brains of mice inoculated with 6 ostrich samples. Nothing was said concerning finding viable *T. gondii*. Of interest is the report of finding *T. gondii*-like oocysts microscopically in soil samples from 5 of 20 paddocks; all were confirmed by PCR. In repeat sampling, *T. gondii*-like oocysts were found microscopically in soil samples from 9 of 20 paddocks and results were confirmed by PCR. It should be noted that only a few oocysts are normally present in soil and their detection is a challenge (J.P.D. own observation).

In marked contrast to the Brazilian report, in a Chinese study, *T. gondii* DNA was not detected in any of the 293 hearts and 77 brains of ostriches (Feng *et al.*, 2017).

A study in Egypt found *T. gondii* DNA in the blood of 9 of 120 ostriches (El-Madawy and Metawea, 2013). These authors also tested tissues of 5 ostriches that had died of toxoplasmosis-like illness; *T. gondii* DNA was found in the brains of 5, hearts of 3 and leg muscle of 1 (El-Madawy and Metawea, 2013) (Comment J.P.D. – tissues of these ostriches should be examined histologically for verification of molecular results).

Other wild avian species

Data on *T. gondii* seroprevalence, viable parasite and DNA characterization are arranged by scientific order of birds, by region and chronologically in Tables 2, 3 and 7.

Carnivorous birds

Prevalence of *T. gondii* in carnivorous birds reflects the prevalence of the parasite in their prey. For example, owls consume hundreds of rodents yearly. In a recent survey of kestrels at an airport site in Italy, *T. gondii* antibodies were detected in 33.3% of kestrel trapped during 2016 and 14.3% of 91 kestrel during 2017; seroprevalence was lower in juveniles than in adults (Iemmi *et al.*, 2020). Based on the remnants of animals in the feces of kestrels, rodents were a major component of the kestrel diet (Iemmi *et al.*, 2020). In a large sample size of raptors in Spain, *T. gondii* antibodies were found in 51.0% of 96 common buzzard, 17.7% of 175 Griffon vulture and 17.0% of Spanish Imperial eagle (Cabezón *et al.*, 2011). Among the raptors, vultures are considered resistant to clinical toxoplasmosis

Table 2. Isolation of viable *Toxoplasma gondii* from wild bird by bioassay in mice and/or cats

Host	Country	Location	No. tested	Tissues	No. isolated	Strain designation	PCR-RFLP genotype (Toxo DB)	Notes	Reference
Accipitriformes									
Bald eagle (<i>Haliaeetus leucocephalus</i>)	USA	Alabama	1	B	1	TgBeAl	1 genotype: #5 (1, TgBeAl)		Yu <i>et al.</i> (2013)
Bald eagle (<i>Haliaeetus leucocephalus</i>)	USA	California	5	H	1	TgHlUs1	1 genotype: #1, Type II (1, TgHlUs1)		Dubey <i>et al.</i> (2011a), Shwab <i>et al.</i> (2014)
Common buzzard (<i>Buteo buteo</i>)	Turkey	Seferihisar, Manisa	25	B, H	9	TgBirdTr_Izmir4,6,7 TgBirdTr_Manisa1-3,5-7	Genotyped by microsatellites. 2 genotypes: Type II, ToxoDB #1 or #3 (6, TgBirdTr_Izmir4, TgBirdTr_Manisa2, TgBirdTr_Manisa3, TgBirdTr_Manisa6, TgBirdTr_Manisa7, TgBirdTr_Izmir7). Type III, ToxoDB #2 (1, TgBirdTr_Manisa1). Mixed types (2, TgBirdTr_Izmir6, TgBirdTr_Manisa5)		Karakavuk <i>et al.</i> (2018)
Ferruginous hawk (<i>Buteo regalis</i>)	USA	Colorado	7	B, H	1	TgBrCoUs1	1 genotype: #1 (1, TgBrCoUs1)		Dubey <i>et al.</i> (2010)
Red-shouldered hawk (<i>Buteo lineatus</i>)	USA	Alabama	1	B	1	TgRshAL	1 genotype: #10, Type I (1, TgRshAL)		Yu <i>et al.</i> (2013)
Red-tailed hawk (<i>Buteo jamaicensis</i>)	USA	Colorado	25	B, H	1	TgBjCoUs1	1 genotype: #1 (1, TgBjCoUs1)		Dubey <i>et al.</i> (2010)
Red-tailed hawk (<i>Buteo jamaicensis</i>)	USA	Wisconsin	1	H	1	TgBjUS1	1 genotype: #15 (1, TgBjUS1)		Dubey <i>et al.</i> (2011a), Shwab <i>et al.</i> (2014)
Roadside hawk (<i>Rupornis magnirostris</i>)	Brazil	Minas Gerais	7	B, H	1	TgWildBrMG1	1 genotype: #108 (1, TgWildBrMG1)		Rêgo <i>et al.</i> (2018)
Rough-legged hawk (<i>Buteo lagopus</i>)	USA	Colorado	4	B, H	1	TgBlCoUs1	1 genotype: #2, Type III (1, TgBlCoUs1)		Dubey <i>et al.</i> (2010)
Swainson's hawk (<i>Buteo swainsoni</i>)	USA	Colorado	13	B, H	2	TgBsCoUs1,2	2 genotypes: #1, Type II (1, TgBsCoUs2), #167 (1, TgBsCoUs1)		Dubey <i>et al.</i> (2010)
Anseriformes									
Canada geese (<i>Branta canadensis</i>)	USA	Maryland	Hunted	H	9	TgGooseUS1-9	5 genotypes #1 (1, TgGooseUS9), #2 (4, TgGooseUS3,5,6,7), #4 (2, TgGooseUS2,8), #266 (1, TgGooseUS1), #267 (1, TgGooseUS4)	One isolate by bioassay in cat	Verma <i>et al.</i> (2016)
Domestic duck (unspecified)	China	Chongqing	12	B, H, K, Li, Lu, Sp	1	ND	PCR (ITS-1)		Zhao <i>et al.</i> (2015)
Domestic duck (unspecified)	Malaysia	Peninsular	23	B, H	4	DK1-4	PCR-RFLP-partial genotyping data	Backyards of homes	Puvanesuaran <i>et al.</i> (2013)
Mallard duck (<i>Anas platyrhynchos</i>)	France	Hunted	2	H	1	ND	Microsatellite markers, 1 genotype: type II, ToxoDB, #1 or #3 (1, one isolate)	Wild	Aubert <i>et al.</i> (2010)

(Continued)

Table 2. (Continued.)

Host	Country	Location	No. tested	Tissues	No. isolated	Strain designation	PCR-RFLP genotype (Toxo DB)	Notes	Reference
Mallard duck (<i>Anas platyrhynchos</i>)	Malaysia	4 states	30 hunted	B, H	4	NS	ND	<i>T. gondii</i> DNA was detected by PCR-RFLP in tissues of inoculated mice	Puvanesuaran <i>et al.</i> (2013)
Mallard duck (<i>Anas platyrhynchos</i>)	Senegal	Saint-Louis	1	B, H	1	TgA117073	Genotyped by 15 microsatellite markers as Type II ToxoDB #1 or #3 (1, TgA117073)		Galal <i>et al.</i> (2019)
Muskovy duck (<i>Cairina moschata</i>)	Senegal	Dakar, Kedougou, Saint-Louis	15	B, H	11	TgA117015, 117017, 117018, 117025, 117032, 117038, 117041, 117054, 117060, 117061, 117070.	12 strains genotyped with 15 microsatellites 4 genotypes: Type II, ToxoDB #1 or #3 (3, TgA117038, 117041, 117054), Type III, ToxoDB #2 (1, TgA117015), Africa 1, ToxoDB #6 (1, TgA117070), Africa 4, ToxoDB #20 (6, TgA117017, 177018, 177025, 177032, 177060, 117061).		Galal <i>et al.</i> (2019)
Mute swan (<i>Cygnus olor</i>)	USA	Great lakes	14	H	3	TgSwanUs1-3	2 genotypes: #2 (2, TgSwanUs1,2), #216 (1, TgSwanUs3)		Dubey <i>et al.</i> (2013)
Charadriiformes									
Yellow-legged gull (<i>Larus michahellis</i>)	Turkey	Balıçova	2	B, H	2	TgBirdTr_Izmir2,3	Genotyped by microsatellite. 2 genotypes: Type II, ToxoDB #1 or #3 (1, TgBirdTr_Izmir2). Type III, ToxoDB #2 (1, TgBirdTr_Izmir3)		Karakavuk <i>et al.</i> (2018)
Columbiformes									
Eared dove (<i>Zenaida auriculata</i>)	Brazil	Paraná	46	Several tissues	12	TgDoveBr1-12	Nine strains typed. 5 genotypes: #1 (4, TgDoveBr1,2,8,9), #6 (1, TgDoveBr7), #17 (1, TgDoveBr11), #65 (1, TgDoveBr12), #182 (2, TgDoveBr6,10)		de Barros <i>et al.</i> (2014)
Rock pigeon (<i>Columba livia</i>)	Mexico	Durango	7	B, H	1	TgPigeonMx1	PCR-RFLP ToxoDB genotype # 9 (1, TgPigeonMx1)	MAT 1:400	Alvarado-Esquivel <i>et al.</i> (2011), Shwab <i>et al.</i> (2014)
Rock pigeon (<i>Columba livia</i>)	Portugal	Lisbon	41	B	24	NS	Genotyping by SAG2 (26 type II, two type III, one type I) and 5 microsatellites markers (12 type II, two type III, one type I and one recombinant)		Waap <i>et al.</i> (2012), Vilares <i>et al.</i> (2014)
Rock pigeon (<i>Columba livia</i>)	Serbia	Belgrade	4	H	3	NS	529 bp, PCR-RFLP-incomplete		Marković <i>et al.</i> (2014)
Falconiformes									
American kestrel (<i>Falco sparverius</i>)	USA	Colorado	5	B, H	1	TgFsCoUs1	1 genotype: #157 (1, TgFsCoUs1)		Dubey <i>et al.</i> (2010)

Southern caracara (<i>Caracara plancus</i>)	Brazil	Minas Gerais	7	B, H	2	TgWildBrMG4,6	2 genotypes: #13 (1, TgWildBrMG6) #290 (1, TgWildBrMG4)	Rêgo <i>et al.</i> (2018)	
Galliformes									
Double-spurred spurfowl (<i>Pternistis bicalcaratus</i>)	Senegal	Kedougou	4	B, H	1	TgA117077	Genotyped with 15 microsatellites. 1 genotype: Africa 1, ToxoDB #6 (1, TgA117077)	Galal <i>et al.</i> (2019)	
Guinea fowl (<i>Numida meleagris</i>)	Brazil	Minas Gerais	2	B, H	1	TgNmBr1	1 genotype: # 1, type II (1, TgNmBr1)	Dubey <i>et al.</i> (2011b)	
Guinea fowl (<i>Numida meleagris</i>)	Senegal	Kedougou	1	B, H	1	TgA117058	Genotyped with 15 microsatellites. 1 genotype: Type Africa 1, ToxoDB genotype #6 (1, TgA117058)	Galal <i>et al.</i> (2019)	
Turkeys (<i>Meleagris gallopavo</i>)	USA	Pennsylvania	20 hunted	H, Sk	5	TgturkeyPa1-5 (designated here)	2 genotypes: #5 (4, TgturkeyPa1-4), #216 (1, TgturkeyPa5) 18NC0055- (TgturkeyPa1) 18-WITU-0002 (TgturkeyPa2) 18-WITU-0012 (TgturkeyPa3) 18-WITU-0015 (TgturkeyPa4) 18-WITU-0019 (TgturkeyPa5)	Cerqueira-Cézar <i>et al.</i> (2019)	
Passeriformes									
Hawaiian crow (<i>Corvus hawaiiensis</i>)	USA	Hawaii	2	B, Li, Sp, Lu	2	TgHcUS1, 2	1 genotype: #177 (2, TgHcUS1, 2)	Work <i>et al.</i> (2000), Shwab <i>et al.</i> (2014)	
Pelecaniformes									
Striated heron (<i>Butorides striata</i>)	Brazil	Pernambuco	2	B, H, Sk	1	TgButstBrPE1	1 genotype: #13 (1, TgButstBrPE1)	Silva <i>et al.</i> (2018b)	
Piciformes									
Campo flicker (<i>Colaptes campestris</i>)	Brazil	Minas Gerais	1	B, H	1	TgWildBrMG3	1 genotype: #11 (1, TgWildBrMG3)	Rêgo <i>et al.</i> (2018)	
Keel-billed toucan (<i>Ramphastos sulfuratus</i>)	Costa Rica	Zoo	1	Sk	1	TgRsCr1	1 genotype: #52 (1, TgRsCr1)	Dubey <i>et al.</i> (2009), Shwab <i>et al.</i> (2014)	
Toco toucan (<i>Ramphastos toco</i>)	Brazil	Minas Gerais	3	B, H	1	TgWildBrMG2	1 genotype: #290 (1, TgWildBrMG2)	Rêgo <i>et al.</i> (2018)	
Psittaciformes									
Peach-Faced Lovebird (<i>Agapornis roseicollis</i>)	Australia	Sydney	1 pet	B	1	AgapornisTg1	1 genotype: #3, Type II variant (1, AgapornisTg1) <i>T. gondii</i> genotyping in B of the bird	Cooper <i>et al.</i> (2015)	
Strigiformes									
Barn owl (<i>Tyto alba</i>)	Turkey	Konak	2	B, H	2	TgBirdTr_Izmir1,5	Genotyped by microsatellites. 2 genotypes: Type II, ToxoDB #1 or #3 (1, TgBirdTr_Izmir5), Type III, ToxoDB #2 (1, TgBirdTr_Izmir1)	Karakavuk <i>et al.</i> (2018)	
Barn owl (<i>Tyto alba</i>)	USA	Colorado	1	B, H	1	TgTaCoUs1	PCR-RFLP ToxoDB genotype # 5 MAT<1:25	Dubey <i>et al.</i> (2010)	
Barred owl (<i>Strix varia</i>)	USA	Alabama	1	B	1	TgSaUS1	1 genotype: #3, Type II variant (1, TgSaUS1) MAT 1:50	Love <i>et al.</i> (2016)	

(Continued)

Table 2. (Continued.)

Host	Country	Location	No. tested	Tissues	No. isolated	Strain designation	PCR-RFLP genotype (Toxo DB)	Notes	Reference
Eurasian eagle-owl (<i>Bubo bubo</i>)	Turkey	Salihli	2	B, H	1	TgBirdTr_Manisa4	Mixed types (1, TgBirdTr_Manisa4)		Karakavuk <i>et al.</i> (2018)
Tropical screech owl (<i>Megascops choliba</i>)	Brazil	Minas Gerais	4	B, H	1	TgWildBrMG5	1 genotype: #8 (1, TgWildBrMG5)		Rêgo <i>et al.</i> (2018)
Struthioniformes									
Ostrich (<i>Struthio camelus</i>)	Brazil	São Paulo	38	B	14	TgOsBr1,2	2 genotypes: # 161 (1, TgOsBr2), #206 (1, TgOsBr1)	Abattoirs	da Silva and Langoni (2016)
Suliformes									
Great cormorant (<i>Phalacrocorax carbo</i>)	Turkey	Çiğli	1	B, H	1	NS	ND		Karakavuk <i>et al.</i> (2018)

B, brain; H, heart; K, kidney; Li, liver; Lu, lung; Sk, skeletal muscle; Sp, spleen; NS, not stated; ND, no data; PCR, polymerase chain reaction; PCR-RFLP, Restriction fragment length polymorphism.

Table 3. *Toxoplasma gondii* DNA from tissues of wild birds

Host	Country	Region	No. tested	Tissue	No. positive	% Positive	Remarks and PCR (gene)	Genotyping	Reference
Accipitriformes									
Black kite (<i>Milvus migrans</i>)	Spain	Several	3	B	1	33.3	N-PCR (529 bp – TOX9,11 primers)	ND	Darwich <i>et al.</i> (2012)
Common buzzard (<i>Buteo buteo</i>)	Turkey	Bergama, Bornova, Çiğli, Kemalpaşa, Konak, Manisa Salihli, Saruhanli, Seferihisar, Turgutlu	25	B, H	23	92.0	RT-PCR, (529 bp – TOX-SE,AS primers). Tg isolated	15 Microsatellite markers – ToxoDB #1 or #3 in 6, #2 in 1 and mixed in 2	Karakavuk <i>et al.</i> (2018)
Eurasian sparrow hawk (<i>Accipiter nisus</i>)	Turkey	Bornova, Çiğli, Karabağlar, Salihli, Saruhanlı	5	B, H	4	80.0	RT-PCR, (529 bp – TOX-SE,AS primers).		Karakavuk <i>et al.</i> (2018)
Griffon vulture (<i>Gyps fulvus</i>)	Spain	Several	105	B	1	0.9	N-PCR (529 bp – TOX9,11 primers)	ND	Darwich <i>et al.</i> (2012)
Anseriformes									
Canada geese (<i>Branta canadensis</i>) Snow geese (<i>Chen caerulescens</i>)	Canada	Nunavik	156	B, G, H, Li, Sk	14	9.0	RT-PCR (529 bp – TOX9,11 primers), highest concentration of parasite DNA (tachyzoite equivalent) in H = 744, B = 300, Sk = 104, Li = 33, G = 8	PCR-RFLP using GRA6 – type II	Bachand <i>et al.</i> (2019)

Eurasian green-winged teal (<i>Anas crecca</i>)	Italy	NS	8	B, H	5	62.5	N-PCR (B1)	ND	Nardoni <i>et al.</i> (2019)
Eurasian teal (<i>Anas crecca</i>)	Italy	Tuscany	3	B	1	33.3	N-PCR (B1)	10 PCR-RFLP markers – genotype incomplete	Mancianti <i>et al.</i> (2013)
Falcated teal (<i>Falcated teal</i>)	China	Jilin	8	H or Lu	1	12.5	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype incomplete	Zhang <i>et al.</i> (2015)
Geese (unspecified)	China	Shandong	42	Sk	2	4.7	N-PCR (B1)		Zou <i>et al.</i> (2017)
Hawaiian geese (<i>Branta sandvicensis</i>)	USA	Hawaii	4	Li, Lu	4	100.0	ND	10 PCR-RFLP markers – ToxoDB genotype #261 (21729-Li, 21774-Lu, 25022-Li), #262 (21850-Li) (DNA from frozen tissues of IHC confirmed toxoplasmosis)	Work <i>et al.</i> (2016)
Mallard (<i>Anas platyrhynchos</i>)	China	Jilin	25	H or Lu	5	20.0	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype incomplete	Zhang <i>et al.</i> (2015)
Mallard (<i>Anas platyrhynchos</i>)	China	Shandong	115	Sk	9	7.8	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #9 in a duck	Zou <i>et al.</i> (2017)
Mallard (<i>Anas platyrhynchos</i>)	Czech Republic	Several	280	B, H, Sk	15	5.4	RT-PCR, (B1,529 bp), <i>Tg</i> genotyped	6 PCR-RFLP markers – 6 type II, 7 type III, 6 type II/III	Skorpikova <i>et al.</i> (2018)
Mallard (<i>Anas platyrhynchos</i>)	Italy	Tuscany	2	B	1	50.0	N-PCR (B1)	10 PCR-RFLP markers – genotype incomplete	Mancianti <i>et al.</i> (2013)
Northern shoveler (<i>Anas clypeata</i>)	Italy	Tuscany	2	B	1	50.0	N-PCR (B1)	10 PCR-RFLP markers – genotype incomplete	Mancianti <i>et al.</i> (2013)
Bucerotiformes									
Southern-Yellow-billed hornbill (<i>Tockus leucomelas</i>)	South Africa	Limpopo	4	B	1	25.0	PCR (B1 – Tg1,2 primers)		Lukášová <i>et al.</i> (2018)
Charadriiformes									
Eurasian stone curlew (<i>Burhinus oedicnemus</i>)	Turkey	Çiğli	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
Little tern (<i>Sternula albifrons</i>)	Turkey	Çiğli	1	B, H	0	0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
Woodcock (<i>Scolopax rusticola</i>)	Greece	Macedonia, Mesolonghi	86	B	4	4.7	PCR (529 bp – Tox -9upAu, Tox-11doAu primers)	ND	Moustakidis <i>et al.</i> (2017)
Yellow-legged gull (<i>Larus michahellis</i>)	Turkey	Balçova, Konak	2	B, H	2	100.0	RT-PCR, (529 bp – TOX-SE,AS primers). <i>Tg</i> isolated	15 Microsatellite markers – ToxoDB #1 or #3 in 1, #2 in 1	Karakavuk <i>et al.</i> (2018)
Ciconiiformes									
Black stork (<i>Ciconia nigra</i>)	Turkey	NS	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
White stork (<i>Ciconia ciconia</i>)	Turkey	Konak	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
Columbiformes									
Laughing dove (<i>Streptopelia senegalensis</i>)	South Africa	Limpopo	4	B	1	25.0	PCR (B1 – Tg1,2 primers)		Lukášová <i>et al.</i> (2018)

(Continued)

Table 3. (Continued.)

Host	Country	Region	No. tested	Tissue	No. positive	% Positive	Remarks and PCR (gene)	Genotyping	Reference
Mourning dove (<i>Zenaida macroura</i>)	USA	Tennessee	186	B	2	1.0	PCR (529 bp, TOX4,5 primers)		Ammar <i>et al.</i> (2020)
Red-eyed dove (<i>Streptopelia semitorquata</i>)	South Africa	Limpopo	5	B	1	20.0	PCR (B1 – Tg1,2 primers)	15 microsatellites markers – genotyping type II	Lukášová <i>et al.</i> (2018)
Rock pigeon (<i>Columba livia</i>)	Iran	Khuzestan	43	B, H	3	6.9	PCR (B1 – TG1,2 primers)	PCR-RFLP using GRA6, 1 type II and 2 type III	Khademvatan <i>et al.</i> (2013)
Rock pigeon (<i>Columba livia</i>)	Pakistan	Punjab	54	H, Sk	19	35.1	PCR (B1 – TOX4,5 primers)	ND	Nazir <i>et al.</i> (2018)
Rock pigeon (<i>Columba livia</i>)	Portugal	Lisbon	41	B	28	68.2	PCR (B1)	Genotyping by SAG2 (26 type II, 2 type III, 1 type I), and 5 microsatellites markers (12 type II, 2 type III, 1 type I, 1 recombinant)	Vilares <i>et al.</i> (2014)
Rock pigeon (<i>Columba livia</i>)	Serbia	Belgrade	7	H	5	71.4	RT-PCR (529 bp)	6 PCR-RFLP markers – 2 type II and 1 type III	Marković <i>et al.</i> (2014)
Wood pigeon (<i>Columba palumbus</i>)	Italy	NS	1	B, H	1	100.0	N-PCR (B1)	ND	Nardoni <i>et al.</i> (2019)
Wood pigeon (<i>Columba palumbus</i>)	Turkey	Konak	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
Falconiformes									
Common kestrel (<i>Falco tinnunculus</i>)	Italy	NS	3	B, H	2	66.6	N-PCR (B1)	ND	Nardoni <i>et al.</i> (2019)
Common kestrel (<i>Falco tinnunculus</i>)	Poland	Several	3	B, H, Li, Sk	1	33.3	PCR (B1)	5 PCR-RFLP markers – type II/III	Sroka <i>et al.</i> (2019)
Common kestrel (<i>Falco tinnunculus</i>)	Turkey	Konak	1	B, H	0	0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
Eurasian jay (<i>Garrulus glandarius</i>)	Slovakia	Tatra National Park	2	Sk	1	50.0	PCR (TGR1E – TGR1E-1,2 primers)	PCR-RFLP using SAG2 – type III	Turčeková <i>et al.</i> (2014)
New Zealand falcon (<i>Falco novaeseelandiae</i>)	New Zealand	Palmerston	35	H, Li, Lu, Sp	2	5.7	N-PCR (Pppk-dhps)	7 PCR-RFLP markers – genotype incomplete	Mirza <i>et al.</i> (2017)
Northern goshawk (<i>Accipiter gentilis</i>)	Slovakia	Tatra National Park	2	Sk	1	50.0	PCR (TGR1E – TGR1E-1,2 primers)	PCR-RFLP using SAG2 – type II	Turčeková <i>et al.</i> (2014)
Peregrine falcon (<i>Falco peregrinus</i>)	Turkey	Çiğli	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)
Galliformes									
Quail (<i>Coturnix coturnix</i>)	China	Shandong	390	Sk	25	6.4	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #9 in 5	Cong <i>et al.</i> (2017b)
Turkeys (<i>Meleagris gallopavo</i>)	Iran	Fars	54	B, Sk, T	33	66.0	N-PCR (B1). Samples from abattoirs	ND	Sarkari <i>et al.</i> (2014)

Gruiformes										
Eurasian coot (<i>Fulica atra</i>)	China	Jilin	25	H or Lu	1	4.0	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype incomplete	Zhang <i>et al.</i> (2015)	
Passeriformes										
Hooded crow (<i>Corvus cornix</i>)	Iran	Tehran	55	B	9	16.3	N-PCR (GRA6)	PCR-RFLP using GRA6 – type III	Abdoli <i>et al.</i> (2018)	
Hooded crow (<i>Corvus cornix</i>)	Israel	Haifa	101	B	1	1.0	PCR (529 bp – TOX4,5 primers)	11 PCR-RFLP markers – type II	Salant <i>et al.</i> (2013)	
House sparrows (<i>Passer domesticus</i>)	Brazil	Bahia, Pernambuco	40	B	10	10.0	N-PCR (ITS1)	ND	Gondim <i>et al.</i> (2010)	
House sparrows (<i>Passer domesticus</i>)	Brazil	Pernambuco	10	B	3	30.0	N-PCR (B1)		Vilela <i>et al.</i> (2011)	
House sparrows (<i>Passer domesticus</i>)	China	Lanzhou	39	B, H, Lu	11	28.2	N-PCR (B1)	9 PCR-RFLP markers – ToxoDB genotype #3 in 3 and a new type in 1	Cong <i>et al.</i> (2013)	
House sparrows (<i>Passer domesticus</i>)	China	Several	22	Sk	1	4.5	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #3	Huang <i>et al.</i> (2012)	
House sparrows (<i>Passer domesticus</i>)	Iran	Khuzestan	64	B, H	17	26.5	PCR (B1 – TG1,2 primers)	PCR-RFLP using GRA6, 1 type II and 16 type III	Khademvatan <i>et al.</i> (2013)	
House sparrows (<i>Passer domesticus</i>)	Iran	Tehran	200	B	17	8.5	LAMP, PCR (529 bp – B3,F3, TOX4,5 primers)	ND	Abdoli <i>et al.</i> (2016)	
Magpie (<i>Pica pica</i>)	Italy	Tuscany	41	Bl, H	15	36.6	N-PCR (B1)	5 PCR-RFLP markers – 8 type II and 7 type III	Mancianti <i>et al.</i> (2020)	
Magpie (<i>Pica pica</i>)	Slovakia	Tatra National Park	3	Sk	1	33.3	PCR (TGR1E – TGR1E-1,2 primers)	PCR-RFLP using SAG2 – type III	Turčeková <i>et al.</i> (2014)	
Magpie (<i>Pica pica</i>)	Spain	Several	33	B	5	12.8	N-PCR (529 bp – TOX9,11 primers)	ND	Darwich <i>et al.</i> (2012)	
Olive-backed pipit (<i>Anthus hodgsoni</i>)	China	Hunan	44	B	1	2.2	N-PCR (B1)		Liu <i>et al.</i> (2019)	
Oriental skylark (<i>Alauda gulgula</i>)	China	Lanzhou, Tianshui	34	B	3	8.8	N-PCR (B1)	10 PCR-RFLP markers – type II variant in 2, ToxoDB genotype #3	Cong <i>et al.</i> (2014)	
Starling (<i>Sturnus vulgaris</i>)	Iran	Khuzestan	39	B, H	5	12.8	PCR (B1 – TG1,2 primers)	PCR-RFLP using GRA6, 2 type II and 3 type III	Khademvatan <i>et al.</i> (2013)	
Thrush nightingale (<i>Luscinia luscinia</i>)	Turkey	Konak	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)	
Tree sparrows (<i>Passer montanus</i>)	China	Hunan	131	B	11	8.4	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #10 in 3 sparrows	Liu <i>et al.</i> (2019)	
Tree sparrows (<i>Passer montanus</i>)	China	Several	35	Sk	1	2.8	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #10	Huang <i>et al.</i> (2012)	
Yellow-breasted bunting (<i>Emberiza aureola</i>)	China	Hunan	26	B	1	3.8	N-PCR (B1)		Liu <i>et al.</i> (2019)	
Phoenicopteriformes										
Great flamingo (<i>Phoenicopterus roseus</i>)	Turkey	Konak	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk <i>et al.</i> (2018)	

(Continued)

Table 3. (Continued.)

Host	Country	Region	No. tested	Tissue	No. positive	% Positive	Remarks and PCR (gene)	Genotyping	Reference
Psittiformes									
Black-headed gull (<i>Chroicocephalus ridibundus</i>)	Italy	NS	4	B, H	4	100.0	N-PCR (B1)	ND	Nardoni et al. (2019)
Black-headed gull (<i>Chroicocephalus ridibundus</i>)	Poland	Several	2	B, H, Li, Sk	1	50.0	PCR (B1)	5 PCR-RFLP markers type II/III	Sroka et al. (2019)
Strigiformes									
Barn owl (<i>Tyto alba</i>)	Turkey	Çiğli, Konak	2	B, H	2	100.0	RT-PCR, (529 bp – TOX-SE,AS primers). Tg isolated	15 Microsatellite markers – ToxoDB #1 or #3 in 1, #2 in 1	Karakavuk et al. (2018)
Eurasian eagle-owl (<i>Bubo bubo</i>)	Turkey	Kemalpaşa, Salihli	2	B, H	2	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)	15 Microsatellite markers – mixed type in 1	Karakavuk et al. (2018)
Little owl (<i>Athene noctua</i>)	Turkey	Çiğli	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk et al. (2018)
Morepork (<i>Ninox novaeseelandiae</i>)	New Zealand	Palmerston	36	H, Li, Lu, Sp	3	8.3	PCR (Pppk-dhps)	7 PCR-RFLP markers – genotype incomplete	Mirza et al. (2017)
Struthioniformes									
Ostrich (<i>Struthio camelus</i>)	Brazil	São Paulo	38	B	25	65.7	PCR (529 bp and 18S rRNA – TOX4,5 and Tg18s48F,359R primers). Tg isolated	10 PCR-RFLP markers – ToxoDB #161 and #206 in 2	da Silva and Langoni (2016)
Ostrich (<i>Struthio camelus</i>)	Egypt	Ismailia	120	Bl	9	7.5	N-PCR (B1)	ND	El-Madawy and Metawea (2013)
Suliformes									
Australasian harrier (<i>Circus approximans</i>)	New Zealand	Palmerston	46	H, Li, Lu, Sp	4	8.7	PCR (Pppk-dhps)	7 PCR-RFLP markers – genotype incomplete	Mirza et al. (2017)
Baikal teal (<i>Anas formosa</i>)	China	Jilin	50	H or Lu	11	22.0	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #9 in 2	Zhang et al. (2015)
Common pheasants (<i>Phasianus colchicus</i>)	China	Several	98	Sk	2	2.0	N-PCR (B1)	10 PCR-RFLP markers – ToxoDB genotype #3	Huang et al. (2012)
Common pheasant (<i>Phasianus colchicus</i>)	Czech Republic	Several	350	B, H, Sk	12	3.4	RT-PCR, (B1,529 bp), Tg genotyped	6 PCR-RFLP markers – 1 type II, 4 type III, 4 type II/III	Skorpikova et al. (2018)
Dalmatian pelican (<i>Pelecanus crispus</i>)	Turkey	Çiğli	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers)		Karakavuk et al. (2018)
Eurasian jays (<i>Garrulus glandarius</i>)	Spain	Several	23	B	5	21.7	N-PCR (529 bp – TOX9,11 primers)	ND	Darwich et al. (2012)
Eurasian siskin (<i>Carduelis spinus</i>)	China	Lanzhou	41	B	5	12.2	N-PCR (B1)	10 PCR-RFLP markers – type II variant in 2, ToxoDB #3	Cong et al. (2014)
Great cormorant (<i>Phalacrocorax carbo</i>)	Turkey	Çiğli	1	B, H	1	100.0	RT-PCR, (529 bp – TOX-SE,AS primers). Tg isolated		Karakavuk et al. (2018)

B, brain; Bl, blood; H, heart; G, gizzard; Li, liver; Lu, lung; Sk, muscle; Sp, spleen; Tg, *Toxoplasma gondii*; ND, not done; NS, not stated; PCR, polymerase chain reaction; N-PCR, nested PCR; RT-PCR, real-time PCR; PCR, polymerase chain reaction; PCR-RFLP, Restriction fragment length polymorphism.

Table 4. Seroprevalence of *Toxoplasma gondii* in domestic ducks

Country	Area	Source	No. tested	No. positive	% Positive	Test	Cut-off	Remarks	Reference
China	Chongqing	NS	635 – FR	84	13.3	MAT	1:25	Tg isolated. Location, type of raising AS	Zhao <i>et al.</i> (2015)
			527 – caged	35	6.6				
China	Guangzhou	Abattoirs	349	56	16.0	MAT	1:5	6 had MAT titers of 1:40 or higher	Yan <i>et al.</i> (2009)
China	Jilin	Farms	268	29	0.8	LAT ^a	1:64		Li <i>et al.</i> (2020)
China	Lanzhou	Abattoirs	111 – caged	7	6.3	MAT	1:5		Cong <i>et al.</i> (2012)
			223 – FR	31	13.9				
China	Liaoning	Farms	268	26	9.7	MAT	1:25		Wang <i>et al.</i> (2014)
China	Shenyang	Abattoirs	146 – caged	11	7.5	MAT	1:25		Yang <i>et al.</i> (2012)
			122 – FR	15	12.3				
Czech Republic	Several	Abattoirs	360	52	14.0	IFA	1:40		Bártová <i>et al.</i> (2009)
Egypt	Behera	Abattoirs	151	21	13.9	MAT	1:25		AbouLaila <i>et al.</i> (2011)
Egypt	Kafr El-Sheikh	Farms	58	32	55.0	IHA ^a	1:80		Harfoush and Tahoon (2010)
Egypt	5 regions	Market	142	15	10.5	ELISA		IHC 3 positive. Region AS	Ibrahim <i>et al.</i> (2018)
Germany	Lower Saxony	61 farms	2534	145	5.7	ELISA – SAG1	1:200	Protection AS to indoors kept animals	Maksimov <i>et al.</i> (2011)
Iraq	Al-Qadisiya	Farms	50	28	56.0	LAT ^a	1:2		Alkhaled <i>et al.</i> (2012)
Iran	Mazandaran	Market	87	40	46.0	MAT	1:20		Amouei <i>et al.</i> (2018)
Malaysia	Johor, Kedah, Melaka, Perak	Farms, FR	205	30	14.6	MAT	1:6	Tg DNA and genotyping	Puvanesaruan <i>et al.</i> (2013)
Poland	Lublin	Farms	33	7	21.2	MAT ^a	1:40		Sroka <i>et al.</i> (2010)
Senegal	Dakar, Saint-Louis	Households	306	16	5.2	MAT	1:20	Tg isolated	Galal <i>et al.</i> (2019)

NS, not stated; ELISA, enzyme-linked immunosorbent assay; IFA, indirect fluorescent antibody test; IHAT, indirect haemagglutination test; LAT, latex agglutination test; MAT, modified agglutination test (Dubey and Desmonts, 1987); FR, free range; Tg, *Toxoplasma gondii*; AS, association; IHC, immunohistochemical.

^aIHA (Toxo-IHA Fumouze Diagnostics, France); LAT (PLASMATECH Co., UK); MAT (Toxo-Screen DA[®], Biomerieux, Lyon, France). This is the same test as MAT.

Table 5. Seroprevalence of *Toxoplasma gondii* in domestic goose

Country	Area	Source	No. tested	No. positive	% Positive	Test	Cut-off	Remarks	Reference
China	Guangdong	Farms	274	41	14.9	MAT	1:5	–	Yan <i>et al.</i> (2011b)
China	Guangdong	Farms	520	9	1.7	MAT		Age, management	Wang <i>et al.</i> (2012)
China	Hainan	Farms	600	102	17.0	IHA ^a	1:64	Presence of cats, hygiene AS. Tg DNA and genotyping	Rong <i>et al.</i> (2014)
China	Jilin	Farms	379	50	13.2	IHA ^a	1:64		Li <i>et al.</i> (2020)
China	Liaoning	Farms	128	9	7.0	MAT	1:25		Wang <i>et al.</i> (2014)
China	Shenyang	Abattoirs	83 – caged	5	6.0	MAT	1:25		Yang <i>et al.</i> (2012)
			45 – FR	4	8.9				
Czech Republic	Bohemia, Olomouc	Abattoirs	178	77	43.0	IFA	1:40		Bártová <i>et al.</i> (2009)
Germany	Lower Saxony	13 farms	373	94	25.2	ELISA-SAG1	1:200	Protection AS to indoors kept animals	Maksimov <i>et al.</i> (2011)
Iran	Mazandaran	Market	5	5	100.0	MAT	1:20	–	Amouei <i>et al.</i> (2018)
USA	Maryland	Hunted	169	12	7.1	MAT	1:25	Canada geese, Tg isolated	Verma <i>et al.</i> (2016)

AS, association; FR, free-range; Tg, *Toxoplasma gondii*; ELISA, enzyme-linked immunosorbent assay; IFA, indirect fluorescent antibody test; IHA, indirect haemagglutination assay; MAT, modified agglutination test (Dubey and Desmonts, 1987).

^aIHA kit (Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, China).

Table 6. Seroprevalence of *Toxoplasma gondii* in ratites

Host	Country	Area	Source	No. tested	No. positive	% positive	Test	Cut-off	Remarks	Reference
Emus (<i>Dromaius novaehollandiae</i>)	Brazil	Goiás	Farm	16	8	50.0	MAT	1:25		Gallo et al. (2019)
Ostriches (<i>Struthio camelus</i>)	Brazil	Mato Grosso, Rio Grande do Sul, São Paulo	3 Farms	46	8	17.4	MAT	1:16		Almeida et al. (2013)
Ostriches (<i>Struthio camelus</i>)	Brazil	Rio de Janeiro	Abattoirs	20	16	80.0	MAT	1:25		Gallo et al. (2019)
Ostriches (<i>Struthio camelus</i>)	Brazil	São Paulo	Farms	195	28	14.3	MAT	1:16	MAT titer 1:16384 in 2 ostriches	Contente et al. (2009)
Ostriches (<i>Struthio camelus</i>)	Brazil	São Paulo	Abattoir, 4 farms	344	38	11.0	MAT	1:8	Tg DNA and parasite isolation, Water source, presence of non-ostrich feces AS	da Silva and Langoni (2016)
Ostriches (<i>Struthio camelus</i>)	China	Hebei, Henan	Abattoirs	315	20	6.4	MAT	1:25	Tg DNA and parasite isolation	Feng et al. (2017)
Ostriches (<i>Struthio camelus</i>)	Egypt	Ismailia	Farm	120	15	12.5	MAT	1:25	Tg DNA. 5 (4.2%) IgM positive	El-Madawy and Metawea (2013)
Rheas (<i>Rhea americana</i>)	Brazil	Espírito Santo, Goiás, São Paulo	Farm	68	18	26.5	MAT	1:25		Gallo et al. (2019)
Rheas (<i>Rhea americana</i>)	Brazil	Rio Grande do Sul, Santa Catarina	2 Farms	20	10	50.0	MAT	1:25		Almeida et al. (2013)

MAT, modified agglutination test (Dubey and Desmonts, 1987); AS, association; Tg, *Toxoplasma gondii*.

Table 7. Serologic prevalence of antibodies to *Toxoplasma gondii* in wild birds

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Accipitriformes									
Bald eagle (<i>Haliaeetus leucocephalus</i>)	Czech Republic	Zoos	1	1	100.0	LAT ^a			Bártová <i>et al.</i> (2018)
Bald eagle (<i>Haliaeetus leucocephalus</i>)	USA	Alabama	13	12	92.3	MAT	1:25		Love <i>et al.</i> (2016)
Bearded vulture (<i>Gypaetus barbatus</i>)	Spain	Several	15	6	42.8	MAT	1:25		Cabezón <i>et al.</i> (2011)
Black kite (<i>Milvus migrans</i>)	Portugal	Central, northern	1	1	100.0	MAT	1:20		Lopes <i>et al.</i> (2011)
Black kite (<i>Milvus migrans</i>)	Senegal	Dakar	2	2	100.0	MAT	1:20		Galal <i>et al.</i> (2019)
Black kite (<i>Milvus migrans</i>)	Spain	Several	17	5	29.4	MAT	1:25		Cabezón <i>et al.</i> (2011)
Bonelli's eagle (<i>Aquila fasciata</i>)	Spain	Several	9	1	11.1	MAT	1:25		Cabezón <i>et al.</i> (2011)
Booted eagle (<i>Hieraaetus pennatus</i>)	Spain	Southern	1	1	100.0	MAT	1:25		Cano-Terriza <i>et al.</i> (2015)
Broad-winged hawk (<i>Buteo platypterus</i>)	USA	Alabama	20	6	30.0	MAT	1:25		Love <i>et al.</i> (2016)
Cinereous vulture (<i>Aegypius monachus</i>)	Czech Republic	Zoos	11	7	64.0	LAT ^a	–		Bártová <i>et al.</i> (2018)
Cinereous vulture (<i>Aegypius monachus</i>)	Spain	Several	23	6	26.0	MAT	1:25		Cabezón <i>et al.</i> (2011)
Common buzzard (<i>Buteo buteo</i>)	Portugal	Central, northern	26	18	69.2	MAT	1:20		Lopes <i>et al.</i> (2011)
Cooper's hawk (<i>Accipiter cooperii</i>)	USA	Alabama	12	6	50.0	MAT	1:25		Love <i>et al.</i> (2016)
Crested goshawk (<i>Accipiter trivirgatus</i>)	Taiwan	Several	41	7	17.0	MAT ^b	1:40		Chen <i>et al.</i> (2015)
Eagle (<i>Aquila</i> spp.)	Mexico	Durango	2	1	50.0	MAT	1:25	Location AS	Alvarado-Esquivel <i>et al.</i> (2011)
Eurasian buzzard (<i>Buteo buteo</i>)	Italy	Northern	12	1	8.3	MAT ^b	1:40		Gazzonis <i>et al.</i> (2018)
Eurasian buzzard (<i>Buteo buteo</i>)	Spain	Several	96	49	51.0	MAT	1:25		Cabezón <i>et al.</i> (2011)
Eurasian griffon (<i>Gyps fulvus</i>)	Czech Republic	Zoos	4	3	75.0	LAT ^a	–		Bártová <i>et al.</i> (2018)
Eurasian sparrow hawk (<i>Accipiter nisus</i>)	Italy	Northern	2	1	50.0	MAT ^b	1:40		Gazzonis <i>et al.</i> (2018)
Golden eagle (<i>Aquila chrysaetos</i>)	Spain	Several	8	5	62.5	MAT	1:25		Cabezón <i>et al.</i> (2011)
Golden eagle (<i>Aquila chrysaetos</i>)	USA	Alabama	1	1	100.0	MAT	1:25		Love <i>et al.</i> (2016)
Griffon vulture (<i>Gyps fulvus</i>)	Israel	4 Areas	101	40	39.6	MAT	1:25		Salant <i>et al.</i> (2013)
Griffon vulture (<i>Gyps fulvus</i>)	Spain	Several	175	31	17.7	MAT	1:25		Cabezón <i>et al.</i> (2011)
Montagu's harrier (<i>Circus pygargus</i>)	Spain	Several	7	1	14.3	MAT	1:25		Cabezón <i>et al.</i> (2011)
Northern goshawk (<i>Accipiter gentilis</i>)	Portugal	Central, northern	3	3	100.0	MAT	1:20		Lopes <i>et al.</i> (2011)
Northern goshawk (<i>Accipiter gentilis</i>)	Spain	Several	5	2	40.0	MAT	1:25		Cabezón <i>et al.</i> (2011)
Osprey (<i>Pandion haliaetus</i>)	Spain	Several	7	2	28.5	MAT	1:25		Cabezón <i>et al.</i> (2011)

(Continued)

Table 7. (Continued.)

Order, common name, (scientific name)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Red kite (<i>Milvus milvus</i>)	Spain	Several	3	1	33.3	MAT	1:25		Cabezón et al. (2011)
Red-shouldered hawk (<i>Buteo lineatus</i>)	USA	Alabama	41	9	21.9	MAT	1:25		Love et al. (2016)
Red tailed hawk (<i>Buteo jamaicensis</i>)	USA	Alabama	71	22	30.9	MAT	1:25		Love et al. (2016)
Roadside hawk (<i>Rupornis magnirostris</i>)	Brazil	São Paulo	1	1	100.0	MAT	1:16		Gonçalves et al. (2013)
Serpent eagle (<i>Spilornis cheela</i>)	Taiwan	Several	43	20	46.5	MAT ^b	1:40		Chen et al. (2015)
Short-toed Snake-eagle (<i>Circaetus gallicus</i>)	Spain	Several	10	5	50.0	MAT	1:25		Cabezón et al. (2011)
Spanish Imperial eagle (<i>Aquila adalberti</i>)	Spain	Several	146	25	17.0	MAT	1:25		Cabezón et al. (2011)
Tawny eagle (<i>Aquila rapax</i>)	Czech Republic	Zoos	1	1	100.0	LAT ^a	–		Bártová et al. (2018)
Western marsh-harrier (<i>Circus aeruginosus</i>)	Italy	Northern	2	1	50.0	MAT ^b	1:40		Gazzonis et al. (2018)
Western marsh-harrier (<i>Circus aeruginosus</i>)	Spain	Several	6	3	50.0	MAT	1:25		Cabezón et al. (2011)
Anseriformes									
Bar-headed goose (<i>Anser indicus</i>)	Spain	Córdoba	11	1	9.1	MAT	1:25		Cano-Terriza et al. (2015)
Barnacle goose (<i>Branta leucopsis</i>)	Europe	Russia, Svalbard, The Netherlands	1087	98	9.0	MAT ^b	1:40	See text	Sandström et al. (2013)
Black swan (<i>Cygnus atratus</i>)	Portugal	Zoo	7	1	14.3	MAT ^b	1:20		Tidy et al. (2017)
Canada goose (<i>Branta canadensis</i>)	Canada	Saskatoon	71	2	2.8	MAT ^b	1:40		Al-Adhami et al. (2016)
Canada goose (<i>Branta canadensis</i>)	Canada	Nunavik	148	16	10.8	MAT ^a	1:25		Bachand et al. (2019)
Canada goose (<i>Branta canadensis</i>)	Europe	The Netherlands	38	3	7.9	MAT ^b	1:40	See text	Sandström et al. (2013)
Common teal (<i>Anas crecca</i>)	Italy	Tuscany	41	3	7.3	MAT ^b	1:10	PCR positive in 1 seropositive	Mancianti et al. (2013)
Coscoroba swan (<i>Coscoroba coscoroba</i>)	Czech Republic	Zoos	2	1	50.0	LAT ^a			Bártová et al. (2018)
Domestic geese (<i>Anser anser</i>)	Europe	The Netherlands	161	13	8.1	MAT ^b	1:40	See text	Sandström et al. (2013)
Emperor goose (<i>Chen canagica</i>)	Spain	Southern	2	1	50.0	MAT	1:25		Cano-Terriza et al. (2015)
Eurasian green-winged teal (<i>Anas crecca</i>)	Iran	Mazandaran	20	12	60.0	MAT	1:20		Amouei et al. (2018)
Garganey (<i>Spatula querquedula</i>)	Senegal	Dakar	28	8	28.5	MAT	1:20		Galal et al. (2019)
Geese (<i>Anser</i> sp.)	Brazil	Paraná	149	27	18.1	IFA	1:25	Higher prevalence in animals from urban parks	Konell et al. (2019)

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Lesser snow geese (<i>Chen caerulescens</i>)	Canada	Karrak Lake Nunavut	121	43	36.0	IFA	1:20	Results higher than MAT ^b at 1:40 dilution	Elmore <i>et al.</i> (2014)
Lesser snow geese (<i>Chen caerulescens</i>)	Canada	Nunavut	233	66	28.3	ELISA			Elmore <i>et al.</i> (2015)
Lesser snow geese (<i>Chen caerulescens</i>)	Canada	Nunavik	8	2	25.0	MAT ^a	1:25		Bachand <i>et al.</i> (2019)
Ross's geese (<i>Chen rossii</i>)	Canada	Nunavut	234	76	32.4	ELISA			Elmore <i>et al.</i> (2015)
Mallard duck (<i>Anas brachyrhynchus</i>)	Europe	Denmark, Svalbard	573	59	10.2	MAT ^b	1:40	See text	Sandström <i>et al.</i> (2013)
Mallard duck (<i>Anas platyrhynchos</i>)	Mexico	Durango	2	1	50.0	MAT	1:25	Location AS	Alvarado-Esquivel <i>et al.</i> (2011)
Mallard duck (<i>Anas platyrhynchos</i>)	Iran	Mazandaran	20	9	45.0	MAT	1:20	–	Amouei <i>et al.</i> (2018)
Mallard duck (<i>Anas platyrhynchos</i>)	Italy	Tuscany	17	2	11.7	MAT ^b	1:10	PCR positive in 1 seropositive	Mancianti <i>et al.</i> (2013)
Mallard duck (<i>Anas platyrhynchos</i>)	Portugal	Zoo	4	3	75.0	MAT ^b	1:20		Tidy <i>et al.</i> (2017)
Mallard duck (<i>Anas platyrhynchos</i>)	Spain	Several	6	2	33.3	MAT	1:25		Cabezón <i>et al.</i> (2011)
Mallard duck (<i>Anas platyrhynchos</i>)	Spain	Southern	4	1	25.0	MAT	1:25		Cano-Terriza <i>et al.</i> (2015)
Mallard duck (<i>Anas platyrhynchos</i>)	Taiwan	Several	1	1	100.0	MAT ^b	1:40		Chen <i>et al.</i> (2015)
Mandarin duck (<i>Aix galericulata</i>)	Portugal	Zoo	6	3	50.0	MAT ^b	1:20		Tidy <i>et al.</i> (2017)
Mexican duck (<i>Anas diazi</i>)	Mexico	Durango	2	1	50.0	MAT	1:25	Location AS	Alvarado-Esquivel <i>et al.</i> (2011)
Mute swan (<i>Cygnus olor</i>)	USA	Northeastern	632	54	8.5	MAT	1:25	Tg isolated	Dubey <i>et al.</i> (2013)
Northern pintail (<i>Anas acuta</i>)	Spain	Several	1	1	100.0	MAT	1:25		Cabezón <i>et al.</i> (2011)
Orinoco (<i>Neochen jubata</i>)	Brazil	Goiás	41	35	85.3	IFA	1:20	<i>N. caninum</i> in 12.1%	André <i>et al.</i> (2019)
Paradise shelduck (<i>Tadorna variegata</i>)	Czech Republic	Zoos	1	1	100.0	LAT ^a	–		Bártová <i>et al.</i> (2018)
Ross's geese (<i>Chen rossii</i>)	Canada	Karrak Lake Nunavut	123	48	39.0	IFA	1:20	Results higher than MAT ^b at 1:40 dilution	Elmore <i>et al.</i> (2014)
Shoveller duck (<i>Anas clypeata</i>)	Italy	Tuscany	11	2	18.1	MAT ^b	1:10	PCR positive in 1 seropositive	Mancianti <i>et al.</i> (2013)
Sudanese duck (<i>Anas platyrhynchos</i>)	Egypt	Several	142	15	10.5	ELISA		Region AS. IHC	Ibrahim <i>et al.</i> (2018)
Swan goose (<i>Anser cygnoides</i>)	Portugal	Zoo	1	1	100.0	MAT ^b	1:20		Tidy <i>et al.</i> (2017)
Tundra swan (<i>Cygnus columbianus</i>)	China	Jiangxi	9	2	22.2	IHA ^b	1:64		Luo <i>et al.</i> (2017)
Wood duck (<i>Aix sponsa</i>)	Portugal	Zoo	7	2	28.6	MAT ^b	1:20		Tidy <i>et al.</i> (2017)
Bucerotiformes									
Eurasian hoopoe (<i>Upupa epops</i>)	Spain	Several	1	1	100.0	MAT	1:25		Cabezón <i>et al.</i> (2011)

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Cariamiformes									
Crested seriema (<i>Cariama cristata</i>)	Brazil	Minas Gerais, São Paulo	1	1	100.0	MAT	1:5		Vitaliano et al. (2014)
Charadriiformes									
Audouin's gull (<i>Larus audouinii</i>)	Spain	Alboran Island	46	1	2.2	MAT	1:25	Age, food source, year AS	Cabezón et al. (2016)
Caracara (<i>Caracara plancus</i>)	Brazil	Pernambuco	115	6	5.2	MAT	1:25		Silva et al. (2018a)
Common kestrel (<i>Falco tinnunculus</i>)	Italy	Northern	18	1	5.5	MAT ^b	1:40		Gazzonis et al. (2018)
Common snipe (<i>Gallinago gallinago</i>)	Italy	Tuscany	8	2	25.0	MAT ^b	1:10		Mancianti et al. (2013)
Common snipe (<i>Gallinago gallinago</i>)	Pakistan	Punjab	4	1	25.0	LAT ^b	1:64	Age, health status AS	Naveed et al. (2019)
Yellow-legged gull (<i>Larus michahellis</i>)	Europe	France, Spain, Tunisia	988 nests	233	23.6	ELISA ^b		ELISA in 1122 egg yolk. Different prevalence between colonies AS	Gamble et al. (2019)
Yellow-legged gull (<i>Larus michahellis</i>)	Spain	Iberian Peninsula	479	109	22.8	MAT	1:25	Age, food source, year AS	Cabezón et al. (2016)
Ciconiiformes									
Black stork (<i>Ciconia nigra</i>)	Spain	Several	1	0	0	MAT	1:25		Cabezón et al. (2011)
White stork (<i>Ciconia ciconia</i>)	Spain	Several	64	9	14.1	MAT	1:25		Cabezón et al. (2011)
Columbiformes									
Dove (<i>Spilopelia</i> sp.)	Senegal	Dakar	1	1	100.0	MAT	1:20		Galal et al. (2019)
Eared dove (<i>Zenaida auriculata</i>)	Brazil	Paraná	206	46	22.3	MAT	1:16	Location AS. Tg isolated	de Barros et al. (2014)
Eurasian collared dove (<i>Streptopelia decaocto</i>)	Pakistan	Punjab	10	2	20.0	LAT ^b	1:64	Age, health status AS	Naveed et al. (2019)
Feral pigeon (<i>Columba livia</i> var. <i>domestica</i>)	Spain	Southern	142	13	9.2	MAT	1:25		Cano-Terriza et al. (2015)
Gray-fronted dove (<i>Leptotila rufaxilla</i>)	Brazil	Paraíba	5	1	20.0	MAT	1:25		Andrade et al. (2016)
Oriental turtle dove (<i>Streptopelia orientalis</i>)	Taiwan	Several	16	4	25.0	MAT ^b	1:40		Chen et al. (2015)
Picazuro pigeon (<i>Patagioenas picazuro</i>)	Brazil	Minas Gerais, São Paulo	3	1	33.3	MAT	1:5		Vitaliano et al. (2014)
Rock pigeon (<i>Columba livia</i>)	Brazil	São Paulo	126	0	0	MAT	1:5	Tg not isolated by bioassay of any of 126 pigeons	de Godoi et al. (2010)
Rock pigeon (<i>Columba livia</i>)	Brazil	São Paulo	120	1	0.8	IFA	1:20		de Sousa et al. (2010)
Rock pigeon (<i>Columba livia</i>)	Brazil	São Paulo	238	12	5.0	MAT	1:8	Tg not isolated	de Lima et al. (2011)
Rock pigeon (<i>Columba livia</i>)	China	Guangdong	275	24	8.7	MAT	1:5		Yan et al. (2011a)
Rock pigeon (<i>Columba livia</i>)	China	Several	963	104	10.8	IHA ^b	1:64	Age, gender AS	Zhang et al. (2019)

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Rock pigeon (<i>Columba livia</i>)	Egypt	Several	310	42	13.5	ELISA		IHC	Ibrahim <i>et al.</i> (2018)
Rock pigeon (<i>Columba livia</i>)	Israel	Several	495	20	4.0	MAT	1:5	Climate, population near to human settlement	Salant <i>et al.</i> (2009)
Rock pigeon (<i>Columba livia</i>)	Mexico	Durango	521	7	1.9	MAT	1:25	Location AS. Tg isolated	Alvarado-Esquivel <i>et al.</i> (2011)
Rock pigeon (<i>Columba livia</i>)	Portugal	Lisbon	1507	39	2.6	MAT ^b	1:20	Tg isolated in cell cultures seeded with 13 of 20 seropositive pigeons	Waap <i>et al.</i> (2012)
Rock pigeon (<i>Columba livia</i>)	Russia	Circus animals	28	10	35.7	ELISA ^a			Sivkova and Neprimerova (2017)
Rock pigeon (<i>Columba livia</i>)	Spain	Córdoba	142	13	9.2	MAT	1:25		Cano-Terriza <i>et al.</i> (2015)
Rock pigeon (<i>Columba livia</i>)	Taiwan	Several	62	1	1.6	MAT ^b	1:40		Chen <i>et al.</i> (2015)
Rock pigeon (<i>Columba livia</i>)	Turkey	Niğde	216	2	0.9	DT	1:16		Karatepe <i>et al.</i> (2011)
Rock pigeon (<i>Columba livia</i>)	USA	Colorado	129	5	3.9	MAT	1:25		Dubey <i>et al.</i> (2010)
Rock pigeon (<i>Columba livia</i>)	USA	Tennessee	1	1	100.0	MAT	1:32		Gerhold <i>et al.</i> (2017)
Spotted-necked dove (<i>Streptopelia chinensis</i>)	Taiwan	Several	17	5	29.4	MAT ^b	1:40		Chen <i>et al.</i> (2015)
Cuculiformes									
Coucal (<i>Centropus</i> sp.)	Senegal	Dakar	1	1	100.0	MAT	1:20		Galal <i>et al.</i> (2019)
Falconiformes									
Common kestrel (<i>Falco tinnunculus</i>)	Italy	Parma	238	62	26.1	MAT ^b	1:25	Age, year sampled, AS	Iemmi <i>et al.</i> (2020)
Common kestrel (<i>Falco tinnunculus</i>)	Spain	Several	13	4	30.7	MAT	1:25		Cabezón <i>et al.</i> (2011)
Crested caracara (<i>Caracara plancus</i>)	Brazil	Minas Gerais, São Paulo	2	1	50.0	MAT	1:5		Vitaliano <i>et al.</i> (2014)
Eurasian hobby (<i>Falco subbuteo</i>)	Italy	Northern	3	1	33.3	MAT ^b	1:40		Gazzonis <i>et al.</i> (2018)
Lesser kestrel (<i>Falco naumanni</i>)	Spain	Several	5	2	40.0	MAT	1:25		Cabezón <i>et al.</i> (2011)
Peregrine falcon (<i>Falco peregrinus</i>)	Spain	Several	4	1	25.0	MAT	1:25		Cabezón <i>et al.</i> (2011)
Yellow-headed caracara (<i>Milvago chimachima</i>)	Brazil	São Paulo	3	2	66.6	MAT	1:16		Gonçalves <i>et al.</i> (2013)
Galliformes									
Common quail (<i>Coturnix coturnix</i>)	China	3 Provinces	620	59	9.5	MAT	1:5		Cong <i>et al.</i> (2017b)
Common quail (<i>Coturnix coturnix</i>)	Mexico	Durango	7	1	14.2	MAT	1:25	Location AS	Alvarado-Esquivel <i>et al.</i> (2011)
Common quail (<i>Coturnix coturnix</i>)	Turkey	Niğde	144	0	0	DT	NS		Kiliç <i>et al.</i> (2017)
Guinea fowl (<i>Numida meleagris</i>)	Brazil	Minas Gerais	10	2	20.0	MAT	1:10	Tg isolated	Dubey <i>et al.</i> (2011a)

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Guinea fowl (<i>Numida meleagris</i>)	Brazil	Rio de Janeiro	114	14	12.3	MAT	1:16		Ferreira et al. (2013)
Guinea fowl (<i>Numida meleagris</i>)	Senegal	Dakar	13	1	7.7	MAT	1:20	Tg isolated	Galal et al. (2019)
Hawaiian goose (<i>Branta sandvicensis</i>)	USA	Hawaii	94	26	27.6	MAT	1:25	48% in Molokai, 23% in Maui, and 21% in Kauai	Work et al. (2016)
Silver pheasant (<i>Lophura nycthemera</i>)	Portugal	Zoo	6	1	16.6	MAT ^b	1:20		Tidy et al. (2017)
Rock ptarmigan (<i>Lagopus muta</i>)	Canada	Saskatoon	25	1	4.0	MAT ^b	1:40		Al-Adhami et al. (2016)
Gruiformes									
Common coot (<i>Fulica atra</i>)	Spain	Several	1	1	100.0	MAT	1:25		Cabezón et al. (2011)
Common moorhen (<i>Gallinula chloropus</i>)	Pakistan	Punjab	5	1	20.0	LAT ^b	1:64	Age, health status AS	Naveed et al. (2019)
Common moorhen (<i>Gallinula chloropus</i>)	Spain	Several	1	1	100.0	MAT	1:25		Cabezón et al. (2011)
Eurasian coot (<i>Fulica atra</i>)	Iran	Mazandaran	10	5	50.0	MAT	1:20		Amouei et al. (2018)
Grey crowned crane (<i>Balearica regulorum</i>)	Portugal	Zoo	4	0	0.0	MAT ^b	1:20		Tidy et al. (2017)
Japanese crane (<i>Grus japonensis</i>)	Portugal	Zoo	2	1	50.0	MAT ^b	1:20		Tidy et al. (2017)
Rail (<i>Rallus</i> sp.)	Senegal	Dakar	2	1	50.0	MAT	1:20		Galal et al. (2019)
Otidiformes									
Great bustard (<i>Otis tarda</i>)	Spain	Several	7	1	14.2	MAT	1:25		Cabezón et al. (2011)
Passeriformes									
Ashy-throated casinornis (<i>Casiornis fuscus</i>)	Brazil	Bahia	1	1	100.0	MAT	1:25		Andrade et al. (2016)
Black drongo (<i>Dicrurus macrocercus</i>)	Pakistan	Punjab	9	1	11.1	LAT ^b	1:64	Age, health status AS.	Naveed et al. (2019)
Buff-browed foliage-gleaner (<i>Syndactyla rufosuperciliata</i>)	Brazil	São Paulo	7	4	57.1	MAT	1:5		Gennari et al. (2014)
Common babbler (<i>Turdoides caudatus</i>)	Pakistan	Punjab	2	1	50.0	LAT ^b	1:64	Age, health status AS.	Naveed et al. (2019)
Common myna (<i>Acridotheres tristis</i>)	Pakistan	Punjab	11	2	18.1	LAT ^b	1:64	Age, health status AS	Naveed et al. (2019)
Common raven (<i>Corvus corax</i>)	Spain	Northeast	113	91	80.5	MAT	1:25	Year sampled AS	Molina-López et al. (2012)
Creamy-bellied thrush (<i>Turdus amaurochalinus</i>)	Brazil	São Paulo	2	1	50.0	MAT	1:5		Gennari et al. (2014)
Curve-billed thrasher (<i>Toxostoma curvirostre</i>)	Mexico	Durango	2	1	50.0	MAT	1:25		Alvarado-Esquivel et al. (2011)
European goldfinch (<i>Carduelis carduelis</i>)	Spain	Several	1	1	100.0	MAT	1:25		Cabezón et al. (2011)
European starling (<i>Sturnus vulgaris</i>)	USA	Colorado	27	4	15.0	MAT	1:50		Dubey et al. (2010)
Formosan blue magpie (<i>Urocissa caerulea</i>)	Taiwan	Several	16	6	37.5	MAT ^b	1:40		Chen et al. (2015)
Golden-crowned warbler (<i>Basileuterus culicivorus</i>)	Brazil	São Paulo	12	2	16.7	MAT	1:5		Gennari et al. (2014)
Great-tailed grackle (<i>Quiscalus mexicanus</i>)	Mexico	Durango	18	5	27.8	MAT	1:25	Location AS	

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
									Alvarado-Esquivel <i>et al.</i> (2011)
Hooded crow (<i>Corvus cornix</i>)	Israel	6 areas	101	48	47.5	MAT	1:25	Geographic location AS	Salant <i>et al.</i> (2013)
Hooded crow (<i>Corvus cornix</i>)	Italy	Pisa	120	4	3.3	MAT ^b	1.25		Mancianti <i>et al.</i> (2020)
House crow (<i>Corvus splendens</i>)	Israel	Eilat	16	2	12.5	MAT	1:25		Salant <i>et al.</i> (2013)
House crow (<i>Corvus splendens</i>)	Pakistan	Punjab	17	6	35.3	LAT ^b	1:64	Age, health status AS	Naveed <i>et al.</i> (2019)
House sparrow (<i>Passer domesticus</i>)	Brazil	Bahia, Pernambuco	293	3	1.0	IHA	1:32	Tg DNA	Gondim <i>et al.</i> (2010)
House sparrow (<i>Passer domesticus</i>)	Brazil	Pernambuco	151	91	60.3	IHA ^a	1:16	Tg DNA	Vilela <i>et al.</i> (2011)
House sparrow (<i>Passer domesticus</i>)	China	Lanzhou	313	39	12.4	MAT	1:5	Tg DNA in 11 tissues of 39 seropositive	Cong <i>et al.</i> (2013)
House sparrow (<i>Passer domesticus</i>)	Pakistan	Punjab	9	1	11.1	LAT ^b	1:64	Age, health status AS	Naveed <i>et al.</i> (2019)
Indian silverbill (<i>Lonchura malabarica</i>)	Pakistan	Punjab	4	2	50.0	LAT ^b	1:64	Age, health status AS	Naveed <i>et al.</i> (2019)
Jackdaw crow (<i>Corvus monedula</i>)	Israel	Ganei Yavne	5	2	40.0	MAT	1:25	–	Salant <i>et al.</i> (2013)
Java sparrows (<i>Lonchura oryzivora</i>)	China	Beijing, Shangqiu	350	120	34.3	MAT	1:20	Gender, colour AS	Huang <i>et al.</i> (2019)
Jungle babbler (<i>Turdoides striatus</i>)	Pakistan	Punjab	6	1	16.6	LAT ^b	1:64	Age, health status AS	Naveed <i>et al.</i> (2019)
Lesser woodcreeper (<i>Xiphorhynchus fuscus</i>)	Brazil	São Paulo	2	1	50.0	MAT	1:5		Gennari <i>et al.</i> (2014)
Magpie (<i>Pica pica</i>)	Italy	Pisa	651	41	6.2	MAT ^b	1:40		Mancianti <i>et al.</i> (2020)
Magpie (<i>Pica pica</i>)	Taiwan	Several	22	10	45.5	MAT ^b	1:40		Chen <i>et al.</i> (2015)
Pale-breasted thrush (<i>Turdus leucomelas</i>)	Brazil	São Paulo	1	1	100.0	MAT	1:5		Gennari <i>et al.</i> (2014)
Pied crow (<i>Corvus albus</i>)	Senegal	Dakar	2	2	100.0	MAT	1:20		Galal <i>et al.</i> (2019)
Plain antvireo (<i>Dysithamnus mentalis</i>)	Brazil	São Paulo	11	5	45.4	MAT	1:5		Gennari <i>et al.</i> (2014)
Red-crowned ant-tanager (<i>Habia rubica</i>)	Brazil	São Paulo	13	8	61.5	MAT	1:5		Gennari <i>et al.</i> (2014)
Red vented bulbul (<i>Pycnonotus cafer</i>)	Pakistan	Punjab	11	3	27.2	LAT ^b	1:64	Age, health AS	Naveed <i>et al.</i> (2019)
Ruby-crowned tanager (<i>Tachyphonus coronatus</i>)	Brazil	São Paulo	6	1	16.7	MAT	1:5		Gennari <i>et al.</i> (2014)
Rufous-bellied thrush (<i>Turdus rufiventris</i>)	Brazil	São Paulo	18	15	83.3	MAT	1:5		Gennari <i>et al.</i> (2014)
Rufous-breasted leaftosser (<i>Sclerurus scansor</i>)	Brazil	São Paulo	1	1	100.0	MAT	1:5		Gennari <i>et al.</i> (2014)
Rufous gnateater (<i>Conopophaga lineata</i>)	Brazil	São Paulo	15	6	40.0	MAT	1:5		Gennari <i>et al.</i> (2014)
Squamate antbird (<i>Myrmeciza squamosa</i>)	Brazil	São Paulo	4	3	75.0	MAT	1:5		Gennari <i>et al.</i> (2014)
White-browed warbler (<i>Basileuterus leucoblepharus</i>)	Brazil	São Paulo	5	1	20.0	MAT	1:5		Gennari <i>et al.</i> (2014)
White-eyed foliage-gleaner (<i>Automolus leucophthalmus</i>)	Brazil	São Paulo	14	3	21.4	MAT	1:5		Gennari <i>et al.</i> (2014)
White-lined tanager (<i>Tachyphonus rufus</i>)	Brazil	Bahia	16	1	6.2	MAT	1:25		Andrade <i>et al.</i> (2016)

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
White-necked thrush (<i>Turdus albicollis</i>)	Brazil	São Paulo	5	5	100.0	MAT	1:5		Gennari et al. (2014)
White-shouldered fire-eye (<i>Pyriglena leucoptera</i>)	Brazil	São Paulo	11	4	36.4	MAT	1:5		Gennari et al. (2014)
White-spotted woodpecker (<i>Veniliornis spilogaster</i>)	Brazil	São Paulo	1	1	100.0	MAT	1:5		Gennari et al. (2014)
White-throated woodcreeper (<i>Xiphocolaptes albicollis</i>)	Brazil	São Paulo	2	1	50.0	MAT	1:5		Gennari et al. (2014)
White-vented myna (<i>Acridotheres javanicus</i>)	Taiwan	Several	1	1	100.0	MAT ^b	1:40		Chen et al. (2015)
White wagtail (<i>Motacilla alba</i>)	Pakistan	Punjab	11	1	9.0	LAT ^b	1:64	Age, health AS	Naveed et al. (2019)
Pelecaniformes									
Black-crowned night heron (<i>Nycticorax nycticorax</i>)	Taiwan	Several	2	2	100.0	MAT ^b	1:40		Chen et al. (2015)
Buff-necked ibis (<i>Theristicus caudatus</i>)	Brazil	Minas Gerais, São Paulo	2	2	100.0	MAT	1:5		Vitaliano et al. (2014)
Cattle egret (<i>Bubulcus ibis</i>)	Brazil	Fernando de Noronha	197	157	79.7	MAT	1:5		Costa et al. (2012)
Cattle egret (<i>Bubulcus ibis</i>)	Taiwan	Several	7	4	57.1	MAT ^b	1:40		Chen et al. (2015)
Eurasian spoonbill (<i>Platalea leucorodia</i>)	Spain	Several	81	5	6.2	MAT	1:25		Cabezón et al. (2011)
Grey heron (<i>Ardea cinerea</i>)	Spain	Several	5	3	60.0	MAT	1:25		Cabezón et al. (2011)
Malay night heron (<i>Gorsachius melanolophus</i>)	Taiwan	Several	21	5	23.8	MAT ^b	1:40		Chen et al. (2015)
Purple heron (<i>Ardea purpurea</i>)	Spain	Several	4	0	0	MAT	1:25		Cabezón et al. (2011)
Roseate spoonbill (<i>Platalea ajaja</i>)	Czech Republic	Zoos	6	1	16.6	LAT ^a	–		Bártová et al. (2018)
Phaethontiformes									
Red-billed tropic bird (<i>Phaeton aethereus</i>)	Brazil	Abrolhos Archipelago	25	7	28.0	MAT	1:5		Gennari et al. (2016b)
Phoenicopteriformes									
Lesser flamingo (<i>Phoenicopterus minor</i>)	Portugal	Zoo	3	1	33.0	MAT ^b	1:20		Tidy et al. (2017)
Piciformes									
Black vulture (<i>Coragyps atratus</i>)	Brazil	São Paulo	121	16	13.2	MAT	1:5		Gennari et al. (2017)
Black vulture (<i>Coragyps atratus</i>)	USA	Alabama	8	1	12.0	MAT	1:25		Love et al. (2016)
Crescent-chested puffbird (<i>Malacoptila striata</i>)	Brazil	São Paulo	3	2	66.7	MAT	1:5		Gennari et al. (2014)
Psittiformes									
Alexandrine parakeets (<i>Psittacula eupatria</i>)	China	Beijing, Weifang	61	4	6.5	MAT	1:5	Season AS	Zhang et al. (2014)
Black-goggled tanager (<i>Lanio melanops</i>)	Brazil	São Paulo	15	5	33.3	MAT	1:5		Gennari et al. (2014)
Black-headed gull (<i>Chroicocephalus ridibundus</i>)	China	Dianchi Lake	659	131	19.9	MAT	1:5		Miao et al. (2014)
Budgerigars (<i>Melopsittacus undulatus</i>)	China	Beijing, Weifang	202	18	8.9	MAT	1:5	Season AS	Zhang et al. (2014)

(Continued)

Table 7. (Continued.)

Order, common name, (scientific name)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Cockatiels (<i>Nymphicus hollandicus</i>)	China	Beijing, Weifang	22	3	13.6	MAT	1:5	Season AS	Zhang <i>et al.</i> (2014)
Lovebirds (<i>Agapornis</i> sp.)	China	Beijing, Weifang	26	1	3.8	MAT	1:5	Season AS	Zhang <i>et al.</i> (2014)
Rainbow lorikeet (<i>Trichoglossus haematodus</i>)	Czech Republic	Zoos	4	1	25.0	LAT ^a	–		Bártová <i>et al.</i> (2018)
Red-tailed Amazon parrot (<i>Amazona brasiliensis</i>)	Brazil	Paraná	51	0	0	IFA	1:16		Sato <i>et al.</i> (2020)
Turquoise-fronted Amazon (<i>Amazona aestiva</i>)	Brazil	São Paulo	71	7	10.0	MAT	1:16		Gonçalves <i>et al.</i> (2013)
Sphenisciformes									
Galapagos penguin (<i>Spheniscus mendiculus</i>)	Republic of Ecuador	Galapagos	298	7	2.3	MAT	1:50	Region AS	Deem <i>et al.</i> (2010)
Humboldt penguin (<i>Spheniscus humboldti</i>)	Czech Republic	Zoos	4	2	50.0	LAT ^a	–		Bártová <i>et al.</i> (2018)
Magellanic penguin (<i>Spheniscus magellanicus</i>)	Brazil	Bahia, Espírito Santo, Rio de Janeiro	145	18	12.4	MAT	1:20	Tg not isolated	Acosta <i>et al.</i> (2018)
Magellanic penguin (<i>Spheniscus magellanicus</i>)	Brazil	Espírito Santo, Santa Catarina, São Paulo	100	28	28.0	MAT	1:20	Captive	Gennari <i>et al.</i> (2016a, 2016b)
Magellanic penguin (<i>Spheniscus magellanicus</i>)	Chile	Magdalena Island	132	57	43.2	MAT	1:20		Acosta <i>et al.</i> (2019)
Struthioniformes									
Greater rhea (<i>Rhea americana</i>)	Portugal	Zoo	1	1	100.0	MAT ^b	1:20		Tidy <i>et al.</i> (2017)
Suliformes									
Brown booby (<i>Sula leucogaster</i>)	Brazil	Abrolhos Archipelago	19	9	47.4	MAT	1:5		Gennari <i>et al.</i> (2016a, 2016b)
Masked booby (<i>Sula dactylatra</i>)	Brazil	Abrolhos Archipelago	23	8	34.8	MAT	1:5		Gennari <i>et al.</i> (2016a, 2016b)
Flightless cormorants (<i>Phalacrocorax harrisi</i>)	Republic of Ecuador	Galapagos	258	6	2.3	MAT	1:50	Tg isolated	Deem <i>et al.</i> (2010)
Strigiformes									
Barn owl (<i>Tyto alba</i>)	Brazil	São Paulo	4	2	50.0	MAT	1:16		Gonçalves <i>et al.</i> (2013)
Barn owl (<i>Tyto alba</i>)	Spain	Several	45	6	13.3	MAT	1:25		Cabezón <i>et al.</i> (2011)
Barred owl (<i>Strix varia</i>)	USA	Alabama	54	25	46.2	MAT	1:25	Tg isolated	Love <i>et al.</i> (2016)
Burrowing owl (<i>Athene cunicularia</i>)	Brazil	Minas Gerais, São Paulo	5	1	20.0	MAT	1:5		Vitaliano <i>et al.</i> (2014)
Collard scops owl (<i>Otus bakkamoena</i>)	Taiwan	Several	74	20	27.0	MAT ^b	1:40		Chen <i>et al.</i> (2015)

(Continued)

Table 7. (Continued.)

Order, common name, (<i>scientific name</i>)	Country	Region	No. tested	No. positive	% Positive	Test	Cut-off	Notes	Reference
Common scops-owl (<i>Otus scops</i>)	Spain	Several	4	1	25.0	MAT	1:25		Cabezón et al. (2011)
Eastern screech owl (<i>Megascops asio</i>)	USA	Alabama	16	2	12.0	MAT	1:25		Love et al. (2016)
Eurasian eagle owl (<i>Bubo bubo</i>)	Portugal	Central, northern	3	1	33.3	MAT ^b	1:20		Lopes et al. (2011)
Eurasian eagle owl (<i>Bubo bubo</i>)	Spain	Several	144	98	68.0	MAT	1:25		Cabezón et al. (2011)
Grass owl (<i>Tyto capensis</i>)	Taiwan	Several	20	3	15.0	MAT ^b	1:40		Chen et al. (2015)
Great horned owl (<i>Bubo virginianus</i>)	USA	Alabama	28	13	46.4	MAT	1:25		Love et al. (2016)
Little owl (<i>Athene noctua</i>)	Italy	Northern	17	4	23.5	MAT ^b	1:40		Gazzonis et al. (2018)
Little owl (<i>Athene noctua</i>)	Spain	Several	19	3	15.7	MAT	1:25		Cabezón et al. (2011)
Long-eared owl (<i>Asio otus</i>)	Spain	Several	9	1	11.1	MAT	1:25		Cabezón et al. (2011)
Long-eared owl (<i>Asio otus</i>)	Taiwan	Several	4	1	25.0	MAT ^b	1:40		Chen et al. (2015)
Short-eared owl (<i>Asio flammeus</i>)	Taiwan	Several	11	1	9.0	MAT ^b	1:40		Chen et al. (2015)
Spotted owl (<i>Athene brama</i>)	Pakistan	Punjab	9	3	33.3	LAT ^b	1:64	Age, health AS	Naveed et al. (2019)
Tawny owl (<i>Strix aluco</i>)	Italy	Northern	12	1	8.3	MAT ^b	1:40		Gazzonis et al. (2018)
Tawny owl (<i>Strix aluco</i>)	Portugal	Central, northern	5	1	20.0	MAT	1:20		Lopes et al. (2011)
Tawny owl (<i>Strix aluco</i>)	Spain	Several	38	5	13.6	MAT	1:25		Cabezón et al. (2011)

ELISA, enzyme-linked immunosorbent assay. Unless stated otherwise, ELISA=ELISA in-house; IFA, indirect fluorescent antibody test; IHA, indirect haemagglutination assay; LAT, latex agglutination test; MAT, modified agglutination test (Dubey and Desmonts, 1987); Tg, *Toxoplasma gondii*; AS, positive association; IHC, immunohistochemical.

^aELISA Vekto-Toxo antibodies kits (Vektor-Best, Russian Federation); IHA (ImmunoHAI-Toxoplasmosis, Wama Diagnostic, São Carlos, SP, Brazil); WAMA Diagnosis. Imuno-HAI; LAT Pastorex toxo, Bio-Rad Laboratories s.r.o., Prague, Czech Republic; MAT (New Life Diagnostic LLC, Carlsbad, CA, USA).

^bELISA ID Screen Avian Toxoplasmosis Indirect (ID VET, France); IHA (Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, China); LAT Latex agglutination test (Antec Diagnostic Products, Bridport, Dorset, UK); MAT (Toxo-Screen DA®, Biomerieux, Lyon, France). This is the same test as MAT.

and other microbial infections in general. In 2 large surveys, *T. gondii* antibodies were found in 17.7% of 175 *Gyps fulvus* in Spain (Cabezón *et al.*, 2011) but twice (39.6% of 101) more were infected in Israel (Salant *et al.*, 2013); the results are comparable because modified agglutination test (MAT) at the serum dilution of 1:25 was used in both studies. Finding of *T. gondii* DNA in tissues of 43 of 48 (89.6%) wild birds (mostly carnivorous) in Turkey suggests a very high prevalence of the parasite in local rodents (Karakavuk *et al.*, 2018). Of 281 raptors from a rehabilitation centre in the USA, 34.5% were seropositive with highest (46.0%) prevalence in Barred owl (Love *et al.*, 2016)

Viable *T. gondii* was isolated (Table 2) and DNA demonstrated (Table 3) from several species of carnivorous birds. The highest prevalence of DNA was in *Buteo buteo* from Turkey; 23 (92.0%) of 25 were infected (Table 3).

Experimental infection of crested caracara with *T. gondii*

An experiment on caracara in Brazil provided useful information. Caracaras are raptors/scavengers with wide distribution. Eight caracaras that could not be released from a rehabilitation centre were used in this experiment. They were serologically negative to *T. gondii* by IFA (cut-off 1:40). Five caracaras were fed rodent (*Calomys callosus*) infected with Me49 strain of *T. gondii*. The birds were euthanized 68 days p.i. Blood was collected for serological examination weekly or more often. Seroconversion occurred between 5 and 14 days p.i. The serological response was erratic. In 1 bird (No.6), transient antibody was observed on day 14 and then at 45 days p.i. The highest antibody titer was 1:650. Antibodies became undetectable at 68 days p.i. in 2 birds. By immunohistochemistry, *T. gondii* was detected in the hearts and muscles of all 5 caracaras and demonstrated by bioassay in the hearts of 2

Table 8. Clinical toxoplasmosis in wild birds

Host	Country	Observations	Diagnostic method	Remarks	Reference
Bar-shouldered dove (<i>Geopelia humeralis</i>)	France	A 6-year-old dove born in the zoo was found dead. Pneumonia was the main lesion and only lungs were studied histologically. Numerous tachyzoites were seen in lesions. Diagnosis was confirmed ultrastructurally	Histopathology and TEM	–	Rigoulet <i>et al.</i> (2014)
Black-footed penguin (<i>Spheniscus demersus</i>)	The Netherlands	3 captive penguins in a zoo died within a day after neurological signs were observed. All 3 had similar lesions. Liver, lungs, intestines, and brain had intralosomal tachyzoites. Diagnosis confirmed by IHC and TEM	Histopathology, IHC, TEM	–	Ploeg <i>et al.</i> (2011)
Guinea fowl (<i>Numida meleagris</i>)	Brazil	Nine of 29 guinea fowl on a farm became ill and 6 died. Two guinea fowl were necropsied. Both birds had disseminated toxoplasmosis with intralosomal <i>T. gondii</i> . One of the 2 sera tested had a MAT titer of 1:1280. Diagnosis was confirmed by IHC	Serology, histopathology, IHC	–	Vielmo <i>et al.</i> (2019)
Kereru (<i>Hemiphaga novaeseelandiae</i>)	New Zealand	2 kererus were found dead in a wildlife park. Pneumonia was the main lesion in both birds. Tachyzoites and tissue cysts were associated with lesions	Histopathology, IHC	Limited genotyping revealed atypical Type II	Howe <i>et al.</i> (2014)
Lovebird (<i>Agapornis roseicollis</i>)	Australia	A female pet bird developed ataxia 1 year after being caged. It had head tilt, tremors and was euthanized. The most important histologic findings were enlargement of spleen, encephalitis and cerebral haemorrhage	IHC, PCR	ToxoDB genotype #3 (Type II) DNA isolated from frozen brain sample	Cooper <i>et al.</i> (2015)
North Island brown kiwi (<i>Apteryx mantelli</i>)	New Zealand	A juvenile hand raised kiwi was found dead. Most important lesion was hepatitis with tissue cysts. Lesions were seen in other organs, but protozoa were not identified	Histopathology, IHC	Limited genotyping revealed atypical Type II	Howe <i>et al.</i> (2014)
North Island kaka (<i>Nestor meridionalis</i>)	New Zealand	1 kaka was found dead in wild. Most important lesion was hepatitis with tissue cysts. Lesions were seen in other organs, but protozoa were not identified	Histopathology, IHC	Limited genotyping revealed atypical Type II	Howe <i>et al.</i> (2014)
Valley quail (<i>Callipepla californica</i>)	Brazil	A captive raised quail recently acquired from another facility died suddenly with clinical signs of diarrhoea and dyspnoea. Disseminated toxoplasmosis involving most organs, many tachyzoites present	IHC, PCR, <i>T. gondii</i> DNA in many tissues	<i>T. gondii</i> DNA from kidney genotyped with 10 PCR-RFLP markers revealed ToxoDB genotype #87	Casagrande <i>et al.</i> (2015)

IHC, immunohistochemistry; PCR, polymerase chain reaction; TEM, transmission electron microscopy; PCR-RFLP, Restriction fragment length polymorphism.

birds. Three control caracaras not inoculated with *T. gondii* remained serologically negative to *T. gondii* (Vitaliano et al., 2010).

Herbivorous/insectivorous birds

Seroprevalence varied depending on the geography, host and the habitat (Table 7). For example, in the ground feeding pigeon (*Columba livia*), seroprevalences varied from 1.6 to 35.7% (Table 7). In a large sample size of pigeons, a low prevalence (1.9% of 521) was reported; the pigeons were from Durango City, Mexico that has a dry climate (Alvarado-Esquivel et al., 2011). This low prevalence is likely related to the effect of climate on oocyst survival. In another survey from cold climate, only 3.9% of pigeons from Colorado, USA were seropositive (Dubey et al., 2010). *Toxoplasma gondii* antibodies were detected in 11.8% of 35 pigeons and 9.5% of 620 quails that were destined for human consumption in China (Cong et al., 2012, 2017a). Finding of *T. gondii* DNA in the muscles of 6.4% of 390 quails indicates that the parasite was present in these birds destined for human consumption (Cong et al., 2017b). Similarly, *T. gondii* DNA was detected in 5.4% of 280 wild ducks and 3.4% of 350 common pheasants hunted for human consumption in the Czech Republic (Skorpikova et al., 2018).

Toxoplasma gondii infection in sea gulls and other scavenging birds indicates contamination of marine/lake waters with oocysts. In one study, 22.8% of 479 yellow-legged gulls (*Larus michahellis*) were seropositive to *T. gondii* (Cabezón et al., 2016). In another investigation, *T. gondii* antibodies (assessed by ELISA) were detected in 233 of 1122 freshly laid *L. michahellis* eggs (Gamble et al., 2019). The freshness of eggs was verified by immersion in water; such eggs do not float in water (Gamble et al., 2019). The occurrence of *T. gondii* antibodies in 4 other species of sea birds (*Sula* spp. and *Phaeton* spp.) from Brazil (Gennari et al., 2016b) and in 19.9% of 659 black-headed gulls (*Chroicocephalus ridibundus*) in China (Miao et al., 2014) indicates that *T. gondii* infection is common in sea birds. *Toxoplasma gondii* oocysts from feline feces can be washed into sewage and freshwater run-off and contaminate marine waters. Antibodies to *T. gondii* were detected in 8.5% of 632 mute swans (*Cygnus olor*) from the USA and viable *T. gondii* was isolated from hearts of 3 (Dubey et al., 2013). Mute swan is an invasive species present in US waters; infection in these hosts is indicative of oocyst contamination.

Reports of viable *T. gondii* and parasite DNA from tissues of wild birds are summarized in Tables 2 and 3, respectively.

Experimental infection of pigeons with *T. gondii*

Little information is available concerning the efficacy of different diagnostic methods for the detection of *T. gondii* infection in wild birds. A diagnostically useful experimental study was conducted in pigeons in Brazil (de Godoi et al., 2010). Sixteen seronegative pigeons (*C. livia*) were inoculated orally with 50 oocysts of the *T. gondii* VEG strain and divided into 4 groups of 4 pigeons each and euthanized at 15, 30, 45 and 60 days p.i. One pigeon died of toxoplasmosis on day 23 p.i. and tissue cysts were found in its brain; other pigeons remained healthy. All pigeons were seronegative by the MAT (cut-off 1:5) and IFA (cut-off 1:4) before feeding oocysts and developed antibody titers of more than 1:4000 by both MAT and IFA. Tissues (brain, heart, muscle) were tested by bioassay in mice and by PCR. Viable *T. gondii* was isolated from 5 of 12 pigeons and DNA was detected by nested PCR in tissues of 7 of 12 pigeons. By serology and bioassay, none of the 160 naturally exposed pigeons were positive for *T. gondii* providing further evidence of the validity of serology (de Godoi et al., 2010).

Migratory birds

Toxoplasma gondii infections in migratory birds are of epidemiological significance because the parasite can be transported with the host and the introduction of *T. gondii* in new geographic locations can disturb the equilibrium (Gennari et al., 2014). For example, ToxoDB genotype #9 (Chinese 1) occurs mainly in China but has been occasionally found in other countries, including the USA and Mexico. Whether migratory birds could have transported the parasite is a possibility. Antibodies to *T. gondii* were detected in Magellanic penguins in Chile and these birds migrate throughout South American coastline (Acosta et al., 2018, 2019).

A study of migratory and non-migratory geese revealed interesting results (Sandström et al., 2013). A total of 2675 birds, both adults and juveniles, of 4 goose species (*Anser anser*, $n = 266$; *A. brachyrhynchus*, $n = 787$; *Branta canadensis*, $n = 79$; *B. leucopsis*, $n = 1543$) at Arctic brood-rearing areas in Russia and on Svalbard, and temperate wintering grounds in the Netherlands and Denmark (migratory populations) as well as temperate brood-rearing grounds (the Netherlands, non-migratory

Table 9. *Toxoplasma gondii* genotypes based on DNA directly from host tissue

Country	Host	No.	10 PCR-RFLP markers ToxoDB genotype	Reference
China	<i>Alauda gulgula</i>	2	#3	Cong et al. (2014)
China	<i>Anas formosa</i>	2	#9	Zhang et al. (2015)
China	<i>Anas</i> sp.	1	#9	Zou et al. (2017)
China	<i>Carduelis spinus</i>	2	#3	Cong et al. (2014)
China	<i>Coturnix coturnix</i>	5	#9	Cong et al. (2017a, 2017b)
China	<i>Passer domesticus</i>	1	#3	Huang et al. (2012)
China	<i>Passer domesticus</i>	4	#3 in 3 and possible new type in 1	Cong et al. (2013)
China	<i>Phasianus colchicus</i>	1	#3	Huang et al. (2012)
China	<i>Passer montanus</i>	1	#10	Huang et al. (2012)
China	<i>Passer montanus</i>	3	#10	Liu et al. (2019)
Israel	<i>Corvus cornix</i>	1	#1	Salant et al. (2013)
South Africa	<i>Streptopelia semitorquata</i>	1	#1 or #3 (15 microsatellites)	Lukášová et al. (2018)
USA	<i>Branta sandvicensis</i>	4	#261 in 3, #262 in 1	Work et al. (2016)

Table 10. Distribution of PCR-RFLP (ToxoDB) *Toxoplasma gondii* genotypes from wild birds from different continents/countries

Continent/country	Total typed	Classic types										Others
		I ToxoDB #10	II ToxoDB #1 or #3	III ToxoDB#2	#4	#5	#6	#9	#11	#20		
Africa												
Senegal	15		5	1			3			6		Galal <i>et al.</i> (2019)
Asia												
China	(22) ^a	4	9					8			1 Possible new type	See Table 9
Israel	(1) ^a		1									Salant <i>et al.</i> (2013)
Oceania												
Australia	1		1									Cooper <i>et al.</i> (2015)
Europe												
France	1		1									Aubert <i>et al.</i> (2010)
Turkey	9		7	2								Karakavuk <i>et al.</i> (2018)
Central America												
Costa Rica	1										1 (#52)	Dubey <i>et al.</i> (2009), Shwab <i>et al.</i> (2014)
South America												
Brazil	19		5				1		1		12 (#8-1, #13-2, #17-1, #65-1, #108-1, #161-1, #182-2, #206-1, #290-2)	da Silva and Langoni (2016), de Barros <i>et al.</i> (2014), Dubey <i>et al.</i> (2011b), Rêgo <i>et al.</i> (2018), Silva <i>et al.</i> (2018a, 2018b)
North America												
Mexico	1							1				Alvarado-Esquivel <i>et al.</i> (2011), Shwab <i>et al.</i> (2014)
USA	32(4) ^a	1	6	4	2	6					13 (#15-1, #157-1, #167-1, #177-2, #216-2, # 61-3, #262-1, #266-1, #267-1)	Cerqueira-Cézar <i>et al.</i> (2019), Dubey <i>et al.</i> (2010), Dubey <i>et al.</i> (2011b), Dubey <i>et al.</i> (2013), Love <i>et al.</i> (2016), Shwab <i>et al.</i> (2014), Verma <i>et al.</i> (2016), Yu <i>et al.</i> (2013)
Grand total	102	5	35	7	2	6	4	9	1	6	27	

^aDNA from tissues. Twenty-seven of the 102 samples were DNA from bird tissues.

populations) were tested for *T. gondii* antibodies (MAT, 1:40). Only adult *B. leucopsis* were seropositive: *T. gondii* antibodies were found in 14.8% of 811 adults from Svalbard and 17.7% of 157 from Russia sampled during summer but not in any of the 456 juveniles sampled in summer (Arctic) and summer and winter in the Netherlands. Similar results were obtained with 3 other species of goose (Sandström *et al.*, 2013). The authors concluded that geese become infected postnatally at wintering groups.

In a study of migratory birds and resident Nearctic brown lemmings (*Lemmus trimucronatus*) from Arctic Canada, *T. gondii* infections were detected only in migratory geese (Table 1) and not in resident lemmings (Elmore *et al.*, 2015).

Clinical toxoplasmosis in wild birds

Little is known of clinical toxoplasmosis among wild birds in nature. Among all avian species, most severe toxoplasmosis has been reported in canaries (*Serinus canaria*), Hawaiian geese (*Branta sandvicensis*) and Hawaiian crows (*Corvus hawaiiensis*); reports between 1988 and 2009 were summarized previously (Dubey, 2010). In the past decade, few cases of fatal toxoplasmosis were documented in captive birds or those from National Parks (Table 8).

Hawaiian crows (*C. hawaiiensis*) and Nene goose (*B. sandvicensis*) are endangered native species in Hawaii. Nene is the largest extant terrestrial bird in Hawaiian Island and the official state bird (Work *et al.*, 2015). Of 300 Nene examined at necropsy, inflammatory conditions were found in 69 and 16.0% of these were thought to be toxoplasmosis (Work *et al.*, 2015). Thus, 11 geese died of toxoplasmosis. The presence of cats in colonies near the native bird sites is thought to be a source of *T. gondii* oocysts for the birds (Lepczyk *et al.*, 2020).

Two episodes of clinical toxoplasmosis were reported in guinea fowl in the USA and Brazil. An owner in Mississippi, USA lost 7 of 20 backyard guinea fowls. Birds were lethargic before death. Two dead birds were necropsied. Severe lesions of multifocal necrosis, fibrin exudation and inflammation of spleen, lung, heart and bone marrow were seen microscopically in 1 and mild lesions in the other guinea fowl. *Toxoplasma gondii* was identified histologically in tissues of both birds and the diagnosis was confirmed by PCR (Jones *et al.*, 2012).

The Brazilian outbreak of clinical toxoplasmosis was reported in guinea fowl on a chicken farm; the farm had 47 chickens (*G. domesticus*) and 29 guinea fowl (Vielmo *et al.*, 2019). Of these 76 birds, 22 (13 chickens and 9 guinea fowl) had clinical signs and 15 (9 chickens, 6 guinea fowl) died. Two guinea fowl were examined at necropsy and both had toxoplasmosis (Table 8).

Genetic diversity of *T. gondii* isolates

PCR-RFLP genetic data based on extraction of DNA from host tissue are summarized in Table 9 and from the live tachyzoites in Table 10. A total of 102 samples from birds were genotyped in this summary (Table 10), including 75 from viable *T. gondii* isolates (Table 2) and 27 from DNA extracted from tissues of birds (Table 10). Overall, genotype distribution follows the global patterns recognized previously (Shwab *et al.*, 2014; Su and Dubey, 2020), with ToxoDB genotypes #1 and #3 (collectively known as Type II), and genotype #2 (known as Type III) being dominant in Africa and Europe. Most genotypes identified in the Americas were diverse and different from those in the Old World. Of interest is the predominance of ToxoDB genotype #9 (Chinese 1) in China and its rare occurrence in Mexico (Alvarado-Esquivel *et al.*, 2011; Schwab *et al.*, 2014). Type I isolates (ToxoDB genotype #10) are considered rare worldwide. It was detected in a hunted turkey in 4 tree sparrows in China (Table 3). As this genotype

is highly virulent to mice and relatively easy to isolate by bioassay, future study to obtain isolates for genotyping is needed to confirm the findings. Also, of interest is the finding of ToxoDB genotypes #4 and #5, together known as Type 12, in North America but their rare frequency from the rest of the world.

Conclusions

Here, we summarized seroprevalence, clinical disease, epidemiology and genetic diversity of *T. gondii* strains isolated from wild birds worldwide for the past decade. It is obvious that *T. gondii* infection in raptors is common and they are excellent sentinels to monitor *T. gondii* in rodents and small other animals. Detection of *T. gondii* antibodies in eggs offers a non-invasive sampling method. In one investigation, *T. gondii* antibodies (assessed by ELISA) were detected in 233 of 1122 freshly laid yellow-legged gulls (*L. michahellis*) eggs (Gamble *et al.*, 2019). Finding antibodies in sea gulls indicates contamination of fresh and marine waters with *T. gondii* oocysts. In general, *T. gondii* infection in herbivorous birds is a good measure of oocyst contamination in the environment. Genetic studies revealed low genetic diversity in Europe, Asia, Africa and the USA, but higher diversity of *T. gondii* in South America. A study of migratory and non-migratory geese at Arctic brood-rearing areas in Russia and on Svalbard, and temperate wintering grounds in the Netherlands and Denmark (migratory populations) revealed that geese become infected postnatally at wintering groups (Sandström *et al.*, 2013).

Acknowledgements. This research was supported in part by an appointment of Camila K. Cerqueira-Cézar and Fernando H. A. Murata to the Agricultural Research Service (ARS) Research Participation administered by the Oak Ridge Institute for Science and Education (ORISE) through an inter-agency agreement between the US Department of Energy (DOE) and the US Department of Agriculture (USDA). ORISE was managed by ORAU under DOE contract number DE-SC 0014664. All opinions expressed in this paper were the authors' and did not necessarily reflect the policies and views of USDA, ARS, DOE or ORAU/ORISE. We thank Dr Yurong Yang for help with Chinese references.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of interest. None.

Ethical standards. Not applicable.

References

- Abdoli A, Dalimi A, Soltanghorae H and Ghaffarifar F (2016) Molecular detection of *Toxoplasma gondii* in house sparrow (*Passer domesticus*) by LAMP and PCR methods in Tehran, Iran. *Journal of Parasitic Diseases* **40**, 1317–1321.
- Abdoli A, Arbabi M, Pirestani M, Mirzaghavami M, Ghaffarifar F, Dalimi A and Sadraei J (2018) Molecular assessment of *Neospora caninum* and *Toxoplasma gondii* in hooded crows (*Corvus cornix*) in Tehran, Iran. *Comparative Immunology, Microbiology and Infectious Diseases* **57**, 69–73.
- AbouLaila M, El-Bahy N, Hilali M, Yokoyama N and Igarashi I (2011) Serodiagnosis of *Toxoplasma gondii* in ducks from Behere Governorate, Egypt. *Journal of Protozoology Research* **21**, 45–49.
- Acosta ICL, Soares RM, Mayorga LFSP, Alves BF, Soares HS and Gennari SM (2018) Occurrence of tissue cyst forming coccidia in Magellanic penguins (*Spheniscus magellanicus*) rescued on the coast of Brazil. *PLoS ONE* **13**, e0209007.
- Acosta ICL, Souza-Filho AF, Muñoz-Leal S, Soares HS, Heinemann MB, Moreno L, González-Acuña D and Gennari SM (2019) Evaluation of antibodies against *Toxoplasma gondii* and *Leptospira* spp. in Magellanic penguins (*Spheniscus magellanicus*) on Magdalena Island, Chile. *Veterinary Parasitology: Regional Studies and Reports* **16**, 100282.

- Al-Adhami BH, Simard M, Hernández-Ortiz A, Boireau C and Gajadhar AA (2016) Development and evaluation of a modified agglutination test for diagnosis of *Toxoplasma* infection using tachyzoites cultivated in cell culture. *Food and Waterborne Parasitology* **2**, 15–21.
- Alkhaled MJA, Yakoob AY and Al-Hamadani AHU (2012) An investigation of toxoplasmosis in free range chickens, industrial chickens and duck in mid Euphrates area of Iraq. *Al-Qadisiya Journal of Veterinary Medicine Sciences* **11**, 17–24.
- Almeida AB, Krindges MM, de Barros LD, Garcia JL, Camillo G, Vogel FSF, Araujo DN, Stefani LM and da Silva AS (2013) Occurrence of antibodies to *Toxoplasma gondii* in rheas (*Rhea americana*) and ostriches (*Struthio camelus*) from farms of different Brazilian regions. *Revista Brasileira de Parasitologia Veterinária* **22**, 437–439.
- Alvarado-Esquivel C, Rajendran C, Ferreira LR, Kwok OCH, Choudhary S, Alvarado-Esquivel D, Rodríguez-Peña S, Villena I and Dubey JP (2011) Prevalence of *Toxoplasma gondii* infection in wild birds in Durango, Mexico. *Journal of Parasitology* **97**, 809–812.
- Ammar S, Purple K and Gerhold R (2020) *Toxoplasma gondii* prevalence in hunter-killed mourning doves (*Zenaida macroura*) and rock pigeons (*Columba livia*) from east Tennessee. *Journal of Wildlife Diseases* **56**, 479–481.
- Amouei A, Sharif M, Hosseini SA, Sarvi S, Mizani A, Salehi S, Gholami S, Jafar-Ramaji T and Daryani A (2018) Prevalence of *Toxoplasma gondii* infection in domestic and migrating birds from Mazandaran province, Northern Iran. *Avian Biology Research* **11**, 12–15.
- Andrade LHM, Lugarini C, Oliveira RAS, Silva LTR, Marvulo MFV, Garcia JE, Dubey JP and Silva JCR (2016) Ocorrência de anticorpos anti-*Toxoplasma gondii* em aves silvestres de três Unidades de Conservação Federais da Paraíba e da Bahia. *Pesquisa Veterinária Brasileira* **36**, 103–107.
- André MR, De Santi M, Luzzi MC, de Oliveira JP, Fernandes SJ, Machado RZ and Werther K (2019) Serological evidence of exposure to *Toxoplasma gondii* and *Neospora caninum* in free-ranging Orinoco goose (*Neochen jubata*) in Brazil. *Revista Brasileira de Parasitologia Veterinária* **28**, 816–820.
- Asgari Q, Sarkari B, Amerinia M, Panahi S, Mohammadpour I and Sarvestani AS (2013) *Toxoplasma* infection in farm animals: a seroepidemiological survey in Fars Province, south of Iran. *Jundishapur Journal of Microbiology* **6**, 269–272.
- Aubert D, Ajzenberg D, Richomme C, Gilot-Fromont E, Terrier ME, de Gevigney C, Game Y, Maillard D, Gibert P, Dardé ML and Villena I (2010) Molecular and biological characteristics of *Toxoplasma gondii* isolates from wildlife in France. *Veterinary Parasitology* **171**, 346–349.
- Ayinmode AB, Obebe OO and Aiki-Raji CO (2017) Detection of *Toxoplasma gondii* antibodies in farmed turkeys (*Meleagris gallopavo*). *Folia Veterinaria* **61**, 5–10.
- Bachand N, Ravel A, Leighton P, Stephen C, Ndao M, Avaré E and Jenkins E (2019) Serological and molecular detection of *Toxoplasma gondii* in terrestrial and marine wildlife harvested for food in Nunavik, Canada. *Parasites & Vectors* **12**, 155.
- Bangoura B, Zöller B, Koethe M, Ludewig M, Pott S, Fehlhaber K, Straubinger RK and Dauschies A (2013) Experimental *Toxoplasma gondii* oocysts infections in turkeys (*Meleagris gallopavo*). *Veterinary Parasitology* **196**, 272–277.
- Bártová E, Sedláčková K and Literák I (2009) Serologic survey for toxoplasmosis in domestic birds from the Czech Republic. *Avian Pathology* **38**, 317–320.
- Bártová E, Lukášová R, Vodička R, Váhala J, Pavlačík L, Budíková M and Sedláčková K (2018) Epizootological study on *Toxoplasma gondii* in zoo animals in the Czech Republic. *Acta Tropica* **187**, 222–228.
- Butty ET (2009) Diagnostic study of *Toxoplasma gondii* in Turkey (*Meleagris gallopavo*) in some regions in Ninewah governorate. *Iraqi Journal of Veterinary Sciences* **23**, 57–62.
- Cabezón O, García-Bocanegra I, Molina-López R, Marco I, Blanco JM, Höfle U, Margalida A, Bach-Raich E, Darwich L, Echeverría I, Obón E, Hernández M, Lavín S, Dubey JP and Almería S (2011) Seropositivity and risk factors associated with *Toxoplasma gondii* infection in wild birds from Spain. *PLoS ONE* **6**, e29549.
- Cabezón O, Cerdà-Cuellar M, Morera V, García-Bocanegra I, González-Solís J, Napp S, Ribas MP, Blanch-Lázaro B, Fernández-Aguilar X, Antilles N, López-Soria S, Lorca-Oró C, Dubey JP and Almería S (2016) *Toxoplasma gondii* infection in seagull chicks is related to the consumption of freshwater food resources. *PLoS ONE* **11**, e0150249.
- Cano-Terriza D, Guerra R, Lecollinet S, Cerdà-Cuellar M, Cabezón O, Almería S and García-Bocanegra I (2015) Epidemiological survey of zoonotic pathogens in feral pigeons (*Columba livia* var. *domestica*) and sympatric zoo species in Southern Spain. *Comparative Immunology, Microbiology and Infectious Diseases* **43**, 22–27.
- Casagrande RA, Pena HFJ, Cabral AD, Rolim VM, de Oliveira LGS, Boabaid FM, Wouters ATB, Wouters F, Cruz CEF and Driemeier D (2015) Fatal systemic toxoplasmosis in valley quail (*Callipepla californica*). *The International Journal for Parasitology: Parasites and Wildlife* **4**, 264–267.
- Cerqueira-Cézar CK, da Silva AF, Murata FHA, Sadler M, Abbas IE, Kwok OCH, Brown JD, Casalena MJ, Blake MR, Su C and Dubey JP (2019) Isolation and genetic characterization of *Toxoplasma gondii* from tissues of wild turkeys (*Meleagris gallopavo*) in Pennsylvania. *Journal of Parasitology* **105**, 391–394.
- Chen JC, Tsai YJ and Wu YL (2015) Seroprevalence of *Toxoplasma gondii* antibodies in wild birds in Taiwan. *Research in Veterinary Science* **102**, 184–188.
- Cong W, Huang SY, Zhou DH, Xu MJ, Wu SM, Yan C, Zhao Q, Song HQ and Zhu XQ (2012) First report of *Toxoplasma gondii* infection in market-sold adult chickens, ducks and pigeons in northwest China. *Parasites & Vectors* **5**, 110.
- Cong W, Huang SY, Zhou DH, Zhang XX, Zhang NZ, Zhao Q and Zhu XQ (2013) Prevalence and genetic characterization of *Toxoplasma gondii* in house sparrows (*Passer domesticus*) in Lanzhou, China. *Korean Journal of Parasitology* **51**, 363–367.
- Cong W, Meng QF, Song HQ, Zhou DH, Huang SY, Qian AD, Su C and Zhu XQ (2014) Seroprevalence and genetic characterization of *Toxoplasma gondii* in three species of pet birds in China. *Parasites & Vectors* **7**, 152.
- Cong W, Chi WB, Sun WW, Shan XF, Kang YH, Meng QF and Qian AD (2017a) First report of *Toxoplasma gondii* infection in common quails (*Coturnix coturnix*) intended for human consumption in three provinces of northeastern China. *Vector-Borne and Zoonotic Diseases* **17**, 351–353.
- Cong W, Ju HL, Zhang XX, Meng QF, Ma JG, Qian AD and Zhu XQ (2017b) First genetic characterization of *Toxoplasma gondii* infection in common quails (*Coturnix coturnix*) intended for human consumption in China. *Infection, Genetics and Evolution* **49**, 14–16.
- Contente APA, Domingues PF and da Silva RC (2009) Prevalence of *Toxoplasma gondii* antibodies in ostriches (*Struthio camelus*) from commercial breeding facilities in the states of São Paulo, Brazil. *Brazilian Journal of Veterinary Research and Animal Science* **46**, 175–180.
- Cooper MK, Šlapeta J, Donahoe SL and Phalen DN (2015) Toxoplasmosis in a pet peach-faced lovebird (*Agapornis roseicollis*). *Korean Journal of Parasitology* **53**, 749–753.
- Costa DGC, Marvulo MFV, Silva JSA, Santana SC, Magalhães FJR, Lima Filho CDF, Ribeiro VO, Alves LC, Mota RA, Dubey JP and Silva JCR (2012) Seroprevalence of *Toxoplasma gondii* in domestic and wild animals from the Fernando de Noronha, Brazil. *Journal of Parasitology* **98**, 679–680.
- Darwich L, Cabezón O, Echeverría I, Pabón M, Marco I, Molina-López R, Alarcía-Alejos O, López-Gatius F, Lavín S and Almería S (2012) Presence of *Toxoplasma gondii* and *Neospora caninum* DNA in the brain of wild birds. *Veterinary Parasitology* **183**, 377–381.
- da Silva RC and Langoni H (2016) Risk factors and molecular typing of *Toxoplasma gondii* isolated from ostriches (*Struthio camelus*) from a Brazilian slaughterhouse. *Veterinary Parasitology* **225**, 73–80.
- de Barros LD, Taroda A, Zulpo DL, da Cunha IAL, Sammi AS, Cardim ST, Miura AC, Su C, Machado RZ, Vidotto O and Garcia JL (2014) Genetic characterization of *Toxoplasma gondii* isolates from eared doves (*Zenaida auriculata*) in Brazil. *Revista Brasileira de Parasitologia Veterinária* **23**, 443–448.
- Deem SL, Merkel J, Ballweber L, Vargas FH, Cruz MB and Parker PG (2010) Exposure to *Toxoplasma gondii* in Galapagos penguins (*Spheniscus mendiculus*) and flightless cormorants (*Phalacrocorax harrisi*) in the Galapagos Islands, Ecuador. *Journal of Wildlife Diseases* **46**, 1005–1011.
- de Godoi FSL, Nishi SM, Pena HFJ and Gennari SM (2010) *Toxoplasma gondii*: diagnosis of experimental and natural infection in pigeons (*Columba livia*) by serological, biological and molecular techniques. *Revista Brasileira de Parasitologia Veterinária* **19**, 238–243.
- de Lima VY, Langoni H, da Silva AV, Pezerico SB, de Castro APB, da Silva RC and Araújo JP (2011) *Chlamydomytila psittaci* and *Toxoplasma gondii*

- infection in pigeons (*Columba livia*) from São Paulo State, Brazil. *Veterinary Parasitology* 175, 9–14.
- de Sousa E, Berchieri AJ, Pinto AA, Machado RZ, de Carrasco AO, Marciano JA and Werther K (2010) Prevalence of *Salmonella* spp. antibodies to *Toxoplasma gondii*, and Newcastle disease virus in feral pigeons (*Columba livia*) in the city of Jaboticabal, Brazil. *Journal of Zoo and Wildlife Medicine* 41, 603–607.
- Dubey JP (2010) *Toxoplasmosis of Animals and Humans*, 2nd Edn. Boca Raton, Florida: CRC Press, pp. 1–313.
- Dubey JP and Desmots G (1987) Serological responses of equids fed *Toxoplasma gondii* oocysts. *Equine Veterinary Journal* 19, 337–339.
- Dubey JP, Velmurugan GV, Morales JA, Arguedas R and Su C (2009) Isolation of *Toxoplasma gondii* from the keel-billed toucan (*Ramphastos sulfuratus*) from Costa Rica. *Journal of Parasitology* 95, 467–468.
- Dubey JP, Felix TA and Kwok OCH (2010) Serological and parasitological prevalence of *Toxoplasma gondii* in wild birds from Colorado. *Journal of Parasitology* 96, 937–939.
- Dubey JP, Passos LMF, Rajendran C, Ferreira LR, Gennari SM and Su C (2011a) Isolation of viable *Toxoplasma gondii* from feral guinea fowl (*Numida meleagris*) and domestic rabbits (*Oryctolagus cuniculus*) from Brazil. *Journal of Parasitology* 97, 842–845.
- Dubey JP, Velmurugan GV, Rajendran C, Yabsley MJ, Thomas NJ, Beckmen KB, Sinnott D, Ruid D, Hart J, Fair PA, McFee WE, Shearn-Bochsler V, Kwok OCH, Ferreira LR, Choudhary S, Faria EB, Zhou H, Felix TA and Su C (2011b) Genetic characterisation of *Toxoplasma gondii* in wildlife from North America revealed widespread and high prevalence of the fourth clonal type. *International Journal for Parasitology* 41, 1139–1147.
- Dubey JP, Choudhary S, Kwok OCH, Ferreira LR, Oliveira S, Verma SK, Marks DR, Pedersen K, Mickley RM, Randall AR, Arsnoe D and Su C (2013) Isolation and genetic characterization of *Toxoplasma gondii* from mute swan (*Cygnus olor*) from the USA. *Veterinary Parasitology* 195, 42–46.
- Dubey JP, Pena HFJ, Cerqueira-Cézar CK, Murata FHA, Kwok OCH, Yang YR, Gennari SM and Su C (2020) Epidemiologic significance of *Toxoplasma gondii* infections in chickens (*Gallus domesticus*): the past decade. *Parasitology* 147, 1263–1289.
- El-Madawy SR and Metawea FY (2013) Serological assay and PCR for detection of *Toxoplasma gondii* infection in an ostrich farm at Ismailia Province, Egypt. *IOSR Journal of Agriculture and Veterinary Science* 2, 56–60.
- Elmore SA, Huyvaert KP, Bailey LL, Milhous J, Alisaukas RT, Gajadhar AA and Jenkins EJ (2014) *Toxoplasma gondii* exposure in Arctic-nesting geese: a multi-state occupancy framework and comparison of serological assays. *International Journal for Parasitology: Parasites and Wildlife* 3, 147–153.
- Elmore SA, Samelius G, Fernando C, Alisaukas RT and Jenkins EJ (2015) Evidence for *Toxoplasma gondii* in migratory vs nonmigratory herbivores in a terrestrial Arctic ecosystem. *Canadian Journal of Zoology* 93, 671–675.
- Feng Y, Lu Y, Wang Y, Zhang L and Yang Y (2017) *Toxoplasma gondii* and *Neospora caninum* in farm-reared ostriches (*Struthio camelus*) in China. *BMC Veterinary Research* 13, 301.
- Ferreira LC, Casartelli-Alves L, Figueiredo FB, da Silva RC, Langoni H, das Neves LB, Nicolau JL, Amendoira MRR and Menezes RC (2013) Ocorrência da infecção por *Toxoplasma gondii* Pela detecção de anticorpos em galinhas-d'angola criadas extensivamente no estado do Rio de Janeiro. *Revista Brasileira de Ciência Veterinária* 20, 140–143.
- Galal L, Sarr A, Cuny T, Brouat C, Coulibaly F, Sembène M, Diagne M, Diallo M, Sow A, Hamidović A, Plault N, Dardé ML, Ajzenberg D and Mercier A (2019) The introduction of new hosts with human trade shapes the extant distribution of *Toxoplasma gondii* lineages. *PLoS Neglected Tropical Diseases* 13, e0007435.
- Gallo SSM, Frazão-Teixeira E, Ederli NB and Oliveira FCR (2019) Prevalence of anti-*Toxoplasma gondii* antibodies in ratites from Brazil. *Journal of Parasitology* 105, 733–737.
- Gamble A, Ramos R, Parra-Torres Y, Mercier A, Galal L, Pearce-Duvel J, Villena I, Montalvo T, González-Solis J, Hammouda A, Oro D, Selmi S and Boulinier T (2019) Exposure of yellow-legged gulls to *Toxoplasma gondii* along the Western Mediterranean coasts: tales from a sentinel. *International Journal for Parasitology: Parasites and Wildlife* 8, 221–228.
- Gazzonis AL, Zanzani SA, Santoro A, Veronesi F, Olivieri E, Villa L, Lubian E, Lovati S, Bottura F, Epis S and Manfredi MT (2018) *Toxoplasma gondii* infection in raptors from Italy: seroepidemiology and risk factors analysis. *Comparative Immunology, Microbiology and Infectious Diseases* 60, 42–45.
- Gennari SM, Ogrzewalska M, Soares HS, Saraiva DG, Pinter A, Labruna MB and Dubey JP (2014) Occurrence of *Toxoplasma gondii* antibodies in birds from the Atlantic Forest, state of São Paulo, Brazil. *Veterinary Parasitology* 200, 193–197.
- Gennari SM, Niemeyer C, Catão-Dias JL, Soares HS, Acosta ICL, Dias RA, Ribeiro JD, Lassalvia C, Maracini P, Kolesnikovas CKM, Mayorga LFSP and Dubey JP (2016a) Survey of *Toxoplasma gondii* antibodies in Magellanic penguins (*Spheniscus magellanicus* Forster, 1781). *Journal of Zoo and Wildlife Medicine* 47, 364–366.
- Gennari SM, Niemeyer C, Soares HS, Musso CM, Siqueira GCC, Catão-Dias JL, Dias RA and Dubey JP (2016b) Seroprevalence of *Toxoplasma gondii* in seabirds from Abrolhos Archipelago, Brazil. *Veterinary Parasitology* 226, 50–52.
- Gennari SM, Raso TF, Guida FJV, Pena HFJ, Soares HS and Dubey JP (2017) Occurrence of antibodies to *Toxoplasma gondii* in scavenging black vultures (*Coragyps atratus*) in Brazil. *Brazilian Journal of Veterinary Research and Animal Science* 54, 197–199.
- Gerhold RW, Saraf P, Chapman A, Zou X, Hickling G, Stiver WH, Houston A, Souza M and Su C (2017) *Toxoplasma gondii* seroprevalence and genotype diversity in select wildlife species from the southeastern United States. *Parasites & Vectors* 10, 508.
- Gonçalves GAM, de Almeida SM, Camossi LG, Langoni H and Andreatti Filho RL (2013) Avaliação sorológica de *Parainfluenzavirus* Tipo 1, *Salmonella* spp., *Mycoplasma* spp. e *Toxoplasma gondii* Em aves silvestres. *Ciência Animal Brasileira* 14, 473–480.
- Gondim LSQ, Abe-Sandes K, Uzêda RS, Silva MSA, Santos SL, Mota RA, Vilela SMO and Gondim LFP (2010) *Toxoplasma gondii* and *Neospora caninum* in sparrows (*Passer domesticus*) in the Northeast of Brazil. *Veterinary Parasitology* 168, 121–124.
- Harfoush M and Tahoon AE (2010) Seroprevalence of *Toxoplasma gondii* antibodies in domestic ducks, free-range chickens, turkeys and rabbits in Kafr El-Sheikh Governorate Egypt. *Journal of the Egyptian Society of Parasitology* 40, 295–302.
- Hotop A, Buschtöns S, Bangoura B, Zöller B, Koethe M, Spekker-Bosker K, Hotop SK, Tenter AM, Däubener W, Straubinger RK and Groß U (2014) Humoral immune responses in chickens and turkeys after infection with *Toxoplasma gondii* by using recombinant antigens. *Parasitology Research* 113, 1473–1480.
- Howe L, Hunter S, Burrows E and Roe W (2014) Four cases of fatal toxoplasmosis in three species of endemic New Zealand birds. *Avian Diseases* 58, 171–175.
- Huang SY, Cong W, Zhou P, Zhou DH, Wu SM, Xu MJ, Zou FC, Song HQ and Zhu XQ (2012) First report of genotyping of *Toxoplasma gondii* isolates from wild birds in China. *Journal of Parasitology* 98, 681–682.
- Huang SY, Fan YM, Chen K, Yao QX and Yang B (2019) Seroprevalence and risk assessment of *Toxoplasma gondii* in Java sparrows (*Lonchura oryzivora*) in China. *BMC Veterinary Research* 15, 129.
- Ibrahim HM, Osman GY, Mohamed AH, Al-Selwi AGM, Nishikawa Y and Abdel-Ghaffar F (2018) *Toxoplasma gondii*: prevalence of natural infection in pigeons and ducks from middle and upper Egypt using serological, histopathological, and immunohistochemical diagnostic methods. *Veterinary Parasitology: Regional Studies and Reports* 13, 45–49.
- Iemmi T, Vismarra A, Mangia C, Zanin R, Genchi M, Lanfranchi P, Kramer LH, Formenti N and Ferrari N (2020) *Toxoplasma gondii* in the Eurasian kestrel (*Falco tinnunculus*) in northern Italy. *Parasites & Vectors* 13, 262.
- Jones KH, Wilson FD, Fitzgerald SD and Kiupel M (2012) A natural outbreak of clinical toxoplasmosis in a backyard flock of guinea fowl in Mississippi. *Avian Diseases* 56, 750–753.
- Karakavuk M, Aldemir D, Mercier A, Atalay Sahar E, Can H, Murat JB, Döndüren Ö, Can S, Özdemir HG, Degirmenci Döskaya A, Pektas B, Dardé ML, Gürüz AY and Döskaya M (2018) Prevalence of toxoplasmosis and genetic characterization of *Toxoplasma gondii* strains isolated in wild birds of prey and their relation with previously isolated strains from Turkey. *PLoS ONE* 13, e0196159.
- Karatepe M, Kiliç S, Karatepe B and Babür C (2011) Prevalence of *Toxoplasma gondii* antibodies in domestic (*Columba livia domestica*) and wild (*Columba livia livia*) pigeons in Niğde region, Turkey. *Turkiye Parazitoloji Dergisi* 35, 23–26.
- Khademvatan S, Saki J, Yousefi E and Abdizadeh R (2013) Detection and genotyping of *Toxoplasma gondii* strains isolated from birds in the south-west of Iran. *British Poultry Science* 54, 76–80.

- Kiliç S, Karatepe M, Babür C and Karatepe B (2017) Seroprevalence of *Toxoplasma gondii* in quails (*Coturnix coturnix japonica*) in Niğde Province, Turkey. *Kocatepe Veterinary Journal* **10**, 129–133, (in Turkish).
- Koethe M, Pott S, Ludewig M, Bangoura B, Zöllner B, Dausgchies A, Tenter AM, Spekker K, Bittame A, Mercier C, Fehlhaber K and Straubinger RK (2011) Prevalence of specific IgG-antibodies against *Toxoplasma gondii* in domestic turkeys determined by kinetic ELISA based on recombinant GRA7 and GRA8. *Veterinary Parasitology* **180**, 179–190.
- Koethe M, Straubinger RK, Pott S, Bangoura B, Geuthner AC, Dausgchies A and Ludewig M (2015) Quantitative detection of *Toxoplasma gondii* in tissues of experimentally infected turkeys and in retail Turkey products by magnetic-capture PCR. *Food Microbiology* **52**, 11–17.
- Konell AL, Sato AP, Stival M, Malaguini NP, dos Anjos A, Ferreira RF and Locatelli-Dittrich R (2019) Serosurvey of *Toxoplasma gondii*, *Sarcocystis* sp. and *Neospora caninum* in geese (*Anser* sp.) from urban parks and captivity. *Revista Brasileira de Parasitologia Veterinária* **28**, 221–228.
- Lélu M, Villena I, Dardé ML, Aubert D, Geers R, Dupuis E, Marnef F, Poulle ML, Gotteland C, Dumètre A and Gilot-Fromont E (2012) Quantitative estimation of the viability of *Toxoplasma gondii* oocysts in soil. *Applied and Environmental Microbiology* **78**, 5127–5132.
- Lepczyk CA, Haman KH, Sizemore GC and Farmer C (2020) Quantifying the presence of feral cat colonies and *Toxoplasma gondii* in relation to bird conservation areas on O'ahu, Hawai'i. *Conservation Science and Practice* **2**, e179.
- Li MH, Yang BT, Yin ZW, Wang W, Zhao Q and Jiang J (2020) A seroepidemiological survey of *Toxoplasma gondii* and *Chlamydia* infection in chickens, ducks, and geese in Jilin Province, Northeastern China. *Vector-Borne and Zoonotic Diseases*. doi: 10.1089/vbz.2020.2614.
- Liu MT, Jiang WX, Gui BZ, Jin YC, Yi JN, Li F, Zheng WB and Liu GH (2019) Molecular prevalence and genetic characterization of *Toxoplasma gondii* in wild birds in Hunan Province, China. *Vector-Borne and Zoonotic Diseases* **19**, 378–383.
- Lopes AP, Sargo R, Rodrigues M and Cardoso L (2011) High seroprevalence of antibodies to *Toxoplasma gondii* in wild animals from Portugal. *Parasitology Research* **108**, 1163–1169.
- Loss SR, Will T and Marra PP (2013) The impact of free-ranging domestic cats on wildlife of the United States. *Nature Communications* **4**, 1396.
- Love D, Kwok OC, Verma SK, Dubey JP and Bellah J (2016) Antibody prevalence and isolation of viable *Toxoplasma gondii* from raptors in the southeastern USA. *Journal of Wildlife Diseases* **52**, 653–656.
- Lukášová R, Kobédová K, Halajian A, Bártová E, Murat JB, Rampedi KM and Luus-Powell WJ (2018) Molecular detection of *Toxoplasma gondii* and *Neospora caninum* in birds from South Africa. *Acta Tropica* **178**, 93–96.
- Luo H, Li K, Zhang H, Gan P, Shahzad M, Wu X, Lan Y and Wang J (2017) Seroprevalence of *Toxoplasma gondii* infection in zoo and domestic animals in Jiangxi Province, China. *Parasite* **24**, 7.
- Maksimov P, Buschtöns S, Herrmann DC, Conraths FJ, Görlich K, Tenter AM, Dubey JP, Nagel-Kohl U, Thoms B, Bötcher L, Kühne M and Schares G (2011) Serological survey and risk factors for *Toxoplasma gondii* in domestic ducks and geese in Lower Saxony, Germany. *Veterinary Parasitology* **182**, 140–149.
- Maksimov P, Basso W, Zerweck J, Schutkowski M, Reimer U, Maksimov A, Conraths FJ and Schares G (2018) Analysis of *Toxoplasma gondii* clonal type-specific antibody reactions in experimentally infected turkeys and chickens. *International Journal for Parasitology* **48**, 845–856.
- Mancianti F, Nardoni S, Mugnaini L and Poli A (2013) *Toxoplasma gondii* in waterfowl: the first detection of this parasite in *Anas crecca* and *Anas clypeata* from Italy. *Journal of Parasitology* **99**, 561–563.
- Mancianti F, Terracciano G, Sorichetti C, Vecchio G, Scarselli D and Perrucci S (2020) Epidemiologic survey on *Toxoplasma gondii* and *Trichinella pseudospiralis* infection in corvids from Central Italy. *Pathogens* **9**, 336.
- Marković M, Ivočić V, Štajner T, Djokić V, Klun I, Bobić B, Nikolić A and Djurković-Djaković O (2014) Evidence for genetic diversity of *Toxoplasma gondii* in selected intermediate hosts in Serbia. *Comparative Immunology, Microbiology and Infectious Diseases* **37**, 173–179.
- Miao Q, Han JQ, Xiang X, Yuan FZ, Liu YZ, Duan G, Zhu XQ and Zou FC (2014) Prevalence of antibody to *Toxoplasma gondii* in black-headed gulls (*Chroicocephalus ridibundus*), Dianchi Lake, China. *Journal of Wildlife Diseases* **50**, 717–719.
- Mirza V, Burrows EB, Gils S, Hunter S, Gartrell BD and Howe L (2017) A retrospective survey into the presence of *Plasmodium* spp. and *Toxoplasma gondii* in archived tissue samples from New Zealand raptors: New Zealand falcons (*Falco novaeseelandiae*), Australasian harriers (*Circus approximans*) and moreporks (*Ninox novaeseelandiae*). *Parasitology Research* **116**, 2283–2289.
- Molina-López R, Cabezon O, Pabón M, Darwich L, Obón E, Lopez-Gatius F, Dubey JP and Almería S (2012) High seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in the common raven (*Corvus corax*) in the Northeast of Spain. *Research in Veterinary Science* **93**, 300–302.
- Moustakidis K, Economou V, Dovas C, Symeonidou I, Papadopoulou E and Papazahariadou M (2017) First report of *Toxoplasma gondii* in the woodcock (*Scolopax rusticola*): preliminary results. *Proceedings of the 8th International Conference on Information and Communication Technologies in Agriculture, Food and Environment (HAICTA 2017)*, Chania, Greece, 21–24 September 2017, pp. 20–27.
- Nardoni S, Rocchigiani G, Varvaro I, Altomonte I, Ceccherelli R and Mancianti F (2019) Serological and molecular investigation on *Toxoplasma gondii* infection in wild birds. *Pathogens* **8**, 58.
- Naveed A, Ali S, Ahmed H, Simsek S, Rizwan M, Kaleem I, Gondal MA, Shabbir A, Pervaiz F, Khan MA, Nadeem MS, Afzaal MS and Umar S (2019) Seroprevalence and risk factors of *Toxoplasma gondii* in wild birds of Punjab Province, Pakistan. *Journal of Wildlife Diseases* **55**, 129–135.
- Nazir MM, Ayaz MM, Ahmed AN, Maqbool A, Ashraf K, Oneeb M, Yasin G, Subhani A, Ali MA, Nazir N and Sajid MA (2018) Prevalence of *Toxoplasma gondii*, *Neospora caninum*, and *Sarcocystis* species DNA in the heart and breast muscles of rock pigeons (*Columbia livia*). *Journal of Parasitology Research* **2018**, 6264042.
- Ploeg M, Ultee T and Kik M (2011) Disseminated toxoplasmosis in black-footed penguins (*Spheniscus demersus*). *Avian Diseases* **55**, 701–703.
- Puvanesuaran VR, Noordin R and Balakrishnan V (2013) Isolation and genotyping of *Toxoplasma gondii* from free-range ducks in Malaysia. *Avian Diseases* **57**, 128–132.
- Rêgo WMF, Costa JGL, Baraviera RCA, Pinto LV, Bessa GL, Lopes REN, Silveira JAG and Vitor RWA (2018) Genetic diversity of *Toxoplasma gondii* isolates obtained from free-living wild birds rescued in Southeastern Brazil. *International Journal for Parasitology: Parasites and Wildlife* **7**, 432–438.
- Rigoulet J, Hennache A, Lagourette P, George C, Longeart L, Le Net JL and Dubey JP (2014) Toxoplasmosis in a bar-shouldered dove (*Geopelia humeralis*) from the Zoo of Clères, France. *Parasite* **21**, 62.
- Rong G, Zhou HL, Hou GY, Zhao JM, Xu TS and Guan S (2014) Seroprevalence, risk factors and genotyping of *Toxoplasma gondii* in domestic geese (*Anser domestica*) in tropical China. *Parasites & Vectors* **7**, 459.
- Sá SG, Lima DCV, Silva LTR, Pinheiro Júnior JW, Dubey JP, Silva JCR and Mota RA (2016) Seroprevalence of *Toxoplasma gondii* among turkeys on family farms in the state of Northeastern Brazil. *Acta Parasitologica* **61**, 401–405.
- Salant H, Landau DY and Baneth G (2009) A cross sectional survey of *Toxoplasma gondii* antibodies in Israeli pigeons. *Veterinary Parasitology* **165**, 145–149.
- Salant H, Hamburger J, King R and Baneth G (2013) *Toxoplasma gondii* prevalence in Israeli crows and Griffon vultures. *Veterinary Parasitology* **191**, 23–28.
- Sandström CAM, Buma AGJ, Høy BJ, Prop J, van der Jeugd H, Voslamber B, Madsen J and Loonen MJJE (2013) Latitudinal variability in the seroprevalence of antibodies against *Toxoplasma gondii* in non-migrant and Arctic migratory geese. *Veterinary Parasitology* **194**, 9–15.
- Sarkari B, Asgari Q, Bagherian N, Ashkani Esfahani S, Kalantari M, Mohammadpour I, Ashrafmansori M, Amerinia M and Sabet Sarvestani F (2014) Molecular and serological evaluation of *Toxoplasma gondii* infection in reared turkeys in Fars Province, Iran. *Jundishapur Journal of Microbiology* **7**, e11598.
- Sato AP, Vaz FF, Konell AL, Koch MO, Ferreira RF, Sipinski EAB and Dittrich RL (2020) Survey of *Toxoplasma gondii*, *Neospora caninum* and *Sarcocystis neurona* antibodies in wild red-tailed Amazon parrots (*Amazona brasiliensis*). *Revista Brasileira de Parasitologia Veterinária* **29**, e017519.
- Shwab EK, Zhu XQ, Majumdar D, Pena HFJ, Gennari SM, Dubey JP and Su C (2014) Geographical patterns of *Toxoplasma gondii* genetic diversity revealed by multilocus PCR-RFLP genotyping. *Parasitology* **141**, 453–461.
- Silva MA, Pena HFJ, Soares HS, Aizawa J, Oliveira S, Alves BF, Souza DS, Melo RPB, Gennari SM, Mota RA and Silva JCR (2018a) Isolation and genetic characterization of *Toxoplasma gondii* from free-ranging and captive birds and mammals in Pernambuco state, Brazil. *Revista Brasileira de Parasitologia Veterinária* **27**, 481–487.

- Silva LTR, Silva JSA, Lima DCV, Rolim VPM, Marvulo MFV, Silva JCR, Mota RA and Oliveira AAF (2018b) Anticorpos anti-*Toxoplasma gondii* em carcarás (*Caracara plancus*) procedentes da região aeroportuária do Recife, Pernambuco, Brasil. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* **70**, 505–510.
- Sivkova TN and Neprimerova TA (2017) Toxoplasmosis in circus animals. *GISAP: Medical Science, Pharmacology* doi: 10.18007/gisap: msp.v0i14.1663.
- Skorpikova L, Reslova N, Lorencova A, Plhal R, Drimaj J, Kamler J and Slany M (2018) Molecular detection of *Toxoplasma gondii* in feathered game intended for human consumption in the Czech Republic. *International Journal of Food Microbiology* **286**, 75–79.
- Sroka J, Wójcik-Fatla A, Szymańska J, Dutkiewicz J, Zajac V and Zwoliński J (2010) The occurrence of *Toxoplasma gondii* infection in people and animals from rural environment of Lublin region – estimate of potential role of water as a source of infection. *Annals of Agricultural and Environmental Medicine* **17**, 125–132.
- Sroka J, Karamon J, Wójcik-Fatla A, Dutkiewicz J, Biliska-Zajac E, Zajac V, Piotrowska W and Cencek T (2019) *Toxoplasma gondii* infection in selected species of free-living animals in Poland. *Annals of Agricultural and Environmental Medicine* **26**, 656–660.
- Su C and Dubey JP (2020) Isolation and genotyping of *Toxoplasma gondii* strains. *Methods in Molecular Biology* **2071**, 49–80.
- Tidy A, Figueiro S, Dubey JP, Cardoso L and Lopes AP (2017) Seroepidemiology and risk assessment of *Toxoplasma gondii* infection in captive wild birds and mammals in two zoos in the North of Portugal. *Veterinary Parasitology* **235**, 47–52.
- Turčeková L, Hurníková Z, Spišák F, Miterpáková M and Chovancová B (2014) *Toxoplasma gondii* in protected wildlife in the Tatra National Park (TANAP), Slovakia. *Annals of Agricultural and Environmental Medicine* **21**, 235–238.
- Verma SK, Calero-Bernal R, Cerqueira-Cézar CK, Kwok OCH, Dudley M, Jiang T, Su C, Hill D and Dubey JP (2016) Toxoplasmosis in geese and detection of two new atypical *Toxoplasma gondii* strains from naturally infected Canada geese (*Branta canadensis*). *Parasitology Research* **115**, 1767–1772.
- Vielmo A, Pena HFJ, Panziera W, Bianchi RM, De Lorenzo C, Oliveira S, Alves BF, Gennari SM, Pavarini SP, de Barros CSL and Driemeier D (2019) Outbreak of toxoplasmosis in a flock of domestic chickens (*Gallus gallus domesticus*) and guinea fowl (*Numida meleagris*). *Parasitology Research* **118**, 991–997.
- Vilares A, Gargaté MJ, Ferreira I, Martins S, Júlio C, Waap H, Ângelo H and Gomes JP (2014) Isolation and molecular characterization of *Toxoplasma gondii* isolated from pigeons and stray cats in Lisbon, Portugal. *Veterinary Parasitology* **205**, 506–511.
- Vilela SMO, Silva JSA, Junior JWP, Moraes ÉPBX, Saukas TN, Gondim LFP and Mota RA (2011) Sparrows (*Passer domesticus* L.) as intermediary hosts of *Toxoplasma gondii* in poultry farms from the 'agreste' region of Pernambuco, Brazil. *Pesquisa Veterinária Brasileira* **31**, 169–172.
- Vitaliano SN, Mineo TWP, André MR, Machado RZ, Mineo JR and Werther K (2010) Experimental infection of crested caracara (*Caracara plancus*) with *Toxoplasma gondii* simulating natural conditions. *Veterinary Parasitology* **172**, 71–75.
- Vitaliano SN, Soares HS, Pena HFJ, Dubey JP and Gennari SM (2014) Serological evidence of *Toxoplasma gondii* infection in wild birds and mammals from southeast Brazil. *Journal of Zoo and Wildlife Medicine* **45**, 197–199.
- Waap H, Cardoso R, Leitão A, Nunes T, Vilares A, Gargaté MJ, Meireles J, Cortes H and Ângelo H (2012) *In vitro* isolation and seroprevalence of *Toxoplasma gondii* in stray cats and pigeons in Lisbon, Portugal. *Veterinary Parasitology* **187**, 542–547.
- Wang XQ, Gao Y, Li GY, He Y, Yuan ZG, Weng YB and Lin RQ (2012) Survey of *Toxoplasma gondii* in geese from Qingyuan City, Guangdong Province. *Chinese Journal of Veterinary Medicine* **48**, 10, (in Chinese).
- Wang DW, Han XH, Mu MY, Yuan GM, Zhang GX, He JB, Yang N and Li HK (2014) Epidemiological survey of toxoplasmosis in some animals from northeastern China. *Heilongjiang Animal Science and Veterinary Medicine* **2014**, 129–131, (in Chinese).
- Work TM, Massey JG, Rideout BA, Gardiner CH, Ledig DB, Kwok OCH and Dubey JP (2000) Fatal toxoplasmosis in free-ranging endangered 'Alala from Hawaii. *Journal of Wildlife Diseases* **36**, 205–212.
- Work TM, Dagenais J, Rameyer R and Breeden R (2015) Mortality patterns in endangered Hawaiian geese (Nene; *Branta sandvicensis*). *Journal of Wildlife Diseases* **51**, 688–695.
- Work TM, Verma SK, Su C, Medeiros J, Kaiakapu T, Kwok OC and Dubey JP (2016) *Toxoplasma gondii* antibody prevalence and two new genotypes of the parasite in endangered Hawaiian geese (nene: *Branta sandvicensis*). *Journal of Wildlife Diseases* **52**, 253–257.
- Yan C, Yue CL, Yuan ZG, He Y, Yin CC, Lin RQ, Dubey JP and Zhu XQ (2009) *Toxoplasma gondii* infection in domestic ducks, free-range and caged chickens in southern China. *Veterinary Parasitology* **165**, 337–340.
- Yan C, Yue CL, Qiu SB, Li HL, Zhang H, Song HQ, Huang SY, Zou FC, Liao M and Zhu XQ (2011a) Seroprevalence of *Toxoplasma gondii* infection in domestic pigeons (*Columba livia*) in Guangdong Province of southern China. *Veterinary Parasitology* **177**, 371–373.
- Yan C, Yue CL, Zhang H, Yin CC, He Y, Yuan ZG, Lin RQ, Song HQ, Zhang KX and Zhu XQ (2011b) Serological survey of *Toxoplasma gondii* infection in the domestic goose (*Anser domestica*) in southern China. *Zoonoses and Public Health* **58**, 299–302.
- Yang N, Mu MY, Li HK, Long M and He JB (2012) Seroprevalence of *Toxoplasma gondii* infection in slaughtered chickens, ducks, and geese in Shenyang, northeastern China. *Parasites & Vectors* **5**, 237.
- Yu L, Shen J, Su C and Sundermann CA (2013) Genetic characterization of *Toxoplasma gondii* in wildlife from Alabama, USA. *Parasitology Research* **112**, 1333–1336.
- Zhang XX, Zhang NZ, Tian WP, Zhou DH, Xu YT and Zhu XQ (2014) First report of *Toxoplasma gondii* seroprevalence in pet parrots in China. *Vector-Borne and Zoonotic Diseases* **14**, 394–398.
- Zhang FK, Wang HJ, Qin SY, Wang ZD, Lou ZL, Zhu XQ and Liu Q (2015) Molecular detection and genotypic characterization of *Toxoplasma gondii* in wild waterfowls in Jilin Province, Northeastern China. *Parasitology International* **64**, 576–578.
- Zhang XX, Qin SY, Li X, Ren WX, Hou G, Zhao Q and Ni HB (2019) Seroprevalence and related factors of *Toxoplasma gondii* in pigeons intended for human consumption in northern China. *Vector-Borne and Zoonotic Diseases* **19**, 302–305.
- Zhao G, Song Z, Wang S, Hassan IA, Wang W, Cheng F and Yang X (2015) A seroepidemiological survey of *Toxoplasma gondii* infection in free-range and caged ducks in Southwest China. *Israel Journal of Veterinary Medicine* **70**, 41–45.
- Zöller B, Koethe M, Ludewig M, Pott S, Dausgchies A, Straubinger RK, Fehlhaber K and Bangoura B (2013) Tissue tropism of *Toxoplasma gondii* in turkeys (*Meleagris gallopavo*) after parenteral infection. *Parasitology Research* **112**, 1841–1847.
- Zou Y, Nie LB, Zhang NZ, Zou FC, Zhu XQ and Cong W (2017) First genetic characterization of *Toxoplasma gondii* infection in poultry meat intended for human consumption in eastern China. *Infection, Genetics and Evolution* **55**, 172–174.