

Event-Related Functional Magnetic Resonance Imaging Changes during Relational Retrieval in Normal Aging and Amnesic Mild Cognitive Impairment

Kelly S. Giovanello,^{1,2} Felipe De Brigard,^{1,3} Jaclyn Hennessey Ford,¹ Daniel I. Kaufer,⁴ James R. Burke,^{5,6} Jeffrey N. Browndyke,^{5,7} AND Kathleen A. Welsh-Bohmer^{5,6,7}

¹Department of Psychology, The University of North Carolina, Chapel Hill, North Carolina

²Biomedical Research Imaging Center, The University of North Carolina, Chapel Hill, North Carolina

³Department of Philosophy, The University of North Carolina, Chapel Hill, North Carolina

⁴Department of Neurology, The University of North Carolina, Chapel Hill, North Carolina

⁵Joseph & Kathleen Bryan Alzheimer's Disease Research Center, Duke University Medical Center, Durham, North Carolina

⁶Division of Neurology, Duke University Medical Center, Durham, North Carolina

⁷Department of Psychiatry & Behavioral Sciences, Duke University Medical Center, Durham, North Carolina

(RECEIVED August 31, 2011; FINAL REVISION April 18, 2012; ACCEPTED April 19, 2012)

Abstract

The earliest cognitive deficits observed in amnesic mild cognitive impairment (aMCI) appear to center on memory tasks that require relational memory (RM), the ability to link or integrate unrelated pieces of information. RM impairments in aMCI likely reflect neural changes in the medial temporal lobe (MTL) and posterior parietal cortex (PPC). We tested the hypothesis that individuals with aMCI, as compared to cognitively normal (CN) controls, would recruit neural regions outside of the MTL and PPC to support relational memory. To this end, we directly compared the neural underpinnings of successful relational retrieval in aMCI and CN groups, using event-related functional magnetic resonance imaging (fMRI), holding constant the stimuli and encoding task. The fMRI data showed that the CN, compared to the aMCI, group activated left precuneus, left angular gyrus, right posterior cingulate, and right parahippocampal cortex during relational retrieval, while the aMCI group, relative to the CN group, activated superior temporal gyrus and supramarginal gyrus for this comparison. Such findings indicate an early shift in the functional neural architecture of relational retrieval in aMCI, and may prove useful in future studies aimed at capitalizing on functionally intact neural regions as targets for treatment and slowing of the disease course. (*JINS*, 2012, 18, 886–897)

Keywords: Mild cognitive impairment, Aging, Memory, Functional MRI, Medial temporal lobe, Parietal lobe

INTRODUCTION

Amnesic mild cognitive impairment (aMCI) is considered to be a transitional stage between healthy aging and Alzheimer's disease (AD; Petersen et al., 1999). Individuals with aMCI exhibit an objective memory impairment, preserved general cognitive abilities, no or minimal decline in activities of daily living, and no dementia (Petersen, 2004; Winblad et al., 2004). Although the classification of aMCI does not guarantee development of dementia, as some individuals meeting the criteria later perform normally on memory testing, there is a significant risk of progressing to clinical AD (e.g., 80% over 6 years;

Petersen et al., 1999; Petersen, 2004). Identifying the cognitive tests that are most sensitive for detecting aMCI in conjunction with associated alterations in the underlying neural architecture will facilitate earlier diagnosis of AD and may prove useful in demonstrating the beneficial effects of future disease modifying treatments.

Individuals with aMCI perform poorly on tests of episodic memory, defined as the encoding and conscious retrieval of contextually-specific information, such as an event that occurred at a particular place and time (Tulving, 1983). Performance on episodic memory tasks relies both on relational memory, the ability to integrate unrelated pieces of information, as well as item memory, which provides the basis for knowing that a stimulus has occurred (Yonelinas, 2001). Several reports over the last decade suggest that the earliest cognitive deficits in aMCI center on memory tasks that

Correspondence and reprint requests to: Kelly S. Giovanello, Department of Psychology, The University of North Carolina, Campus Box 3270, Chapel Hill, NC 27713. E-mail: kgio@unc.edu

require relational memory (e.g., paired-associate learning, cued-recall, and associative recall). For example, Fowler, Saling, Conway, Semple, and Louis (2002) conducted longitudinal neuropsychological assessments in control participants, individuals with “questionable dementia,” and individuals with probable early AD and found that performance on a paired associate learning test best identified the onset of progressive memory decline in the questionably demented individuals, all of whom went on to fulfill NINCDS-ADRDA criteria for probable AD over a 2-year period. In another study, Ivanoiu et al. (2005) evaluated cued-recall performance in individuals with subjective memory complaints, MCI, and mild probable AD and showed that a cued-recall task correctly classified 88% of the MCI participants and was a good predictor of MCI and mild AD status. More recently, Troyer et al. (2008) used standardized neuropsychological tasks thought to tap relational memory processes and found that tests of associative recall were particularly sensitive to early cognitive change in aMCI. Finally, Anderson and colleagues (2008) reported age-related reductions in recollection, a mnemonic process that influences performance on relational memory tasks (Hockley and Consoli, 1999). Notably, the reductions in recollection were greater among aMCI individuals than cognitively normal subjects. Taken together, these neuropsychological findings suggest that tests of relational memory may be among the most sensitive measures for detecting cognitive changes associated with aMCI.

A prominent neural correlate of aMCI is volume loss in the medial temporal lobe (MTL), particularly the hippocampus and entorhinal cortex (Convit et al., 1997; Dickerson et al., 2001; Xu et al., 2000), with increasing atrophy in these structures from normal aging to aMCI to AD (Du et al., 2001; Penner et al., 2004). Longitudinal studies of aMCI patients have revealed that diminished baseline hippocampal and entorhinal volume is associated with an increased likelihood of progressing to clinical dementia (De Santi et al., 2001; Grundman et al., 2002; Jack et al., 1999; Kaye et al., 2005; Killiany et al., 2000). Memory decline is the primary cognitive consequence of atrophy in these MTL structures and, in general, hippocampal and entorhinal volumes correlate with performance on memory tasks (Rodrigue and Raz, 2004; Rosen et al., 2003). Several lines of research suggest that structures within the MTL, particularly the hippocampus, make a critical contribution to relational memory processing, by linking or binding the elements of a to-be-remembered episode (Eichenbaum, Yonelinas, & Ranganath, 2007). Indeed, Rajah, Kromas, Han, and Pruessner (2010) recently demonstrated that volumetric reductions in anterior hippocampus related to poorer retrieval of spatial and temporal context information with age. As such, relational memory deficits in aMCI likely reflect pathological alterations, at least in part, in the MTL.

Atrophic changes in parietal cortex (PC), particularly in the posterior region of medial parietal cortex, referred to as precuneus, have also been documented in MCI (Buckner et al., 2005). Studies of resting glucose metabolism have demonstrated

hypometabolism in the inferior parietal lobule, that progresses with the disease and correlates with mental status (e.g., Herholz et al., 2002; Minoshima et al., 1997), and is present in individuals at genetic risk for AD (Reiman et al., 1996). More recently, fMRI studies using analysis of intrinsic activity correlations have reported functional changes in parietal regions in aMCI (e.g., Celone et al., 2006; Rombouts, Goekoop, Stam, Barkhof, & Scheltens, 2005; Wang et al., 2007), with alteration of neural activity in medial and lateral parietal regions directly related to loss of MTL functional integrity, notably during tasks involving relational encoding (e.g., learning face-name pairs; Celone et al., 2006).

Although prior fMRI studies have assessed neural activity during the encoding of relational information (e.g., Celone et al., 2006; Dickerson et al., 2005; Petrella, Prince, Wang, Hellegers, & Doraiswamy, 2007), or during retrieval of item information (e.g., Lenzi et al., 2011; Machulda et al., 2009; Trivedi, et al., 2008), to our knowledge no study has directly compared retrieval of item and relational information in aMCI. Such a comparison offers three advantages. First, successful retrieval of relational information requires recovery of more specific, detailed information than does successful item retrieval which can be based on a general sense of prior occurrence (i.e., familiarity). As such, a direct comparison of the neural underpinnings of item and relational memory in aMCI provides the opportunity to pinpoint whether, and in what manner, alterations in the functional neural architecture are present in aMCI based on the *specificity* of information retrieved. Although the term “specificity” has been used broadly (see Schacter, Gallo, & Kensinger, 2007), here we use the term “specificity” to denote differences in the degree to which specific, detailed information is required at retrieval for successful performance (e.g., retrieval of a novel link formed between unrelated words). Second, a comparison of item and relational memory in aMCI, coupled with event-based fMRI, allows for the analysis of *correct* memory trials only, eliminating potential confounds due to differences in retrieval success across levels of specificity. Finally, this aspect of the current design, in turn, allowed us to identify those regions recruited by aMCI participants during successful relational retrieval. Given the importance of capitalizing on functionally intact neural regions as targets for treatment and slowing of the disease course (Dickerson & Sperling, 2009; Golde, 2006), elucidating the neural regions that may contribute to relational memory is a valuable research endeavor.

Here, we used a paradigm in which encoding stimuli and encoding tasks were held constant, and then subsequently compared successful retrieval of item information and relational information in aMCI individuals and cognitively normal, age- and education-matched participants. In light of prior behavioral reports documenting relational memory deficits in aMCI, as well the neural alterations observed in the MTL and PC in this group, we had three hypotheses. First, we hypothesized that the aMCI group would perform worse on the relational memory task as compared to the CN group. Second, we hypothesized that the aMCI group, as compared to the CTL group, would not

recruit regions traditionally shown to be engaged during relational memory (i.e., the MTL and PC). Rather, under conditions of successful relational retrieval, we hypothesized that aMCI participants, as compared to the CN group, would recruit regions outside of the MTL and PC, particularly lateral temporal regions during relational retrieval, due to the nature of the encoding task (i.e., generation of encoding sentences).

METHODS

Participants

Twelve cognitively normal (CN) and 12 individuals with aMCI were recruited for this study through the Bryan Alzheimer's Disease Research Center (ADRC) at Duke Medical Center and the University of North Carolina at Chapel Hill (UNC-CH) Memory Disorders Clinic. This study was approved by the UNC-CH and Duke Medical Center Institutional Review Boards. Informed consent was obtained from each participant. All subjects were paid for their participation.

The classification of CN and aMCI was based on the input of two sources: the neurologist's (J.R.B. or D.I.K.) clinical opinion based on their interview and examination of the participants and cognitive test results interpreted by the neuropsychologist (see below).

MCI Subjects

Amnesic MCI was defined with the following criteria: (1) memory complaint corroborated by an informant, (2) not normal for age (as determined by the neurologists' and neuropsychologists' clinical judgment), (3) not demented, (4) mild cognitive impairment, (5) essentially normal functional activities, and (6) memory was the only cognitive domain mildly impaired relative to normal comparison.

Subjects

Cognitively normal elderly met the following criteria: (1) no cognitive complaints, (2) no active neurological or psychiatric illness, (3) independently functioning community dwellers, (4) normal neurological and neuropsychological exam, and (5) not taking any medications in doses that would impact cognitive performance.

Exclusion Criteria

Exclusion criteria were: (1) diagnosis other than CN or aMCI, (2) left-handedness, (3) non-native English speaker, (3) dementia, (4) medical contra-indications for MRI, (5) structural abnormalities (e.g., infarctions), and (6) concurrent illnesses interfering with cognitive function other than aMCI (i.e., heart/liver/renal failure, psychiatric disorders, and substance abuse).

Neuropsychological Testing

Neuropsychological testing was completed within 5 months of participation in the study. The battery used is one that has been used in longitudinal studies at the Bryan ADRC

(e.g., Tschanz et al., 2006) and includes all the requisite measures of the National Alzheimer's Disease Coordinating Center (NACC; see Hayden et al., 2011). Narrative memory was assessed by performance on Story A from the Wechsler Memory Scale – Revised (Wechsler, 1987) Logical Memory Immediate and Delay subtests, as well as by scores on subtests of the Consortium to Establish a Registry for Alzheimer's Disease (CERAD) list learning task (i.e., word list learning, recall intrusions, perseverations, recall, recognition, constructional praxis recall, and constructional praxis recognition; Morris et al., 1989). Language tests measured object naming (30 item version of the Boston Naming Test, Kaplan, Goodglass, & Weintraub, 1983), phonemic fluency (Controlled Oral Word Association Test; COWAT), and category fluency (animals, Morris et al., 1989 and vegetables). Attention and executive tests include the Trail-making test Part A and B (Spreen & Strauss, 1991) and both the Digit Span and Digit Symbol subtests from the Wechsler Adult Intelligence Scale – Revised (Wechsler, 1981). Additional tests included the AD8 (a screening test that assesses memory, orientation, executive functioning, and interest in activities; 2005, Washington University, St. Louis, MO), the Shipley Vocabulary Test (as an estimate of premorbid function and intelligence; Shipley, 1967), the Mini-mental Status Exam (MMSE; Folstein, Folstein, & McHugh, 1975), the Geriatric Depression Scale, and the Hachinski Ischaemia Questionnaire (Hachinski et al., 1975).

Functional MRI Tasks

Stimuli were 288 one- to three-syllable unrelated nouns (M Freq = 56.3; SD = 63.5). Following extensive practice outside the scanner, participants received four study/retrieval runs. During the unscanned study phase, participants studied two words (e.g., HIGHWAY CAFÉ) on each of a total of 96 trials, with 24 trials per list. Participants were instructed to covertly create a short sentence that incorporated the two words, ensuring that the word on the left side of the screen was used in the sentence before the word on the right side of the screen (e.g., "The highway was in front of the café"). All participants indicated *via* button press that they had created and encoded a sentence for each trial. During the scanned retrieval phase, which started immediately following the study phase, participants performed one of two recognition tasks: Relational or Item. In the *Relational* task, participants saw pairs of words that were previously seen together (Intact Pair – IP), pairs of words that were previously seen but not together (Recombined Pair – RP), and pairs of novel words that were not seen previously (New Pair – NP). Participants were asked to indicate whether the two words were previously seen together; they were instructed to press "1" if they were or "2" if they were not seen together previously. In the *Item* task, participants saw pairs of words previously seen but not together (Recombined items – RI), pairs consisting of one old word and one new word (Old/New Items – ONI), and pairs consisting of two new words (New Items – NI). Participants were asked to indicate whether both words of a pair were previously seen; they were instructed to press "1"




Fig. 1. Event-related task design with alternating blocked task periods of relational retrieval (“together previously?”) and item retrieval (“both old?”). There were four study/retrieval phases. Imaging data was acquired during the retrieval phase only. IP = intact pair; RP = rearranged pair; NP = new pair; RI = rearranged items; OI = old/new items; and NI = new items.

if they were both seen previously at all or “2” if they were not. Four task blocks alternated between relational memory and item memory (Figure 1). Each block consisted of nine trials drawn from each of the task-appropriate experimental conditions types, as well as three control trials during which participants viewed ampersands and number signs, and were instructed to indicate on which side of the screen the ampersands had appeared. Control trials were used to introduce jitter during each scanner run. Trials were randomized within each task block. Starting task and stimulus conditions were counterbalanced across participants.

MR Image Acquisition

All imaging data were acquired at the UNC-CH’s Biomedical Research Imaging Center on a Siemens 3 Tesla Allegra head-only imaging system equipped for echo planar imaging (EPI; Siemens Medical Systems, Iselin, NJ) using a three-axis gradient head coil. For each participant, an anatomical scan was acquired using a high resolution T1-weighted MPRAGE sequence (repetition time [TR] = 1700 ms; echo time [TE] = 4.38 ms; flip angle = 8°; field of view [FOV] = 280 × 320; 160 slices; matrix = 224 × 256, 1.25 × 1.25 × 1.25 mm resolution). After the anatomical scan, four functional runs were acquired for each participant during the test phase. For the functional runs, imaging was performed using a T2*-weighted EPI sequence (TR = 3000 ms; TE = 30 ms; flip angle = 80°). Each brain volume was composed of 50 slices (FOV = 243 × 243; matrix = 64 × 64, 3 × 3 × 3 mm resolution, slices were oriented along the long axis of the hippocampus, collected interleaved, inferior to superior). For all functional runs, data from the first two volumes were discarded to allow for stabilization of magnetic fields. Stimuli were back-projected onto a screen and viewed on an MR-compatible mirror mounted above the participant’s head. Responses were recorded *via* a response box. Head motion was restricted with a pillow and foam inserts. Subjects requiring vision correction were given MRI-compatible glasses with prescriptions matching their own. The task was presented using MacStim software (CogState Ltd, Melbourne, Australia).

Imaging Analysis

Imaging data were processed using SPM 8 (Wellcome Department of Cognitive Neurology, London) run within Matlab (Matlab Mathwork, Inc., Natick, MA). For pre-processing, fMRI data were slice-time corrected for acquisition order (referenced to the first slice), then realigned and unwarped to correct for motion across runs. Next, the images were spatially normalized by warping each participant’s anatomical scan to MNI (Montreal Neurological Institute) defined standardized brain space, and then applying that algorithm to the EPI data. Finally, the EPI images were spatially smoothed with a Gaussian kernel of 8 mm FWHM. The time series were then high pass filtered at 128 s.

Statistical analyses were performed using the general linear model for event-related designs in SPM 8. For each participant, a whole-brain voxel-wise analysis was conducted in which in which instances of a particular event type were modeled through the convolution with a canonical hemodynamic response function. Each retrieval trial (6 s in duration) was modeled as three 2-s TRs. Because our interest centered on neural recruitment during successful retrieval, all memory conditions were modeled for correct decisions only. Effects for each event type were estimated using a subject-specific, fixed effects model. These data were then entered into a second order, random effects analysis. Analyses contrasted activation as a function of recognition type (relational *vs.* item) using the appropriate trial types. For all random effects analyses, we used combined intensity and cluster size thresholds of $p < .005$ with a minimum cluster size of 20 contiguous voxels ($k \geq 20$) to balance between types I and II error rates (Lieberman & Cunningham, 2009). Conjunction analyses examined what neural regions were commonly activated by CN and MCI participants during item and relational retrieval. For conjunction analyses, the threshold for each contrast was set at $p < .0225$ (such that the conjoint probability of the conjunction analysis, using Fisher’s estimate was $p < .005$; Fisher, 1950; Lazar et al., 2002). Finally, we examined regions differentially activated by CN or MCI participants during item and relational retrieval using two-sample *t* tests. Voxel coordinates are reported in Montreal Neurological Institute (MNI) coordinates and reflect the most significant voxel within the cluster.

RESULTS

Sample Characteristics

Demographic and neuropsychological data are presented in Table 1. Pairwise *t* tests comparing CN and aMCI participants across each measure showed no difference in the demographic variables of age and education, vascular risk (Hachinski Score), mood depression, nor any differences on measures of vocabulary, naming, or generative fluency (animals, vegetables, and COWAT). The two groups did differ significantly as expected on global measures of cognition (MMSE and AD8), episodic learning and memory (Logical Memory Immediate and

Table 1. Demographic and mean neuropsychological data

	Normal <i>n</i> = 12	aMCI <i>n</i> = 12
Age (<i>SD</i>), years	72.6 (5.9)	75.2 (4.3)
Men/women	5/6	5/6
Education (<i>SD</i>), years	15.6 (3.1)	16.3 (2.9)
MMSE	29.5 (0.9)	27.8 (1.7)*
Hachinski Score	1.2 (1.0)	1.6 (0.5)
AD8	.5 (0.8)	3.8 (1.2)*
Shipley Vocabulary Test	34.5 (7.4)	33.8 (4.2)
Digit Span (WAIS-R) Total	16.7 (3.3)	14.8 (2.5)
Logical Memory Immediate (WMS-R)†	16.2 (2.7)	7.9 (4.8)*
Logical Memory Delay (WMS-R)†	15.5 (2.2)	3.8 (5.2)*
CERAD		
– Word List Learning	22.7 (2.0)	17.3 (2.2)*
– Recall Intrusions	<1 (.03)	<1 (1.1)
– Perseverations	<1 (1.1)	<1 (1.0)
– Recall	8.2 (1.2)	3.1 (2.0)*
– Recognition Correct Yes	9.9 (0.3)	8.7 (1.7)*
– Recognition Correct No	10.0 (0.0)	8.8 (2.1)
– Immediate Constructional Praxis	10.4 (1.1)	10.0 (1.3)
– Delay Constructional Praxis Recall	8.9 (1.5)	4.5 (2.1)*
– Delay Constructional Praxis Recognition	4.0 (0.0)	3.5 (0.5)*
Trails A errors	0.0 (0.0)	0.0 (0.0)
Trails A Time	26.1 (9.2)	39.4 (6.5)*
Trails B errors	<1 (0.5)	<1 (0.8)
Trails B Time	68.6 (7.7)	95.0 (6.0)*
Digit Symbol (WAIS-R)	49.5 (6.6)	35.3 (3.6)*
Boston Naming Test	27.7 (2.4)	25.6 (3.4)
Animal Fluency	19.5 (5.6)	16.8 (5.0)
Vegetable Fluency	14.0 (4.8)	13.0 (4.0)
COWAT	38.9 (7.2)	33.8 (8.1)
Geriatric Depression Scale	<1 (0.8)	1.14 (2.2)

Note. Standard deviations are presented in parentheses. WMS-R = Wechsler Memory Scale-Revised; WAIS-R = Wechsler Adult Intelligence Scale – Revised; sec. = seconds.

*Indicates a significant difference between the two groups at $p < .05$.

†Scores are for Story A only.

Delayed, CERAD word list learning and recall, and CERAD delayed recall and recognition of constructional praxis figures), as well as on measures of speeded motor performance (Digit Symbol and Trail Making). Such results were consistent with our recruitment of aMCI participants with single-domain memory impairment.¹

Behavioral Performance During Scanning

The proportion of studied and unstudied stimuli endorsed as “old” are shown in Table 2. Relational recognition accuracy was calculated as the difference between “old” judgments to *intact pairs* (hits) and “old” judgments to *recombined pairs*

¹ It should be noted that the aMCI group performed more slowly on Trails A and B (i.e., non-memory tasks) than the control group. These results are in-line with a recent study reporting additional cognitive impairments in aMCI, particularly on tasks of fluency and executive function, even when the criteria for aMCI are fairly narrow (Kramer et al., 2006).

Table 2. Behavioral accuracy during retrieval of relational information and item information

	Control	aMCI
Relational Memory		
Hits	.84 (.09)	.63 (.16)
False Alarms	.20 (.17)	.28 (.12)
Corrected Recognition	.64 (.22)	.35 (.11)
Item Memory		
Hits	.54 (.25)	.46 (.19)
False Alarms	.12 (.10)	.14 (.08)
Corrected Recognition	.42 (.23)	.32 (.23)

(false alarms), while item recognition was calculated as the difference in “old” judgments to *recombined items* (hits) and “old” judgments to *new items* (false alarms). An analysis of variance (ANOVA) with memory type (item, relational) as a within-subjects factor and group (normal control, aMCI) as a between-subjects factor, revealed a main effect of group ($F(1,22) = 10.31$; $p < .05$), indicating that aMCI participants accuracy was lower than that of CN participants, as well as a main effect of memory task ($F(1,22) = 5.91$; $p < .05$), indicating that relational retrieval accuracy was higher than item retrieval accuracy. The interaction of memory task \times group did not reach significance ($F(1,22) = 2.96$; $p > .05$). As such, the fMRI analysis was not confounded with group by task differences.

Functional Neuroimaging Data

Direct contrasts between item memory and relational memory conditions were conducted to pinpoint alterations in the functional neural architecture based on the specificity of the information retrieved.² The neuroimaging data were analyzed for within-group (CTL or aMCI) differences between relational memory and item memory (i.e., Relational Memory > Item Memory; and vice versa) to specify neural regions differentially activated by each task. Between-group analyses were conducted to identify differential neural activity for each group for each task. Finally, neural regions commonly engaged by CN and aMCI participants were assessed for each task.

Neural regions differentially associated with accurate retrieval of relational information and item information in normal control subjects

We assessed regions differentially activated by CN participants during relational and item retrieval (see Table 3). First, we contrasted activity for all accurate relational memories compared to all accurate item memories (IP + RP + NP > RI + ONI + NI).

² We contrasted all relational memory versus all item memory conditions to assess alterations in functional activity based on the specificity of the information retrieved (i.e. relational vs. item). This approach allowed us to model all conditions (and available data) in the study, which afforded more statistical power. Single condition contrasts were also conducted (e.g., Intact Pair > Recombined). The results were highly similar to those reported in Table 3, except with lower *t*-values.

Table 3. Neural activity during accurate retrieval of relational information and item information in normal control and aMCI participants

Contrast	Region of Activation	Hemisphere	BA	MNI Coordinates				
				x	y	z	t	k
<i>Normal Control</i>								
Relational > Item	Inferior Parietal Lobule	R	40	60	-28	20	6.68	148
	Angular Gyrus	L	39	-48	-62	14	6.33	69
	Parahippocampal Cortex	R	36	30	-34	-8	4.34	71
Item > Relational	Inferior Frontal Gyrus	R	47	40	22	-6	4.34	53
	Superior Frontal Gyrus	R	10	30	60	8	4.18	45
<i>aMCI</i>								
Relational > Item	Inferior Temporal Gyrus	L	20	-50	-16	-22	6.03	26
	Middle Frontal Gyrus	L	9	-32	52	24	5.28	44
	Superior Temporal Gyrus	L	22	-46	-11	4	5.13	58
		R	22	54	-8	2	5.13	27
Item > Relational	Inferior Frontal Gyrus	R	45	60	22	16	6.00	59
	Precuneus	L	7	-18	-40	52	4.98	27
	Superior Parietal Lobule	L	7	-24	-58	50	4.23	20
<i>Normal Control > aMCI</i>								
Relational > Item	Precuneus	L	7	-16	-56	38	4.35	195
	Posterior Cingulate	R	31	18	-44	28	4.21	43
	Parahippocampal Cortex	R	36	30	-34	-10	4.12	50
	Angular Gyrus	L	39	-48	-62	16	4.10	79
Item > Relational	Inferior Frontal Gyrus	R	47	32	22	-6	4.74	77
	Superior Frontal Gyrus	R	10	20	64	6	4.38	86
	Inferior Frontal Gyrus	R	47	30	12	-14	4.14	28
<i>aMCI > Normal Control</i>								
Relational > Item	Superior Temporal Gyrus	L	22	-50	-16	2	5.26	74
	Supramarginal Gyrus	R	40	64	-28	20	4.14	52
Item > Relational	Anterior Cingulate	R	24	4	24	2	4.46	49
	Inferior Frontal Gyrus	R	45	60	22	18	4.39	64
<i>Normal Control AND aMCI</i>								
Relational > Item	Superior Temporal Gyrus	R	21	40	2	-12	7.80	49
	Item > Relational	Middle Frontal Gyrus	R	10	46	46	-12	6.33

Note. Regions significant at $p < .005$, $k > 20$. T = *t*-value; R = right; L = left; PHC = parahippocampal cortex; BA = approximate Brodmann area based on Montreal Neurological Institute coordinates.

This contrast showed activity in left inferior parietal lobule (Brodmann Area, BA 40), left angular gyrus (BA 39), and right parahippocampal cortex (BA 36). The reverse contrasts, comparing activity for all accurate item memories compared to all accurate relational memories (IP + RP + NP < RI + ONI + NI) showed activity in right inferior (BA 47) and right superior frontal (BA 10) regions.

Neural regions differentially associated with accurate retrieval of relational information and item information in aMCI subjects

Next, we examined regions differentially activated by aMCI participants during relational and item retrieval (see Table 3). We contrasted activity during all accurate relational memories compared to activity during all accurate item memories. This contrast showed activity in bilateral superior temporal

gyrus (BA 22), left inferior temporal gyrus (BA 20), and left middle frontal gyrus (BA 9). The reverse contrasts, comparing activity for all accurate item memories versus activity for all accurate relational memories showed activity in right inferior frontal gyrus (BA 45), left precuneus, and left superior parietal lobule (BA 7).

Neural regions associated with retrieval of relational information as a function of group

We hypothesized that the aMCI group, as compared to the CTL group, would not recruit regions traditionally shown to be engaged during relational memory (i.e., the MTL and PC). Rather, under conditions of successful relational retrieval, we predicted that aMCI participants, as compared to the CN group, would recruit regions outside of the MTL and PC, particularly lateral temporal regions during relational retrieval.




Fig. 2. Neural activity during relational retrieval greater than item retrieval in (a) the normal control group relative to the amnesic mild cognitive impairment (aMCI) group and (b) the aMCI group relative to the normal control group.

To test this hypothesis, we assessed regions activated by the CN (i.e., CN > aMCI) or the aMCI (i.e., aMCI > CN) group (relative to the other group) for retrieval of relational, as compared to item, information. To do so, we contrasted activity during all accurate relational memories compared to activity during all accurate item memories. The comparison showed that CN (*vs.* aMCI) activated left precuneus (BA 7), left angular gyrus (BA 39), right posterior cingulate (BA 31), and right parahippocampal cortex (BA 36), while aMCI (*vs.* CN) activated superior temporal gyrus (BA 22) and supramarginal gyrus (BA 40; see Figure 2).

Neural regions associated with retrieval of item information as a function of group

To identify the neural regions associated with item memory as a function of group, we assessed regions activated by the CN (i.e., CN > aMCI) or the aMCI (i.e., aMCI > CN) group (relative to the other group) for retrieval of item, as compared to relational, information. As such, we contrasted activity during accurate item memory compared to activity during accurate relational memory. The comparison showed that CN (*vs.* aMCI) activated right inferior (BA 47) and superior frontal gyrus (BA 10), while aMCI (versus CN) activated anterior cingulate and inferior frontal gyrus (BA 45; see Figure 3).

Neural regions commonly engaged by normal controls and aMCI

Finally, we examined shared regions of activation across CN and aMCI participants during retrieval of relational, as compared to item, memories (see Figure 4), as well as during retrieval of item, as compared to relational, memories. To do so, we conducted a conjunction analysis to identify regions

that were more active during the relational memory than during item memory for both CN and aMCI groups. This analysis revealed activity in superior temporal gyrus (BA 21). Next, we conducted the same type of conjunction analysis to identify regions that were more active during the item memory than during relational memory for both CN and aMCI groups. This analysis showed activity in right anterior frontal gyrus (BA 10).

DISCUSSION

We compared the neural underpinnings of item and relational retrieval in aMCI and CN participants, using event-based fMRI, while holding constant the stimuli at encoding and the encoding task for the two retrieval conditions (item and relational). The use of event-related fMRI allowed us to limit the analysis to *correct* memory trials only, eliminating potential confounds due to differences in retrieval success. Accuracy was numerically lower in the aMCI, relative to the CN, group. However, the group by memory task interaction did not reach significance, indicating that the aMCI group showed no disproportionate impairment in relational memory performance. This result may have arisen from the encoding task used in the current study, in which participants were shown two words and instructed to create a meaningful sentence. Such deep (elaborative) semantic processing, coupled with generative (and likely self-referential) processing, leads to favorable encoding conditions for older adults during retrieval of relational information (Giovanello et al., 2012; Glisky et al., 2001). It should be noted, however, that the aMCI group did not benefit to the same extent as the control group. Future studies should elucidate the conditions under which aMCI participants do and do not benefit from encoding support during tasks of relational memory. Nonetheless, the




Fig. 3. Neural activity during item retrieval greater than relational retrieval in (a) the normal control group relative to the amnesic mild cognitive impairment (aMCI) group and (b) the aMCI group relative to the normal control group.

group by memory task interaction was not significant in the current study, indicating that the fMRI data were not confounded by differential success rates on the two memory tasks across the two groups.

An examination of the neural regions commonly recruited by CN and aMCI groups showed significant activity in right

STG during relational, as compared to item, retrieval, and right anterior frontal gyrus during item, as compared to relational, retrieval. These findings highlight the notion that not all neural regions are functionally affected in aMCI and provide insight into the nature of the cognitive processes that may be commonly used in CN and aMCI participants. For example, activity in right STG has been reported in studies using word stimuli during tasks of semantic retrieval (e.g., Dalla Barba, Parlato, Jobert, Samson, & Pappata, 1998). Additionally, right superior temporal activity has been observed during reduced attentional allocation (Iidaka, Anderson, Kapur, Cabeza, & Craik, 2000) and during gist-based processing (Dennis, Kim, & Cabeza, 2008), that is, the retention of the general meaning of a concept or an event (Brainerd & Reyna, 1990). As such, right STG activity in the current study may reflect retrieval of gist-based, semantic information (e.g., partial recovery of the sentences formed at encoding) that aids in successful relational memory retrieval. The activity in anterior frontal cortex was observed in both groups during item, compared to relational, retrieval. Anterior frontal cortex (BA10) has been associated with internally directed attention (Burgess, Simons, Dumontheil, & Gilbert, 2005; Simons, Gilbert, Owen, Fletcher, & Burgess, 2005; Simons, Owen, Fletcher, & Burgess, 2005), as well as with retrieval mode (Düzel et al., 1999; Lepage, Ghaffar, Nyberg, & Tulving, 2000; Velanova et al., 2003), a sustained cognitive set associated with perceiving stimuli as cues to elicit memory (Nyberg et al., 1995; Tulving, 1983). Thus, recruitment of anterior frontal cortex may reflect a sustained, internally directed retrieval mechanism that is common to both groups. The preservation of such retrieval mechanisms in aMCI could prove useful for intervention or training techniques aimed at capitalizing on intact cognitive processes in these individuals.




Fig. 4. Neural activity common to both normal control and aMCI groups during (a) relational retrieval relative to item retrieval and (b) item retrieval relative to relational retrieval.

In addition to the neural regions commonly activated by the two groups, however, several group differences in brain activity were observed. During accurate relational retrieval, CN adults showed significantly greater activity than aMCI individuals in left precuneus, right posterior cingulate, right parahippocampal cortex, and left angular gyrus. Several of these regions, particularly those located along the midline (i.e., precuneus, posterior cingulate, and parahippocampal cortex) have been characterized as being part of the default mode network (DMN) – defined as a set of functionally connected brain regions that exhibit task-induced deactivation and increase activity at rest (Buckner, Andrews-Hanna, Schacter, 2008; Raichle & Snyder, 2007). While the role of the DMN in cognition is unclear, increased activity in midline DMN regions has been reported to predict subsequent successful memory when a subjective, social, or self-referential judgment is made at encoding (Dobbs & Wagner, 2005; Kelley et al., 2002; Mitchell, Macrae, & Banaji, 2004). The current study used an encoding task that required generation of a meaningful sentence that related the two word stimuli appearing on each trial. It is possible that CN adults generated sentences that engendered a higher degree of social or self-referential processing, than did the aMCI group. This suggestion, however, is purely speculative and should be tested empirically by either inclusion of an alternative encoding task or administration of a post-test debriefing questionnaire targeted at social and self-referential processing.

Beyond to the involvement of parahippocampal cortex and precuneus in the DMN, these regions also have been frequently observed during memory retrieval studies (Cabeza & Nyberg, 2000). Evidence from neuropsychological, neuroimaging, and neurophysiological studies indicate that parahippocampal cortex and precuneus regions contribute to the mnemonic processes of recollection and familiarity. Whereas recollection is a conscious, attention demanding process, in which prior contextual aspects of an experience are retrieved, familiarity, in contrast, is thought to be an unconscious, automatic process that arises when fluent processing of a stimulus is attributed to prior experience with that stimulus. Recollection and familiarity are thought to be functionally distinct and to rely on different neural underpinnings. One influential memory model postulates that the parahippocampal cortex contributes to recollection, possibly *via* the representation and retrieval of contextual information, with the hippocampus serving to bind such contextual information with other components of an event (for a review, see Eichenbaum et al., 2007). More recently, the precuneus (located in dorsal parietal cortex) has been implicated during retrieval of familiar, yet low-confident decisions (for a review see Cabeza, Ciaramelli, Olson, & Moscovitch, 2008).

Indeed, the role of PC in general has received considerable attention recently, and functionally distinct regions within PC have been proposed to mediate performance during episodic memory retrieval (for reviews, see Cabeza et al., 2008; Wagner, Shannon, Kahn, & Buckner, 2005). Traditionally, the PC has been associated with aspects of attention. For example, Corbetta and Shulman (2002) suggest that the

dorsal parietal cortex (DPC) is involved in top-down, voluntary attention, whereas the ventral parietal cortex (VPC) is involved in bottom-up, involuntary attention. Support for this theory comes from studies showing that the DPC is preferentially engaged during the cuing period, when participants are actively and voluntarily searching for a target, whereas the VPC is primarily recruited during target detection (Corbetta, Kincade, Ollinger, McAvoy, & Shulman, 2000). This distinction has recently been extrapolated to the domain of memory. Specifically, it has been hypothesized that DPC and VPC differentially contribute to episodic memory retrieval, where DPC contributes top-down attentional processes guided by retrieval goals, and VPC mediates bottom-up attentional processes captured by retrieval output (Cabeza, 2008; Ciaramelli, Grady, Levine, Ween, & Moscovitch, 2010). Additionally, recollection has been associated with VPC, while familiarity has been associated with DPC (Wheeler & Buckner, 2004; Yonelinas, Otten, Shaw, & Rugg, 2005). We observed activity in both VPC (angular gyrus BA 39) and DPC (precuneus, BA7) in the CN group, compared to the aMCI group, during successful relational retrieval. Taken together, the observed activity in parahippocampal cortex, precuneus (DPC), and angular gyrus (VPC) in CN individuals, as compared to aMCI subjects, suggests that the CN group engaged both recollection and familiarity based processing to a greater extent than did aMCI subjects to support successful relational retrieval.

An examination of the regions showing greater activity in aMCI subjects, as compared to the CN group, revealed activity in right supramarginal gyrus (BA 40) and left superior temporal gyrus (STG; BA 22). The role of left STG in language is well-documented, particularly with regard to semantic processing (e.g., Friederici, Makuuchi, & Bahlmann, 2009). This finding, coupled with the activity observed in right STG for both CN and aMCI groups, suggests that aMCI subjects may have attempted recovery of encoding sentences which aided in successful relational memory judgments about the words pairs. Indeed, Lenzi et al. (2011) conducted an fMRI investigation of multiple cognitive domains in aMCI and found that right STG was activated in aMCI, compared to CN, individuals, and correlated significantly with neuropsychological scores of Story Recall. Recruitment of right supramarginal gyrus (BA 40), a region in VPC, is noteworthy given the evidence of its role in recollection-based processing. Given the contribution of recollection-based processing to relational memory (Hockley & Consoli, 1999), the engagement of this region points to aMCI-related alterations in the neural network engaged during successful retrieval of relational information.

During accurate item retrieval, CN adults showed significantly greater activity than aMCI individuals in right inferior (BA 47) and superior frontal gyri (BA 10), whereas aMCI participants showed significantly greater activity than CN adults in right cingulate (BA 24) and right inferior frontal gyrus (BA 45). Such engagement of frontal regions, in the context of under recruitment of posterior regions (i.e., MTL and PC) during memory retrieval, has been documented in CN subjects and is known as the posterior to anterior shift in aging (PASA; Davis, Dennis, Daselaar, Fleck, & Cabeza, 2008).

The current findings indicate that the PASA pattern is also observed in aMCI.

In summary, we directly compared the neural underpinnings of successful relational retrieval in aMCI and CN groups using event-related fMRI. The data showed that the CN, compared to the aMCI, group activated left precuneus (BA 7), left angular gyrus (BA 39), right posterior cingulate (BA 31), and right parahippocampal cortex (BA 36) during relational retrieval. These regions have been implicated in the DMN, as well as during memory retrieval, particularly with regard to the mnemonic processes of recollection and familiarity. Specifically, recruitment of parahippocampal cortex, VPC, and DPC by CN individuals suggests that control participants used multiple regions associated with recollection, as well as regions thought to support familiarity to achieve successful relational retrieval. In contrast, the aMCI group, compared to the CN, group activated superior temporal gyrus (BA 22) and supramarginal gyrus (BA 40) for this comparison, indicating an early shift in the functional neural architecture of relational retrieval in aMCI. These findings may prove useful in future studies aimed at capitalizing on functionally intact neural regions as targets for treatment and slowing of the disease course.

ACKNOWLEDGEMENTS

This research was supported, in part, by NIH/NIA grants K01-AG028774 (KSG), L30-AG029001 (JNB), and P30-AG028377 (KWB).

The authors report no conflicts of interest.

REFERENCES

- Anderson, N.D., Ebert, P.L., Jennings, J.M., Grady, C.L., Cabeza, R., & Graham, S.J. (2008). Recollection- and familiarity-based memory in healthy aging and amnesic mild cognitive impairment. *Neuropsychologia*, *22*, 177–187.
- Brainerd, C.J., & Reyna, V.F. (1990). Gist is the gist: The fuzzy-trace theory and new intuitionism. *Developmental Review*, *10*, 3–47.
- Buckner, R.L., Andrews-Hanna, J.R., & Schacter, D.L. (2008). The brain's default network: Anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*, *1124*, 1–38.
- Buckner, R.L., Snyder, A.Z., Shannon, B.J., LaRossa, G., Sachs, R., Fotenos, A.F., ... Mintun, M.A. (2005). Molecular, structural, and functional characterization of Alzheimer's disease: Evidence for a relationship between default activity, amyloid, and memory. *Journal of Neuroscience*, *25*, 7709–7717.
- Burgess, P.W., Simons, J.S., Dumontheil, I., & Gilbert, S.J. (2005). The gateway hypothesis of rostral prefrontal cortex (area 10) function. In J. Duncan, L. Phillips, P. McLeod (Eds), *Measuring the mind: Speed, control, and age* (pp. 217–248). Oxford: Oxford University Press.
- Cabeza, R. (2008). Role of parietal regions in episodic memory retrieval: The dual attentional processes hypothesis. *Neuropsychologia*, *46*, 1813–1827.
- Cabeza, R., Ciaramelli, E., Olson, I.R., & Moscovitch, M. (2008). The parietal cortex and episodic memory: An attentional account. *Nature Reviews Neuroscience*, *9*, 613–625.
- Cabeza, R., & Nyberg, L. (2000). Neural bases of learning and memory: Functional neuroimaging evidence. *Current Opinion in Neurology*, *13*, 415–421.
- Celone, K., Calhoun, V., Dickerson, B., Atri, A., Chua, E., Miller, S., ... Sperling, R. (2006). Alterations in memory networks in mild cognitive impairment and Alzheimer's disease: An independent component analysis. *The Journal of Neuroscience*, *26*, 10222–10231.
- Ciaramelli, E., Grady, C.L., Levine, B., Ween, J., & Moscovitch, M. (2010). Top-down and bottom-up attention to memory are dissociated in posterior parietal cortex: Neuroimaging and neuropsychological evidence. *Journal of Neuroscience*, *30*, 4943–4956.
- Convit, A., de Leon, M.J., Tarshishi, C., De Santri, S., Tsui, W., Rusinek, H., & George, A.E. (1997). Specific hippocampal volume reductions in individuals at risk for Alzheimer's disease. *Neurobiology of Aging*, *18*, 131–138.
- Corbetta, M., Kincade, J.M., Ollinger, J.M., McAvoy, M.P., & Shulman, G.L. (2000). Voluntary orienting is dissociated from target detection in human posterior parietal cortex. *Nature Neuroscience*, *3*, 292–297.
- Corbetta, M., & Shulman, G.L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience*, *3*, 201–215.
- Dalla Barba, G., Parlato, V., Jobert, A., Samson, Y., & Pappata, S. (1998). Cortical networks implicated in semantic and episodic memory: Common or unique? *Cortex*, *34*, 547–561.
- Davis, S.W., Dennis, N.A., Daselaar, S.M., Fleck, M.S., & Cabeza, R. (2008). Qué PASA? The posterior-anterior shift in aging. *Cerebral Cortex*, *18*, 1201–1209.
- De Santi, S., de Leon, M.J., Rusinek, H., Convit, A., Tarshish, C.Y., Roche, A., ... Fowler, J. (2001). Hippocampal formation glucose metabolism and volume losses in MCI and AD. *Neurobiology of Aging*, *22*, 529–539.
- Dennis, N.A., Kim, H., & Cabeza, R. (2008). Age-related differences in brain activity during true and false memory retrieval. *Journal of Cognitive Neuroscience*, *20*, 1390–1402.
- Dickerson, B.C., Goncharova, I., Sullivan, M.P., Forchetti, C., Wilson, R.S., Bennett, D.A., ... deToledo-Morrell, L. (2001). MRI-derived entorhinal and hippocampal atrophy in incipient and very mild Alzheimer's disease. *Neurobiology of Aging*, *22*, 747–754.
- Dickerson, B.C., Salat, D.H., Greve, D.N., Chua, E.F., Rand-Giovannetti, E., Rentz, D.M., ... Sperling, R.A. (2005). Increased hippocampal activation in mild cognitive impairment compared to normal aging and AD. *Neurology*, *65*, 404–411.
- Dickerson, B.C., & Sperling, R.A. (2009). Large-scale functional brain network abnormalities in Alzheimer's disease: Insights from functional neuroimaging. *Behavioral Neurology*, *21*, 63–75.
- Dobbins, I.G., & Wagner, A.D. (2005). Domain-general and domain-sensitive prefrontal mechanisms for recollecting events and detecting novelty. *Cerebral Cortex*, *15*, 1768–1778.
- Du, A.T., Schuff, N., Amend, D., Laakso, M.P., Hsu, Y.Y., Jagust, W.J., ... Weiner, M.W. (2001). Magnetic resonance imaging of the entorhinal cortex and hippocampus in mild cognitive impairment and Alzheimer's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, *71*, 441–447.
- Düzel, E., Cabeza, R., Picton, T.W., Yonelinas, A.P., Scheich, H., Heinze, H.J., & Tulving, E. (1999). Task-related and item-related brain processes of memory retrieval. *Proceedings of the National Academy of Sciences of the United States of America*, *96*, 1794–1799.

- Eichenbaum, H., Yonelinas, A.P., & Ranganath, C. (2007). The medial temporal lobe and recognition memory. *Annual Review of Neuroscience*, *30*, 123–152.
- Fisher, R.A. (1950). *Statistical methods for research workers*. London: Oliver and Boyd.
- Folstein, M.F., Folstein, S.E., & McHugh, P.R. (1975). “Minimal mental state”. A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatry Research*, *12*, 189–198.
- Fowler, K.S., Saling, M.M., Conway, E.L., Semple, J.M., & Louis, W.J. (2002). Paired associate performance in the early detection of DAT. *Journal of the International Neuropsychological Society*, *8*, 58–71.
- Friederici, A.D., Makuuchi, M., & Bahlmann, J. (2009). The role of the posterior superior temporal cortex in sentence comprehension. *Neuroreport*, *20*, 563–568.
- Giovanello, K.S., & Schacter, D.L. (2012). Reduced specificity of hippocampal and posterior ventrolateral prefrontal activity during relational retrieval in normal aging. *Journal of Cognitive Neuroscience*, *24*, 159–170.
- Glisky, E.L., Rubin, S.R., & Davidson, P.S. (2001). Source memory in older adults: An encoding or retrieval problem? *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *27*, 1131–1146.
- Golde, T.E. (2006). Disease modifying therapy for AD? *Journal of Neurochemistry*, *99*, 689–707.
- Grundman, M., Sencakova, D., Jack, C.R., Jr., Petersen, R.C., Kim, H.T., Schultz, A., ... Thal, L.J. (2002). Brain MRI hippocampal volume and prediction of clinical status in a mild cognitive impairment trial. *Journal of Molecular Neuroscience*, *19*, 23–27.
- Hachinski, V.C., Iliff, L.D., Zilhka, E., Du Boulay, G.H., McAllister, V.L., Marshall, J., ... Symon, L. (1975). Cerebral blood flow in dementia. *Archives of Neurology*, *32*, 632–637.
- Hayden, K.M., Jones, R.N., Zimmer, C., Plassman, B., Browndyke, J.N., Pieper, C., ... Welsh-Bohmer, K.A. (2011). Factor structure of the National Alzheimer’s Coordinating Centers Uniform Dataset Neuropsychological Battery: An evaluation of invariance between and within diagnostic groups over time. *Alzheimer’s and Association Disorders*, *25*, 128–137.
- Herholz, K., Salmon, E., Perani, D., Baron, J.C., Holthoff, V., Frölich, L., ... Heiss, W.D. (2002). Discrimination between Alzheimer dementia and controls by automated analysis of multicenter FDG PET. *Neuroimage*, *17*, 302–316.
- Hockley, W.A., & Consoli, C. (1999). Familiarity and recollection in item and associative recognition. *Memory and Cognition*, *27*, 657–664.
- Iidaka, T., Anderson, N.D., Kapur, S., Cabeza, R., & Craik, F.I. (2000). The effect of divided attention on encoding and retrieval in episodic memory revealed by positron emission tomography. *Journal of Cognitive Neuroscience*, *12*, 267–280.
- Ivanou, A., Adam, S., Van der Linden, M., Salmon, E., Juillerat, A.C., Mulligan, R., & Seron, X. (2005). Memory evaluation with a new cued recall test in patients with mild cognitive impairment and Alzheimer’s disease. *Journal of Neurology*, *252*, 47–55.
- Jack, C.R., Jr., Petersen, R.C., Xu, Y.C., O’Brien, P.C., Smith, G.E., Ivnik, R.J., ... Kokmen, E. (1999). Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment. *Neurology*, *52*, 1397–1403.
- Kaplan, E., Goodglass, H., & Weintraub, S. (1983). *Boston Naming Test*. Philadelphia, PA: Lea and Febiger.
- Kaye, J.A., Moore, M.M., Dame, A., Quinn, J., Camicioli, R., Howieson, D., ... Sexton, G. (2005). Asynchronous regional brain volume losses in presymptomatic to moderate AD. *Journal of Alzheimer’s Disease*, *8*, 51–56.
- Kelley, W.M., Macrae, C.N., Wyland, C.L., Caglar, S., Inati, S., & Heatherton, T.F. (2002). Finding the self? An event-related fMRI study. *Journal of Cognitive Neuroscience*, *14*, 785–794.
- Killiany, R.J., Gomez-Isla, T., Moss, M., Kikinis, R., Sandor, T., Jolesz, F., ... Albert, M.S. (2000). Use of structural magnetic resonance imaging to predict who will get Alzheimer’s disease. *Annals of Neurology*, *47*, 430–439.
- Kramer, J.H., Nelson, A., Johnson, J.K., Yaffe, K., Glenn, S., Rosen, H.J., & Miller, B.L. (2006). Multiple cognitive deficits in amnesic mild cognitive impairment. *Dementia and Geriatric Cognitive Disorders*, *22*, 306–311.
- Lazar, N.A., Luna, B., Sweeney, J.A., & Eddy, W.F. (2002). Combining brains: a survey of methods for statistical pooling of information. *Neuroimage*, *16*, 538–550.
- Lenzi, D., Serra, L., Perri, R., Pantano, P., Lenzi, G.L., Paulesu, E., ... Macaluso, E. (2011). Single domain amnesic MCI: A multiple cognitive domains fMRI investigation. *Neurobiology of Aging*, *32*, 1542–1557.
- Lepage, M., Ghaffar, O., Nyberg, L., & Tulving, E. (2000). Prefrontal cortex and episodic memory retrieval mode. *Proceedings of the National Academy of Sciences of the United States of America*, *97*, 506–511.
- Lieberman, M.D., & Cunningham, W.A. (2009). Type I and Type II error concerns in fMRI research: re-balancing the scale. *Social Cognitive and Affective Neuroscience*, *4*, 423–428.
- Machulda, M.M., Senjem, M.L., Weigand, S.D., Smith, G.E., Ivnik, R.J., Boeve, B.F., ... Jackie, C.R. (2009). Functional magnetic resonance imaging changes in amnesic and nonamnesic mild cognitive impairment during encoding and recognition tasks. *Journal of the International Neuropsychological Society*, *15*, 372–382.
- Minoshima, S., Giordani, B., Berent, S., Frey, K.A., Foster, N.L., & Kuhl, D.E. (1997). Metabolic reduction in the posterior cingulate cortex in very early Alzheimer’s disease. *Annals of Neurology*, *42*, 85–94.
- Mitchell, J.P., Macrae, C.N., & Banaji, M.R. (2004). Encoding-specific effects of social cognition on the neural correlates of subsequent memory. *Journal of Neuroscience*, *24*, 4912–4917.
- Morris, J.C., Heyman, A., Mohs, R.C., Hughes, J.P., van Belle, G., Fillenbaum, G., ... Clark, C. (1989). The Consortium to Establish a Registry for Alzheimer’s Disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimer’s disease. *Neurology*, *39*, 1159–1165.
- Nyberg, L., Tulving, E., Habib, R., Nilsson, L.G., Kapur, S., Houle, S., ... McIntosh, A.R. (1995). Functional brain maps of retrieval mode and recovery of episodic information. *Neuroreport*, *7*, 249–252.
- Petrella, J.R., Prince, S.E., Wang, L., Hellegers, C., & Doraiswamy, P.M. (2007). Prognostic value of posteromedial cortex deactivation in mild cognitive impairment. *PLoS One*, *2*, e1104.
- Pennanen, C., Kivipelto, M., Tuomainen, S., Hartikainen, P., Hänninen, T., Laakso, M.P., ... Soininen, H. (2004). Hippocampus and entorhinal cortex in mild cognitive impairment and early AD. *Neurobiology of Aging*, *25*, 303–310.
- Petersen, R.C. (2004). Mild cognitive impairment as a diagnostic entity. *Journal of Internal Medicine*, *256*, 183–194.
- Petersen, R.C., Smith, G.E., Waring, S.C., Ivnik, R.J., Tangalos, E., & Kokmen, E. (1999). Mild cognitive impairment: Clinical

- characterization and outcome. *Archives of Neurology*, 56, 303–308.
- Raichle, M.E., & Snyder, A.Z. (2007). A default mode of brain function: A brief history of an evolving idea. *Neuroimage*, 37, 1083–1090.
- Rajah, M.N., Kromas, M., Han, J.E., & Pruessner, J.C. (2010). Group differences in anterior hippocampal volume and in the retrieval of spatial and temporal context memory in healthy young versus older adults. *Neuropsychologia*, 48, 4020–4030.
- Reiman, E.M., Caselli, R.J., Yun, L.S., Chen, K., Bandy, D., Minoshima, S., ... Osborne, D. (1996). Preclinical evidence of Alzheimer's disease in persons homozygous for the epsilon 4 allele for apolipoprotein E. *New England Journal of Medicine*, 334, 752–758.
- Rodrigue, K.M., & Raz, N. (2004). Shrinkage of the entorhinal cortex over five years predicts memory performance in healthy adults. *Journal of Neuroscience*, 24, 956–963.
- Rombouts, S.A., Goekoop, R., Stam, C.J., Barkhof, F., & Scheltens, P. (2005). Delayed rather than decreased BOLD response as a marker for early Alzheimer's disease. *Neuroimage*, 26, 1078–1085.
- Rosen, A.C., Prull, M.W., Gabrieli, J.D., Stoub, T., O'Hara, R., Friedman, L., ... deToledo-Morrell, L. (2003). Differential associations between entorhinal and hippocampal volumes and memory performance in older adults. *Behavioral Neuroscience*, 117, 1150–1160.
- Schacter, D.L., Gallo, D.A., & Kensinger, E.A. (2007). The cognitive neuroscience of implicit and false memories: Perspectives on processing specificity. In J.S. Nairne (Ed.), *The foundations of remembering: Essays in honor of Henry L. Roediger III* (pp. 353–378). New York: Psychology Press.
- Shipley, W.S. (1967). *Shipley Institute of Living Scale*. Los Angeles, CA: Western Psychological Services.
- Simons, J.S., Gilbert, S.J., Owen, A.M., Fletcher, P.C., & Burgess, P.W. (2005). Distinct roles for lateral and medial anterior prefrontal cortex in contextual recollection. *Journal of Neurophysiology*, 94, 813–820.
- Simons, J.S., Owen, A.M., Fletcher, P.C., & Burgess, P.W. (2005). Anterior prefrontal cortex and the recollection of contextual information. *Neuropsychologia*, 43, 1774–1783.
- Spreen, O., & Strauss, E. (1991). *A compendium of neuropsychological tests: Administration, norms, and commentary*. New York: Oxford University Press.
- Trivedi, M.A., Murphy, C.M., Goetz, C., Shah, R.C., Gabrieli, J.D., Whitfield-Gabrieli, S., ... Stebbins, G.T. (2008). fMRI activation changes during successful episodic memory encoding and recognition in amnesic mild cognitive impairment relative to cognitively healthy older adults. *Dementia and Geriatric Cognitive Disorders*, 26, 123–137.
- Troyer, A.K., Murphy, K.J., Anderson, N.D., Hayman-Abello, B.A., Craik, F.I., & Moscovitch, M. (2008). Item and associative memory in amnesic mild cognitive impairment: performance on standardized memory tests. *Neuropsychology*, 22, 10–16.
- Tschanz, J.T., Welsh-Bohmer, K.A., Lykestos, C.G., Corcoran, C., Green, R.C., Norton, M.C., ... Breitner, J.C.S. and the Cache County Investigators (2006). Conversion to dementia from mild cognitive disorder: The Cache County Study. *Neurology*, 67, 229–234.
- Tulving, E. (1983). *Elements of episodic memory*. Oxford: Clarendon Press.
- Velanova, K., Jacoby, L.L., Wheeler, M.E., McAvoy, M.P., Peterson, S.E., & Buckner, R.L. (2003). Functional-anatomic correlates of sustained and transient processing components engaged during controlled retrieval. *The Journal of Neuroscience*, 23, 8460–8470.
- Wagner, A., Shannon, B., Kahn, I., & Buckner, R.L. (2005). Parietal lobe contributions to episodic memory retrieval. *Trends in Cognitive Sciences*, 9, 445–453.
- Wang, K., Liang, M., Wang, L., Tian, L., Zhang, X., Li, K., & Jiang, T. (2007). Altered functional connectivity in early Alzheimer's disease: a resting-state fMRI study. *Human Brain Mapping*, 28, 967–978.
- Wechsler, D. (1981). *Wechsler Adult Intelligence Scale – Revised*. San Antonio, TX: The Psychological Corporation.
- Wechsler, D. (1987). *Wechsler Memory Scale – Revised*. San Antonio, TX: The Psychological Corporation.
- Wheeler, M.E., & Buckner, R.L. (2004). Functional-anatomic correlates of remembering and knowing. *Neuroimage*, 21, 1337–1349.
- Winblad, B., Palmer, K., Kivipelto, M., Jelic, V., Fratiglioni, L., Wahlund, L.O., ... Petersen, R. (2004). Mild cognitive impairment – Beyond controversies, towards a consensus: Report of the International Working Group on Mild Cognitive Impairment. *Journal of Internal Medicine*, 256, 240–246.
- Xu, Y.C., Jack, C.R., O'Brien, P.C., Kokmen, E., Smith, G.E., Ivnik, R.J., ... Petersen, R.C. (2000). Usefulness of MRI measures of entorhinal cortex versus hippocampus in AD. *Neurology*, 54, 1760–1767.
- Yonelinas, A.P. (2001). Components of episodic memory: the contribution of recollection and familiarity. *Philosophical Transactions Royal Society London B Biological Science*, 356, 1363–1374.
- Yonelinas, A.P., Otten, L.J., Shaw, K.N., & Rugg, M.D. (2005). Separating the brain regions involved in recollection and familiarity in recognition memory. *Journal of Neuroscience*, 25, 3002–3008.