

# Angiostrongylus cantonensis in urban populations of terrestrial gastropods and rats in an impoverished region of Brazil

## Research Article

\*These authors contributed equally to this work.

**Cite this article:** Souza FN *et al* (2021). *Angiostrongylus cantonensis* in urban populations of terrestrial gastropods and rats in an impoverished region of Brazil. *Parasitology* **148**, 994–1002. <https://doi.org/10.1017/S0031182021000597>

Received: 28 October 2020

Revised: 4 April 2021

Accepted: 5 April 2021

First published online: 12 April 2021












### Key words:

Angiostrongyliasis; helminth; one health; public health; *Rattus norvegicus*; risk factors; slug; snail; zoonoses

### Author for correspondence:

Fábio N. Souza,

E-mail: [fabionevesouza@gmail.com](mailto:fabionevesouza@gmail.com)

Fábio N. Souza<sup>1,2\*</sup> , Máisa Aguiar Santos<sup>1,2,\*</sup> , Daniele Almeida Alves<sup>1,2</sup> , Leyva Cecília Vieira de Melo<sup>3</sup>, Dan Jessé Gonçalves da Mota<sup>3</sup>, Arsinoê Cristina Pertile<sup>1,2</sup>, Ricardo Gava<sup>3</sup>, Pedro Luiz Silva Pinto<sup>3</sup> , Max T. Eyre<sup>4</sup> , Caio Graco Zeppelini<sup>1</sup> , Mitermayer G. Reis<sup>1,2,5</sup> , Albert I. Ko<sup>5</sup> , Mike Begon<sup>6</sup> , Thiago C. Bahiense<sup>1</sup>, Federico Costa<sup>1,2,5,6</sup> , and Ticiania Carvalho-Pereira<sup>1,2</sup> 

<sup>1</sup>Universidade Federal da Bahia, UFBA, Salvador, Brazil; <sup>2</sup>Instituto Gonçalo Moniz, Fundação Oswaldo Cruz, Ministério da Saúde, Salvador, Brazil; <sup>3</sup>Instituto Adolfo Lutz, São Paulo, Brazil; <sup>4</sup>Centre for Health Informatics, Computing, and Statistics, Lancaster University Medical School, Lancaster, UK; <sup>5</sup>Department of Epidemiology of Microbial Disease, Yale School of Public Health, New Haven, USA and <sup>6</sup>Institute of Integrative Biology, University of Liverpool, Liverpool, UK

### Abstract

The nematode *Angiostrongylus cantonensis* is the most common cause of neuroangiostrongyliasis (manifested as eosinophilic meningitis) in humans. Gastropod molluscs are used as intermediate hosts and rats of various species are definitive hosts of this parasite. In this study, we identified several environmental factors associated with the presence and abundance of terrestrial gastropods in an impoverished urban region in Brazil. We also found that body condition, age and presence of co-infection with other parasite species in urban *Rattus norvegicus*, as well as environmental factors were associated with the probability and intensity of *A. cantonensis* infection. The study area was also found to have a moderate prevalence of the nematode in rodents (33% of 168 individuals). Eight species of molluscs (577 individuals) were identified, four of which were positive for *A. cantonensis*. Our study indicates that the environmental conditions of poor urban areas (presence of running and standing water, sewage, humidity and accumulated rain and accumulation of construction materials) influenced both the distribution and abundance of terrestrial gastropods, as well as infected rats, contributing to the maintenance of the *A. cantonensis* transmission cycle in the area. Besides neuroangiostrongyliasis, the presence of these hosts may also contribute to susceptibility to other zoonoses.

### Introduction

The rat lungworm *Angiostrongylus cantonensis* (Nematoda: Metastrongyloidea) is a parasitic nematode that causes neuroangiostrongyliasis, because of its larvae tropism for the central nervous system (CNS). This can manifest as eosinophilic meningitis with possible severe and fatal outcomes (Wang *et al.*, 2008; Cowie, 2013; Morassutti *et al.*, 2014). Humans are accidental hosts, with infection occurring after ingestion of terrestrial gastropods (slugs and snails) infected by third-stage larvae (L3), including accidental consumption of these invertebrates in vegetables; or through the ingestion of paratenic hosts containing L3 (e.g. freshwater shrimp, crabs, planarians) (Thiengo *et al.*, 2013). Neuroangiostrongyliasis has expanded from southeastern Asia and Pacific Islands, where the disease is considered endemic, to countries and islands around the world (Cowie, 2013), advancing from tropical to temperate zones (Červená *et al.*, 2019). The disease is well documented in South America (Valente *et al.*, 2018) and since the first reported case in Brazil in 2007 it has continued to emerge in the north (Barbosa *et al.*, 2020), northeastern (Thiengo *et al.*, 2010) and southeastern regions (Caldeira *et al.*, 2007) of the country. Infected intermediate and definitive hosts are found in almost all regions of the country (Carvalho *et al.*, 2012; Moreira *et al.*, 2013; Carvalho-Pereira *et al.*, 2018).

The transmission cycle involves gastropods as the intermediate host of *A. cantonensis* and rodents as definitive hosts. Infected rodents release first-stage larvae in their feces which then infect molluscs by ingestion or, perhaps less importantly, by penetrating the body wall or the respiratory pore (Thiengo *et al.*, 2013). The larvae then develop to the third stage within the molluscs and infect rodents when they ingest infected molluscs. In rats, these larvae develop to the subadult stage (L5) in the CNS, subsequently migrating to the pulmonary arteries, where they develop into adult nematodes, reproduce and complete the cycle (Thiengo *et al.*, 2013).

In impoverished tropical urban areas, both rats and certain widespread gastropods thrive because of the highly favourable environmental conditions (Panti-May *et al.*, 2016; Onyishi

*et al.*, 2018). However, the relationship between specific environmental variables and gastropod abundance and diversity has not been studied in detail in this context. The life cycle of *A. cantonensis* can be maintained by a broad range of intermediate hosts (Kim *et al.*, 2014) but the introduced (and invasive) snail species *Achatina fulica* Bowdich, 1822 and the slug *Sarasinula marginata* Semper, 1885 are the most commonly found intermediate host species in Brazil (Caldeira *et al.*, 2007; Carvalho *et al.*, 2012). The lack of adequate sanitation and refuse collection offers a habitat with plentiful food sources which enables definitive host populations for *A. cantonensis* – such as one of its main definitive hosts, *Rattus norvegicus* Berkenhout, 1769 (Valente *et al.*, 2018) – to proliferate (Masi *et al.*, 2010). This diversity of abundant host species is therefore likely to facilitate *A. cantonensis* infection in residents of impoverished urban areas already affected by several other zoonotic agents (Costa *et al.*, 2014; Carvalho-Pereira *et al.*, 2018).

A recent study in an impoverished urban area of Brazil estimated that the local population of *R. norvegicus* had an *A. cantonensis* prevalence of around 40%, indicating that residents were at risk of exposure to this nematode (Carvalho-Pereira *et al.*, 2018). However, information on the abundance and diversity of intermediate host species and the prevalence of *A. cantonensis* within these host populations is scarce for impoverished urban settings, complicating efforts to understand the risk of exposure in human populations. Consequently, the main objective of this study was to describe the occurrence and abundance of terrestrial gastropods present in a Brazilian impoverished urban area and to identify the associated environmental factors. Our secondary objective was to identify whether environmental and demographic factors, body condition or coinfection with other helminths were associated with the presence and intensity of *A. cantonensis* infection in *R. norvegicus*, using long-term monitoring data. This study seeks to identify factors facilitating the maintenance of this helminth's life cycle, which can be used to identify high-risk areas and to guide future targeting of public health interventions.

## Materials and methods

### Study area

We conducted our study in the neighbourhood of Pau da Lima in the municipality of Salvador-Bahia (northeastern Brazil), in an area (0.17 km<sup>2</sup>) geographically stratified in three valleys, with approximately 3717 inhabitants (Panti-May *et al.*, 2016) (Fig. 1). The community lives at the intersection of poverty and the environment, with poor-quality housing and inadequate sanitation infrastructure (Panti-May *et al.*, 2016; Carvalho-Pereira *et al.*, 2018). Transmission of other zoonotic diseases is well-documented within the community (Kikuti *et al.*, 2015; Hagan *et al.*, 2016), for example, leptospirosis, a rodent-borne zoonosis had an incidence of 35.4 cases per 1000 people/year (Hagan *et al.*, 2016).

### Study design

Sampling sites were selected from 108 locations that had been established for a previous study in the area with a minimum distance between locations of 20 m (Panti-May *et al.*, 2016). From these 108 locations, we randomly selected 40 gastropod sampling sites and 45 *R. norvegicus* sampling sites, 17 of which coincided, stratifying by the three valleys to ensure a similar number of sites in each valley. The sampling area for both taxa at each site consisted of a 15 m radius circle centred at the geotagged location.

Molluscs were sampled during a single sampling period between November 2016 and January 2017. Rat data were

obtained from four sampling periods spanning 2014–2016: two periods in the rainy season (March–July 2014 and March–April 2016) and two in the dry season (October–December 2014 and November–December 2015).

### Terrestrial gastropod sampling

Exhaustive mollusc sampling was performed by visual searching of the sampling sites, following the protocol developed by the Brazilian Ministry of Health (Ministério da Saúde, 2008). Sampling was carried out in the early morning (08:00–10:00) because the molluscs are more active and exposed during this time of day. The individuals collected were taken to the laboratory, where they were then identified using morphological characteristics, based on the literature (Salgado and dos Santos Coelho, 2003; Simone, 2006; Colley, 2012; Breure and Ablett, 2014).

### Processing of terrestrial mollusc samples and larval extraction

At each site, gastropods of the same species with a length of <2 cm were pooled for analysis (Carvalho *et al.*, 2012), while individuals >2 cm were dissected and analysed separately. Gastropods from each site were processed separately. The animals (or pools) were macerated and digested following Wallace and Rosen (1969), and adaptations of the technique of Rugai *et al.* (1954) for faecal samples, as it follows: the mollusc macerate was divided into portions of approximately 3 g, laid in six layers of gauze and distributed in sedimentation cones, for sedimentation with pepsin/HCl solution (4% pepsin 0.7% hydrochloric acid, in de-chlorinated water) at 42°C; then, the cones were incubated at 37°C for 2 h; after incubation, the sediment in the sediment cones was aspirated using a pipette, transferred to a Petri dish and observed under a binocular microscope at 40×. All larvae found were counted and fixed in 70% ethanol, with a subsample stored at –20°C for molecular analysis by PCR-RFLP (polymerase chain reaction-restriction fragment length polymorphism).

### Morphometrics and identification of *A. cantonensis* larvae by PCR-RFLP


Larvae with metastrongyloid characteristics [based on their morphology, mainly the tail conformation (Ash, 1970)] were identified, with 10% randomly selected for morphometric analysis. The following measurements were taken for each specimen: total body length, length of oesophagus, distance between excretory pore and anterior extremity, distance between genital primordium and tail, length of genital primordium and distance between anus and tail. The measurements were compared with the measurements from reference specimens of *A. cantonensis* and *A. costaricensis*, in Instituto Adolfo Lutz (IAL) by calculating the percentage of similar samples. All samples were sent for molecular confirmation (PCR-RFLP) of species identification as *A. cantonensis*, *A. costaricensis* or *A. vasorum*, following the method of Caldeira *et al.* (2003).

### Rat sampling and determination of *A. cantonensis* infection

Rats were captured using live rat-trapping, with two traps left at each location for 24 h on two consecutive days following previously established protocols (Panti-May *et al.*, 2016; Carvalho-Pereira *et al.*, 2018).

Demographic data (sex, age, reproductive status) and body condition [presence and severity of wounds and amount of fat deposits, as well as the Scaled Mass Index (SMI) (Peig and Green, 2009) – which accounts for the effect of age] were recorded.

Feces were collected directly from the large intestine and stored in 10% formaldehyde for subsequent identification of first-stage larvae of *A. cantonensis* following the protocol of



**Fig. 1.** Distribution of the presence of molluscs and intensity of *Angiostrongylus cantonensis* in rats, in Brazil. (A) The northeastern region of Brazil, where the state of Bahia and the city of Salvador are located. (B) Study area, health district of Pau da Lima, which has an elevated topography divided into three geographic valleys. B also represents the distribution of the presence of terrestrial molluscs and the intensity of *Angiostrongylus cantonensis* found in rats. *Angiostrongylus cantonensis* larvae were confirmed in terrestrial gastropods by molecular tests.

Carvalho-Pereira *et al.* (2018). Faecal sample analyses consisted of a qualitative stage followed by a quantitative stage. During the qualitative stage, we employed an adapted sedimentation method (Hoffman *et al.*, 1934), with larval identification based on morphological characteristics observed under light microscopy, with probable metastrongyloid nematodes indicated by observing the tail conformation, a particular characteristic of this group (Anderson, 1978). We also identified eggs of other parasites based on the morphology described in the literature (Vicente *et al.*, 1997). To confirm the species of identified parasite, we also searched for adult worms by collecting target tissues from the rodents captured during one of the sampling periods. Samples were taken from the intestines, stomach, heart and lungs, fixated in formalin–acetic acid–alcohol at 50°C and stored at 4°C. This allowed for the confirmation of occurrence of adults of *A. cantonensis* in captured *R. norvegicus* pulmonary arteries, as well as the other helminth species included in the study (see Additional file 1: Table S2). For the quantitative analysis, larvae per gram of feces were counted for *A. cantonensis* and eggs per gram of feces were counted for all other helminths. This was conducted using an adapted flotation method (Gordon *et al.*, 1939), using a saturated zinc sulphate solution with a density of 1.18 g cm<sup>-3</sup> (Faust *et al.*, 1938). Co-infection variables were created as the intensity of infection for each parasite species.

### Environmental variables

Environmental variables were selected for the survey by considering factors associated with mollusc abundance that reflects their physiological and ecological needs (Nunes and Santos, 2012). At all mollusc and rat sampling locations, environmental data were recorded. We observed data on the presence of food access, garbage access, rubbish, construction material, open sewage and water (standing or running). Variables related to habitat were

also collected: type of ground (permeable/impermeable) and vegetation (absent/herbaceous/shrubby). Additionally, during the capture of rodents, we recorded whether food and harbour sites for rats (pet food, human garbage, water sources and vegetation coverage by number of trees) were present within a radius of 10 m of the site. Additional information including the valley in which the samples were collected, the number of rat collection events and mollusc collection days (sampling effort: 2 or 3 days by two field technicians) was included as variables in this study. Daily pluviometry data were obtained from the station in the study area, managed by the Institute of Environment and Hydric Resources of Bahia (INEMA).

### Statistical analysis

The occurrence and abundance of mollusc species at each study site were determined, as was the prevalence and intensity (number of larvae) of *A. cantonensis* in each mollusc species identified. The prevalence of *A. cantonensis* in rats was then estimated for each sampling period, with comparisons between periods made by  $\chi^2$  tests.

To model the association between gastropod abundance and environmental factors, we used zero-inflated negative binomial models using the *R* function (R Core Team, 2013) *pscl* (Loeys *et al.*, 2012) to account for zero-inflation in the data. This analysis was performed separately for each gastropod species. Univariate analyses were performed to explore the relationship between each environmental variable and the abundance outcome, with variables statistically significant at the  $P \leq 0.1$  level included in the final multivariate model for each species. To account for the difference in the sampling effort between sites, the mollusc collection days was included in the models as a fixed effect.

To identify factors associated with the intensity of *A. cantonensis* in rats, we conducted a model selection process for the

three groups of (i) environmental, (ii) demographic and (iii) co-infection variables separately (see Additional file 1: Table S2). We fitted a zero-inflated generalized linear model with a *Gamma* error distribution to the intensity response variable, transformed to the base 2 logarithm to facilitate the interpretation of results (Carvalho-Pereira *et al.*, 2018). Modelling followed three steps: (1) univariate analysis of the environmental variables with variables included in the multivariate model if  $P \leq 0.1$ ; (2) then each rat demography and body condition variable was added one at a time into the multivariate environmental model and the same process of variable selection was repeated; (3) finally, each helminth co-infection intensity variable was added into the multivariate model selected in (2) and the same process was repeated for these variables. Final models of each step were defined for terrestrial molluscs and rats using model selection criteria based on Akaike's Information Criterion corrected for small samples (AICc) (Zuur and Ieno, 2016), with the help of the *MuMIn* package, which ranks models by AICc based on the importance of each explanatory variable included (Bartón, 2013). Thus, it was possible to identify all plausible models ( $\Delta\text{AICc} \leq 2.00$ ) and the most parsimonious, this being the model with the fewest explanatory variables among the plausible models.

## Results

We collected 577 gastropods, widely distributed in all valleys of the area studied, representing eight species from seven families. The most common species were *Subulina octona* (47.5%), *S. marginata* (30%), *A. fulica* (25%) and *Bulimulus tenuissimus* (22.5%) with the co-occurrence of the four species common in the sampled areas. Only these four species were found to be infected with *A. cantonensis*. *Bradybaena similaris*, *Drymaeus papyraceus*, *Helicina* sp. and a representative of the family Streptaxidae occurred at fewer sites and in lower abundance, with an average of five individuals per occurrence site.

The prevalence of *A. cantonensis* was 33, 11, 8 and 5% in *A. fulica*, *B. tenuissimus*, *S. marginata* and *S. octona*, respectively. *Achatina fulica* exhibited the highest number of larvae per individual, with an average of  $22.7 \pm 22.5$ . The remaining species exhibited lower levels of infection, with only one individual of *S. marginata* infected, with four larvae. *Bulimulus tenuissimus* and *S. octona*, predominantly analysed in pooled samples, exhibited the means of 0.5 and 0.3 larvae per individual, respectively.

Occurrence and abundance of mollusc species in the study area were associated with several local environmental variables, with the presence of water an important factor (see Additional file 1: Tables S3 and S4). Humidity and accumulated rain exhibited positive associations with the abundance of *A. fulica* and *S. octona*, respectively. The presence of construction materials was positively correlated with the number of *S. marginata*, which was significantly less abundant in valleys 2 and 3 compared to valley 1. In the univariate models (see Additional file 1: Table S3), the presence of herbaceous vegetation was positively correlated with the abundance of *S. octona*. On the other hand, areas with shrub vegetation exhibited higher *B. tenuissimus* numbers. The model including solely sampling effort was significantly associated with the presence of *B. tenuissimus*, indicating that a 3-day sampling effort was associated with a higher chance of finding individuals of *B. tenuissimus*, compared to 2-day efforts (Table 1). Only seven of the gastropod samples were infected with *A. cantonensis*. However, the infected samples occurred among four species (*A. fulica*, *B. tenuissimus*, *S. marginata* and *S. octona*) and in five sampling sites. Given the small number of infected gastropod samples, analysis of the risk factors associated with the presence and intensity of infection by *A. cantonensis* in the gastropod

species was disregarded. All 17 sites sampled for both rats and molluscs were positive for both taxa. In 45% of these sites, rats were positive for *A. cantonensis* (Fig. 1B).

In total, 168 *R. norvegicus* were trapped during the four sampling periods. Among them, sub-adults (103; 61.3%) were captured more frequently than the young (35; 20.8%) and adults (30; 17.9%), with more females (90; 53.9%) than males (77; 46.1%); and a similar number of captures in dry (85; 50.6%) and rainy seasons (83; 49.4%). First-stage *A. cantonensis* larvae were detected in 56 (33.3%) individuals, presenting an average of 90.00 (11.00) (Fig. 1) and with no significant difference between sampling periods. In addition to *A. cantonensis*, another seven helminth species already recorded by Carvalho-Pereira *et al.* (2018) were identified. Of these, the soil-transmitted nematode *Strongyloides* spp. (160; 95%) and *Nippostrongylus brasiliensis* (52; 30.9%) were the most prevalent.

Accumulated rainfall in the previous 2 weeks, as well as the presence of construction materials, was positively associated with a higher prevalence of *A. cantonensis* in *R. norvegicus* (Table 2). These relations were maintained when controlling for demographic and body condition variables (Table 2 – step 2; accumulated rain: rate 1.000, IC 95% 0.999–1.001; construction materials: 1.333, IC 95% 1.056–1.693). In this model, rat body condition (SMI) was significantly negatively associated with the probability of infection by *A. cantonensis*. Contrary to what was expected, however, the increase of age (in days) was negatively associated with infection intensity (Table 2). The final model, including coinfection variables, retained body condition as the sole significantly associated factor linked to the probability of infection. However, in terms of *A. cantonensis* infection intensity in rats, besides the accumulated rain, which was positively associated with greater *A. cantonensis* intensity, the intensity of *A. cantonensis* exhibited a negative association with that of *N. brasiliensis* and a positive association with that of *Strongyloides* spp.

## Discussion

This study demonstrated that in the impoverished urban area, studied environmental conditions that are correlated with the occurrence of terrestrial gastropods (intermediate hosts) are also indicators that rats (definitive hosts) are infected with *A. cantonensis*. The transmission cycle of *A. cantonensis* can be established and maintained in these contexts because these environmental conditions (available water and accumulation of inorganic matter such as construction materials) create microhabitats that favour both mollusc and rat abundance, with molluscs a food source for rats within these shared environments. The occurrence of several mollusc species in most of the studied area and lack of seasonality in rat prevalence across the two study years suggest that transmission of *A. cantonensis* is stable within the study area. This demonstrates the need for preventive action given the severity of human neuroangiostrongyliasis, in spite of the current lack of recorded cases of the disease in the community. This may, in turn, be a result of misdiagnosis because the main symptoms (e.g. headache, pain in the neck and fever) (Morassutti *et al.*, 2014) can be easily mistaken for other illnesses that co-occur in impoverished urban areas with abundant rats and molluscs.

Many mollusc species have been reported as susceptible to *A. cantonensis* infection and capable of maintaining transmission (Kim *et al.*, 2014). However, usually only a few species are considered as the primary intermediate hosts in a given region (Wang *et al.*, 2008). The presence of the species *S. octona*, *A. fulica*, *S. marginata*, *B. tenuissimus* and *B. similaris* (all characterized as *A. cantonensis* hosts in the literature) may indicate the occurrence of

**Table 1.** Final zero-inflated negative binomial regression models for terrestrial molluscs

Species	Predictors	IRR	CI	Sig.
	(Intercept)	$1.11 \times 10^{-7}$	$3.8 \times 10^{-13}$ – $1.2 \times 10^{-2}$	*
	Humidity	<b>1.245</b>	<b>1.070–1.456</b>	**
<i>Achatina fulica</i>	<b>Zero-inflated model</b>			
	(Intercept)	1.276	$7.2 \times 10^{-6}$ –5.484	
	Humidity	0.987	0.835–1.167	
	Sample effort			
	(Intercept)	4.823	0.418–9.296	***
	2 days (Ref.)	–	–	
	3 days	1.000	0.418–2.389	
<i>Bulimulus tenuissimus</i>	<b>Zero-inflated model</b>			
	Sample effort			
	(Intercept)	6.235	2.146–18.114	***
	2 days (Ref.)	–	–	
	3 days	<b>0.149</b>	<b>0.027–0.794</b>	*
	(Intercept)	4.003	2.296–6.977	***
	Construction material	<b>4.696</b>	<b>2.680–8.227</b>	***
	Geographic valley			
	Valley 1 (Ref.)	–	–	
<i>Sarasinula marginata</i>	Valley 2	<b>0.522</b>	<b>0.317–0.858</b>	**
	Valley 3	<b>0.012</b>	<b>0.000–0.446</b>	**
	<b>Zero-inflated model</b>			
	(Intercept)	2.190	0.518–9.264	
	Construction material	0.386	0.047–3.176	
	Geographic valley			
	Valley 1 (Ref.)	–	–	
	Valley 2	0.465	0.072–3.015	
	Valley 3	0.611	$0.000–2.97 \times 10^4$	
	(Intercept)	8.683	5.911–12.756	***
	Cumulative rain	<b>1.426</b>	<b>1.087–1.871</b>	*
	Sample effort			
	2 days (Ref.)	–	–	
	3 days	1.054	0.568–1.955	
<i>Subulina octona</i>	<b>Zero-inflated model</b>			
	(Intercept)	1.732	0.789–3.802	
	Cumulative rain	2.283	0.461–11.313	
	Sample effort			
	2 days (Ref.)	–	–	
	3 days	0.014	0.000–1.132	

IRR, incidence rate ratios; CI, confidence intervals; Sig., significance codes: 0; '\*\*\*\*' 0.001; '\*\*\*' 0.01; '\*\*' 0.05; '#' 0.1; '.' 1. The significant variables are shown in bold.

more contamination foci, given that these species can occupy different habitats. This is the first record of *B. tenuissimus* as a natural intermediate host of *A. cantonensis* in the state of Bahia. *Bradybaena similaris*, although previously reported as a host in Brazil (Carvalho et al., 2012), was not positive for *A. cantonensis* infection in this study, which is consistent with the findings of Kim et al. (2014). Overall, the study demonstrated a diversity of positive intermediate host species and that the co-occurrence of more than one species infected by *A. cantonensis* in a sampling

site is common. This could favour the development of larvae to the infective phase, which could promote continuous dissemination.

We observed in this study similarity in the distribution of highly infected rats and of infected molluscs, although geospatial analyses are still needed. Other characteristics of each study valley that were not measured could also contribute to the uneven distribution of mollusc species. For example, compared to valleys 2 and 3, valley 1 exhibited (based on unquantified observation) more poor sanitary conditions and more potential faunal refuges

**Table 2.** Final zero-inflated regression models with Gamma family for *A. cantonensis* infection in *Rattus norvegicus*

	Predictors	IRR	CI	Sig.
Environment ( <i>n</i> = 129)				
	(Intercept)	4.260	2.820–6.436	***
	Water			
	Absent (Ref.)	–	–	
	Running water	1.049	0.764–1.440	
	Standing water	0.797	0.542–1.171	
	Cumulative rain	<b>1.001</b>	<b>1.000–1.002</b>	*
Step 1	Construction materials	<b>1.359</b>	<b>1.060–1.743</b>	**
	<b>Zero-inflated model</b>			
	(Intercept)	0.578	0.181–1.840	
	Water			
	Absent (Ref.)	–	–	
	Running water	<b>3.515</b>	<b>1.366–9.042</b>	**
	Standing water	<b>4.849</b>	<b>1.468–16.018</b>	**
	Cumulative rain	1.000	0.997–1.003	
	Construction materials	0.957	0.404–2.263	
Step 1 + demography and body condition ( <i>n</i> = 129)				
	(Intercept)	6.519	3.909–10.872	***
	Age (days)	<b>0.995</b>	<b>0.992–0.999</b>	**
	Cumulative rain	<b>1.000</b>	<b>0.999–1.001</b>	*
	Construction materials	<b>1.333</b>	<b>1.050–1.693</b>	**
Step 2	Smi	1.000	0.997–1.002	
	<b>Zero-inflated model</b>			
	(Intercept)	0.591	0.104–3.354	
	Age (days)	0.993	0.984–1.002	
	Cumulative rain	0.997	0.994–1.000	
	Construction materials	0.800	0.346–1.848	
	Smi	<b>1.009</b>	<b>1.001–1.017</b>	*
Step 2 + co-infection ( <i>n</i> = 119)				
	(Intercept)	2.501	1.219–5.130	**
	Age (days)	<b>0.995</b>	<b>0.991–0.999</b>	*
	Cumulative rain	<b>1.001</b>	<b>1.000–1.001</b>	*
Step 3 – final model	Smi	1.000	0.997–1.002	
	Db <i>N. brasiliensis</i> intensity	<b>0.871</b>	<b>0.801–0.947</b>	***
	Db <i>Strongyloides</i> sp. intensity	<b>1.164</b>	<b>1.100–1.232</b>	***
	Db <i>N. brasiliensis</i> intensity: male (Ref.)	–	–	
	Db <i>N. brasiliensis</i> intensity: female	<b>1.122</b>	<b>1.026–1.226</b>	*
	<b>Zero-inflated model</b>			
	(Intercept)	0.217	0.016–2.836	
	Age (days)	0.992	0.982–1.003	
	Cumulative rain	0.997	0.994–1.000	
	Smi	<b>1.011</b>	<b>1.002–1.020</b>	**
	Db <i>N. brasiliensis</i> intensity	1.041	0.864–1.255	
	Db <i>Strongyloides</i> sp. intensity	1.072	0.884–1.301	
	Db <i>N. brasiliensis</i> intensity: male (Ref.)	–	–	
	Db <i>N. brasiliensis</i> intensity: female	0.906	0.719–1.143	

IRR, incidence rate ratios; CI, confidence intervals; Db, 'double'; Sig., significance codes: 0; '\*\*\*\*' 0.001; '\*\*\*' 0.01; '\*\*' 0.05; '#' 0.1; '.' 1. The significant variables are shown in bold.

(e.g. construction material), which could contribute to the mollusc's abundance in the environment. *Achatina fulica*, a generalist species, was the most abundant and an important host of *A. cantonensis*, as found in other studies (Barratt *et al.*, 2016; Thiengo *et al.*, 2010). The abundance of *A. fulica* may be due to it being a generalist invasive species in Brazil. Humid habitats may contribute to its increased abundance (Miranda *et al.*, 2015), which may increase the chances of infection by *A. cantonensis*.

The prevalence of *A. cantonensis* in our samples of the *R. norvegicus* in our study (33% of 116 specimens) was moderate, similar to a previous study in the area (Carvalho-Pereira *et al.*, 2018), but additionally stable among seasons. Other studies in Brazil recorded prevalences of 11% of 19 rats in the state of Pará (Moreira *et al.*, 2013) and of 71% of 114 rats in the state of Rio de Janeiro (Simões *et al.*, 2014). Stable prevalences between seasons indicate that there may be continuous contamination of the environment with *A. cantonensis* through rat feces, and potential year-round transmission, similar to the results of Simões *et al.* (2014) – which considered a larger sampling scale. Both regions of Bahia and Rio de Janeiro have tropical climates, with limited variation in temperature and rainfall throughout the year (Climate-Data.org), which may explain both the prevalences found and their stability between the dry and rainy seasons.

The majority of relationships found in the demographic and body condition analysis were expected (Carvalho-Pereira *et al.*, 2018, 2019). However with regard to the association with age, as the present study used cross-sectional data, a possibility is that older rats had been previously exposed to the nematode, and, consequently, had developed the capacity to modulate the parasite load when re-infected, thus exhibiting a lower infection rate (Au and Ko, 1979). In terms of coinfection associations, there was a different pattern than that previously found given that, when analysing rats stratified by sex, females presented with a positive association between the intensity of *N. brasilienses* and *A. cantonensis* (see Carvalho-Pereira *et al.*, 2019). This may be because pregnant or lactating females allocate fewer resources to fending off parasites, thus being more prone to infections and coinfections (Spickett *et al.*, 2017; Castillo and Paller, 2018).

This study was carried out in an impoverished urban community, which is an environment characterized by precarious sanitation services, a lack of urban infrastructure such as pavement, and the presence of points with rubbish containing construction materials and trash. Although the accumulation of construction materials was not retained in the final model for *A. cantonensis* infection in rats, this was considered a factor associated with gastropod abundance, probably by fostering the formation of preferred shelter. The absence of pavement was not identified as a risk factor for rat infection or for the abundance of molluscs, despite it being previously described as an important factor for the establishment of rodent populations, given their habit of digging burrows in soil (Cavia *et al.*, 2009). The absence of pavement can also contribute to the accumulation of rainwater in the soil, a risk factor detected by our study, which may explain why accumulated rain of the previous 2 weeks was an important predictor of infection intensity by *A. cantonensis* in rats and abundance of molluscs. This is likely to be because it creates humid, favourable environments for the molluscs. This could be related to a potential contribution of this phenomenon to the viability of *A. cantonensis* larvae in the environment and may favour infection in intermediate hosts (by contact with water contaminated with rodent feces) and rats (by ingestion of contaminated water or infected molluscs).

The integration of ecological mollusc and rodent data may be the key to understanding the establishment and transmission of *A. cantonensis* in human populations. The variation in environmental conditions within the urban landscape, especially in

areas with precarious sanitation, poor living and infrastructure conditions, may facilitate the existence of heterogeneous microhabitats (Sólymos *et al.*, 2009; Astor *et al.*, 2017) affecting the composition, abundance and distribution of the terrestrial mollusc assemblage. Microhabitats that favour an elevated abundance of molluscs may contribute to an increase in the invertebrate hosts' contact with the parasite, and consequently increase the number of infected rats. In contrast, if environmental changes occur, the equilibrium of the microhabitats may be upset, affecting the target animal populations (Feng and Himsworth, 2014), and probably resulting in reduced *A. cantonensis* transmission. This highlights the importance of carrying out structural and sanitation interventions to help prevent the transmission of *A. cantonensis* between hosts and to the people of the area. Local and short-term measures (e.g. cleaning yards and empty lots, reduction of construction material) and long-term actions (e.g. structural urban interventions that reduce water accumulation in the environment, such as street paving) may disrupt the parasite cycle already established in areas lacking urban infrastructure and with poor sanitary conditions.

One potential limitation of this study is the time interval between the collection of rat and mollusc data. However, we believe that this will have had a limited impact on our results because rat abundance did not vary seasonally (Panti-May *et al.*, 2016). Additionally, other studies suggest that mollusc abundance does not vary through time in the tropical urban context (Oliveira *et al.*, 2015). Another factor that could have limited the identification of environmental factors associated with the mollusc species – among the ones defined in this study – is that some of the collected species are generalists, occupying several habitats that vary from humid to dry and hot, while others may depend on specific variables that were not accounted for in this study (Hylander *et al.*, 2005; Dias *et al.*, 2007; Nunes and Santos, 2012).

In conclusion, this study indicates that the transmission cycle of the rat lungworm, *A. cantonensis*, is well established in an impoverished urban community in Brazil with evidence of its presence in definitive hosts and five species of intermediate hosts. Overall, our findings indicate the necessity for the involvement of government entities in the execution of structural and sanitation interventions, as well as the involvement of public health and zoonoses control agencies that can inform and guide residents with educational and participative actions regarding the risks associated with *A. cantonensis* and the potential of molluscs and rats as hosts of an array of zoonotic parasites. Thus, not only would neuroangiostrongyliasis, but also other relevant zoonosis, such as leptospirosis (transmitted by rat urine) and schistosomiasis (transmitted by snails) be prevented.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182021000597>

**Acknowledgements.** We thank the residents of the Pau da Lima community, Salvador, Bahia, Brazil, and anonymous reviewers for their valuable recommendations.

**Financial support.** This research was supported by projects funded by the Oswaldo Cruz Foundation and Secretariat of Health Surveillance, Brazilian Ministry of Health; the Federal University of Bahia; the National Institutes of Health of the United States (grant numbers F31 AI114245, R01 AI052473, U01 AI088752, R01 TW009504 and R25 TW009338); Medical Research Council (MR/P024084/1 and MR/T029781/1); and by the Wellcome Trust (102330/Z/13/Z and 218987/Z/19/Z). F.N.S. and C.G.Z. participated in this study under a FAPESB doctorate scholarship.

**Conflict of interest.** None.

**Ethical standards.** The rodent sampling component of the study used protocol 003/2012, which was approved by the Ethical Committee of the Animal Use (CEUA), of the Gonçalo Moniz Institute (IGM) – Oswaldo

Cruz Foundation (Fiocruz). The license for the collection of terrestrial molluscs in an urban area of Salvador-Bahia was provided by the Chico Mendes Institute for Biodiversity Conservation (ICMbio), a federal agency linked to the Brazilian Ministry of the Environment. None of the specimens collected belong to threatened or protected species.

## References

- Anderson RC (1978) Keys to genera of the superfamily Metastrongyloidea. CIH keys to the nematode parasites of vertebrates, No. 5. Commonwealth Agricultural Bureau, Farnham Royal, Bucks, UK.
- Ash LR (1970) Diagnostic morphology of the third-stage larvae of *Angiostrongylus cantonensis*, *Angiostrongylus vasorum*, *Aelurostrongylus abstrusus*, and *Anaflaroides rostratus* (Nematoda: Metastrongyloidea). *The Journal of Parasitology* **56**, 249–253.
- Astor T, Proschwitz T, Strengbom J, Berg MP and Bengtsson J (2017) Importance of environmental and spatial components for species and trait composition in terrestrial snail communities. *Journal of Biogeography* **44**, 1362–1372.
- Au ACS and Ko RC (1979) Changes in worm burden, haematological and serological response in rats after single and multiple *Angiostrongylus cantonensis* infections. *Zeitschrift für Parasitenkunde* **58**, 233–242.
- Barbosa TA, Thiengo SC, Fernandez MA, Graeff-Teixeira C, Morassutti AL, Mourão FRP, Miranda COS, Jorge MM, Costa LF and Gomes SR (2020) Infection by *Angiostrongylus cantonensis* in both humans and the snail *Achatina (Lissachatina) fulica* in the city of Macapá, in the Amazon Region of Brazil. *Memorias do Instituto Oswaldo Cruz* **115**, e200115.
- Barratt J, Chan D, Sandaradura I, Malik R, Spielman D, Lee R, Marriott D, Harkness J, Ellis J and Stark D (2016) *Angiostrongylus cantonensis*: a review of its distribution, molecular biology and clinical significance as a human pathogen. *Parasitology* **143**, 1087–1118.
- Bartón K (2013) Model selection and model averaging based on information criteria (AICc and alike). *The Comprehensive R Archive Network* **1**, 13.
- Breure ASH and Ablett JD (2014) Annotated type catalogue of the Bulimulidae (Mollusca, Gastropoda, Orthalicoidea) in the Natural History Museum, London. *ZooKeys* **392**, 1–367. doi: 10.3897/zookeys.392.6328
- Caldeira RL, Carvalho OS, Mendonça CL, Graeff-Teixeira C, Silva MCF, Ben R, Maurer R, Lima WS and Lenzi HL (2003) Molecular differentiation of *Angiostrongylus costaricensis*, *A. cantonensis*, and *A. vasorum* by polymerase chain reaction-restriction fragment length polymorphism. *Memorias do Instituto Oswaldo Cruz* **98**, 1039–1043.
- Caldeira RL, Mendonça CL, Goveia CO, Lenzi HL, Graeff-Teixeira C, Lima WS, Mota EM, Pecora IL, de Medeiros AMZ and Carvalho OS (2007) First record of molluscs naturally infected with *Angiostrongylus cantonensis* (Chen, 1935) (Nematoda: Metastrongylidae) in Brazil. *Memorias do Instituto Oswaldo Cruz* **102**, 887–889.
- Carvalho-Pereira T, Souza FN, Santos LRN, Walker R, Pertile AC, de Oliveira DS, Pedra GG, Minter A, Rodrigues MG, Bahiense TC, Reis MG, Diggle PJ, Ko AI, Childs JE, da Silva EM, Begon M and Costa F (2018) The helminth community of a population of *Rattus norvegicus* from an urban Brazilian slum and the threat of zoonotic diseases. *Parasitology* **145**, 797–806.
- Carvalho-Pereira TSA, Souza FN, Santos LRN, Pedra GG, Minter A, Bahiense TC, Reis MG, Ko AI, Childs JE, Silva EM, Costa F and Begon M (2019) Coinfection modifies carriage of enzootic and zoonotic parasites in Norway rats from an urban slum. *Ecosphere (Washington, D.C.)* **10**(10), 1–14. doi: 10.1002/ecs2.2887
- Carvalho ODS, Scholte RGC, de Mendonça CLF, Passos LKJ and Caldeira RL (2012) *Angiostrongylus cantonensis* (Nematode: Metastrongyloidea) in molluscs from harbour areas in Brazil. *Memorias do Instituto Oswaldo Cruz* **107**, 740–746.
- Castillo DSC and Paller VGV (2018) Occurrence of *Angiostrongylus cantonensis* in rodents from the rice granary of the Philippines and associated risk factors for zoonotic transmission. *Journal of Parasitic Diseases: Official Organ of the Indian Society for Parasitology* **42**, 350–356.
- Cavia R, Cueto GR and Suárez OV (2009) Changes in rodent communities according to the landscape structure in an urban ecosystem. *Landscape and Urban Planning* **90**, 11–19.
- Červená B, Modrý D, Fecková B, Hrazdilová K, Foronda P, Alonso AM, Lee R, Walker J, Niebuhr CN and Malik R (2019) Low diversity of *Angiostrongylus cantonensis* complete mitochondrial DNA sequences from Australia, Hawaii, French Polynesia and the Canary Islands revealed using whole genome next-generation sequencing. *Parasites & Vectors* **12**, 241.
- Climate-Data.org. Salvador: Temperatura, Tempo e Dados climatológicos Salvador. Temperatura da água Salvador. Available at <https://pt.climate-data.org/america-do-sul/brasil/bahia/salvador-854/> (Accessed 19 June 2020).
- Colley E (2012) Molluscos terrestres e a malacologia paranaense: histórico e importância no cenário nacional. *Estudos de Biologia* **34**, 1–34. doi: 10.7213/estud.biol.6127
- Costa F, Porter FH, Rodrigues G, Farias H, de Faria MT, Wunder EA, Osikowicz LM, Kosoy MY, Reis MG, Ko AI and Childs JE (2014) Infections by *Leptospira interrogans*, *Seoul virus*, and *Bartonella* spp. among Norway rats (*Rattus norvegicus*) from the urban slum environment in Brazil. *Vector Borne and Zoonotic Diseases* **14**, 33–40.
- Cowie RH (2013) Biology, systematics, life cycle, and distribution of *Angiostrongylus cantonensis*, the cause of rat lungworm disease. *Hawai'i Journal of Medicine & Public Health*, **72**(6 suppl. 2), 6.
- Dias RJP, Bessa ECA and D'Ávila S (2007) Influence of substrate humidity on desiccation resistance capacity in *Subulina octona* (Mollusca, Subulinidae). *Brazilian Archives of Biology and Technology* **50**, 137–139.
- Faust EC, D'Antoni JS, Odom V, Miller MJ, Peres C, Sawitz W, Thomen LF, Tobie J and Walker HJ (1938) A critical study of clinical laboratory techniques for the diagnosis of protozoan cysts and helminth eggs in feces. *The American Journal of Tropical Medicine and Hygiene* **s1–18**, 169–183.
- Feng AYT and Himsforth CG (2014) The secret life of the city rat: a review of the ecology of urban Norway and black rats (*Rattus norvegicus* and *Rattus rattus*). *Urban Ecosystems* **17**, 149–162.
- Gordon HM and Whitlock HV (1939) A new technique for counting nematode eggs in sheep faeces. *Journal of the Council for Scientific and Industrial Research* **12**, 50–52.
- Hagan JE, Moraga P, Costa F, Capian N, Ribeiro GS, Wunder EA Jr, Felzemburgh RDM, Reis RB, Nery N, Santana FS, Fraga D, Dos Santos BL, Santos AC, Queiroz A, Tassinari W, Carvalho MS, Reis MG, Diggle PJ and Ko AI (2016) Spatiotemporal determinants of urban leptospirosis transmission: four-year prospective cohort study of slum residents in Brazil. *PLoS Neglected Tropical Diseases* **10**, e0004275.
- Hoffman WA, Pons JA and Janer JL (1934) The sedimentation-concentration method in *Schistosomiasis mansoni*. *Puerto Rico Journal of Public Health and Tropical Medicine* **9**, 283–298.
- Hylander K, Nilsson C, Jonsson GB and Göthner T (2005) Differences in habitat quality explain nestedness in a land snail meta-community. *Oikos* **108**, 351–361.
- Kikuti M, Cunha GM, Paploski IAD, Kasper AM, Silva MMO, Tavares AS, Cruz JS, Queiroz TL, Rodrigues MS, Santana PM, Lima HCAV, Calcagno J, Takahashi D, Gonçalves AHO, Araújo JMG, Gauthier K, Diuk-Wasser MA, Kitron U, Ko AI, Reis MG and Ribeiro GS (2015) Spatial distribution of dengue in a Brazilian urban slum setting: role of socioeconomic gradient in disease risk. *PLoS Neglected Tropical Diseases* **9**, e0003937.
- Kim JR, Hayes KA, Yeung NW and Cowie RH (2014) Diverse gastropod hosts of *Angiostrongylus cantonensis*, the rat lungworm, globally and with a focus on the Hawaiian Islands. *PLoS ONE* **9**, e94969.
- Loeys T, Moerkerke B, De Smet O and Buysse A (2012) The analysis of zero-inflated count data: beyond zero-inflated Poisson regression. *The British Journal of Mathematical and Statistical Psychology* **65**, 163–180.
- Masi E, Pino FA, Santos MGS, Genehr L, Albuquerque JOM, Bancher AM and Alves JCM (2010) Socioeconomic and environmental risk factors for urban rodent infestation in Sao Paulo, Brazil. *Journal of Pest Science* **83**, 231–241.
- Ministério da Saúde (2008) *Vigilância e Controle de Moluscos de Importância Epidemiológica: Diretrizes Técnicas: Programa de Vigilância e Controle da Esquistossomose (PCE)*. Ministério da Saúde-Brasil.
- Miranda MS, Fontenelle JH and Pecora IL (2015) Population structure of a native and an alien species of snail in an urban area of the Atlantic Rainforest. *Journal of Natural History* **49**, 19–35.
- Morassutti AL, Thiengo SC, Fernandez M, Sawanyawisuth K and Graeff-Teixeira C (2014) Eosinophilic meningitis caused by *Angiostrongylus cantonensis*: an emergent disease in Brazil. *Memorias do Instituto Oswaldo Cruz* **109**, 399–407.
- Moreira VLC, Giese EG, Melo FTV, Simões RO, Thiengo SC, Maldonado A and Santos JN (2013) Endemic angiostrongyliasis in the Brazilian Amazon: Natural parasitism of *Angiostrongylus cantonensis* in *Rattus rattus* and *Rattus norvegicus*, and sympatric giant African land snails, *Achatina fulica*. *Acta Tropica* **125**, 90–97.



- Nunes GKM and Santos SB** (2012) Environmental factors affecting the distribution of land snails in the Atlantic Rain Forest of Ilha Grande, Angra dos Reis, RJ, Brazil. *Brazilian Journal of Biology* **72**, 79–86.
- Oliveira AP, Gentile R, Maldonado Júnior A, Lopes Torres EJ and Thiengo SC** (2015) *Angiostrongylus cantonensis* infection in molluscs in the municipality of São Gonçalo, a metropolitan area of Rio de Janeiro, Brazil: role of the invasive species *Achatina fulica* in parasite transmission dynamics. *Memorias do Instituto Oswaldo Cruz* **110**, 739–744.
- Onyishi GC, Aguzie IO, Okoro JO, Nwani CD, Ezenwaji N, Oluah NS and Okafor FC** (2018) Terrestrial snail fauna and associated helminth parasites in a tropical semi-urban zone, Enugu State, Nigeria. *Pakistan Journal of Zoology* **50**, 1080–1085.
- Panti-May JA, Carvalho-Pereira TSA, Serrano S, Pedra GG, Taylor J, Pertile AC, Minter A, Airam V, Carvalho M, Júnior NN, Rodrigues G, Reis MG, Ko AI, Childs JE, Begon M and Costa F** (2016) A two-year ecological study of norway rats (*Rattus norvegicus*) in a Brazilian Urban Slum. *PLoS ONE* **11**, e0152511.
- Peig J and Green AJ** (2009) New perspectives for estimating body condition from mass/length data: the scaled mass index as an alternative method. *Oikos* **118**, 1883–1891.
- R Core Team** (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <http://www.R-project.org/>.
- Rugai E, Mattos T and Brisola AP** (1954) Nova técnica para isolar larvas de nematóides das fezes-modificação do método de Baermann. *Revista do Instituto Adolfo Lutz* **14**, 5–8.
- Salgado NC and dos Santos Coelho AC** (2003) Moluscos terrestres do Brasil (Gastrópodes operculados ou não, exclusive Veronicellidae, Milacidae e Limacidae). *Revista de Biología Tropical* **51**, 149–189.
- Simões RO, Júnior A, Olifiers N, Garcia JS, Bertolino AVFA and Luque JL** (2014) A longitudinal study of *Angiostrongylus cantonensis* in an urban population of *Rattus norvegicus* in Brazil: the influences of seasonality and host features on the pattern of infection. *Parasites & Vectors* **7**, 100.
- Simone LRL** (2006) *Land and Freshwater Molluscs of Brazil: An Illustrated Inventory on the Brazilian Malacofauna, Including Neighbor Regions of the South America, Respect to the Terrestrial and Freshwater Ecosystems*. Museu de Zoologia, Universidade de São Paulo.
- Sólymos P, Farkas R, Kemencei Z, Páll-Gergely B, Vilisics F, Nagy A, Kisfali M and Hornung E** (2009) Micro-habitat scale survey of land snails in dolines of the Alsó-hegy, Aggtelek National Park, Hungary. *Mollusca* **27**, 167–171.
- Spickett A, Junker K, Krasnov BR, Haukisalmi V and Matthee S** (2017) Helminth parasitism in two closely related South African rodents: abundance, prevalence, species richness and impinging factors. *Parasitology Research* **116**, 1395–1409.
- Thiengo SC, Maldonado A, Mota EM, Torres EJM, Caldeira R, Carvalho OS, Oliveira APM, Simões RO, Fernandez MA and Lanfredi RM** (2010) The giant African snail *Achatina fulica* as natural intermediate host of *Angiostrongylus cantonensis* in Pernambuco, northeast Brazil. *Acta Tropica* **115**, 194–199.
- Thiengo SC, de Oliveira Simões R, Fernandez MA and Júnior AM** (2013) *Angiostrongylus cantonensis* and rat lungworm disease in Brazil. *Hawai'i Journal of Medicine & Public Health* **72**(6 suppl. 2), 18.
- Valente R, Robles MDR, Navone GT and Diaz JI** (2018) *Angiostrongylus* spp. in the Americas: geographical and chronological distribution of definitive hosts versus disease reports. *Memorias do Instituto Oswaldo Cruz* **113**, 143–152.
- Vicente JJ, Rodrigues HDO, Gomes DC and Pinto RM** (1997) Nematóides do Brasil. Parte V: nematóides de mamíferos. *Revista Brasileira de Zoologia* **14**, 1–452.
- Wallace GD and Rosen L** (1969) Techniques for recovering and identifying larvae of *Angiostrongylus cantonensis* from molluscs. *Malacologia* **7**, 427–438.
- Wang Q-P, Lai D-H, Zhu X-Q, Chen X-G and Lun Z-R** (2008) Human angiostrongyliasis. *The Lancet Infectious Diseases* **8**, 621–630.
- Zuur AF and Ieno EN** (2016) *Beginner's Guide to Zero-Inflated Models with R*. United Kingdom: Highland Statistics Limited.