


Comparative assessment of DNA extraction procedures for *Ascaris* spp. eggsI.D. Amoah¹ , G. Singh¹, K. Troell², P. Reddy³, T.A. Stenström¹ and F. Bux¹

Research Paper

Cite this article: Amoah ID, Singh G, Troell K, Reddy P, Stenström TA, Bux F (2020).

Comparative assessment of DNA extraction procedures for *Ascaris* spp. eggs. *Journal of Helminthology* **94**, e78, 1–7. <https://doi.org/10.1017/S0022149X19000683>

Received: 5 March 2019

Revised: 14 June 2019

Accepted: 5 July 2019

Key words:

Soil-transmitted helminths; *Ascaris suum*; qPCR; NanoDrop; Qubit; PCR inhibitors; DNA extraction kits

Author for correspondence:

I.D. Amoah, E-mail: amoahkid@gmail.com

¹Institute for Water and Wastewater Technology, Durban University of Technology, PO Box 1334, Durban 4000, South Africa; ²Department of Microbiology, National Veterinary Institute, SE-751 89, Uppsala, Sweden and

³Department of Community Health Studies, Faculty of Health Sciences, Durban University of Technology, PO Box 1334, Durban 4000, South Africa

Abstract

A central and critical step in the molecular detection of soil-transmitted helminths from environmental sources is the extraction of DNA from the eggs. In this study, we investigated the yield of DNA extracted from known quantities (500, 100, 50, 20, 10 and 5) of *Ascaris suum* eggs, as well as directly from wastewater and sludge samples containing *Ascaris* spp. eggs, using six commercial DNA extraction kits. The amount of DNA extracted was quantified with NanoDrop, Qubit and Ct values from quantitative polymerase chain reaction (qPCR) assay using CFX96 Touch™ real-time PCR equipment. The PowerLyzer Ultraclean Microbial DNA isolation kit and PowerSoil DNA isolation kit gave the highest yield of DNA based on the NanoDrop, Qubit and Ct values. However, the qPCR results indicate that in some of the kits, PCR inhibitors may have been carried over to the PCR reaction. DNA extraction kits that incorporate a bead-beating step as well as other mechanical eggshell disruption steps were superior in extracting DNA from *Ascaris* spp. eggs. Additionally, for the accurate quantification of extracted DNA, the use of Ct values from qPCR and Qubit readings gives better results compared to the NanoDrop readings. For efficient downstream applications, the use of DNA extraction kits with superior inhibitor removal technology is essential, in addition to a high yield of DNA.

Introduction

The main route of infection by soil-transmitted helminths (STHs) is through exposure to contaminated soil, water and food (Grimes *et al.*, 2016; Rostami *et al.*, 2016). The common conventional methods for their detection in environmental samples are based on sedimentation and/or flotation, aimed to quantitatively separate and concentrate the eggs before final microscopic identification and quantification (Collender *et al.*, 2015; Amoah *et al.*, 2017). Microscopic examination of eggs is prone to errors – for instance, in samples with low egg concentrations, false negatives or under-estimation of the egg counts could occur (Verweij *et al.*, 2007). This lack of sensitivity in microscopy is also due to the difficulty in morphological differentiation and speciation of STH eggs (Valero *et al.*, 2009; Ai *et al.*, 2010) – for instance, hookworm eggs of different genera and species are indistinguishable based on egg morphology (Gordon *et al.*, 2015). Another challenge is the presence of debris from the sample on the microscope slide that may interfere with the viewing.

The advent of polymerase chain reaction (PCR) methods has led to a more sensitive detection of different microorganisms, including STHs in environmental samples (Basuni *et al.*, 2011). The method is highly sensitive and specific in detecting *Strongyloides stercoralis* and hookworms as compared to microscopy (Verweij *et al.*, 2007, 2009). Despite the advantages with nucleic-acid-based methods for the detection and quantification of STH eggs in environmental samples, the main challenge is the extraction of consistent quantities and good quality of nucleic acid material from these eggs. A low-yielding DNA extraction method may also lead to false negative results, especially in environmental samples where low egg concentrations occur (Salonen *et al.*, 2010). The main challenge associated with nucleic acid extraction from helminthic eggs is due to the tough protective eggshell. For instance, the eggshell of *Ascaris* spp. has four layers, composed of a uterine layer (a glycoprotein), followed by a thin vitelline proteinaceous layer, a chitinaceous layer and then the innermost lipid layer (termed the ‘ascaroside membrane’) (Quilès *et al.*, 2006). These combined layers make the extraction of the nucleic material difficult. To overcome the protective barrier of the tough eggshell, harsh extraction conditions (such as sonication or bead beating) may sometimes be incorporated, but this may also lead to the shearing of the extracted DNA, negatively impacting downstream applications such as PCR. Commercial kits are commonly used for extracting nucleic acid from STH eggs (Pecson *et al.*, 2006; Gyawali *et al.*, 2017; Luo *et al.*, 2017). These are, however, not optimized explicitly for helminth eggs and may vary between manufacturers

in respect to cost, type of sample, sample processing time, amount of sample required, their ability to recover nucleic acids (Boesenberg-Smith *et al.*, 2012) and removal of PCR inhibitors (Miller *et al.*, 1999; Lakay *et al.*, 2006). Extraction of DNA from environmental samples, such as wastewater and sludge, are further affected by the complex sample matrices. A variable mixture of components may be present, including different proteins, lipids and humic acids, which may have an inhibitory effect on downstream analysis such as quantitative PCR (qPCR) if they follow-through in the extraction (Hall *et al.*, 2013; Josefsen *et al.*, 2015).

In this study, DNA from *Ascaris* spp. eggs was extracted using six commercial kits, and the amount of DNA was quantified using NanoDrop, Qubit and Ct values from qPCR performed on the extracted DNA. *Ascaris* spp. eggs were used as a surrogate for STHs because eggs of this helminth are the most resilient (Naidoo *et al.*, 2016) and have been applied as an indicator pathogen in wastewater and sludge. In addition, morphologically and structurally, the eggs of the two most common *Ascaris* spp. (*A. lumbricoides* and *A. suum*) are indistinguishable, whereby similar challenges are encountered in extracting DNA from the eggs of both species.

Materials and methods

Preparation of egg concentrations

Ascaris suum eggs were purchased from Excelsior Sentinel Inc. (Ithaca, NY). These eggs were initially stored in formalin to inhibit their development. A working solution with approximately 500 eggs was prepared in distilled water. These were counted microscopically under $\times 100$ magnification (Leica DM1000, Leica Microsystems). The stock was then divided into six subsamples, resulting in approximately 83 eggs per extraction method/kit, and used for the comparison of DNA extraction kits. The limit of detection of the three best-performing extraction kits was determined by preparing eggs of different concentrations – 100, 50, 20, 10 eggs – which was done with a newly prepared stock solution and concentrations determined microscopically. DNA was also extracted from raw wastewater and sludge samples ($n = 9$ for each sample) from a conventional activated sludge wastewater treatment plant. The raw wastewater was taken at the influent point and sludge samples from fresh material on the sludge drying beds. These samples were assumed to contain the human parasite *Ascaris lumbricoides*, whereby the performance of the kits was assessed for their applicability in extracting DNA from this species as well. These samples were processed using a conventional STH egg-detection method, involving filtration and flotation (using Zinc sulphate with a specific gravity of 1.30). Briefly, 1 l of wastewater was filtered through sieves of sizes 100 and 20 μm ; for the sludge samples, 20 g was weighed and mixed thoroughly with saturated ammonium bicarbonate and filtered similarly to the wastewater samples. Deposits collected on the 20 μm sieve were then washed into a test tube and flotation with zinc sulphate was carried out. Eggs were then collected after flotation and concentrated eggs washed thoroughly with distilled water before DNA extraction. The eggs were extracted from each sample, and pooled together to get the total egg concentration per wastewater and sludge. The concentration and stage of development of the *Ascaris* spp. eggs in the wastewater and sludge samples was determined prior to DNA extraction through microscopic examination of the eggs. Table 1 shows the stage of development of the eggs that were used.

DNA extraction and quantification

DNA was extracted from the different egg concentrations in triplicate using the six commercial DNA extraction kits (listed in table 2) and extractions were done three times (nine extractions in all per extraction kit) each time with fresh set of quantified eggs, according to the manufacturer's instructions. These kits were selected based on a thorough literature search to find the most commonly used commercial kits for the extraction of DNA from STH eggs in environmental samples. Extracted DNA was quantified using the NanoDrop ND-1000 (Thermo Scientific, Waltham, MA, USA) according to the manufacturer's instructions using 1 μl of DNA sample. The Qubit 2.0 Fluorometer (Carlsbad, California, USA) was also used to quantify the extracted DNA, using the Qubit dsDNA High Sensitivity Assay Kit and a 10 μl sample volume.

qPCR

The forward primer 5'-GTA ATA GCA GTC GGC GGT TTC TT-3' and reverse primer 5'-GCC CAA CAT GCC ACC TAT TC-3' targeting the first Internal transcribed spacer (ITS-1) region of *Ascaris* spp. were adopted from Wiria *et al.*, (2010). The qPCR assay was performed using the CFX96 Touch™ Real-Time PCR Detection System (BIO-RAD, Hercules, CA, USA). Briefly, the reaction mixture contained Maxima SYBR Green/ROX qPCR Master Mix (12.5 μl), primers (0.4 μM each) and 5 μl DNA template in a final volume of 25 μl . Water was used instead of template in the negative control.

The qPCR amplification protocol for the targeted genes consists of initial denaturation for 5 min at 95°C, followed by 45 cycles of three steps consisting of 10 s at 95°C, 20 s at 58°C and 20 s at 72°C. The fluorescence signals were measured at the end of each extension step. Serial dilutions of the DNA were made up to 1:100 to check for inhibition of enzymatic reaction and all qPCR reactions were repeated at least three times (in triplicates). These dilutions were done to determine if we had any PCR inhibition in our reactions; therefore, only a few of these eggs were diluted and the test was repeated three times to give accurate information on the presence or otherwise of inhibitors based on the change in Ct values.

Statistical analysis

Statistical analysis on the total DNA yields was performed in GraphPadPrism version 7.0 (GraphPad Software Inc.) using one-way analysis of variance followed by Tukey's multiple comparisons test. A *P*-value of less than 0.05 was considered significant.

Results

Comparison of DNA extraction kits and methods of DNA quantification

All six DNA extraction kits yielded DNA of varying concentrations, as confirmed by the Ct values from qPCR reactions, as well as the NanoDrop and Qubit results (table 3). The qPCR reactions showed different quantities of template DNA as extracted by the six kits. PowerSoil and PowerLyzer Ultraclean DNA isolation kits gave the least Ct values, indicating higher template DNA concentrations than the other extraction kits, followed by the PowerLyzer PowerSoil DNA isolation kit. The least quantity of DNA was extracted with the QIAamp Fast DNA Stool Mini Kit (Ct value of 30.11 ± 0.08). Considering all three methods of

Table 1. The percentage of *Ascaris* spp. eggs per each extraction at different stages of development.

	Total egg count	Stage of development of eggs ^a				
		Percentage of eggs at 1 cell (<i>n</i>)	Percentage of eggs at 2–3 cells (<i>n</i>)	Percentage of eggs at 4–6 cells (<i>n</i>)	Percentage of eggs with ≥7 cells (<i>n</i>)	Percentage of eggs with larvae (<i>n</i>) ^b
Stock	500	92.8 (464)	1.6 (8)	2.4 (12)	1.5 (7.5)	1.7 (8.5)
Sludge	94	21.9 (20.6)	1.0 (0.9)	45.9 (43.2)	11.2 (10.5)	20.0 (18.8)
Wastewater	343	22.0 (75.5)	2.0 (6.9)	45.0 (154.3)	12.1 (41.5)	18.9 (64.8)

^aDetermined microscopically prior to DNA extraction.

^bIt is estimated that each larvated egg will contain about 600 cells (Pecson *et al.*, 2006).

DNA quantification, the PowerLyzer Ultraclean DNA, PowerSoil DNA and PowerLyzer PowerSoil DNA isolation kits were the best in descending order. However, it is worth noting that the PowerLyzer Ultraclean DNA isolation kit had an elution solution of 50 µl, which is half the amount required in comparison with the other extraction kits.

The Ct values after qPCR were statistically significant between the different kits, showing a variation in the amount of template DNA extracted. However, this statistical difference could also be attributed to the uneven distribution of larvated eggs in the samples. Eggs undergoing embryonation could have high copies of nucleic material and, therefore, influence the amount of DNA extracted. With the incorporation of a freeze and thaw step with extractions using the QIAamp Fast DNA Stool Mini Kit, there was an increase in the amount of DNA extracted, with NanoDrop readings of 7.1 ± 2.1 ng/µl and Qubit readings of 2.32 ± 0.24 ng/µl as compared to the 2.1 (± 1.6) and 0.18 (± 0.05) recorded.

Relationship between *A. suum* egg concentration and DNA yield

The three best DNA extraction kits (PowerLyzer Ultraclean DNA, PowerSoil DNA and PowerLyzer PowerSoil DNA isolation kits) have similar extraction steps and reagents (see table 2). Therefore, in the further extraction tests, the worst performing of these three (PowerLyzer PowerSoil DNA isolation kit) was replaced with the QIAamp Fast DNA Stool Mini Kit (Qiagen Inc.). This was to enable us to determine if the different extraction steps and reagents, based on difference in manufacturers (at the time of this study, Qiagen Inc. and MO BIO Laboratories were separate companies), would result in different amounts of extracted DNA when applied to a decreasing number of eggs, as well as in different sample matrices.

As expected, the measured DNA decreased when the egg concentrations were reduced. Large differences did occur between the different extraction kits, as can be seen in table 4. However, the removal of assumed inhibitors through dilution resulted in comparable good performance by the PowerSoil DNA isolation kit (table 4). The recorded CT values for the ten- and 100-fold dilutions for the 50 eggs using the PowerLyzer Ultraclean Microbial DNA isolation kit and the QIAamp Fast DNA Stool Mini Kit did not show the expected increase in CT value. Dilution of the extracted DNA from the 100 and 50 eggs of *A. suum* showed that with an increase in dilution, the amount of template DNA decreases with a corresponding decrease in PCR inhibitors, as based on the Ct values for the PowerLyzer Ultraclean isolation kit, and especially for the QIAamp Fast DNA Mini Kit (table 4).

Extraction of *Ascaris* spp. DNA from wastewater and sludge samples

The wastewater and sludge samples had mean *Ascaris* spp. egg concentrations of 94.6 (± 25.79) eggs/l and 343.4 (± 25.79) eggs/g, respectively (table 1). As expected, based on the number of eggs in these samples and given the similar percentage of these eggs being embryonated (table 1), the DNA yield from the sludge samples was higher than the yield from the wastewater samples, which was seen for all three of the extraction protocols (table 5). Ct values from the qPCR were as expected for the extraction protocols, except the QIAamp Fast DNA Stool Mini Kit, where a lower Ct value in the wastewater samples than in the sludge samples was found, as shown in table 5. In contrast to the comparisons made on the commercially purchased eggs, dilutions of DNA extracted from the wastewater and sludge samples showed an absence of enzymatic inhibition, which could rule out problems with inhibitor carryover from extraction to qPCR. However, the increase in the Ct values was above the expected 2–3 values. The PCR reactions were run three times and each time with freshly extracted DNA from the eggs.

Discussion

All six of the commercial kits were successful in extracting DNA from the *A. suum* eggs. Using the NanoDrop or Qubit to quantify the extracted DNA, the PowerLyzer Ultraclean Microbial DNA isolation kit gave the highest quantity of DNA yield. However, Ct values that directly correlate with the amount of template DNA in the sample were consistently lower, i.e. more template, in the samples extracted with the PowerSoil DNA isolation kit. The volume of elution solution used with the PowerLyzer Ultraclean Microbial DNA isolation kit was half that recommended for the other extraction kits. Therefore, the final DNA extracts will be more concentrated in the PowerLyzer Ultraclean Microbial DNA isolation kit due to the smaller volume of elution solution. Basing on the Ct values, the PowerSoil DNA isolation kit was the best in extracting the template DNA. It was observed that the type of DNA extraction method did not significantly affect the quantity of DNA extracted, with almost all the extraction kits resulting in low yield. However, kits that have a bead-beating step yielded slightly higher DNA quantities – for instance, the PowerSoil, PowerWater and PowerLyzer Ultraclean DNA isolation kits all make use of beads and these reported the highest yield of DNA. Several studies have established that DNA extraction kits that incorporate bead beating result in higher DNA yield as well as an increase in technical reproducibility (Ariefdjohan *et al.*, 2010; Salonen *et al.*, 2010; Josefsen *et al.*, 2015). In addition, Josefsen *et al.* (2015) showed that a bead-

Table 2. List of DNA extraction kits used in this study and their critical steps/reagents.

Parameter	Commercial kit					
	PowerWater DNA isolation kit (MO BIO Laboratories)	PowerSoil DNA isolation kit (MO BIO Laboratories)	PowerLyzer PowerSoil DNA isolation kit (MO BIO Laboratories)	PowerLyzer Ultraclean Microbial DNA isolation kit (MO BIO Laboratories)	PowerFecal DNA isolation kit (MO BIO Laboratories)	QIAamp Fast DNA Stool Mini Kit (Qiagen Inc.)
Cell lysis solution	Sodium dodecyl sulphate	Sodium dodecyl sulphate	Sodium dodecyl sulphate	Sodium dodecyl sulphate	Sodium dodecyl sulphate	InhibitEX buffer and proteinase K
Cell lysis technique	Vortex at maximum speed for 10 min with beads	Vortex at maximum speed for 10 min with beads	Vortex at maximum speed for 10 min with beads	Vortex at maximum speed for 10 min with beads	Vortex at maximum speed for 10 min with beads	Vortex at maximum speed for 10 min and incubation at 95°C
Protein removal	Patented inhibitor removal technology	Patented inhibitor removal technology	Patented inhibitor removal technology	Solution MD2	Patented inhibitor removal technology	InhibitEX buffer
Humic acid removal	Patented inhibitor removal technology	Patented inhibitor removal technology	Patented inhibitor removal technology	Inhibitor removal solution	Patented inhibitor removal technology	InhibitEX buffer
DNA precipitation	High-concentration salt solution	High-concentration salt solution	High-concentration salt solution	High-concentration salt solution	High-concentration salt solution	Ethanol
DNA purification	Spin filter with silica membrane	Spin filter with silica membrane	Spin filter with silica membrane	Spin filter with silica membrane	Spin filter with silica membrane	Spin filter with silica membrane
Elution buffer	10 mM Tris-HCl, pH 8.0	10 mM Tris-HCl, pH 8.0	10 mM Tris-HCl, pH 8.0	10 mM Tris-HCl, pH 8.0	10 mM Tris-HCl, pH 8.0	Buffer TAE (Buffer containing a mixture of Tris base, acetic acid and EDTA)
Final elution volume (µl)	100	100	100	50	100	100

Table 3. Mean concentration (\pm standard deviation) of extracted DNA (using 83 eggs) as measured by NanoDrop, Qubit and qPCR ($n=9$).

Protocol	DNA concentration		
	NanoDrop (ng/ μ l)	Qubit (ng/ μ l)	qPCR (Ct values)
PowerWater DNA isolation kit	3.73 (\pm 2.5)	0.26 (\pm 0.04)	27.54 (\pm 0.41)
PowerLyzer PowerSoil DNA isolation kit	2.83 (\pm 0.97)	1.18 (\pm 0.10)	26.33 (\pm 0.45)
PowerSoil DNA isolation kit	1.97 (\pm 0.26)	0.86 (\pm 0.15)	25.25 (\pm 0.09)
PowerLyzer Ultraclean Microbial DNA isolation kit	6.07 (\pm 0.51)	2.04 (\pm 0.08)	25.84 (\pm 0.05)
PowerFecal DNA isolation kit	0	0.71 (\pm 0.04)	28.66 (\pm 0.19)
QIAamp Fast DNA Stool Mini Kit	2.1 (\pm 1.6)	0.18 (\pm 0.05)	30.11 (\pm 0.08)

Table 4. Mean concentration (\pm standard deviation) of *Ascaris suum* DNA extracted from varying egg concentrations, as determined by NanoDrop (ng/ μ l) and qPCR (Ct values) after a series of dilutions ($n=9$).

Egg concentration	PowerSoil DNA isolation kit		PowerLyzer Ultraclean Microbial DNA isolation kit		QIAamp Fast DNA Stool Mini Kit	
	ng/ μ l	Ct value	ng/ μ l	Ct value	ng/ μ l	Ct value
100	4.2 (\pm 1.2)		12.47 (\pm 2.5)		1.37 (\pm 0.15)	
1:1		22.60 (\pm 0.24)		24.72 (\pm 0.63)		28.90 (\pm 0.26)
1:10		26.08 (\pm 0.23)		28.21 (\pm 0.21)		29.04 (\pm 0.45)
1:100		28.60 (\pm 0.34)		28.55 (\pm 0.52)		27.32 (\pm 0.41)
50	3.93 (\pm 2.5)		12 (\pm 1.4)		0.97 (\pm 0.35)	
1:1		23.12 (\pm 0.25)		25.55 (\pm 0.32)		28.28 (\pm 0.45)
1:10		26.06 (\pm 0.87)		28.03 (\pm 0.21)		28.90 (\pm 0.21)
1:100		28.55 (\pm 0.64)		28.87 (\pm 0.45)		28.92 (\pm 0.12)
20	0 (\pm 0)	23.72 (\pm 0.45)	10.57 (\pm 1.3)	26.36 (\pm 0.52)	0.47 (\pm 0.32)	30.77 (\pm 0.45)
10	0 (\pm 0)	24.53 (\pm 0.35)	10.1 (\pm 1.8)	26.31 (\pm 0.23)	0.6 (\pm 0.20)	31.73 (\pm 0.31)
5	0 (\pm 0)	22.48 (\pm 0.29)	7.4 (\pm 0.50)	26.46 (\pm 0.15)	0 (\pm 0)	32.02 (\pm 0.41)

beating step in the DNA extraction protocol using the PowerLyzer PowerSoil and PowerFecal DNA isolation kits resulted in twice the amount of total DNA from the bacterium *Campylobacter jejuni* cells, although the resistance of the cell wall was lower here. However, prolonged bead beating may result in the shearing of DNA extracts, which may affect downstream applications, especially for bacterial cells (Josefsen *et al.*, 2015). The bead-beating step also introduces an extra step, which increases the duration of extraction. When an additional cell disruption step using liquid nitrogen to freeze and thaw the eggs was introduced, the quantity of DNA extracted using the QIAamp Fast DNA Stool Mini Kit increased. Therefore, it can be concluded that the eggs are tough and may require an extra mechanical egg disruption step to increase DNA yield. Additionally, the difference in the concentration of DNA extracted using these kits could be attributed to the developmental stage of the eggs. In reference to table 1, some of the purified eggs purchased from a commercial supplier (1.7%) had embryonated, which significantly results in an increase in the cell counts and an increase in gene copies as a result (Pecson *et al.*, 2006). With such a low percentage of embryonated eggs, it is possible that a few of the samples may have contained larvated eggs, thus resulting in an uneven DNA quantity thereafter. One embryonated egg with larvae contained

approximately 600 cells (Pecson *et al.*, 2006) as compared to the majority (92.8%) of the eggs, which only had a single cell. Naturally, one of these embryonated cells will largely influence the outcome of the extraction in terms of template DNA for the qPCR. To overcome this problem, we performed each extraction three times, in triplicates, and the mean value was used.

Our results show that the choice of DNA extraction kit or protocol may play a critical role in the amount of DNA extracted or the carryover inhibitors, both being crucial for further applications such as qPCR. Although the NanoDrop approach has been used extensively for the quantification of nucleic acids, it is limited for very low nucleic acid concentrations as seen in this study. Therefore, for accurate quantification, the use of the Ct values gives better results. In addition, the NanoDrop does not differentiate between amplifiable or non-amplifiable DNA (Thermo Scientific, 2010). Therefore, relying on DNA concentrations quantified using the NanoDrop to determine the best extraction kit/method may result in inaccurate conclusions. For instance, based on the NanoDrop, the best DNA extraction kit was the PowerLyzer Ultraclean DNA isolation kit; the use of the Ct values, however, showed that the PowerSoil DNA extraction kit extracts the highest quantity of template DNA. In contrast, the Qubit technique, together with the Ct values, may give better

Table 5. Mean concentration (\pm standard deviation) of *Ascaris* spp. DNA, as measured by NanoDrop (ng/ μ l) and qPCR (Ct values) in wastewater and sludge ($n = 9$).

Sample matrix	PowerSoil DNA isolation kit		PowerLyzer Ultraclean Microbial DNA isolation kit		QIAamp Fast DNA Stool Mini Kit	
	ng/ μ l	Ct value	ng/ μ l	Ct value	ng/ μ l	Ct value
Wastewater						
1:1	0.89 (± 0.12)	22.69 (± 0.49)	1.24 (± 0.34)	23.71 (± 0.44)	0.8 (± 0.18)	24.90 (± 0.32)
1:10		28.29 (± 0.65)		28.46 (± 0.78)		29.69 (± 0.50)
1:100		32.57 (± 1.50)		35.15 (± 0.61)		35.96 (± 0.69)
Sludge						
1:1	2.38 (± 0.24)	20.53 (± 0.39)	8.91 (± 0.45)	23.37 (± 0.44)	0.97 (± 0.17)	28.45 (± 0.68)
1:10		24.54 (± 0.28)		26.28 (± 0.86)		31.24 (± 0.67)
1:100		27.74 (± 0.59)		29.65 (± 0.71)		32.38 (± 0.91)

quantification. The Qubit technique is based on the binding of intercalating dyes to double-stranded DNA molecules. The fluorescence of these bound dyes is then measured to determine the DNA concentration (Olson & Morrow, 2012). However, the DNA quantified using the Qubit method may not be amplifiable, but the Ct values gives a direct correlation with the amplifiable DNA content. Based on these Ct values, Qubit readings and the amount of PCR inhibitors, as determined through the dilution of the DNA extracts, the PowerSoil DNA isolation kit was seen to give the best result for the samples used in this study.

STH egg concentration in the environment is generally low, unless in endemic areas. Therefore, any DNA extraction method used must be able to extract DNA from a low number of eggs. Only the PowerLyzer Ultraclean DNA isolation kit yielded measurable DNA from as low as five eggs (the least number of eggs studied). However, qPCR was positive from the extracted samples, which clearly showed the presence of template DNA in all samples, which means all kits successfully extracted amplifiable DNA. The concentrations of STHs in wastewater vary from 0 eggs/l (Forsslund *et al.*, 2010; Abreu-Acosta & Vera, 2011) to 16,000 eggs/l (Yen-Phi *et al.*, 2010) depending on geographical location, and are an indication of the infection status of the populations. In sludge samples, concentrations are generally higher, although variable (Scaglia *et al.*, 2014). Therefore, the ability to extract DNA from eggs must cover a wide egg-concentration range.

The main challenge with the molecular detection of STH eggs from environmental samples is the presence of inhibitors that might interfere with the nucleic acid extraction and subsequent PCR applications. The Ct values from the qPCR indicates the presence of inhibitors in the DNA extracts, as can be seen from the results presented in table 4. These inhibitors can be from different sources and vary in type; however, they generally have an impact on the amplification of the DNA. The Ct values reported for the diluted DNA extracts in table 4 could most likely be due to DNA cross-linking and modification caused by the preservation of the eggs with formalin. These eggs were preserved with formalin and this has been reported to affect PCR reactions due to nucleic acid cross-linking, modification and decreased DNA quality (Gilbert *et al.*, 2007). DNA cross-linking may lead to susceptibility of oligonucleotides to mechanical stress and decreased accessibility to DNA polymerase (Dietrich *et al.*, 2013). Sludge and wastewater samples contain a lot of substances and materials,

including proteins, lipids, humic acids, etc., which have been shown to be PCR inhibitors (Tebbe & Vahjen, 1993; Sørensen *et al.*, 2002; Zhou & Thompson, 2002; Fortin *et al.*, 2004; Hall *et al.*, 2013). Surprisingly less, or no, inhibition was observed in the sludge and wastewater samples. The increase in average Ct values beyond the 2–3 folds expected may be attributed to the impact of an embryonated egg/s on the quantity of DNA extracted. In addition to being able to extract higher quantities of DNA, DNA extracts from the PowerSoil DNA extraction kit also had the least carryover of PCR inhibitors.

Conclusion

It can be concluded that DNA extraction kits that make use of bead-beating techniques (PowerSoil, PowerWater and PowerLyzer Ultraclean DNA isolation kits) have a higher yield of DNA, which can most likely be attributed to the mechanical effect of the beads on the eggshell. The worst-performing kit was the QIAamp Fast DNA Stool Mini Kit, which is based solely on enzymatic reactions. In addition, PCR inhibitor impact was less in the DNA extracts from the PowerSoil DNA extraction kit, attributed to a superior inhibitor removal technology. For the quantification of DNA extracted from *Ascaris* spp. eggs, and most probably all helminth eggs, the use of Ct values was also seen to be a better approach, compared to the NanoDrop.

Acknowledgements. We are grateful to the Institute for Water and Wastewater Technology and the Faculty of Health Sciences of the Durban University of Technology for their support.

Financial support. This work was funded by the Bill and Melinda Gates Foundation (grant number OPP1122681) and the Sweden (STINT)/South Africa (National Research Foundation) Bilateral Research Collaboration Programme for the support (grant number STINT150831140656). Additional funding was provided by the South African Research Chairs Initiative (SARChI) of the Department of Science and Technology and National Research Foundation of South Africa.

Conflicts of interest. None.

References

Abreu-Acosta N and Vera L (2011) Occurrence and removal of parasites, enteric bacteria and faecal contamination indicators in wastewater natural

- reclamation systems in Tenerife-Canary Islands, Spain. *Ecological Engineering* 37, 496–503.
- Ai L, Dong SJ, Zhang WY, Elsheikha HM, Mahmmod YS and Lin RQ** (2010) Specific PCR-based assays for the identification of *Fasciola* species: their development, evaluation and potential usefulness in prevalence surveys. *Annals of Tropical Medicine and Parasitology* 104, 65–72.
- Amoah ID, Singh G, Stenström TA and Reddy P** (2017) Detection and quantification of soil-transmitted helminths in environmental samples: a review of current state-of-the-art and future perspectives. *Acta Tropica* 169, 187–201.
- Ariefdjohan MW, Savaiano DA and Nakatsu CH** (2010) Comparison of DNA extraction kits for PCR-DGGE analysis of human intestinal microbial communities from fecal specimens. *Journal of Nutrition* 9, 23–31.
- Basuni M, Muhi J, Othman N, et al.** (2011) A pentaplex real-time polymerase chain reaction assay for detection of four species of soil-transmitted helminths. *American Journal of Tropical Medicine and Hygiene* 84, 338–343.
- Boesenberg-Smith KA, Pessaraki MM and Wolk DM** (2012) Assessment of DNA yield and purity: an overlooked detail of PCR troubleshooting. *Clinical Microbiology Newsletter* 34, 1–5.
- Collender PA, Kirby AE, Addiss DG, Freeman MC and Remais JV** (2015) Methods for quantification of soil-transmitted helminths in environmental media: current techniques and recent advances. *Trends in Parasitology* 31, 623–639.
- Dietrich D, Uhl B, Sailer V, Holmes EE and Jung M** (2013) Improved PCR performance using template DNA from formalin-fixed and paraffin-embedded tissues by overcoming PCR inhibition. *PLoS One* 8, e77771.
- Forslund A, Ensink JHJ, Battilani A, et al.** (2010) Faecal contamination and hygiene aspect associated with the use of treated wastewater and canal water for irrigation of potatoes (*Solanum tuberosum*). *Agricultural Water Management* 98, 440–450.
- Fortin N, Beaumier D, Lee K and Greer CW** (2004) Soil washing improves the recovery of total community DNA from polluted and high organic content sediments. *Journal of Microbiology Methods* 56, 181–191.
- Gilbert MT, Haselkorn T, Bunce M, Sanchez JJ and Lucas SB** (2007) The isolation of nucleic acids from fixed, paraffin-embedded tissues-which methods are useful when? *PLoS One* 2, e537.
- Gordon CA, McManus DP, Acosta LP, Olveda RM, Williams GM, Ross AG, Gray DJ and Gobert GN** (2015) Multiplex real-time PCR monitoring of intestinal helminths in humans reveals widespread polyparasitism in Northern Samar, the Philippines. *International Journal of Parasitology* 45, 477–483.
- Grimes JET, Tadesse G, Mekete K, et al.** (2016) School water, sanitation, and hygiene, Soil-Transmitted Helminths, and Schistosomes: national mapping in Ethiopia. *PLoS Neglected Tropical Diseases* 10(3), e0004515.
- Gyawali P, Ahmed W, Sidhu JP, Jagals P and Toze S** (2017) Quantification of hookworm ova from wastewater matrices using quantitative PCR. *Journal of Environmental Science* 57, 231–237.
- Hall TA, Zovanyi MA, Christensen DR, Koehler JW and Devins MT** (2013) Evaluation of Inhibitor-Resistant Real-Time PCR methods for diagnostics in clinical and environmental samples. *PLoS One* 8(9), e73845.
- Josefsen MH, Andersen SC, Christensen J and Hoorfar J** (2015) Microbial food safety: potential of DNA extraction methods for use in diagnostic metagenomics. *Journal of Microbiology Methods* 114, 30–34.
- Lakay FM, Botha A and Prior BA** (2006) Comparative analysis of environmental DNA extraction and purification methods from different humic acid-rich soils. *Journal of Applied Microbiology* 102, 265–273.
- Luo HQ, Zhang H, Li K, et al.** (2017) Molecular characterization of ascaris from Tibetan pigs by three mitochondrial markers of nad1, cox1 and cox2. *Tropical Biomedicine* 34, 576–582.
- Miller DN, Bryant JE, Madsen EL and Ghiorse WC** (1999) Evaluation and optimization of DNA extraction and purification procedures for soil and sediment samples. *Applied and Environmental Microbiology* 65, 4715–4724.
- Naidoo D, Archer C, Louton B and Rodda N** (2016) Testing household disinfectants for the inactivation of helminth eggs on surfaces and in spills during pit latrine emptying. *Water SA* 42, 560–570.
- Olson ND and Morrow JB** (2012) DNA extract characterization process for microbial detection methods development and validation. *BMC Research Notes* 5, 668–682.
- Pecson BM, Barrios JA, Johnson DR and Nelson KL** (2006) A realtime PCR method for quantifying viable *Ascaris* eggs using the first internally transcribed spacer region of ribosomal DNA. *Applied and Environmental Microbiology* 72, 7864–7872.
- Quilès F, Balandier JY and Capizzi-Banas S** (2006) In situ characterisation of a microorganism surface by Raman microspectroscopy: the shell of *Ascaris* eggs. *Analytical and Bioanalytical Chemistry* 386, 249–255.
- Rostami A, Ebrahimi M, Mehvarar S, Omrani VF, Fallahi S and Behniafar H** (2016) Contamination of commonly consumed raw vegetables with soil transmitted helminth eggs in Mazandaran province, Northern Iran. *International Journal of Food Microbiology* 225, 54–58.
- Salonen A, Nikkilä J, Jalanka-Tuovinen J, Immonen O, Rajilić-Stojanović M, Kekkonen RA, Palva A and de Vos WM** (2010) Comparative analysis of fecal DNA extraction methods with phylogenetic microarray: effective recovery of bacterial and archaeal DNA using mechanical cell lysis. *Journal of Microbiology Methods* 81, 127–134.
- Scaglia B, D'Imporzano G, Garuti G, Negri M and Adani F** (2014) Sanitation ability of anaerobic digestion performed at different temperature on sewage sludge. *Science of the Total Environment* 466–467, 888–897.
- Sørensen SJ, Müller AK, Hansen LH, Rasmussen LD, Liphthay JR and Barkay T** (2002) Molecular methods for assessing and manipulating the diversity of microbiological populations and processes. pp. 289–294 in Burns RG, Dick RP (Eds) *Enzymes in the environment*. New York, Marcel Dekker.
- Tebbe CC and Vahjen W** (1993) Interference of humic acids and DNA extracted directly from soil in detection and transformation of recombinant-DNA from bacteria and a yeast. *Applied and Environmental Microbiology* 59, 2657–2665.
- Thermo Scientific** (2010) *Thermo scientific NanoDrop spectrophotometers: nucleic acid. Product description*. MA, USA, Thermo Fisher Scientific Inc.
- Valero MA, Perez-Crespo I, Periago MV, Khoubbane M and Mas-Coma S** (2009) Fluke egg characteristics for the diagnosis of human and animal fascioliasis by *Fasciola hepatica* and *F. gigantica*. *Acta Tropica* 111, 150–159.
- Verweij JJ, Brienen EA, Ziem J, Yelifari L, Polderman AM and Van Lieshout L** (2007) Simultaneous detection and quantification of *Ancylostoma duodenale*, *Necator americanus*, and *Oesophagostomum bifurcum* in fecal samples using multiplex real-time PCR. *American Journal of Tropical Medicine and Hygiene* 77, 685–690.
- Verweij JJ, Canales M, Polman K, Ziem J, Brienen EA, Polderman AM and van Lieshout L** (2009) Molecular diagnosis of *Strongyloides stercoralis* in fecal samples using real-time PCR. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 103, 342–346.
- Wiria AE, Prasetyani MA, Hamid F, et al.** (2010) Does treatment of intestinal helminth infections influence malaria? Background and methodology of a longitudinal study of clinical, parasitological and immunological parameters in Nangapanda, Flores, Indonesia (ImmunoSPIN Study). *BMC Infectious Diseases* 10, 1–12.
- Yen-Phi VT, Rechenburg A, Vinnerås B, Clemens J and Kistemann T** (2010) Pathogens in septage in Vietnam. *Science of the Total Environment* 408, 2050–2053.
- Zhou JZ and Thompson DK** (2002) Challenges in applying microarrays to environmental studies. *Current Opinions in Biotechnology* 13, 204–207.