

Predictors of Neuropsychological Improvement Following Cognitive Rehabilitation in Patients with Gliomas

Karin Gehring,¹ Neil K. Aaronson,^{2,3} Chad M. Gundy,² Martin J.B. Taphoorn,^{4,5} AND Margriet M. Sitskoorn¹

¹CoRPS, Faculty of Social and Behavioural Sciences, Tilburg University, The Netherlands

²Division of Psychosocial Research and Epidemiology, Netherlands Cancer Institute, The Netherlands

³Department of Clinical Psychology, University of Amsterdam, The Netherlands

⁴Department of Neurology, Medical Center Haaglanden, The Hague, The Netherlands

⁵Department of Neurology, VU University Medical Center Amsterdam, The Netherlands

(RECEIVED April 17, 2010; FINAL REVISION November 11, 2010; ACCEPTED November 16, 2010)

Abstract

This study investigated the specific patient factors that predict responsiveness to a cognitive rehabilitation program. The program has previously been demonstrated to be successful at the group level in patients with gliomas, but it is unclear which patient characteristics optimized the effect of the intervention at the individual level. Four categories of possible predictors of improvement were selected for evaluation: sociodemographic and clinical variables, self-reported cognitive symptoms, and objective neuropsychological test performance. Hierarchical logistic regression analyses were conducted, beginning with the most accessible (sociodemographic) variables and ending with the most difficult (baseline neuropsychological) to identify in clinical practice. Nearly 60% of the participants of the intervention were classified as reliably improved. Reliable improvement was predicted by age ($p = .003$) and education ($p = .011$). Additional results suggested that younger patients were more likely to benefit specifically from the cognitive rehabilitation program ($p = .001$), and that higher education was also associated with improvement in the control group ($p = .024$). The findings are discussed in light of brain reserve theory. A practical implication is that cognitive rehabilitation programs should take the patients' age into account and, if possible, adapt programs to increase the likelihood of improvement among older participants. (*JINS*, 2011, 17, 256–266)

Keywords: Neuropsychological rehabilitation, Brain tumor, Prediction, Individual change, Reliable change, Cognitive impairment

INTRODUCTION

Patients with a glioma, the most common primary brain tumor, may experience deficits in several cognitive domains, including attention, memory, and executive functioning (Klein et al., 2001; Laack et al., 2005; Meyers & Brown, 2006). The peak age of onset for low-grade (less malignant but not benign) gliomas lies between 30 and 40 years (DeAngelis, 2001). Both the tumor itself and its treatment can cause these deficits. Deficits in patients with a low-grade glioma are usually not severe (Anderson, Damasio, & Tranel, 1990; Correa et al., 2008; Laack et al., 2005), but they can have a substantial impact on daily life (Meyers, Berger, & Prados, 2005). Particularly during the disease-free period, which can last for years, when patients are attempting to

resume their normal work and social activities, these deficits can play a significant role. Moreover, subjective cognitive symptoms are among the most common problems reported by patients with brain tumors (Lidstone et al., 2003; Mukand, Blackinton, Crincoli, Lee, & Santos, 2001).

In a recent study (Gehring et al., 2009), we reported the results of a randomized controlled trial of cognitive rehabilitation in 140 patients with a glioma, in which positive group effects were observed. In particular, the intervention group performed significantly better than the control group on neuropsychological tests of attention and verbal memory 6 months after undergoing the cognitive rehabilitation program.

Since group results can mask the variability in individual responses to cognitive rehabilitation, it is of interest to further investigate the specific patient factors that are associated with a more (or less) positive outcome. Undergoing a cognitive rehabilitation program can be quite time-consuming and burdensome for patients. Identification of specific patient characteristics that predict a positive response to cognitive

Correspondence and reprint requests to: Karin Gehring, PhD, Tilburg University, Prisma Building; Room P 512, P.O. Box 90153, 5000 LE Tilburg, The Netherlands. E-mail: k.gehring@uvt.nl

rehabilitation could allow us to reassure patients who possess these characteristics that undergoing this type of intervention is almost certainly worth the effort. Conversely, if we are able to identify patients who are less likely to benefit from the currently available cognitive rehabilitation programs, we may advise them accordingly. Perhaps more importantly, we can use the knowledge gained to adapt cognitive rehabilitation programs to yield optimal benefit for a larger percentage of the target population.

Sociodemographic (e.g., age, sex, and education), clinical (e.g., radiotherapy), subjective (e.g., self-reported cognitive symptoms), and/or neuropsychological (pre-intervention neuropsychological functioning) factors may play a role in predicting the outcome of cognitive rehabilitation. From a practical point of view, it would be desirable to be able to identify patients who are most likely to respond to cognitive rehabilitation on the basis of information that can be collected with relatively little effort (e.g., sociodemographics, basic clinical data). However, use of other patient-reported outcomes (e.g., cognitive symptoms) and performance indicators (neuropsychological test results) may also provide useful in maximizing such predictions.

Identifying responsiveness to a cognitive intervention, or change over time at the individual patient level is complicated by the fact that the reliability of (neuro)psychological measures can be compromised by such factors as practice effects and regression to the mean. In recent years, the Reliable Change Index (RCI) has been introduced as a possible means of addressing this problem. The RCI represents a measure of improvement at the individual level in the context of observed changes over time in a control group (Evans, Margison, & Barkham, 1998); it reflects the individual change beyond that which can be attributed to measurement error, and practice effects. To date, very few studies have used RCIs to evaluate cognitive rehabilitation at the individual level (e.g., Medalia & Richardson, 2005).

The present study investigated the potential predictors of individual responsiveness to a cognitive rehabilitation program that has previously been demonstrated to be successful at the group level in patients with gliomas. We examined whether individual improvement, based on the RCI, is predicted by sociodemographic and clinical variables, self-reported cognitive symptoms, and objective neuropsychological test performance.

METHOD

Study Sample and Design

For a detailed description of the randomized clinical trial and of the cognitive rehabilitation program, the reader is referred to Gehring et al. (2009). Briefly, 366 adult patients with (primarily low-grade) glioma and favorable prognostic factors, whose disease was in remission, were screened for the presence of cognitive symptoms. Those who screened positive and expressed potential interest in participating in a

cognitive rehabilitation program completed a battery of neuropsychological tests of attention, memory and executive functioning. Patients who scored at least one standard deviation below the mean of a healthy comparison group ($n = 294$; van Boxtel et al., 1998) on at least 4 of 20 neuropsychological variables were considered eligible. These subjective and objective cognitive eligibility criteria were used to identify patients who would both be motivated to participate in the program, and would potentially benefit from its content.

Eligible patients were randomized to an intervention group ($n = 70$) or to a (waiting list) control group ($n = 70$). To evaluate the effect of the rehabilitation program, the battery of neuropsychological tests and self-report questionnaires that was administered at baseline was repeated directly following cognitive rehabilitation (or an equivalent time point for the control group), and at 6-months follow-up.

The trial was approved by the institutional review boards of all participating hospitals and all patients provided written informed consent.

The Cognitive Rehabilitation Condition and the Control Condition

The cognitive rehabilitation program (Gehring, Aaronson, Taphoorn, & Sitskoorn, accepted for publication; Gehring et al., 2009; for a more detailed description) consisted of six weekly, individual sessions of 2 hr. The intervention incorporated both cognitive retraining and strategy training. For the retraining component of the program, a computer program (C-Car; Gehring & Sitskoorn, 2004) was developed which consisted of a series of hierarchically graded tasks, designed to strengthen various aspects of attention, based on the patient's needs. The strategy training consisted of six integrated psycho-education sessions addressing attention, memory and executive function. These sessions included both didactic and practical elements aimed at teaching patients to compensate for impaired cognitive functions. Patients were also given weekly "homework" assignments to supplement both the retraining and strategy elements of the program. The waiting list control group received usual care (i.e., regular medical follow-up; no cognitive interventions), and was given the opportunity to follow the program at completion of the trial.

Results at the Group Level

For a detailed description of the trial results, the reader is referred to Gehring et al. (2009). To summarize, at immediate post-treatment, the intervention group reported significantly fewer cognitive symptoms and less symptom burden than the control group. At 6-months follow-up, the intervention group performed significantly better than the control group on a total of six neuropsychological test variables of attention and verbal memory, and reported significantly less mental fatigue. Group differences in other subjective outcomes were no longer statistically significant at 6-months follow-up.

Study Measures

The neuropsychological and self-report measures of cognitive symptoms that were used in this report are summarized in Table 1.

Outcome Measures

For the current analyses, the six neuropsychological measures of attention and verbal memory for which statistically significant group differences were observed at 6-months follow-up (Gehring et al., 2009) (see Neuropsychological measures in Table 1) were selected for further investigation at the individual level using a standardized regression-based RCI (McSweeney, Naugle, Chelune, & Luders, 1993). One global dichotomous measure of improvement was calculated (see Statistical Methods), based on the RCIs of these six measures, that distinguished between patients who improved reliably on at least one of the six measures (“improvers”) and those who did not (“non-improvers”).

Predictors

Potential predictors of outcome (Table 2) that were included in the analyses were: sociodemographics, clinical variables, self-reported cognitive symptoms, and neuropsychological test results. As all patients attended all sessions, “dose of therapy” was not a relevant predictor variable.

Although we considered calculating a composite baseline neuropsychological score based on Z-scores, this was not possible because normative data were not available for all six neuropsychological measures that were also used for the global RCI outcome.

Statistical Methods

For each program participant, change in the six neuropsychological performance measures was calculated by the Reliable Change Index (RCI), reflecting change at the individual level in the context of observed changes for the control group. While several variants of the RCI have been proposed

Table 1. Details of neuropsychological and subjective cognitive measures

Neuropsychological measures (tests)		
Attention		
SCWT	Stroop Color-Word Test (Hammes, 1971; Stroop, 1935), subtest: Card III (time in seconds) –	Attentional inhibition of a dominant response
DS	Digit Span of the WAIS-R (Luteijn & Van der Ploeg, 1983), subtests: Forward (span: 0–8) + Backward (span: 0–7) +	Immediate verbal recall Working memory
LDST	Letter Digit Substitution Test (Jolles, Houx, Van Boxtel, & Ponds, 1995), subtest: 90 Sec Writing (number correct: 0–125) +	Psychomotor speed and speed of information processing
Verbal Memory		
VVLT	Visual Verbal Learning Test, direct and delayed recall (Brand & Jolles, 1987), subscores: Trial 1 (number correct: 0–15) + Delayed Recall (number correct: 0–15) +	Immediate verbal span Verbal memory after an interval
Subjective cognitive measures (questionnaires)		
Cognitive symptoms		
CFS	Cognitive Functioning Scale of the Medical Outcomes Study (MOS) Health Survey questionnaire battery (Stewart, Ware, Sherbourne, & Wells, 1992) Total score (6–36) +	Self-reported frequency of cognitive symptoms
Burden	Study-specific measure Total (3–18) + of three questions on the impact of the cognitive complaints on daily life, worry about the cognitive complaints, being troubled by the cognitive complaints	Self-reported burden of the CFS symptoms
CFQ	Cognitive Failure Questionnaire (CFQ) (Broadbent, Cooper, FitzGerald, & Parkes, 1982; Ponds, Rozendaal, & Jolles) Total score (0–100) –	Self-reported cognitive failures in daily life

Note. + = Higher score indicates better (performance) score; – = Higher score: worse (performance) score; In parentheses: Score/scale ranges.

Table 2. Candidate predictors of improvement: Descriptives and correlations (Pearson’s correlation and Phi coefficient) with ‘reliable improvement on at least one of the six neuropsychological measures’ in the intervention group

Predictor variables		Descriptives	Correlation	<i>p</i>
Demographic variables (personal interview)	Age	Mean: 41.8; SD: 9.5	-.385	.002**
	Sex (male/female)	M: 36 (56%) F: 28 (44%)	.088	.488
	Education (low-medium/high)	L-M: 32 (50%) H: 32 (50%)	.318	.010*
	Currently employed (no/yes)	N: 25 (39%) Y: 39 (61%)	.185	.142
Medical variables (medical record)	Disease duration (median: 4.8; range 0.8–38.9; dichotomized: 0.5–5 years/>5 years)	<5: 33 (52%) >5: 31 (48%)	-.307	.014*
	Years since last tumor treatment (log)	Median: 2.6; Range: 0.6–21.2	-.210	.106
	Hemisphere of lesion (left/right; 2 bilateral lesions excluded)	L: 33 (52%) R: 29 (45%)	.076	.557
	Tumor classification (astrocytic/oligodendroglial; 4 without histopathological diagnosis excluded)	A: 30 (47%) O: 30 (47%)	.068	.605
	Radiotherapy in history (no/yes)	N: 25 (39%) Y: 39 (61%)	-.010	.936
	Epileptic seizures in last year (no/yes)	N: 27 (42%) Y: 37 (58%)	.260	.038*
	Comorbidity [‡] (no/yes)	N: 24 (38%) Y: 40 (62%)	-.312	.012*
	Tumor grade (low-grade/anaplastic glioma) – excluded from analyses due to small cell sizes	L: 54 (84%) A: 10 (16%)		
	Anti-epileptic drug use (no/yes) – excluded from analyses due to small cell sizes	N: 10 (16%) Y: 54 (84%)		
Chemotherapy (no/yes) – excluded from analyses due to small cell sizes	N: 57 (89%) Y: 7 (11%)			
Subjective variables (questionnaires)	Total Cognitive Functioning Scale (CFS)	Mean: 20.9; SD: 4.2	.070	.588
	Burden of the cognitive symptoms	Mean 9.8; SD: 2.9	.092	.472
	Total Cognitive Failures Questionnaire (CFQ)	Mean: 48.5; SD: 10.3	-.067	.602
Baseline neuropsychological variables (neuropsychological assessment)	Stroop Color-Word Test Card III (rec)	Median: 98.5; Range: 52–254	-.102	.422
	Digit Span Forward	Mean: 5.3; SD: 1.0	-.034	.792
	Digit Span Backward	Mean: 4.6; SD: 1.1	-.180	.156
	Letter Digit Substitution Test 90 Sec Writing	Mean: 43.9; SD: 8.4	.263	.036*
	Visual Verbal Learning Test Trial 1	Mean: 5.8; SD: 2.3	.119	.351
	Visual Verbal Learning Test Delayed Recall	Mean: 9.5; SD: 3.0	.022	.866
	One (1) cognitively disturbed domain versus multiple (≥1) domains disturbed at baseline	1: 49 (77%) ≥1: 15 (23%)	.082	.519

Note. **p* < .05, two-tailed. ***p* < .01, two-tailed; log = logarithmically transformed, rec = reciprocally transformed; [‡]other concurrent diseases or disease symptoms not related to the brain tumor varying from chronic eczema to diabetes or cardiovascular disease.

(Maassen, Bossema, & Brand, 2009), all reflect the ratio of an estimate of the true change in the numerator and a corresponding (measurement) error in the denominator (Maassen et al., 2009). For the current study, a standardized regression-based approach to the RCI was used (McSweeney et al., 1993), that uses the available information of the control

Table 3. Reliably improved individuals in the intervention group (*N* = 64) and in the control group (*N* = 63)

Neuropsychological test variable	Intervention group		Control group	
	<i>N</i> improved	% improved	<i>N</i> improved	% improved
Stroop Color-Word Test Card III	4	6%	5	8%
Digit Span Forward	13	20%	3	5%
Digit Span Backward	17	27%	5	8%
Letter Digit Substitution Test 90-Sec Writing	8	13%	1	2%
Visual Verbal Learning Test Trial 1	13	20%	4	6%
Visual Verbal Learning Test Delayed Recall	11	17%	4	6%
Reliably improved in at least one of the six measures	38	59%	16	25%

Note. Number and percentage of improvers at six months follow-up per neuropsychological test variable, and number and percentage of patients who improved reliably in at least one of the six neuropsychological measures (global RCI outcome).

group more thoroughly than other types of RCIs (Maassen et al., 2009). Practice effect is estimated by means of a regression of the post-test on the pre-test score in the control group; subsequently the relationship between practice effect and the initial score of an individual is used. The standard error of prediction of the control group is used as the standard error in the denominator. The index that is calculated is compared to the critical value for a one-sided *t* test with $n = 64$. The reader is referred to the Appendix for further details of the RCI formula used in this study.

Finally, a global, dichotomous measure of reliable improvement was created to distinguish between patients who improved reliably on at least one of the six measures (“improvers”) and those who did not (“non-improvers”). This global measure was used as the dependent variable in the analyses.

Because of the large number of possible predictors (21; see Table 2) and the relatively small sample size ($n = 64$), univariate correlations (Pearson’s correlation and Phi coefficient) between (both dichotomous and continuous) predictor variables and the global RCI were first calculated. Independent variables that correlated with the global RCI outcome at the $p < .05$ level were included in subsequent multivariate analyses. Dichotomous predictor variables with cell sizes that were too small were excluded from the analyses. Variables that were not normally distributed (an assumption for Pearson’s correlations) were transformed to normalize distributions, or were dichotomized.

Clusters of one or more variables that correlated significantly with the global RCI outcome at the univariate level were included in the logistic regression analyses according to a pre-specified hierarchy: sociodemographic variables, clinical variables, self-reported cognitive symptoms, and baseline neuropsychological test results. This hierarchical ordering was chosen to reflect the ease with which these data could be collected in daily clinical practice. That is, sociodemographic and clinical data can be relatively easily extracted from the medical records, while patient self-reported symptoms and objective neuropsychological test results require additional and, in the case of the neuropsychological tests, quite substantial data collection efforts. The final model was composed of those variables that contributed significantly to predicting reliable improvement in at least one of the six neuropsychological measures (the global RCI outcome).

To investigate whether the factors identified as being multivariate predictors of improvement within the intervention group might also be predictive of improvement in neuropsychological functioning in general, an additional logistic regression analysis was conducted for the entire study sample (the intervention and the control group combined) in which, in addition to the identified predictor variables, both the group variable and group \times predictor interaction terms were included in the model.

Assumptions concerning (absence of) outliers/influential cases, empty cells and multicollinearity were tested for all of the logistic regression models. Variables that had been transformed or dichotomized for the univariate analyses

remained modified for the logistic regression analyses. For all logistic regression analyses, a p value $< .05$ was considered statistically significant. Nagelkerke’s pseudo R^2 was used to provide an indication of the substantive significance of the logistic regression model (analogous to a R^2 value as reported with multiple regression; Field, 2005).

Because of the relatively small sample size, a bootstrap sampling estimation was performed in which 2000 versions of the training group were generated to estimate regression coefficients and confidence intervals with a minimum of assumptions.

Finally, supplemental data on the area under the curve, sensitivity, specificity, positive (PPV) and negative (NPV) predictive values, and classification accuracy were calculated for both predictors, education and age, in the intervention group, using information from a receiver operating characteristic (ROC) curve. As age is a continuous measure, determination of these data depends on the cutoff used for age. This cutoff was determined on the basis of the smallest number of misclassifications (Youden’s Index; Youden, 1950), that is, on the optimal combination of sensitivity and specificity, by summing these two values for each value of age in the sample (a table provided by the SPSS ROC curve output).

For all statistical tests, the Statistical Package for the Social Sciences (SPSS; SPSS Inc. Chicago, IL) version 17.0 was used.

RESULTS

Follow-up data were available of 64 intervention group patients. There were no statistically significant differences in sociodemographic, clinical, self-reported cognitive symptoms or baseline neuropsychological test scores between these 64 patients and the 63 control group patients whose baseline and follow-up neuropsychological test scores were used for calculating the RCI.

RCI Results per Measure and Global RCI-Score

Fifty-nine percent of the patients in the intervention group, as compared to 25% in the control group ($\chi^2(1) = 14.997$; $p < .001$), improved reliably (i.e., corrected for measurement errors and practice effects) on at least one of the six neuropsychological measures (the global RCI outcome; see Table 3, also for the number and percentage of reliably changed individuals in both groups per neuropsychological test variable).

Variables Associated Significantly with Reliable Change in Neuropsychological Functioning at the Univariate Level

Variables that were not normally distributed were transformed (logarithmically for “years since last tumor treatment”, or reciprocally for “Stroop color-word test”; see Table 2) to normalize distributions. One variable (“disease duration”) was dichotomized to solve problems with normality even after transformation, with the cutoff based on the median disease

duration of 5 years. It should be noted that, since the majority of patients had a low-grade glioma, received anti-epileptic drugs and had not received chemotherapy, variables regarding tumor grade, anti-epileptic drugs and chemotherapy were not included in the analyses as the numbers of patients in these cells were too small.

The following variables were found to correlate significantly with reliable improvement in neuropsychological functioning over time (see also Table 2): From the socio-demographic cluster, *age* (younger age) and *education* (higher level); from the clinical cluster, *disease duration* (less than 5 years), *epileptic seizures* in last year (yes), and *comorbidity* (no). None of the self-reported cognitive symptom variables were correlated significantly with outcome. Of the baseline neuropsychological tests, only *letter-digit substitution score* (higher) was associated significantly with reliable improvement.

Logistic Regression Analyses

In the first logistic regression analysis (see Table 4), the demographic variables age and education were included as possible predictors. (Younger) age and (higher) education were associated with reliable improvement (model $\chi^2(2) = 17.190$; $p < .001$).

Addition of the clinical variables disease duration, epileptic seizures and comorbidity did not result in a significant improvement of the model (step/block $\chi^2(3) = 6.582$; $p = .086$). Thus the clinical variables did not contribute significantly to the model and therefore these variables were not included. Finally, inclusion of the letter-digit substitution test to the existing model did not result in an improvement of the model (step/block $\chi^2(1) = .012$; $p = .991$).

The final model (model $\chi^2(2) = 17.190$; $p < .001$) predicting reliable change in the participants that had followed the cognitive rehabilitation program six months earlier was similar to the initial model and included age ($p = .003$) and education ($p = .011$) only. A Nagelkerke's pseudo R^2 of .318 indicated moderate explanatory power of the model. The odds of improvement over time increased 0.9 times per year of age (range, 23 to 59 years), and were 4.7 times greater for higher (vs. low to medium) education level. The bootstrap sampling estimation confirmed these results.

For education, the area under the ROC curve in the intervention group was .66 (asymptotic 95% confidence interval: .585–.850). The sensitivity was 63% and the specificity was 69%. The overall classification accuracy was 65%. The PPV (i.e., given a higher education the chance to improve) was 75%, and the NPV (i.e., given a lower education, the chance of no improvement) was 56%. For age, the area under the ROC curve was .72 (asymptotic 95% confidence interval: .525–.799). At the optimal cutoff of age 50, sensitivity was 95%, specificity was 50%, and the overall classification accuracy was 77%. The PPV (i.e., given a younger age the chance to improve) was 73% and the NPV (i.e., given an older age the chance of no improvement) was 87%.

To investigate whether age and education specifically predicted intervention-related improvement, or were also involved in more general improvement, an additional logistic regression analysis was conducted for the entire study group ($n = 127$) with inclusion of the variables group, age, education, and the two-way interaction terms of group \times age, and group \times education. A high correlation was observed between the group variable and the interaction term group \times age ($r(125) = .95$), resulting in a problem with multicollinearity. For this reason, age was centered around the total group mean age of 41.8 years for use in the interaction term.

This model (model $\chi^2(5) = 41.545$; $p < .001$; Table 4) confirmed that the intervention ($p = .014$) was associated significantly with improvement in the entire group, as was (higher) education ($p = .024$). (Centered) age, however, was only significant when incorporated in the interaction term ($p = .001$), indicating that (younger) age moderated improvement in neuropsychological test performance for the intervention group, but not for the control group. The odds of reliable improvement were 5.3 for the patients in the intervention group (versus the control group), and 4.5 for patients with higher education (versus low to medium education, irrespective of group). The odds ratio for group \times centered age was .86, indicating that for every year above the mean age of 41.8 years, the odds of improvement decreased by a factor of 0.86.

DISCUSSION

The current study investigated the patient characteristics that predicted individual neuropsychological improvement in glioma patients 6 months after undergoing a cognitive rehabilitation program (Gehring et al., 2009). According to the criteria used, with measurement errors and practice effects taken into account, nearly 60% of the 64 patients in the intervention group had improved reliably on at least one of the six neuropsychological measures for which statistically significant intervention effects had been demonstrated at the group level. Four categories of candidate predictors of improvement in neuropsychological test performance were examined at the univariate level and, subsequently, in the context of a hierarchical logistical regression model. Of all of the variables examined, only two sociodemographic variables, age and education, significantly predicted individual improvement in the intervention group. Younger patients and those with a higher education had the highest likelihood of improvement over time. Additional results suggested that age was a specific, intervention-related predictor of improvement in neuropsychological functioning; that is, it did not predict such improvement among those patients not exposed to the cognitive rehabilitation program. Education, however, appeared to be a more general predictor of improvement in neuropsychological functioning, irrespective of whether patients were exposed to the program.

It should be noted that in calculating the RCI, data on practice effects as measured in the control group are taken into account. However, these data are based on "mean"

Table 4. Logistic regression models

Model	χ^2	df	p	R ² (Nagelkerke)	Variables in the equation	B	SE	Wald	df	Sig.	Exp(B)	95% CI for Exp(B)	
												lower	upper
1 Sociodemographic variables	17.190	2	.000	.318	Age	-.103	.035	8.673	1	.003	.902	.842	.966
					Education (dich)	1.552	.611	6.457	1	.011	4.721	1.426	15.628
					Constant	4.055	1.498	7.329	1	.007	57.692		
2 Sociodemographic + clinical variables	23.773	5	.000	.419	Age	-.085	.038	4.922	1	.027	.919	.853	.990
					Education (dich)	1.236	.657	3.535	1	.060	3.441	.949	12.481
					Comorbidity (dich)	-1.129	.678	2.773	1	.096	.323	.086	1.221
					Disease duration (dich)	-1.101	.631	3.051	1	.081	.332	.097	1.144
					Epileptic seizures (dich)	.473	.654	.523	1	.469	1.605	.446	5.778
					Constant	4.393	1.818	5.840	1	.016	80.865		
3 Sociodemographic + neuropsychological variables	17.203	3	.001	.318	Age	-.102	.038	7.117	1	.008	.903	.839	.973
					Education (dich)	1.527	.650	5.513	1	.019	4.603	1.287	16.463
					Letter digit substitution	.004	.040	.012	1	.911	1.005	.928	1.087
					Constant	3.801	2.722	1.951	1	.163	44.764		
4 Final model	17.190	2	.000	.318	Age	-.103	.035	8.673	1	.003	.902	.842	.966
					Education (dich)	1.552	.611	6.457	1	.011	4.721	1.426	15.628
					Constant	4.055	1.498	7.329	1	.007	57.692		
Entire sample	41.545	5	.000	.375	Group	1.673	.681	6.039	1	.014	5.327	1.403	20.228
					Age (centered)	.052	.030	2.936	1	.087	1.053	.993	1.117
					Education (dich)	1.500	.666	5.071	1	.024	4.482	1.215	16.536
					Group X education (dich)	.052	.904	.003	1	.954	1.053	.179	6.192
					Group X age (centered)	-.155	.046	11.220	1	.001	.857	.782	.938
					Constant	-2.045	.556	13.527	1	.000	.129		

Note. dich = dichotomized.

practice effects for the control group. This means that some patients exhibit more than average, and others exhibit less than average practice effects. The 25% of the control group patients that improved “reliably” (a percentage that is significantly different from that observed in the intervention group) may be those patients with greater than average improvement. The results of the total group analyses suggest that education may (also) have moderated neuropsychological improvement in the control group, which is probably related to practice effects.

Furthermore, although the criterion of reliable improvement on at least one of six neuropsychological measures may seem minimal, it should be noted that the RCI is a very stringent index, excluding measurement errors and subjecting the adapted score to statistical testing (a *t* test). This resulted in clearly defined groups that could be used for analyses with reliable results. The difference in proportion of reliably improved patients in the intervention (59%) versus control (25%) group was statistically significant ($\chi^2(1) = 14.997$; $p < .001$). The results of the logistic regression analyses based on the current criteria with group sizes of 38 reliably improved versus 26 not (reliably) improved patients were supplemented by other statistical measures (in particular, the PPV and NPV of age of 73% and 87%, respectively), and yielded different predictor variables for the intervention group as compared to the control group.

The prediction of “success” of the program by age may reflect the nature and content of the program itself. The game-like retraining on a notebook computer may have been more attractive, and familiar to, younger patients. However, in studies on outcome of patients suffering from acquired brain injury (without interventions specifically aimed at treating cognitive deficits), sociodemographic variables such as age and education/intelligence, have also been predictive of functioning, in addition to medical injury severity variables (Chu et al., 2007; Malec & Basford, 1996; Robertson & Murre, 1999; Sherer et al., 2006). Both older age (e.g., Chu et al., 2007; Hukkelhoven et al., 2003; Robertson & Murre, 1999; Sherer et al., 2006) and lower education (Brooks, McKinlay, Symington, Beattie, & Campsie, 1987; Grafman, Salazar, Weingartner, Vance, & Amin, 1986; Robertson & Murre, 1999; Sherer et al., 2006) have been associated with poorer neuropsychological outcome in many of these studies.

The finding that age plays a significant role in predicting (cognitive) recovery may be related to the concept of brain reserve. There is both epidemiological and biological evidence that individual differences in the brain allow some people to cope better than others with brain damage (Fratiglioni & Wang, 2007; Stern, 2009). These individual differences stem from quantitative factors such as brain size, or number of neurons or synapses. Furthermore, brain anatomy can be influenced by life experiences, via neurogenesis and angiogenesis that promote resistance to apoptosis and indirectly promote neural plasticity (Stern, 2009). As aging has been related to losses in gray and white matter in medial-temporal, parietal and frontal areas (Charlton et al., 2006;

Gordon et al., 2008), it has been suggested that these diffuse age-related losses results in a larger threshold for recovery to take place (Robertson & Murre, 1999), decreasing the brain's reserve (Green et al., 2008; Stern, 2009).

Moreover, brain reserve factors may not only buffer the outcome of brain damage, and influence the initial cognitive status after the injury, they may also play a supporting role in restitution of function or functional reorganization during recovery (Green et al., 2008). Some of the few studies on recovery trajectories conducted in brain-injured populations have suggested that age (but not education) can indeed moderate improvement in cognitive functioning in patients with traumatic brain injury (Green et al., 2008; Zwaagstra, Schmidt, & Vanier, 1996). However, in other studies, factors such as age (or premorbid intelligence) only predicted the level of cognitive outcome, rather than influencing the recovery trajectory (Chu et al., 2007; Green et al., 2008).

With regard to improvement after cognitive rehabilitation in particular, age has also been reported to play a moderating role (e.g., Langbaum, Rebok, Bandeen-Roche, & Carlson, 2009; Rohling, Faust, Beverly, & Demakis, 2009; Verhaeghen, de Mey, Helsen, van Assel, & Vanwijnsberghe, 1992; Yesavage, Sheikh, Friedman, & Tanke, 1990). Langbaum and colleagues (2009) studied the different patterns of response to memory training in 619 healthy older adults and concluded that higher education and higher baseline cognitive functioning were predictors of (patterns of) memory improvement. Age had also a moderating effect, although the relationship to response pattern was possibly not linear. However, other studies did not find moderating effects of age (or education) (Medalia & Richardson, 2005; Neely & Backman, 1995).

Thus, although it is well-known that age (and also education) influence (static) performance on neuropsychological tests, this study demonstrates that these variables may also moderate the (dynamic) response to cognitive rehabilitation. In line with this reasoning above, in the current study, younger patients may have a larger brain reserve capacity for dealing with the pathological effects of the brain tumor.

It would have been interesting if we, in addition to a proxy measure of brain reserve (age) also had used some proxy measure for *cognitive reserve* in the analyses. Although we collected data on the DART (the Dutch version of the NART), a measure of premorbid intelligence, we decided not to use these data in the analyses as we observed that they were confounded by hemisphere of lesion. Unfortunately we had no other measure for cognitive reserve available.

The order in which the variables were analyzed, according to our pre-specified clinical hierarchy, in combination with the relatively small power of the study, may have precluded some other possibly relevant variables from reaching statistical significance. The clinical variables that were tested in the multivariate model yielded statistical trends only. The finding that baseline neuropsychological functioning was not predictive of improvement following the program runs contrary to other evidence in the literature (e.g., Fiszdon, Cardenas, Bryson, & Bell, 2005; Langbaum et al., 2009;

Strangman et al., 2008). This may, in part, be due to the fact that we used only one global measure of cognitive improvement, which was also dichotomized, to investigate the relationship with each of the six baseline neuropsychological test variables.

Furthermore, we cannot exclude the possibility that the positive results of the cognitive rehabilitation are not (only) attributable to the cognitive treatment itself, but might also be due, at least in part, to other non-specific factors such as the attention given to the patients. The RCI does not correct for these effects. We considered it unethical, impractical, and not credible to include a “placebo” condition in which patients would be required to attend a series of 2-hr sessions (comparable to the intervention group) in which no substantive rehabilitation program would be offered. However, non-specific treatment effects as the sole or even primary explanation for the improvements in objective test performance is less likely, considering the initially equal improvement in neuropsychological performance for both groups, and the intervention effect observed after a 6-month interval, in which possible placebo effects may be assumed to be absent.

The results of this study suggest that younger patients with a glioma are more likely to benefit from the cognitive rehabilitation program. As younger age at diagnosis is a major prognostic factor in brain cancer (Schiff, Brown, & Giannini, 2007), it is useful to know that these patients are also most likely to benefit from the intervention. This should not, however, suggest that cognitive rehabilitation programs should not be offered to older patients. Rather, greater effort should be devoted to adapting our current training program to increase its effectiveness among older patients. Brooks and colleagues (Brooks, Friedman, Pearman, Gray, & Yesavage, 1999), in a study of 268 community-dwelling adults over the age of 55 years, reported that older participants benefited from increased training time coupled with a comprehensive pre-training regimen in memory (mnemonic) rehabilitation. Thus, pre-training, combined with more sessions over time, may facilitate program success in older patients.

A secondary, but relevant, finding of the study was that more highly educated patients were more likely to exhibit improved cognitive functioning over time, regardless of whether they participated in the program. This suggests that (clinical) neuropsychologists or (other) researchers should be aware that patients with a higher level of education are likely to show improvement over time after multiple neuropsychological test assessments.

Future cognitive rehabilitation approaches, in patients with brain tumors or other diseases that cause cognitive impairment should take the age of participants into account, and investigate whether adapted intervention characteristics (e.g., pretraining, and splitting up sessions) may facilitate enhanced program effectiveness in older participants.

ACKNOWLEDGMENTS

We thank Gerard Maassen for his advice on the use of the Reliable Change Index, Wobbe P. Zijlstra for his statistical advice, and Ruth

E. Nieuwenhuis-Mark for editing an earlier version of this draft of this manuscript. This study was supported by the Dutch Cancer Society (UU2003-2783) and Health Insurers Innovation Foundation (file 504). There are no conflicts of interest.

REFERENCES

- Anderson, S.W., Damasio, H., & Tranel, D. (1990). Neuropsychological impairments associated with lesions caused by tumor or stroke. *Archives of Neurology*, *47*(4), 397–405.
- Brand, N., & Jolles, J. (1987). Information processing in depression and anxiety. *Psychological Medicine*, *17*(1), 145–153.
- Broadbent, D.E., Cooper, P.F., FitzGerald, P., & Parkes, K.R. (1982). The Cognitive Failures Questionnaire (CFQ) and its correlates. *The British Journal of Clinical Psychology*, *21*(Pt 1), 1–16.
- Brooks, J.O., III, Friedman, L., Pearman, A.M., Gray, C., & Yesavage, J.A. (1999). Mnemonic training in older adults: Effects of age, length of training, and type of cognitive pretraining. *International Psychogeriatrics*, *11*(1), 75–84.
- Brooks, N., McKinlay, W., Symington, C., Beattie, A., & Campsie, L. (1987). Return to work within the first seven years of severe head injury. *Brain Injury*, *1*(1), 5–19.
- Charlton, R.A., Barrick, T.R., McIntyre, D.J., Shen, Y., O’Sullivan, M., Howe, F.A., ... Markus, H.S. (2006). White matter damage on diffusion tensor imaging correlates with age-related cognitive decline. *Neurology*, *66*(2), 217–222. doi:10.1212/01.wnl.0000194256.15247.83
- Chu, B.C., Millis, S., Arango-Lasprilla, J.C., Hanks, R., Novack, T., & Hart, T. (2007). Measuring recovery in new learning and memory following traumatic brain injury: a mixed-effects modeling approach. *Journal of Clinical and Experimental Neuropsychology*, *29*(6), 617–625. doi:10.1080/13803390600878893
- Correa, D.D., Shi, W., Thaler, H.T., Cheung, A.M., DeAngelis, L.M., & Abrey, L.E. (2008). Longitudinal cognitive follow-up in low grade gliomas. *Journal of Neuro-Oncology*, *86*(3), 321–327.
- DeAngelis, L.M. (2001). Brain tumors. *New England Journal of Medicine*, *344*(2), 114–123.
- Evans, C., Margison, F., & Barkham, M. (1998). The contribution of reliable and clinically significant change methods to evidence-based mental health. *Evidence-Based Mental Health*, *1*(3), 70–72.
- Field, A. (2005). *Discovering statistics using SPSS* (2nd ed.). Thousand Oaks, CA: Sage Publications, Inc.
- Fiszdon, J.M., Cardenas, A.S., Bryson, G.J., & Bell, M.D. (2005). Predictors of remediation success on a trained memory task. *The Journal of Nervous and Mental Disease*, *193*(9), 602–608.
- Fratiglioni, L., & Wang, H.X. (2007). Brain reserve hypothesis in dementia. *Journal of Alzheimer’s Disease*, *12*(1), 11–22.
- Gehring, K., Aaronson, N.K., Taphoorn, M.J.B., & Sitskoorn, M.M. (accepted for publication). A description of a cognitive rehabilitation program evaluated in brain tumor patients with mild to moderate cognitive deficits.
- Gehring, K., & Sitskoorn, M.M. (2004). *Handleiding C-Car [Manual C-Car]*.
- Gehring, K., Sitskoorn, M.M., Gundy, C.M., Sikkes, S.A., Klein, M., Postma, T.J., ... Aaronson, N.K. (2009). Cognitive rehabilitation in patients with gliomas: A randomized, controlled trial. *Journal of Clinical Oncology*, *27*(22), 3712–3722. doi:10.1200/JCO.2008.20.5765
- Gordon, B.A., Rykhlevskaia, E.I., Brumback, C.R., Lee, Y., Elavsky, S., Konopack, J.F., ... Fabiani, M. (2008). Neuro-anatomical correlates of aging, cardiopulmonary fitness level, and

- education. *Psychophysiology*, 45(5), 825–838. doi:10.1111/j.1469-8986.2008.00676.x
- Grafman, J., Salazar, A., Weingartner, H., Vance, S., & Amin, D. (1986). The relationship of brain-tissue loss volume and lesion location to cognitive deficit. *The Journal of Neuroscience*, 6(2), 301–307.
- Green, R.E., Colella, B., Christensen, B., Johns, K., Frasca, D., Bayley, M., ... Monette, G. (2008). Examining moderators of cognitive recovery trajectories after moderate to severe traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 89(12 Suppl), S16–S24. doi:10.1016/j.apmr.2008.09.551
- Hammes, J. (1971). *De Stroop Kleur-Woord Test: Handleiding [The Stroop Color-Word Test: Manual]*. Amsterdam, The Netherlands: Harcourt.
- Hukkelhoven, C.W., Steyerberg, E.W., Rampen, A.J., Farace, E., Habbema, J.D., Marshall, L.F., ... Maas, A.I. (2003). Patient age and outcome following severe traumatic brain injury: An analysis of 5600 patients. *Journal of Neurosurgery*, 99(4), 666–673. doi:10.3171/jns.2003.99.4.0666
- Jolles, J., Houx, P.J., Van Boxtel, M.P.J., & Ponds, R.W.H.M. (1995). *Maastricht Aging Study: Determinants of Cognitive Aging*. Maastricht, The Netherlands: Neuropsych Publishers.
- Klein, M., Taphoorn, M.J., Heimans, J.J., Van Der Ploeg, H.M., Vandertop, W.P., Smit, E.F., ... Aaronson, N.K. (2001). Neurobehavioral status and health-related quality of life in newly diagnosed high-grade glioma patients. *Journal of Clinical Oncology*, 19(20), 4037–4047.
- Laack, N.N., Brown, P.D., Ivnik, R.J., Furth, A.F., Ballman, K.V., Hammack, J.E., ... Buckner, J.C. (2005). Cognitive function after radiotherapy for supratentorial low-grade glioma: A North Central Cancer Treatment Group prospective study. *International Journal of Radiation Oncology, Biology, Physics*, 63(4), 1175–1183.
- Langbaum, J.B., Rebok, G.W., Bandeen-Roche, K., & Carlson, M.C. (2009). Predicting memory training response patterns: Results from ACTIVE. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 64(1), 14–23. doi:10.1093/geronb/gbn026
- Lidstone, V., Butters, E., Seed, P.T., Sinnott, C., Beynon, T., & Richards, M. (2003). Symptoms and concerns amongst cancer outpatients: Identifying the need for specialist palliative care. *Palliative Medicine*, 17(7), 588–595.
- Luteijn, F., & Van der Ploeg, F.A.E. (1983). *Handleiding Groninger Intelligentie Test (GIT). [Manual GIT]*. Lisse: Swets and Zeitlinger.
- Maassen, G.H., Bossema, E., & Brand, N. (2009). Reliable change and practice effects: Outcomes of various indices compared. [Comparative Study]. *Journal of Clinical and Experimental Neuropsychology*, 31(3), 339–352.
- Malec, J.F., & Basford, J.S. (1996). Postacute brain injury rehabilitation. *Archives of Physical Medicine and Rehabilitation*, 77(2), 198–207.
- McSweeney, A., Naugle, R.I., Chelune, G.J., & Luders, H. (1993). “T Scores for Change”: An illustration of a regression approach to depicting change in clinical neuropsychology. *The Clinical Neuropsychologist*, 7(3), 300–312. doi:10.1080/13854049308401901
- Medalia, A., & Richardson, R. (2005). What predicts a good response to cognitive remediation interventions? *Schizophrenia Bulletin*, 31(4), 942–953.
- Meyers, C.A., Berger, M.S., & Prados, M. (2005). Functional outcomes. *Textbook of neuro-oncology* (pp. 101–104). Philadelphia: Elsevier Saunders.
- Meyers, C.A., & Brown, P.D. (2006). Role and relevance of neurocognitive assessment in clinical trials of patients with CNS tumors. *Journal of Clinical Oncology*, 24(8), 1305–1309.
- Mukand, J.A., Blackinton, D.D., Crincoli, M.G., Lee, J.J., & Santos, B.B. (2001). Incidence of neurologic deficits and rehabilitation of patients with brain tumors. *American Journal of Physical Medicine & Rehabilitation*, 80(5), 346–350.
- Neely, A.S., & Backman, L. (1995). Effects of multifactorial memory training in old age: Generalizability across tasks and individuals. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 50(3), P134–P140.
- Ponds, R.W.H.M., Rozendaal, N., & Jolles, J. (1999). *The Cognitive Failure Questionnaire: Factor structure, effects of age, sex and education and the relation with cognitive performance and psychosocial variables*. Maastricht, The Netherlands: Neuropsych Publishers.
- Robertson, I.H., & Murre, J.M. (1999). Rehabilitation of brain damage: Brain plasticity and principles of guided recovery. *Psychological Bulletin*, 125(5), 544–575.
- Rohling, M.L., Faust, M.E., Beverly, B., & Demakis, G. (2009). Effectiveness of cognitive rehabilitation following acquired brain injury: A meta-analytic re-examination of Cicerone et al.’s (2000, 2005) systematic reviews. *Neuropsychology*, 23(1), 20–39. doi:10.1037/a0013659
- Schiff, D., Brown, P.D., & Giannini, C. (2007). Outcome in adult low-grade glioma: The impact of prognostic factors and treatment. *Neurology*, 69(13), 1366–1373. doi:10.1212/01.wnl.0000277271.47601.a1
- Sherer, M., Stouter, J., Hart, T., Nakase-Richardson, R., Olivier, J., Manning, E., & Yablon, S.A. (2006). Computed tomography findings and early cognitive outcome after traumatic brain injury. *Brain Injury*, 20(10), 997–1005. doi:10.1080/02699050600677055
- Stern, Y. (2009). Cognitive reserve. *Neuropsychologia*, 47(10), 2015–2028. doi:10.1016/j.neuropsychologia.2009.03.004
- Stewart, A.L., Ware, J.E., Sherbourne, C.D., & Wells, K.B. (1992). Psychological distress/well-being and cognitive functioning measures. *Measuring functioning and well-being: The medical outcomes study approach* (pp. 102–142). Durham, NC: Duke University.
- Strangman, G.E., O’Neil-Pirozzi, T.M., Goldstein, R., Kelkar, K., Katz, D.I., Burke, D., ... Glenn, M.B. (2008). Prediction of memory rehabilitation outcomes in traumatic brain injury by using functional magnetic resonance imaging. *Archives of Physical Medicine and Rehabilitation*, 89(5), 974–981. doi:10.1016/j.apmr.2008.02.011
- Stroop, J.R. (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, 18(6), 643–662.
- van Boxtel, M.P., Buntinx, F., Houx, P.J., Metsemakers, J.F., Knotterus, A., & Jolles, J. (1998). The relation between morbidity and cognitive performance in a normal aging population. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 53(2), M147–M154.
- Verhaeghen, P., de Mey, G., Helsen, Z., van Assel, A., & Vanwijnsberghe, L. (1992). *Omgaan met het geheugen. Handboek voor hulpverleners en cursusbegeleiders. [Managing memory. Handbook for therapists and course supervisors.]*. Leuven, Belgium/Amersfoort, The Netherlands: Acco.
- Yesavage, J.A., Sheikh, J.I., Friedman, L., & Tanke, E. (1990). Learning mnemonics: Roles of aging and subtle cognitive impairment. *Psychology and Aging*, 5(1), 133–137.
- Youden, W.J. (1950). Index for rating diagnostic tests. *Cancer*, 3(1), 32–35.

Zwaagstra, R., Schmidt, I., & Vanier, M. (1996). Recovery of speed of information processing in closed-head-injury patients. *Journal of Clinical and Experimental Neuropsychology*, 18(3), 383–393.

APPENDIX

The formula of the Reliable Chang Index (Maassen et al., 2009; McSweeney et al., 1993) is as follows:

$$RCI = (D_i - D_c + (1 - b_c)(X_i - X_c)) / (S_y(1 - r_{XY}^2)^{1/2})$$

D_i = difference score: post-test score – pre-test score of the individual in the intervention group

D_c = difference score: mean of post scores of the control group – mean of pre scores of the control group

b_c = regression coefficient of post-test on pre-test in the control group

X_i = pre-test score of the individual in the intervention group

X_c = mean of pre-test scores of the control group

S_y = standard deviation of the post-test scores of the control group

r_{XY} = test–retest reliability of the pre- and post-test scores in the control group

$RCI > 1.67$ (one-sided t-test ($N = 64$) with $p < 0.05$) indicates reliably improved