

Twin Studies in Human Genetics

SIGNIFICANCE OF SECULAR BODILY CHANGES OF THE JAPANESE FROM THE VIEWPOINT OF A TWIN STUDY

HISASHI SUZUKI

*Department of Anthropology, National Science Museum,
Tokyo, Japan*

The shapes of the skull of the common people in Tokyo-Yokohama area have undergone secular changes from the protohistoric age to the present day.

The common people in Japan are considered to be the descendants of protohistoric farmers who inhabited and cultivated the same locality from generation to generation. Nevertheless, the mentioned changes can be regarded as gradual ones if skeletal materials of the intermediate ages (5th to 19th century A.D.) are examined.

Some features, such as cranial form, facial form, nasal form, and elevation of the nasal root, have changed significantly, whereas other traits, such as the sum of cranial length and breadth, cranial module, horizontal circumference, and mediosagittal arc of the cranium, have undergone little change.

The results of a twin study are supposed to elucidate the probable reason for these age changes.

Anthropological measurements were made on an unselected sample of 286 MZ and 83 same-sexed DZ twin pairs aged 12 to 14 years, all living in Tokyo Metropolitan area. A total of 18 measurements of the head and face were made. Von Verschuer's percentage deviation method was applied for the data analysis.

The mean intrapair difference is maximum in the sum of the head length and breadth, horizontal circumference, module of the head; medium in bizygomatic breadth; minimum in frontal breadth, etc.; maximum in facial height, length-breadth index of the head, nasal height and breadth, elevation of the nasal root.

The results suggest that features showing

minimum intrapair difference are stable to the environment, whereas features showing maximum intrapair difference are unstable to the environment.

Therefore, of the various changes in the cranial features which have taken place since protohistoric times, the most conspicuous ones correspond to such features that are supposed to be most unstable to the environment. On the contrary, the chronologically little changed ones correspond to the stable features.

Prof. H. Suzuki, Department of Anthropology, 3-23-1 Hyakunin-cho, Shinjuku-ku, Tokyo, Japan

TWINS: GENETIC EFFECTS ON PHYSICAL GROWTH FROM BIRTH TO FOUR YEARS

RONALD S. WILSON

*Louisville Twin Study, School of Medicine, University of
Louisville, Kentucky, USA*

Height and weight growth curves have been developed for twins, based on a sample of 584 twins that were repeatedly measured from birth to four years of age. Sex differences in growth rate were evident at several ages: males grew more rapidly in the first 6 months, whereas females grew more rapidly in the third year. Compared to singletons, the size deficit for twins was greatest at birth, but there was a marked recovery in the first six months, followed by only minor changes thereafter. Weight was somewhat more affected by growth suppression than height; the twins remained 0.45 standard deviations below the singletons for weight, whereas they nearly equalled the singletons for height at age 4. The results suggest that prenatal adipose-cell replication may be restricted by the nutritional demands of a twin pregnancy.

When the sample was separated into MZ twin pairs and DZ same-sex pairs, the MZ twins were found to be less alike for birth