



# Presence of *Echinococcus* eggs in the environment and food: a review of current data and future prospects

Roxanne Barosi<sup>1,2</sup> and Gérald Umhang<sup>1</sup>

<sup>1</sup>ANSES LRFSN, National Reference Laboratory for Echinococcus spp., Malzéville, France and <sup>2</sup>University of Reims Champagne-Ardenne, University of Rouen Normandie, ESCAPE, Reims, France

## Review Article

**Cite this article:** Barosi R, Umhang G (2024). Presence of *Echinococcus* eggs in the environment and food: a review of current data and future prospects. *Parasitology* 1–16. <https://doi.org/10.1017/S0031182024000945>

Received: 15 April 2024

Revised: 18 July 2024

Accepted: 27 July 2024

### Keywords:

dispersion; *Echinococcus granulosus*; *Echinococcus multilocularis*; food; soil; survival; water

### Corresponding author:

Gérald Umhang;

Email: [gerald.umhang@anses.fr](mailto:gerald.umhang@anses.fr)

### Abstract

Cystic and alveolar echinococcosis are considered the second and third most significant foodborne parasitic diseases worldwide. The microscopic eggs excreted in the feces of the definitive host are the only source of contamination for intermediate and dead-end hosts, including humans. However, estimating the respective contribution of the environment, fomites, animals or food in the transmission of *Echinococcus* eggs is still challenging. *Echinococcus granulosus* and *E. multilocularis* seem to have a similar survival capacity regarding temperature under laboratory conditions. In addition, field experiments have reported that the eggs can survive several weeks to years outdoors, with confirmation of the relative susceptibility of *Echinococcus* eggs to desiccation. Bad weather (such as rain and wind), invertebrates and birds help scatter *Echinococcus* eggs in the environment and may thus impact human exposure. Contamination of food and the environment by taeniid eggs has been the subject of renewed interest in the past decade. Various matrices from endemic regions have been found to be contaminated by *Echinococcus* eggs. These include water, soil, vegetables and berries, with heterogeneous rates highlighting the need to acquire more robust data so as to obtain an accurate assessment of the risk of human infection. In this context, it is essential to use efficient methods of detection and to develop methods for evaluating the viability of eggs in the environment and food.

### Introduction

Cystic and alveolar echinococcosis (CE & AE) are considered the second and third most significant foodborne parasitic diseases worldwide (FAO and WHO, 2014). *Echinococcus granulosus sensu lato* (*s.l.*), the causative agent of CE, is found throughout the world, whereas AE, caused by *E. multilocularis*, is generally limited to the Northern Hemisphere (Deplazes *et al.*, 2017; Massolo *et al.*, 2022, Fig. 1). The highest AE disease burden is in Asia, especially China, while that for CE is more widely distributed in parts of Eurasia, Africa and South America (Deplazes *et al.*, 2017). The mortality rates in humans are higher for AE (2–5% in Western and Central Europe and North America, 10–30% in Eastern Europe, and around 100% elsewhere) than for CE (1–2%) (Torgerson *et al.*, 2015). Advances during recent decades have led to a better diagnosis and treatment of echinococcosis notably long term albendazole therapy strongly associated to very low mortality rate but the accessibility to them is still remaining disparate between countries (Torgerson *et al.*, 2008). One study estimates an annual combined figure of 200 000 new human cases of either CE or AE (Kern, 2010; Torgerson *et al.*, 2015). Most research in the past has been focused on detection in definitive and intermediate hosts (DHs & IHs), including dead-end hosts such as humans, mainly for the purpose of determining and describing endemic areas. For instance, the diagnosis of human CE and AE is mainly based on imaging and then may be attested with human blood and tissue samples analysis. Parasitic forms (e.g. eggs, worms, metacestodes and protoscolices) from feces and tissue samples of animal hosts are used to study the epidemiology of the parasite in animals. Indeed, *Echinococcus* eggs, transmitted to humans through the fecal-oral route, are initially shed in DH stools. In Europe, red foxes are considered to be the main definitive host of *E. multilocularis* based on high population densities, high susceptibility to infection, high worm burden and high prevalence of infection reaching more than 50% in high endemic areas (Oksanen *et al.*, 2016; Romig *et al.*, 2017). When the prevalence in other sympatric species notably domestic carnivores is far much lower. Despite a high susceptibility to infection of dogs, their low exposure to infected rodent and occasional deworming limit their global contribution to environmental contamination in Europe (Hegglin and Deplazes, 2013). However, the contribution of dogs in environmental contamination in Central Asia and China is significant due to higher prevalence (Wang *et al.*, 2021; Abdykerimov *et al.*, 2024). At the opposite, if the population densities of cats can be very important with frequent predation of rodents (Forin-Wiart *et al.*, 2018), the low worm establishment and the few eggs excreted which are considered to be non-infective result to an insignificant role of cats in *E. multilocularis* transmission (Kapel *et al.*, 2006). Therefore, according to population densities, prevalence and biotic potential, DHs with the

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.

foremost contribution to *E. multilocularis* environmental contamination by eggs in rural and urban areas are foxes (81–96%), followed by dogs (4–19%) and, to a lesser extent, cats (0–0.2%) (Hegglin and Deplazes, 2013). Thus, the biotic potential of overall of *E. multilocularis* is estimated at between more than 57 000 to several million eggs/km<sup>2</sup> depending on the level of endemicity and urbanization (Hegglin and Deplazes, 2013). Based on the number of eggs produced by *E. granulosus* s.l. worms (~1503 eggs per parasite during its lifetime) and the mean number of worms in infected dogs (~202), the biotic potential in DHs seems to be close between *E. granulosus* s.s. (~303 606 eggs) and *E. multilocularis* (~279 910 eggs) (Thompson and Eckert, 1982; Gemmell, 1990; Kapel *et al.*, 2006). Due to the high density of *E. granulosus* DHs (i.e. domestic and stray dogs), we could expect similar or higher values of environmental contamination by *E. granulosus* eggs in endemic areas than those for *E. multilocularis* in foxes.

The environmental contamination by *Echinococcus* eggs may therefore be high in endemic regions where DH prevalence is high, leading to a risk of human infection. Nevertheless, both food and environmental contamination by *Echinococcus* eggs has been neglected for a long time, despite the fact that the oral uptake of infective eggs from these matrices is the main source of human contamination. It is difficult to estimate the potential contribution of each matrix, such as the environment (e.g. soil, water or fomites, including DH hairs), animals or food as a vehicle for the transmission of *Echinococcus* eggs. The assessments obtained by meta-analysis or based on expert knowledge elicitation ultimately give an insight into the risk factors rather than information on the relative contribution of each compartment in human diseases. Indeed, the confidence intervals (CIs) are usually wide and thus uninformative. For instance, based on the WHO (2015) report, the greatest source of human contamination for *E. granulosus* appears to be contact with animals [52% (CI95%: 22 to 75%)], whereas it might be food for *E. multilocularis* [50% (CI95%: 0 to 88%)]. Meta-analysis results suggest that contaminated water could be responsible for up to 29.4% (CI95%: 12 to 52%) of CE infections in humans, while for AE, the major pathway of transmission could be contact with dogs [34.4% (CI95%: 21 to 48%)] (Torgerson *et al.*, 2020). However, in endemic regions within Europe, the canine prevalence of *E. multilocularis* is clearly lower (0 to 1.5%) (Hegglin and Deplazes, 2013) than that of *E. granulosus* (0 to 19%) (Carmena and Cardona, 2013). On the other hand, in Asia, dogs can play an important role in the *E. multilocularis* life cycle, especially in China or Kyrgyzstan, where high canine prevalence and incidences of human infection have been reported (Budke *et al.*, 2005; Ziadinov *et al.*, 2008; Abydykerimov *et al.*, 2024). Instead of attempting to evaluate the relative contribution of contamination sources, some publications have focused on potential echinococcosis risk factors. Contact with animals, especially DHs, is generally significant for CE and AE (Possenti *et al.*, 2016; Conraths *et al.*, 2017; Torgerson *et al.*, 2020). Depending on the regions and the disease, having a kitchen garden or living in a rural area may also have an impact on *Echinococcus* infection rates. It has also been highlighted that the possibility of ingesting eggs from food (e.g. wild fruits, raw/unwashed vegetables and fruits) or water exists, but these factors do not significantly increase the risk of human infection (Possenti *et al.*, 2016; Conraths *et al.*, 2017). However, in Europe, it appears in the light of case-control studies (Torgerson *et al.*, 2020) that food may be a major source of contamination for AE. The direct link between water or unwashed vegetables/fruits and the human disease is still unclear due to the long incubation time (2–15 years) and in the absence of scientific evidence of food-borne transmission. As a result, it is necessary to clarify these

causal links in order to implement prevention plans and reduce the risk of infection.

Currently, we only have an approximate overview of matrices that play a role in the transmission of *Echinococcus* eggs to humans. After excretion in carnivore feces, a huge quantity of eggs must withstand harsh environmental conditions before they are finally ingested by intermediate or dead-end hosts to pursue their life cycle. Time and fluctuating or extreme field conditions may damage egg integrity and so hamper both viability and infectiousness. So how long do they actually remain viable on average in the environment or on food? What are the contamination rates for the environment and food? Answering these questions with any precision is challenging even today, because despite constant, expanding research on the parasite since the 19<sup>th</sup> century, there are still large gaps in the field (Eckert and Thompson, 2017). It is necessary to determine the presence and viability of eggs on various matrices of interest (e.g. soil, food, or water) in order to establish source attributions and risk factors. In this context, this review summarizes current knowledge on (i) the survival properties of *Echinococcus* spp. eggs; (ii) their capacity for dispersal in the environment; (iii) environmental and (iv) food contamination and finally, (v) gaps and future prospects in the field of interest. In light of the lack of data for *Echinococcus* on some topics, literature on other taeniid may be cited as a model.

### Survival of *Echinococcus* eggs

Eggs, the only free-living stage of the parasite, are the only source of contamination for both intermediate and dead-end hosts. Eggs are infectious, and therefore viable, from their moment of excretion. However, harsh environmental conditions can damage their ability to survive. Thus, the survival capacity of *Echinococcus* eggs were a major focus of studies during the last century, in order firstly to find a way of making them inoperative or killing them, and secondly to propose appropriate control strategies.

### Laboratory conditions

In the early days, various laboratory conditions were tested, such as the effect of temperature (negative and positive), humidity and specific chemicals. As this review focuses on the conditions favourable to egg survival in the environment, resistance to chemicals will not be addressed. Only a few authors have estimated the viability of *E. multilocularis* and *E. granulosus* eggs at different temperatures. However, a wide temperature range (−196°C to 100°C) has been applied to *in vivo* tests (Table 1). According to this literature, it appears that *E. granulosus* and *E. multilocularis* have a similar survival capacity regarding the temperature factor. Although the assay conditions (e.g. time of exposure, inoculation method, host species) may differ, an overall profile may be deduced. The eggs are freeze-resistant to −50°C, but they are killed by very low (<−70°C) and high (>65°C) temperatures. Among these articles, two also compared the survival of *E. multilocularis* eggs in water and air with various humidity rates (Veit *et al.*, 1995; Federer *et al.*, 2015). Eggs seem to be more resistant to positive temperatures in water than in air, even with high relative humidity (rh). For example, at 45°C, eggs remain viable 478 days in water compared with 75 days in air with 85–95% rh (Veit *et al.*, 1995). At 65°C, Federer *et al.* (2015) noted that eggs remained infectious in water up to 1 h, while they were killed after 30 minutes with 70% rh. These results show a susceptibility of *Echinococcus* eggs to desiccation, as already observed for *Taenia* species in the literature. Overall, humidity appears to be more critical to taeniid egg survival than temperature. A low humidity rate, under 34%, hampers egg survival (Jansen *et al.*,

**Table 1.** *In vivo* evaluation of the resistance of *E. multilocularis* and *E. granulosus* eggs to temperature

Temperature (°C)	<i>E. granulosus</i> eggs			<i>E. multilocularis</i> eggs		
	Time of exposure	Survival ( <i>in vivo</i> test)	References	Time of exposure	Survival ( <i>in vivo</i> test)	References
-196	-	-	-	20 h	No (voles)	Veit <i>et al.</i> (1995)
-83	-	-	-	48 h	No (voles)	Veit <i>et al.</i> (1995)
-70	24 h (w)	No (jirds)	Colli and Williams (1972)	4 d (96 h)	No (mice)	Hildreth <i>et al.</i> (2004)
-50	24 h (w)	Yes (mice)	Colli and Williams (1972)	24 h (nsf)	Yes (voles)	Schiller (1955)
-30	24 h (w)	Yes (mice)	Colli and Williams (1972)	24 h (nsf)	Yes (voles)	Schiller (1955)
-26	-	-	-	21 h–54 d (nsf)	Yes (voles)	Schiller (1955)
-20	24 h (w)	Yes (jirds)	Colli and Williams (1972)	-	-	-
-18	-	-	-	106, 240 d (w)	Yes (voles)	Veit <i>et al.</i> (1995)
-15	-	-	-	8 d (tissue)	No (mice)	Hildreth <i>et al.</i> (2004)
-10	24 h (w)	Yes (mice)	Colli and Williams (1972)	-	-	-
2	-	-	-	2.5 y	Yes (mice & squirrels)	Thomas and Babero (1956)
4	24 h (w)	Yes (mice & jirds)	Colli and Williams (1972)	478 d (w) 43, 75 d (85–95% rh)	Yes (voles) Yes (voles)	Veit <i>et al.</i> (1995)
6	45 d 7.5 mo	Yes (squirrel) Yes (chipmunk)	Sweatman and Williams (1963)	-	-	-
45	5 min (w)	Yes (mice)	Colli and Williams (1972)	30 min, 1 h, 2 h, 3 h (w&rh) 1 h, 2 h (85–95% rh) 3 h, +6 h (85–95% rh)	Yes (mice) Yes (voles) No (voles)	Federer <i>et al.</i> (2015) Veit <i>et al.</i> (1995) Veit <i>et al.</i> (1995)
50	5 min (w) >1 h (w)	Yes (mice) No (sheep)	Colli and Williams (1972) Nosik (1952)	30 min, 1 h, 2 h, 3 h (w & rh)	Yes (mice)	Federer <i>et al.</i> (2015)
55	5 min (w)	Yes (mice)	Colli and Williams (1972)	30 min, 1 h, 2 h(w) 2 h (rh), 3 h (w & rh)	Yes (mice) No (mice)	Federer <i>et al.</i> (2015)
60	5 min (w)	No (mice)	Colli and Williams (1972)	30 min (w) 30 min (70% rh)	Yes (mice) No (mice)	Federer <i>et al.</i> (2015)
65	5 min (w)	No (mice)	Colli and Williams (1972)	30 min and 1 h (w) 30 min, 1 h (rh) and 2 h, 3 h	Yes (mice) No (mice)	Federer <i>et al.</i> (2015)
70	5 min (w)	No (mice)	Colli and Williams (1972)	30 min	No (mice)	Federer <i>et al.</i> (2015)
75	5 min (w)	No (mice)	Colli and Williams (1972)	15 min	No (mice)	Federer <i>et al.</i> (2015)
80	5 min (w)	No (mice)	Colli and Williams (1972)	7.5 min	No (mice)	Federer <i>et al.</i> (2015)
100	>20 s (w)	No (sheep)	Nosik (1952)	-	-	-

w, water; rh, relative humidity (70%); d, day; mo, month; y, year; nsf, natural semi-fluid.

2021). A high humidity rate combined with a low or high temperature appears to affect infectiousness more than survival.

### Outdoor conditions

Laboratory conditions have been used to evaluate the great resistance of *Echinococcus* eggs under control and steady conditions. Technically, the temperature thresholds listed above allow the eggs to survive on food or in the environment and then to be transmitted to intermediate and dead-end hosts. Nevertheless, is survival possible under fluctuating outdoor conditions? To answer this question, some scientists have dropped eggs onto the ground, either in feces or nylon bags, and then checked their viability as the seasons passed *via in vivo* (Sweatman and Williams, 1963; Abbasov, 1965; Veit *et al.*, 1995; Thevenet *et al.*, 2005) or *in vitro* tests based on hook motility and the eosin exclusion dye test (Wachira *et al.*, 1991). The maximum survival time of *E. multilocularis* and *E. granulosus* eggs in natural environmental conditions has thus been appraised (Table 2). Temperature ranges are very heterogeneous and vary between regions. No real trend stands out regarding the effect of seasons. Indeed, in the same temperature range (~–16°C to ~10°C), *E. multilocularis* eggs may survive around 3 months (96 days) in Germany (Veit *et al.*, 1995) but less than 1 month for *E. granulosus* in Azerbaijan (Abbasov, 1965). However, the maximum survival time for *E. multilocularis* eggs observed in Germany was 240 days during the autumn/winter and 78 days in summer (Veit *et al.*, 1995). It is therefore very complicated to attempt to summarize or draw conclusions about the maximum survival time due to the difference in timescale, the lack of information in some old research papers and, more generally, the scarcity of studies. Nevertheless, in Kenya, Wachira *et al.* (1991) reached the same conclusion about the harmful effect of desiccation during their field experiments in ‘open ground’, ‘house’ and ‘water hole’. Their results showed that *E. granulosus* eggs are viable but not motile for a longer period (19 days) in the Maasai environment than in Turkana (2 days). This may be explained by the difference in climate between these 2 localities: cool, humid conditions in Maasailand *vs* a semi-arid environment in Turkana. In addition, only eggs in a water hole remained viable and motile for 12 days. This tends to prove that low humidity hampers egg survival, whereas water and high humidity favour survival.

However, the field study in New Zealand led by Sweatman and Williams (1963) pointed out the fact that ‘heavy rainfall was a more important limiting factor in the availability and infectivity of the eggs than was meteorological dryness’. Succinctly, the survival of *E. granulosus* eggs was assessed in 2 localities with different climates: high temperatures and winds, very low rainfall (Waipiata) *vs* extreme rainfall (Lower Kokatahi, LK). Eggs were sprayed over a pasture and then, after either 4 (group 1) or 10 months (group 2), lambs were allowed to graze the pasture for several days. One of the 6 lambs from group 1 became weakly infected in Waipiata compared with 2 of the 6 lambs in LK. None of the lambs from group 2 developed cysts after 10 months. These results appear to show no significant difference between the 2 localities. Nevertheless, the heavy rainfall affected the depth dispersal of the eggs, as described below in the dispersal section.

In addition, the outdoor survival of *E. granulosus* eggs has been compared with some species of *Taenia*, including *T. hydatigena* (Sweatman and Williams, 1963; Wachira *et al.*, 1991). Their life cycles are similar, sharing livestock and dogs as intermediate and definitive hosts, respectively. The *in vivo* study by the aforementioned authors revealed no significant difference in the survival time between eggs of these 2 tapeworms, whatever the climate (Sweatman and Williams, 1963). However, the rate of infection after 4 months was higher for *T. hydatigena* (100% in

W and 83.3% in LK) than *E. granulosus* (16.7% in W and 33.3% in LK). After a year’s exposure to an oceanic climate, most of the eggs seemed to be unavailable and/or were no longer infective. Even though the eggs’ viability had been estimated *in vitro*, the results from Wachira *et al.* (1991) appeared to lead to the same conclusion. By contrast, under various controlled humidity conditions (0–80%), Laws (1968) observed higher *in vitro* survival rates for *T. hydatigena* and *T. ovis* than for *T. pisiformis* and *E. granulosus*. Within these taeniid species, and despite a similar life cycle, there appears to be a difference in their resistance capacity. In the literature, taeniid eggs have globally the same resistance capacity (Jansen *et al.*, 2021 table 1; Table 1) but egg survival rates appear to be heterogeneous and to vary according to outdoor conditions, even for the same species.

### Age effect on *Echinococcus* egg survival

Other factors, such as the age or dispersion of eggs, play a role in the survival and infectivity capacity. For instance, Batham (1957) not only studied the impact of humidity on viability, but also the influence of the age of *E. granulosus* eggs. Briefly, one batch was kept in tap water, and the second in air, both in covered petri dishes at room temperature (between 10°C and 21°C) for 1 year. At various timepoints, the *E. granulosus* eggs were injected intraperitoneally into mice, which were autopsied 3 months post infestation. Unexpectedly, results tended to show that eggs kept in air, unlike eggs kept in water, could remain viable for 1 year. Although no mice developed cysts, ‘spots’ (probably indicating dead remains of cestode cysts) were observed in viscera among mice infected with eggs kept in air. In another study, a drop in egg infectivity was noticed after 41 months, as a small number of cysts (2–7/individual) had developed in the 6 sheep (Thevenet *et al.*, 2005) compared with infection using fresh eggs (24.3 cysts/individual) (Lightowers *et al.*, 1999).

To conclude this part about survival of *Echinococcus* eggs, laboratory and field experiments tend to confirm the relative susceptibility of the eggs to desiccation. Under field conditions, egg survival may depend on specific outdoor conditions and perhaps on the *Echinococcus* species, even if they seem to have a similar resistance profile under stable laboratory conditions. Some papers assert that the eggs are sensitive to freezing conditions, occasionally accompanied by a drop in egg longevity (Slepnev, 1982). However, agricultural practices such as irrigation and food storage could help eggs survive the critical summer months.

The comparison with other taeniid regarding the eggs’ resistance capacities seems reasonable under lab conditions, but is it reasonable to use a *Taenia* species as a model for studying the outdoor survival of an *Echinococcus* species? The answer probably depends on the purpose of the study, because it is hard to compare laboratory and field conditions then draw definitive conclusions. However, the morphology of taeniid eggs is the same for all genera within the Taeniidae family. Therefore, eggs from different *Taenia* species — particularly non-zoonotic ones — can be easily used in the study of dispersal.

### Dispersal of *Echinococcus* eggs in the environment

As seen above in the study of Sweatman and Williams (1963), factors favouring egg survival can also lessen their infectivity by affecting availability in the environment. Indeed, the eggs were dispersed quite differently on each location, as reported for *Taenia* species with the same protocol. Heavy rainfall buried *Taenia hydatigena* and *T. pisiformis* eggs deeper into the soil (11–13 cm) at LK and spread them farther apart than in the near-continent climate of W (0–0.64 cm). Although other authors

**Table 2.** Estimation of the survival of *E. multilocularis* or *E. granulosus* eggs under natural conditions

Reference	Locality	Temperature				Precipitation	Climate	Species	Echinococcus Eggs			
		winter	summer	Maximum survival time					Viability test	Results		
				winter	summer							
Veit <i>et al.</i> (1995)	Southwestern Germany	-15 to 28°C	1.4 to 40°C		~800 mm/year <sup>a</sup>	Temperate continental	<i>E. multilocularis</i>	240 d	78 d	<i>In vivo</i> (voles)	(w) 7/31 (s) 6/11	
Slepnev (1982)	Belorussian SSR	-12°C/-17°C	NC 26°C		~700 mm/year <sup>a</sup>	Cold temperate continental*	<i>E. granulosus</i>	<24 h	13 mo 30 d	NC	90 mm depth topsoil	
Abbasov (1965)	Azerbaijan (Nakhichevan ASSR)	-16 to 10°C	NC		~400 mm/year <sup>a</sup>	Cold semi-arid <sup>a</sup>	<i>E. granulosus</i>	<1 mo	<1 mo	<i>In vivo</i> (lambs)	NC	
Thevenet <i>et al.</i> (2005)	South Argentina, Sarmiento	-3 to 37°C (mean annual = 10°C)			<300 mm/year	Inferior arid	<i>E. granulosus</i>	41 mo (Oct 1996 to Feb 2000)		<i>In vivo</i> (sheep)	4/4 (2-7 cysts/ind)	
Wachira <i>et al.</i> (1991)	Kenya:	-					<i>E. granulosus</i>	-		<i>In vitro</i> <sup>b</sup>		
	Turkana		28-69°C		~500 mm/year <sup>a</sup>	Hot semi-arid			2 d			34% viable
	Maasailand		~20°C		~1000 mm/year <sup>a</sup>	Mediterranean <sup>a</sup>			19 d			33% viable
Sweatman and Williams (1963)	New Zealand:						<i>E. granulosus</i>			<i>In vivo</i> (lambs)		
	Waipiata	-5 to 30°C			229 mm (10 mo)	Quasicontinental			~4 mo			1/6 <sup>c</sup> & 0/6 <sup>d</sup>
	Lower Kokatahi	-3.9 to 14.4°C			2584 mm (10 mo)	Oceanic			~4 mo			2/6 <sup>c</sup> & 0/6 <sup>d</sup>

<sup>a</sup>Information not indicated in the article (pulled from climate-data.org);

<sup>b</sup>Viability was assessed using hook motility and eosin exclusion dye test;

<sup>c</sup>4 months after eggs were sprayed over the pasture;

<sup>d</sup>10 months after eggs were sprayed over the pasture.

d, day; mo, month; NC, not communicated in the abstract; (w), winter; (s), summer.

have studied *E. granulosus*, they have never estimated the eggs' burial depth. However, given the similarity between the morphology of *Taenia* spp. and *Echinococcus* spp. eggs, this observation of vertical dispersal might be relevant. Extreme rainfall can scatter taeniid eggs across the soil and even bury them. Dispersal is both boosted by a linear movement over the topsoil and hampered due to eggs being buried. At the same time, these buried eggs can be protected from harmful environmental conditions and then reemerge when the soil is tilled.

### Role of insects and birds

Furthermore, among animals, invertebrates are considered as potential egg vectors. Several species of wild invertebrates have been found to carry parasitic organisms, including *Taenia* spp. eggs. These include cockroaches (1/820) in Thailand (Chamavit *et al.*, 2011), dung beetles (5/309) in Peru (Vargas-Calla *et al.*, 2018) and flies in various countries (Saelens *et al.*, 2022, table 4). On the other hand, no eggs were observed being carried by flies caught in either Mexico ( $n = 600$  and  $n = 1187$ ) (Keilbach *et al.*, 1989; Martinez *et al.*, 2000) or Iran ( $n = 210$ ) (Hemmati *et al.*, 2018). The most experimental studies currently available on the spreading of *Echinococcus* eggs by invertebrates are focused on flies, with only two on beetles (Benelli *et al.*, 2021, table 2). Diptera and Coleoptera from these studies have successfully transmitted ingested *Echinococcus* eggs to intermediate hosts, which developed infection. Eggs are mainly detected in the invertebrates' gut and occasionally on the body.

Data for taeniids such as *T. hydatigena* are more abundant and widespread (Gemmell, 1976; Gemmell *et al.*, 1978; Benelli *et al.*, 2021) than those on *Echinococcus*, but similarities include egg dispersal. In a field experiment in New Zealand, 24.3% (183/754) of wild blowflies, allowed to feed for 2 to 3 minutes on infested dog feces, contained taeniid eggs in their digestive tract (0–860 eggs/fly) (Lawson and Gemmell, 1985). Only one external attachment of an egg was observed in this study. Most of the eggs tested were still viable after their ingestion by flies, as attested by the development of cysts in naive lambs that were fed these flies (ten flies/lamb). In laboratory conditions, blowflies fed for 2 h on dog feces can ingest up to 5411 taeniid eggs per blowfly. No data are available on the resistance of taeniid eggs in the blowfly digestive tract, nor about their excretion in droppings. However, the viability of taeniid eggs ingested into beetles' digestive tract has been proved (Bilý *et al.*, 1978; Gomez-Puerta *et al.*, 2014). Considering that taeniid eggs excreted by flies after ingestion are still viable, the impact on egg dispersal could be significant. Indeed, according to studies, the daily dispersal capabilities of blowflies in temperate and subtropical regions are 0.10–0.15 km and 1.25–2.35 km, respectively (Lee *et al.*, 2023; Table 1). Moreover, the observation of sheep commonly infected by *T. hydatigena* despite the absence of a definitive host on St Kilda, an island 60 km off the Scottish coast, suggests that eggs may be transported by birds or insects from the nearest coastland (Torgerson *et al.*, 1995). Indeed, some birds may be involved in the dispersal of tapeworm eggs, such as herring gulls for *T. saginata* (Crewe and Crewe, 1969) or coprophagous birds like black-billed magpies for *E. multilocularis* (Raymond and St. Clair, 2023). Furthermore, viable helminth eggs have been found in bird droppings (Crewe, 1967; Crewe and Owen, 1978). The eating behaviour of these birds, combined with their proximity to areas inhabited by humans, may increase the contamination risk for humans.

### Role of climatic factors

Wind can help invertebrates to move greater distances, but does it have a direct impact on the dispersal of *Echinococcus* eggs? No

papers have investigated the role of wind in *Echinococcus* egg dispersal either in the field or in laboratory conditions. Regarding taeniid eggs, Lawson and Gemmell (1985) placed feces contaminated by eggs of *T. pisiformis* 3 cm away from 4 trays filled with water (from 1 cm to 2.6 m). A fan was used ( $23.4 \text{ km h}^{-1}$ ) to simulate wind action for 10 days. Four rabbits were then given the water from the trays to drink, and none became infected, failing to prove in this context a potential effect of wind. However, a mathematical model has theoretically shown that *E. multilocularis* eggs can be spread by wind (Siegert and Neumann, 2022). The average flight distance of eggs in forest areas could vary from 1.3 m to 17 m, depending on factors such as wind speed and altitude. Another study using *T. laticollis* eggs as a model for *Echinococcus* observed dispersal of between 0 m and 12 m from fox feces to wild berries in a boreal forest in Finland (Malkamäki *et al.*, 2022). In Argentina, *E. granulosus* eggs were found 10 m from pens housing experimentally infected dogs. This dispersal distance and area matched the direction of prevailing winds (Sánchez Thevenet *et al.*, 2020). It appears that the dispersal distance also depends on the layout of the studied site. It is reasonable to assume that the distance travelled by an egg is probably higher in open areas than in a forest.

According to the abovementioned literature, bad weather (e.g. rainfall, wind), invertebrates and birds may play a role in the scattering of eggs in the environment. Even if the contribution of each factor in the dispersal of eggs is still unclear and hard to quantify, each one increases the contamination risk for intermediate and dead-end hosts. To a lesser extent, domestic and wild animals can also increase the risk of transmission by carrying eggs on their body (Matoff and Kolev, 1964; Okolo, 1986; Nagy *et al.*, 2011). As DHs are the original source of environmental contamination, they obviously play an important role in the dispersal of taeniid eggs through defecation at different places. The population density of DHs, their movements and behaviour are all parameters that significantly impact the distribution of eggs, so they all need to be further investigated in order to shed light on the presence of *Echinococcus* eggs in the environment and food.

### Environmental and food contamination by *Echinococcus* eggs

Since the early 2000s, there has been a significant increase in the production of data on environmental and food contamination by taeniid eggs as reviewed by Saelens *et al.* (2022). The specific detection of *Echinococcus* species follows the same trend, with a majority of articles published in the last 10 years. *Echinococcus* species cannot be specifically identified by microscopic examination of taeniid eggs, so a molecular method was needed, leading to the tedious and much less sensitive microscopic observation being gradually discarded. Nowadays, environmental and food contamination by *Echinococcus* spp. is mainly detected by targeting parasite DNA, assumed to originate from eggs isolated after filtration and/or flotation methods. As early as 1988, Craig *et al.*, specifically identified for the first time *E. granulosus* among the taeniid eggs isolated from soil and water samples from Turkana in Kenya using an immunological technique. It was not until the early 2000s that new studies on environmental contamination were published, this time systematically using molecular biology to identify *Echinococcus* eggs, with the exception of one study that used an immunological test (Sánchez Thevenet *et al.*, 2003). Investigations into environmental contamination mainly focus on 2 types of matrices: water and soil. In all, 17 studies have been published covering a wide range of geographical areas, from Asia to South America, as well as Europe and Africa, and targeting either *E. multilocularis* or *E. granulosus*.

## Water

There are currently 7 studies that investigate the specific detection of *Echinococcus* eggs in water. Four of them focus on the detection of *E. multilocularis* in water, while the others (all concerning *E. granulosus*) took place in a broader context with other matrices such as soil and feces also being tested (Table 3). The isolation method for eggs in water typically involves filtration, though a flotation method has also been successfully used once. However, the performance of these methods using 10 mL to 50 L of water has been evaluated only for the 2 methods used by Lass *et al.* (2019, 2020) resulting in high limits of detection, with 100 and 200 eggs for 10 L. According to the studies available, *Echinococcus* detection rates in water vary mainly between 0 and 6%. Higher values were obtained in only 2 particular situations with a very small number of samples tested: 60% ( $n=5$ ) of samples collected from open water sources in Kenya (Craig *et al.*, 1988) and 20% ( $n=10$ ) in water samples corresponding to the washing of cattle on a farm on the Tibetan plateau (Lass *et al.*, 2020). The marked heterogeneity of the sampling, both in terms of water type and number of samples collected, complicates the interpretation of contamination rates. However, it appears (as expected) that eggs were detected mainly in stagnant water (lakes, waterholes) rather than flowing water (rivers, canals). The presence of eggs in contaminated water can contribute to the maintenance of the parasitic life cycle, but except in certain areas of Africa and Asia, such water is not intended for human consumption. However, it may contribute to soil contamination and may also contaminate vegetables when used for irrigation, which may indirectly cause human cases. Currently there is no specific data about contamination of wastewater by *Echinococcus* eggs, even though the presence of taeniid eggs has already been asserted in some studies (Saelens *et al.*, 2022). There has been a recent increase in interest in waterborne *Echinococcus* detection, motivating 6 of the 7 studies carried out in the last 5 years. With this new interest, it is likely that more studies will follow, providing a better understanding of the role of water in human contamination.

## Soil

Detection of *Echinococcus* in soil has been more frequently reported than in water. A total absence of detection was only reported for 2 of the 10 studies published (Table 4). The first of these 2 studies collected only 2 samples of 36 kg each from the car deck of ferries in order to investigate the potential transfer of *E. multilocularis* eggs by cars from the endemic island of Hokkaido to a bigger island in mainland Japan (Matsuo *et al.*, 2003). The second one analysed 33 samples of 400 g each from educational farms on the island of Sardinia, Italy (Serra *et al.*, 2022), which is highly endemic for *E. granulosus*. For both studies, the sample mass was significantly higher than the other 8, which ranged only between 5 g and 40 g of soil. Unfortunately, the performance of the 9 different methods used by the various studies was evaluated only for 2. One of these reports an increase in the detection limit from 10 g to 20 g (Umhang *et al.*, 2017) with an increase in the soil quantity analysed. According to the studies available, detection rates ranged from 2.7 to 25% for *E. granulosus* and 10.4 to 11.7% for *E. multilocularis*. Contamination has also been detected both inside the huts of the Turkana community, a nomadic tribe in Kenya (Craig *et al.*, 1988), and in vegetable gardens such as in Kazakhstan for *E. granulosus* (Shaikenov *et al.*, 2004) and in Europe for *E. multilocularis* (Szostakowska *et al.*, 2014; Umhang *et al.*, 2017; Da Silva *et al.*, 2021), which can lead to human contamination. These results indicate relatively significant soil contamination, especially when considering it represents contamination by at least one egg from a very small

amount of soil, demonstrating both the eggs' ability to disperse and their high numbers in the environment.

## Food

*Echinococcus* species have only recently been sought in food, despite *E. multilocularis* and *E. granulosus* being considered food-borne parasites for several decades. In Europe, the detection of *Echinococcus* spp. in food mainly focuses on *E. multilocularis*, probably due to its greater impact on human health (Devleeschauwer *et al.*, 2017). The parasite is found on salad vegetables and berries (e.g. strawberries and blueberries), with contamination rates globally ranging from 0 to 4% and 0 to 7%, respectively (Table 5). Among the ten European endemic countries surveyed in the 9 papers concerned, *E. multilocularis* was identified in at least one food item from 8 countries. Contaminated food items were mainly identified in the historical geographical focus, France and Switzerland, but also in Eastern Europe with Estonia, Latvia and Poland (Lass *et al.*, 2015; Guggisberg *et al.*, 2020; Umhang *et al.*, 2024). Nevertheless, detections were also reported in countries considered endemic but at low rates, such as Denmark and the Netherlands (Umhang *et al.*, 2024), and unexpectedly in locally produced mixed salad vegetables from southern Italy, an area not currently known to be endemic (Barlaam *et al.*, 2021). Thus, while the contamination of food by *E. multilocularis* appears to mainly affect highly endemic areas, the risk of contamination cannot be ruled out for endemic areas with lower rates due to the patchy distribution of the parasite. The level of *E. multilocularis* contamination of lettuces from endemic areas in Europe is globally estimated to be 1% according to the 3 studies focusing on this food item (Guggisberg *et al.*, 2020; Barlaam *et al.*, 2021; Umhang *et al.*, 2024). For other types of vegetables (mainly chards, parsley and spinach), a similar global value of 1.8% was found in the multicentre study of the OHEJP Meme project (Umhang *et al.*, 2024). However, these values are much lower than the value of 30.7% obtained for various vegetables, excluding lettuces, in a highly endemic area in Poland (Lass *et al.*, 2015). This latter study revealed the first detection of *Echinococcus* spp. in food. The high levels of contamination by *E. multilocularis* found in various types of samples have been questioned and debated in several papers (Lass *et al.*, 2016; Robertson *et al.*, 2016; Torgerson, 2016). The authors had suggested that choosing a sampling site in a hyper-endemic zone could help explain the high contamination levels, which would probably not be the same as those observed more globally in a given region. While the contamination rate for vegetables other than salad vegetables is very different from that obtained subsequently, it is not currently possible to compare mushrooms (36%) as this item was not targeted in other studies. However, it is possible to compare berries. Lass *et al.* (2015) reported contamination rates of 9.4% for forest fruits (raspberries, cranberries, blueberries, cowberries) collected in the forest and 20% for cultivated raspberries. No contamination by *E. multilocularis* has been reported on berries in some studies (Malkamäki *et al.*, 2019; Temesgen *et al.*, 2022) and especially on berries ( $n=30$ ) or salad vegetables ( $n=74$ ) from areas with a high and low fox prevalence in Poland (Umhang *et al.*, 2024). However, during this same multicentre study (Umhang *et al.*, 2024), an overall contamination rate by *E. multilocularis* of 5.4% for strawberries and 7.3% for blueberries was obtained for the 7 endemic European countries concerned. The highest proportions were found in the 2 Balkan countries sampled, with 13.3% on strawberries ( $n=30$ ) and 13.3% on blueberries ( $n=30$ ) in Latvia, and 16.7% on strawberries ( $n=30$ ) in Estonia. When few strawberries were collected ( $n=21$ ) during this study, the only positive batch was identified among the 4 from Switzerland. Consequently, recent data

**Table 3.** Data from studies concerning the detection of *Echinococcus* species in water

<i>Echinococcus</i> species targeted	Location of sampling	Type and number of water samples	Proportion of contamination	Volume of one sample	Method for isolating eggs	Method of identification	Limit of detection	Reference
<i>E. granulosus</i>	Kenya: Turkana	Open waterholes: 5	60% (3)	200 mL	Filtration	Immunofluorescence test	Not evaluated	Craig <i>et al.</i> (1988)
<i>E. multilocularis</i>	Poland: Warmia-Masuria and Pomerania Provinces	Lake: 49 rivers and canals: 24 groundwater: 32	4.1% (2) 0 0	50 L	Filtration	Nested PCR and sequencing, real-time PCR	100 eggs for 10 L	Lass <i>et al.</i> (2019)
<i>E. multilocularis</i>	China: Qinghai-Tibet Plateau	Rivers: 26 wastewater treatment plants: 137 slaughterhouse: 49 cattle farm: 10	0 5.8% (8) 6.1% (3) 20% (2)	10 L	Flocculation filtration	Nested PCR and sequencing, real-time PCR, LAMP PCR	200 eggs for 10 L	Lass <i>et al.</i> (2020)
<i>E. granulosus</i> , <i>E. multilocularis</i> <sup>a</sup>	Nigeria: Ibadan	Open water sources: 50	2% (4)	10 mL	Flotation	Microscopy and multiplex PCR	Not evaluated	Awosanya <i>et al.</i> (2022)
<i>E. granulosus</i> , <i>E. multilocularis</i> <sup>a</sup>	Italy: Sardinia	Well: 15 cistern: 8 public supply: 10 stream: 2	0	1 L	Filtration	Multiplex PCR	Not evaluated	Serra <i>et al.</i> (2022)
<i>E. multilocularis</i>	Japan: Hokkaido	Stream water: 128	0.8% (1)	900 mL	Filtration	Real-time PCR	Not evaluated	Mori <i>et al.</i> (2023)
<i>E. multilocularis</i>	Armenia	Raw water: 24 sediments: 2	0 0	10 L	Filtration	Multiplex real-time PCR <sup>a</sup>	Not evaluated	Shcherbakov <i>et al.</i> (2024)

<sup>a</sup>Studies assuming the non-endemic status for *E. multilocularis* in the country but detection conducted in the context of a multiplex PCR including other parasite species.



**Table 4.** Data from studies concerning the detection of *Echinococcus* species in soil

<i>Echinococcus</i> species targeted	Location of sampling	Type and number of soil samples	Proportion of contamination	Mass of one sample	Method for isolating eggs	Method of identification	Limit of detection	Reference
<i>E. granulosus</i>	Kenya: Turkana	Inside huts and surroundings: 28	25% (3)	10 g	Filtration	Immunofluorescence test	Not evaluated	Craig <i>et al.</i> (1988)
<i>E. multilocularis</i>	Japan: Aomori	From car deck on ferry-boats: 2	0	36 kg	Filtration and flotation	Microscopy	Not evaluated	Matsuo <i>et al.</i> (2003)
<i>E. granulosus</i>	Kazakhstan: Almaty Oblast	Kitchen gardens: 120	4.1% (5)	5 g	Filtration and flotation	Microscopy and PCR	Not evaluated	Shaikenov <i>et al.</i> (2004)
<i>E. multilocularis</i>	Poland: Varmia-Masuria Province	Wild areas: 29 kitchen garden: 18 farm yards: 15	10.3% (3) 11.1% (2) 13.3% (2)	40 g	Filtration and flotation	Microscopy, PCR and sequencing	10 eggs	Szostakowska <i>et al.</i> (2014)
<i>E. granulosus</i>	Argentina: Sarmiento	Enclosures of experimentally infected dogs: 48 surrounding: 152	40% (19) <sup>b</sup> 0.7% (1) <sup>b</sup>	2 g	Filtration	Microscopy and western blot	Not evaluated	Sánchez Thevenet <i>et al.</i> (2020)
<i>E. multilocularis</i>	France: Moselle and Ardennes	Kitchen garden: 126	11.7% (26)	10 g	Filtration and flotation	Real-time PCR	10 eggs	Umhang <i>et al.</i> (2017)
<i>E. multilocularis</i>	France: Doubs and Ardennes	Urban kitchen garden: 213 rural kitchen garden: 250	0 10.4% (26)	10 g	Umhang <i>et al.</i> (2017)	Umhang <i>et al.</i> (2017)	10 eggs	Da Silva <i>et al.</i> (2021)
<i>E. granulosus</i>	China: Xizang	Kennel: 37	2.7% (1)	15 g	Filtration	PCR and sequencing	Not evaluated	Liu <i>et al.</i> (2022)
<i>E. granulosus</i> , <i>E. multilocularis</i> <sup>a</sup>	Nigeria: Ibadan	Increasing distances from slaughterhouse (5 m to 10 km): 200	8% (16)	20 g	Flotation	Microscopy and multiplex PCR	Not evaluated	Awosanya <i>et al.</i> (2022)
<i>E. granulosus</i> , <i>E. multilocularis</i> <sup>a</sup>	Italy: Sardinia	Farm: 33	0	400 g	Filtration and flotation	Microscopy and multiplex PCR	Not evaluated	Serra <i>et al.</i> (2022)

The presence of a citation for the egg isolation and identification methods means that it is the same method used in the study cited.

<sup>a</sup>A study that assumed the non-endemic status for *E. multilocularis* in the area but detection conducted in the context of a multiplex PCR including other parasite species.

<sup>b</sup>Data from four collections 0, 41, 70 and 84 months after the experimental infection of dogs were pooled.

**Table 5.** Data from studies concerning detection of *Echinococcus* species in food

<i>Echinococcus</i> species targeted	Location of production	Purchasing entity	Type and number of food items sampled	Proportion of contamination	mass of one sample	Method for isolating eggs	Method of identification	Limit of detection	Reference
<i>E. multilocularis</i>	Poland: Varmia-Masuria Province	From the field	Mushrooms: 25 forest fruits: 32 vegetables: 26 raspberries: 20	36% (9) 9.4% (3) 30.7% (8) 20% (4)	Berries: 300–500 g vegetables: 500 g	Tween washing, filtration (up to 50 µm), flotation	Nested PCR and sequencing	100 eggs	Lass <i>et al.</i> (2015)
<i>E. multilocularis</i> , <i>E. granulosus s.l.</i>	Switzerland: Basel	From the field	Mainly salad vegetables and varying quantities of other vegetables and fruits: 40	0%	50 kg of vegetables plus 10 kg of fruits	Tap water washing, filtration (up to 21 µm)	Multiplex PCR and sequencing	Not evaluated	Federer <i>et al.</i> (2015)
	Various European countries	Local markets	Mainly salads and varying quantities of other vegetables and fruits: 46	<i>E. multilocularis</i> : 0% <i>E. granulosus s.l.</i> : 4.3% (2)					
<i>E. multilocularis</i>	Poland: Pomerania province	From the field	Mushrooms: 32 forest fruits: 17 vegetables: 21	15.6% (5) 0% 0%	Berries: 300–500 g vegetables: 500 g	Lass <i>et al.</i> (2015)	Lass <i>et al.</i> (2015)	100 eggs	Lass <i>et al.</i> (2017)
<i>E. multilocularis</i> , <i>E. canadensis</i>	Finland	Supermarkets	Bilberries 21	0%	250 g	Tween washing, filtration (up to 20 µm)	Real time PCR	3 eggs	Malkamäki <i>et al.</i> (2019)
	Estonia	Unknown	Bilberries 21	0%					
<i>E. granulosus s.l.</i>	Tunisia: Sousse, Mahdia, Monastir, and Gafsa	Local markets	Lettuce: 40 parsley: 40 chard: 40 spinach: 40 celery: 40 mint: 40	2.5% (1) 0% 5% (2) 0% 0% 0%	100 g	NaCl washing, sedimentation, sucrose flotation	PCR and sequencing	Not evaluated	M'Rad <i>et al.</i> (2020)
<i>E. multilocularis</i> , <i>E. granulosus s.l.</i>	Italy: Sardinia	Local farms	Mainly broad-leaved: 23	0%	350–500 g	Water washing, filtration (up to 50 µm)	Multiplex PCR and sequencing	Not evaluated	Serra <i>et al.</i> (2022)
<i>E. multilocularis</i>	Norway, but mostly imported from Poland, Belgium, Sweden, Netherlands and 8 non endemic countries	Local and supermarkets	Blueberries: 274 raspberries: 276 strawberries: 270	0%	30 g	Alconox washing, sedimentation	Real-time PCR	5 eggs	Temesgen <i>et al.</i> (2022)
<i>E. multilocularis</i> <sup>a</sup>	Colombia: Bogota	Local and supermarkets	Strawberries: 240	0%	30 g	Temesgen <i>et al.</i> (2019)	Real-time PCR	5 eggs	Ortiz Pineda <i>et al.</i> (2020)
<i>E. multilocularis</i> , <i>E. granulosus s.l.</i>	Switzerland	Local and supermarkets	Lettuce: 157	1.3% (2)	300 g	Tween washing, filtration (up to 20 µm)	Multiplex PCR and sequencing	2 eggs	Guggisberg <i>et al.</i> (2020)
<i>E. multilocularis</i>	Italy	Supermarkets	Ready-to-eat mixed salad vegetables: 324	0.15% (1)	100 g	Temesgen <i>et al.</i> (2019)	Real-time PCR with in parallel microscopic observation (FLOTAC)	5 eggs but not evaluated for FLOTAC	Barlaam <i>et al.</i> (2021)
			Raspberries: 108	0%					
	Mexico		Blackberries: 108	0%					
	Peru		Blueberries: 108	0%					

<i>E. multilocularis</i> <sup>b</sup> , <i>E. granulosus s.l.</i> <sup>b</sup>	Europe: Denmark, Estonia, Finland, France, Germany, Italy, Latvia, the Netherlands, Norway, Poland, Portugal, Switzerland	From the field, local and supermarkets	Lettuce: 570 other vegetables: 71 strawberries: 202 blueberries: 82 other berries: 38	<i>E. multilocularis</i> : 1.2% (7) 1.8% (1) 5.4% (11) 7.3% (6) 2.6% (1)	Lettuce: 300 g  berries: 100–250 g	Guggisberg <i>et al.</i> (2020)	Real-time PCR	3 eggs	Umhang <i>et al.</i> (2024)
			Lettuces: 942 others vegetables: 71 strawberries: 284 blueberries: 105 others berries: 50	<i>E. granulosus s.s.</i> : 1.3% (10) 0% 1.5% (3) 1.3% (1) 0%					
	Tunisia	Local markets	Lettuces: 75 strawberries: 16	<i>E. granulosus s.s.</i> : 12% (9) 81.3% (13)					
Pakistan	Local markets	Lettuces: 100 blueberries: 25 lettuces: 100 blueberries: 25	<i>E. granulosus s.l.</i> : 4% (4) 12% (3) <i>E. multilocularis</i> : 2% (2) 56% (14)						

<sup>a</sup>Studies that assumed the non-endemic status for *E. multilocularis* in the country but detection conducted in the context of a multiplex PCR including other parasite species.

<sup>b</sup>Proportions were calculated considering only those European countries for which areas endemic for *E. multilocularis* have been sampled (Denmark, Estonia, France, Germany, Latvia, the Netherlands, Poland and Switzerland) and European countries where a domestic lifecycle is established for *E. granulosus* (Estonia, France, Italy, Latvia, Norway, Poland, Portugal).

The presence of a citation for the egg isolation and identification methods means that it is the same method used in the study cited.

acquisitions appear to confirm a high rate of *E. multilocularis* contamination among berries in some areas (Umhang *et al.*, 2024), as initially observed in Poland (Lass *et al.*, 2015). However, this does not seem to be the case for vegetables other than salad vegetables, contrary to the high values obtained from the initial study in Poland. The observation of a significant difference in the rate of contamination between salad vegetables and berries is also supported by data from Pakistan, where *E. multilocularis* contamination affected 2% of salad vegetables ( $n=100$ ) and 56% of blueberry batches ( $n=25$ ) mainly from the same region (Umhang *et al.*, 2024).

From these same samples from Pakistan, a similar significant trend in favour of the higher contamination of berries than lettuces was observed for *E. granulosus s.l.* There was a 4% contamination rate for salad vegetables ( $n=100$ ) and a 12% contamination rate for blueberries ( $n=25$ , all *E. granulosus s.s.*, except for one case of *E. canadensis*). A significant difference in contamination between salad vegetables and berries was also described in Tunisia, a hyper-endemic area for *E. granulosus s.s.*, with 12% of salad vegetables ( $n=75$ ) contaminated and 81.3% ( $n=16$ ) of strawberries (Umhang *et al.*, 2024). However, in this specific situation, it is important to note that the strawberries all came from the region of Cap Bon in the northeast of the country, whereas the lettuces came from other parts of Tunisia that had a similar rate of contamination. Previously, *E. granulosus s.s.* eggs were only observed in 2% of the salad vegetables ( $n=40$ ) and 5% of the chards ( $n=40$ ) out of the various vegetables tested from local markets in 4 Tunisian cities (M'rad *et al.*, 2020). These lower proportions may be explained by the use of microscopic egg screening prior to molecular identification, a method whose performance was not evaluated in the study. In Europe, the average proportion of *E. granulosus s.l.* contamination of salad vegetables can be estimated at 1%, with similar values for strawberries (1.1%) and blueberries (1%) according to the 5 studies concerned (Federer *et al.*, 2015; Malkamäki *et al.*, 2019; Guggisberg *et al.*, 2020; Serra *et al.*, 2022; Umhang *et al.*, 2024). The cases reported concern only identified endemic countries in the south (Italy and Portugal) and east of Europe (Latvia, Estonia and Poland), although uncertainty remains about the exact origin within Europe of the cases detected by Federer *et al.* (2016) in lettuces bought in Switzerland. While several European countries with low or very low endemic levels were involved without resulting in detection, the proportion reaches higher values for the countries from highly endemic areas, with an average proportion of 1.9% ( $n=469$ ). For instance, in Italy a total proportion of 3.2% of contaminated salad vegetables ( $n=232$ ) was reported, with detections ranging from 1.3% in Abruzzo to 3.8% in Sardinia and 6.5% in Campania (Umhang *et al.*, 2024).

To sum up, it is only in the past decade that most data on the contamination of food by *Echinococcus* spp. have been produced. The detection rate of *E. multilocularis* and *E. granulosus s.l.* DNA in food mainly ranges from 0 to 5%, but can occasionally reach higher values, especially in berries. The large variation in the few available data reflects the heterogeneity of the studies carried out in terms of the variety of matrices, epidemiological context, number of samples tested per type of matrix and methods used. The food samples tested include various types of berries and a wide range of leafy vegetables, mainly lettuces. Four studies focused on one or two types of food items, which generally allows a high number of samples to be collected, resulting in an accurate estimation of the presence or absence of contamination, as opposed to sampling several different types of vegetables, often in small quantities for each. For now, it may be more appropriate to prioritize the production of robust data for specific food items rather than testing a large panel but with very few samples of

each, as it would be difficult to draw relevant conclusions. The quantities of vegetable and berry matrices tested varied mostly between 30 g and 500 g, which can greatly impact the result depending on the method used. In this way, out of the 11 available studies, only six different methods were employed with detection limits ranging from 3 to 100 eggs, although this limit was not estimated for half of the methods used. However, based on current data, the contamination rate by *E. multilocularis* and *E. granulosus s.s.* appears to be significantly higher on berries than on salad vegetables and probably on other vegetables. Further studies conducted in various epidemiological situations will be necessary to clarify this preliminary trend.

## Current gaps and future prospects

### The need to use efficient detection methods

Various studies conducted over the past decade have provided much-needed data on the contamination of the environment and food by *Echinococcus* eggs. Despite the heterogeneity of the acquired data, it is possible to obtain a first overview of the contamination level for the different matrices tested. Overall, the contamination of soil, where eggs are initially deposited via feces, appears to be generally greater than in water or food following their dispersion. However, there is a need to consolidate these data to obtain more accurate estimates of contamination in different matrices. The multiplication of detection methods to detect taeniid eggs in the same matrix complicates the comparison of results (Saelens *et al.*, 2022). To achieve the goal of using standardised methods validated by an inter-laboratory ring trial (Hazards (BIOHAZ) *et al.*, 2018) it is currently crucial to employ only performance-evaluated methods for each matrix type taking great care to prevent contamination of samples during molecular analyses. The limit of detection may vary for the same method, for example, between leafy vegetables and berries, or between different types of water or soil. A limit of detection that is too high will result in an underestimation of the level of contamination. Currently, there are limited data available on the number of eggs detected in environmental or food samples, which is typically in the range of several dozen eggs per sample (Craig *et al.*, 1988; M'rad *et al.*, 2020; Temesgen *et al.*, 2022). Although smaller quantities (<10 eggs) have also been observed (Shaikenov *et al.*, 2004; Barlaam *et al.*, 2021), the lower sensitivity of microscopic observation compared with molecular methods suggests that these high quantities of eggs observed do not correspond to the majority of contaminations, but only to the few most significant ones. There are currently no data on the minimum number of eggs that could lead to human infection (Alvarez Rojas *et al.*, 2018). It may be difficult to systematically detect a single egg, as evidenced by the most effective methods currently requiring 2–5 eggs from food matrices even though fewer eggs may be detected but with a lower frequency. The precise quantification of the number of eggs present in soil, water and food samples is currently unknown but necessary to define relevant detection limits for methods. The amount of material tested for each sample is also important. Large quantities lead to methodological constraints in handling large volumes and generally result in a loss of sensitivity. This drawback can be compensated for by increasing the number of samples, probably giving a more relevant result but leading to a longer analysis time and higher cost.

### Pursuing the acquisition of data on environmental and food contamination

In the coming years, the use of efficient methods will lead to more information being gathered on the contamination of soil, water, vegetables and berries. Conducting studies simultaneously seeking

*Echinococcus* spp. and taeniid eggs in different matrices should shed light on the relative contamination of soil, water and food, as previously undertaken in some studies (Craig *et al.*, 1988; Awosanya *et al.*, 2022; Serra *et al.*, 2022). It will also be necessary to investigate more specifically the differences in contamination between various types of leafy vegetables and berries. For instance, a difference in the attachment of *Taenia laticollis* eggs has been described between 2 types of berries (bilberries and cowberries) in the boreal forests of Finland (Malkamäki *et al.*, 2022). Such studies could potentially identify foods that are more at risk. Furthermore, understanding these differences in interactions between *Echinococcus* spp. eggs and the surfaces of different foods could lead to an adaptation of domestic and industrial washing methods. Currently, the effectiveness of washing methods has never been evaluated, despite washing being one of the main recommendations for preventing *Echinococcus* food contamination. In the current context of global trading, there is also the question of the risk to humans of contamination due to the importation of vegetables or berries from hyper-endemic areas. *E. granulosus* has been detected on salad vegetables purchased in Switzerland but originating from other European countries (Federer *et al.*, 2016). Similarly, contamination by protozoa from imported berries has also been described in Italy and Norway (Barlaam *et al.*, 2021; Temesgen *et al.*, 2022). In addition, the favourable storage conditions of these exported foods promote the survival of eggs by maintaining a high level of humidity.

Furthermore, most of the articles have extended the identification of parasites by molecular biology beyond *Echinococcus* spp. to include other species of the Taeniidae family (genus *Taenia*, *Hydatigera* and *Versteria*) as well as protozoa (such as *Toxoplasma*, *Cyclospora* or *Cryptosporidium*). Only 2 studies conducted by the same group focus exclusively on one species (i.e. *E. multilocularis*) (Lass *et al.*, 2015, 2017). Although the chosen washing methods can isolate a wide range of nematodes, cestodes and protozoa, it is important to note that the following steps of flotation and/or filtration and especially the molecular tests usually focus on only a few priority species due to their zoonotic potential. However, conducting a more comprehensive screening of parasite species can provide valuable insights into the contamination of food by parasite eggs, oocysts and cysts. To address the limitations and challenges of food sampling, a multi-species approach that encompasses more than one species or parasitic group would be more appropriate. This approach is already used by some groups notably using multiplex PCR to identify species in the Taeniidae family (Trachsel *et al.*, 2007) or to identify a larger spectrum, covering *Cyclospora* sp., *T. gondii* and *E. multilocularis* (Temesgen *et al.*, 2019). In regards to *Echinococcus*, detecting *E. granulosus* s.l. may always appear relevant due to its global distribution. While not systematically testing for *E. multilocularis* may be justified on non-endemic continents (i.e. Africa and South America), it would appear to be of primary interest elsewhere, even in areas not previously known to be endemic, as it may involve imported food or incidental detection, as in the case of locally produced salad vegetables in southern Italy (Barlaam *et al.*, 2021).

Studying seasonal variations in environmental contamination rates would provide a better understanding of the dynamics of the presence of eggs in the environment. For instance, in Europe, the prevalence of *E. multilocularis* in foxes is generally higher in autumn and winter than in spring and summer, indirectly reflecting the proportion of rodents in the fox's diet (Burler *et al.*, 2011; Otero-Abad *et al.*, 2017). Does this supposed increase in environmental contamination still contribute to the risk of food contamination, which is mainly associated with warmer periods corresponding to food production? While egg dispersion can lead to their presence in cultivated areas, it is important to

know whether eggs are still infectious after several months in the environment. Although available data demonstrate that this is possible, it is by no means systematic, as evidenced by the success rates of vole infestations ranging from 0 to 54% after infection by *E. multilocularis* eggs that had spent several months outdoors in experiments conducted by Veit *et al.* (1995). To more accurately assess environmental contamination, it appears necessary to estimate the proportion of eggs that are still viable rather than simply detecting their presence.

### Essential need to assess the viability of *Echinococcus* eggs

Currently, data on environmental and food contamination by *Echinococcus* spp. are based on detecting parasitic DNA from eggs. However, it has not been possible to assess the viability of these eggs, which is a major issue in evaluating the risk of human contamination. Out of all the available studies, only one attempted – unsuccessfully – to evaluate the viability of eggs detected on lettuces (Guggisberg *et al.*, 2020). This currently represents a significant methodological barrier, whether using *in vitro* or *in vivo* methods. Regarding the *in vitro* approach, hatching and activation of taeniid eggs in artificial gastric fluid may overestimate infectivity as eggs first lose their infective ability when they age (Coman and Rickard, 1977; Hazards (BIOHAZ) *et al.*, 2018). On the other hand, the comparison between oral, intraperitoneal and subcutaneous infection of mice showed that the latter route was the most sensitive, allowing for systematic parasite development (4/4) of up to 20 *E. multilocularis* eggs (Federer *et al.*, 2015). However, this detection threshold still appears too high when it comes to environmental or food contamination, bearing in mind that this threshold has not yet been established for *in vitro* methods. In addition to ethical considerations regarding animal welfare, replacing *in vivo* by *in vitro* methods would be appropriate, especially as some limitations can be successfully managed in this case (Alvarez Rojas *et al.*, 2018). A similar *in vitro* method already exists for protozoa such as *T. gondii* and *Cryptosporidium parvum*, using CC-qPCR based on the development of oocysts in cell culture followed by real-time PCR detection to indirectly demonstrate parasitic proliferation (Rousseau *et al.*, 2019; Géba *et al.*, 2021; Kubina *et al.*, 2021). The availability of a suitable method to assess the viability of *Echinococcus* spp. eggs is essential to be able to fully exploit the detection data from environmental and food matrices so as to more accurately evaluate the risk of human contamination.

### Conclusion/future directions

During the last decade, there has been a renewed interest in studying environmental and food contamination by *Echinococcus* spp. eggs, which has provided an overview of the level of contamination in different matrices such as water, soil, berries and raw vegetables. Given the heterogeneity of the available data, it is necessary to continue acquiring information from different sources and in different epidemiological situations. However, it is essential to use methods whose performance has been evaluated and is relevant to the number of contaminated eggs and the type of sample involved. The quantitative data on egg contamination per sample is currently unknown, and must be estimated. Additionally, the development of standardised methods, ideally not limited to the *Echinococcus* genus, would facilitate the comparison of results on parasitic contamination. In the meantime, it is necessary to have accessible and efficient methods for producing data in hyper-endemic areas where economic resources are often limited. However, even in the presence of sensitive and robust methods for environmental or food detection, surveillance/monitoring of *Echinococcus* spp. via feces, where eggs are

concentrated in large quantities, seems far more relevant (Alvarez Rojas *et al.*, 2018). A better understanding of environmental contamination would also require additional data on the mechanisms involved in the dispersion of eggs, and especially the ability to estimate their viability on different matrices with a view to assessing the risk of human echinococcosis infection. If data are more and more available about the understanding and level of environmental and food contamination by *Echinococcus* eggs, there is still a long road to obtain an accurate estimation of the relative contribution of the different transmission pathway to human infection.

**Data availability statement.** Data will be made available on request.

**Author contributions.** G. U. and R. B. conceived and designed the study. G. U. and R. B. conducted data gathering. G. U. and R. B. wrote the article.

**Financial support.** This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

**Competing interests.** None.

**Ethical standards.** Not applicable.

## References

- Abbasov MT (1965) Periods of survival on pasture of the oncospheres and bladder-worms of *Multiceps multiceps*, *Echinococcus granulosus* and *Taenia hydatigena*. *Veterinariia* **9**, 46–47. Available at <https://eurekama.com/research/014/601/014601145.php> (accessed 19 January 2024).
- Abdykerimov KK, Kronenberg PA, Isaev M, Paternoster G, Deplazes P and Torgerson PR (2024) Environmental distribution of Echinococcus- and Taenia spp.-contaminated dog feces in Kyrgyzstan. *Parasitology* **151**, 84–92.
- Alvarez Rojas CA, Mathis A and Deplazes P (2018) Assessing the contamination of food and the environment with Taenia and Echinococcus eggs and their zoonotic transmission. *Current Clinical Microbiology Reports* **5**, 154–163.
- Awasanya EJ, Olagbaju A, Peruzzo A, Masu G, Masala G and Bonelli P (2022) Detection of *Echinococcus granulosus* sensu lato in Environmental Samples from Ibadan, Oyo State, South West Nigeria. *Veterinary Sciences* **9**, 679.
- Barlaam A, Temesgen TT, Tysnes KR, Rinaldi L, Ferrari N, Sannella AR, Normanno G, Cacciò SM, Robertson LJ and Giangaspero A (2021) Contamination of fresh produce sold on the Italian market with *Cyclospora cayatanensis* and *Echinococcus multilocularis*. *Food Microbiology* **98**, 103792.
- Batham EJ (1957) Notes on viability of hydatid cysts and eggs. *New Zealand Veterinary Journal* **5**, 74–76.
- Benelli G, Wassermann M and Brattig NW (2021) Insects dispersing taeniid eggs: who and how? *Veterinary Parasitology* **295**, 109450.
- Bilý S, Stěrba J and Dyková I (1978) Results of an artificial feeding of eggs of *Taenia saginata* Goeze, 1782 to various beetle species. *Folia Parasitologica* **25**, 257–260.
- Budke CM, Jiamin Q, Craig PS and Torgerson PR (2005) Modeling the transmission of *Echinococcus granulosus* and *Echinococcus multilocularis* in dogs for a high endemic region of the Tibetan plateau. *International Journal for Parasitology* **35**, 163–170.
- Burlet P, Deplazes P and Hegglin D (2011) Age, season and spatio-temporal factors affecting the prevalence of *Echinococcus multilocularis* and *Taenia taeniaeformis* in *Arvicola terrestris*. *Parasites & Vectors* **4**, 6.
- Carmena D and Cardona GA (2013) Canine echinococcosis: global epidemiology and genotypic diversity. *Acta Tropica* **128**, 441–460.
- Chamavit P, Sahaisook P and Niamnuay N (2011) The majority of cockroaches from the Samutprakarn province of Thailand are carriers of parasitic organisms. *EXCLI Journal* **10**, 218–222.
- Colli CW and Williams JF (1972) Influence of temperature on the infectivity of eggs of *Echinococcus granulosus* in laboratory rodents. *The Journal of Parasitology* **58**, 422–426.
- Coman BJ and Rickard MD (1977) A comparison of *in vitro* and *in vivo* estimates of the viability of *Taenia pisiformis* eggs aged under controlled conditions, and their ability to immunise against a challenge infection. *International Journal for Parasitology* **7**, 15–20.
- Conraths FJ, Probst C, Possenti A, Boufana B, Saulle R, La Torre G, Busani L and Casulli A (2017) Potential risk factors associated with human alveolar echinococcosis: systematic review and meta-analysis. *PLoS Neglected Tropical Diseases* **11**, e0005801.
- Craig PS, Macpherson CNL, Watson-Jones DL and Nelson GS (1988) Immunodetection of Echinococcus eggs from naturally infected dogs and from environmental contamination sites in settlements in Turkana, Kenya. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **82**, 268–274.
- Crewe SM (1967) Worm eggs found in gull droppings. *Annals of Tropical Medicine & Parasitology* **61**, 358–359.
- Crewe W and Crewe SM (1969) Possible transmission of bovine cysticercosis by gulls. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **63**, 17.
- Crewe W and Owen R (1978, January 1) 750,000 eggs a day – 750,000 (Pounds) a year. Available at <https://natlib.govt.nz/records/30745126> (accessed 5 March 2024).
- Da Silva AM, Bastien M, Umhang G, Boué F, Bastid V, Boucher J-M, Caillot C, De Garam CP, Renault C, Faisse M, Courquet S, Scalabrino V, Millon L, Knapp J and Poulle M-L (2021) Soil contamination by *Echinococcus multilocularis* in rural and urban vegetable gardens in relation to fox, cat and dog faecal deposits. *Parasite* **28**, 74.
- Deplazes P, Rinaldi L, Alvarez Rojas CA, Torgerson PR, Harandi MF, Romig T, Antolova D, Schurer JM, Lahmar S, Cringoli G, Magambo J, Thompson RCA and Jenkins EJ (2017) Chapter Six – global distribution of alveolar and cystic echinococcosis. In Thompson RCA, Deplazes P and Lymbery AJ (eds), *Advances in Parasitology Echinococcus and Echinococcosis part A*, Vol. **95**. Academic Press, pp. 315–493. <https://doi.org/10.1016/bs.apar.2016.11.001>
- Develeeschauer B, Bouwknegt M, Dorny P, Gabriël S, Havelaar AH, Quoilin S, Robertson LJ, Speybroeck N, Torgerson PR, van der Giessen JWB and Trevisan C (2017) Risk ranking of foodborne parasites: state of the art. *Food and Waterborne Parasitology* **8–9**, 1–13.
- Eckert J and Thompson RCA (2017) Chapter One – historical aspects of echinococcosis. In Thompson RCA, Deplazes P and Lymbery AJ (eds), *Advances in Parasitology Echinococcus and Echinococcosis Part A*, Vol. **95**. Academic Press, pp. 1–64. <https://doi.org/10.1016/bs.apar.2016.07.003>
- FAO and WHO (2014) *Multicriteria-based ranking for risk management of food-borne parasites: report of a joint FAO/WHO Expert Meeting, 3-7 September 2012, FAO Headquarters, Rome, Italy*, Boireau P (ed). Rome, Italy: Geneva, Switzerland: Food and Agriculture Organization of the United Nations; World Health Organization.
- Federer K, Armua-Fernandez MT, Hoby S, Wenker C and Deplazes P (2015) In vivo viability of *Echinococcus multilocularis* eggs in a rodent model after different thermo-treatments. *Experimental Parasitology* **154**, 14–19.
- Federer K, Armua-Fernandez MT, Gori F, Hoby S, Wenker C and Deplazes P (2016) Detection of taeniid (*Taenia* spp., *Echinococcus* spp.) eggs contaminating vegetables and fruits sold in European markets and the risk for metacestode infections in captive primates. *International Journal for Parasitology: Parasites and Wildlife* **5**, 249–253.
- Forin-Wiart M-A, Poulle M-L, Piry S, Cosson J-F, Larose C and Galan M (2018) Evaluating metabarcoding to analyse diet composition of species foraging in anthropogenic landscapes using Ion Torrent and Illumina sequencing. *Scientific Reports* **8**, 17091.
- Géba E, Rousseau A, Le Guernic A, Escotte-Binet S, Favennec L, La Carbona S, Gargala G, Dubey JP, Villena I, Betoulle S, Aubert D and Bigot-Clivot A (2021) Survival and infectivity of *Toxoplasma gondii* and *Cryptosporidium parvum* oocysts bioaccumulated by *Dreissena polymorpha*. *Journal of Applied Microbiology* **130**, 504–515.
- Gemmell MA (1976) Factors regulating tapeworm populations: estimations of the build-up and dispersion patterns of eggs after the introduction of dogs infected with *Taenia hydatigena*. *Research in Veterinary Science* **21**, 220–222. Available at <https://pubmed.ncbi.nlm.nih.gov/968189/?dopt=Abstract> (accessed 5 March 2024)
- Gemmell MA (1990) Australasian contributions to an understanding of the epidemiology and control of hydatid disease caused by *Echinococcus granulosus*—past, present and future. *International Journal for Parasitology* **20**, 431–456.
- Gemmell MA, Johnstone PD and Boswell CC (1978) Factors regulating tapeworm populations: dispersion patterns of *Taenia hydatigena* eggs on pasture. *Research in Veterinary Science* **24**, 334–338.
- Gomez-Puerta LA, Lopez-Urbina MT, Garcia HH and Gonzalez AE (2014) Longevity and viability of *Taenia solium* eggs in the digestive system of the

- beetle *Ammophorus rubripes*. *Revista Brasileira de Parasitologia Veterinaria=Brazilian Journal of Veterinary Parasitology: Orgao Oficial Do Colegio Brasileiro de Parasitologia Veterinaria* 23, 94–97.
- Guggisberg AR, Alvarez Rojas CA, Kronenberg PA, Miranda N and Deplazes P** (2020) A sensitive, one-way sequential sieving method to isolate helminths' eggs and protozoal oocysts from lettuce for genetic identification. *Pathogens (basel, Switzerland)* 9, 624.
- Hazards (BIOHAZ) EP on B, Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, Davies R, De Cesare A, Herman L, Hilbert F, Lindqvist R, Nauta M, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Cacciò S, Chalmers R, Deplazes P, Devleeschauwer B, Innes E, Romig T, van der Giessen J, Hempen M, Van der Stede Y and Robertson L** (2018) Public health risks associated with food-borne parasites. *EFSA Journal* 16, e05495.
- Heggin D and Deplazes P** (2013) Control of *Echinococcus multilocularis*: strategies, feasibility and cost–benefit analyses. *International Journal for Parasitology* 43, 327–337.
- Hemmati S, Afshar AA, Mohammadi MA, Afgar A, Nasibi S and Harandi MF** (2018) Experimental and field investigation of non-biting flies as potential mechanical vectors of *Echinococcus granulosus* eggs. *Experimental Parasitology* 189, 43–48.
- Hildreth MB, Blunt DS and Oaks JA** (2004) Lethal effects of freezing *Echinococcus multilocularis* eggs at ultralow temperatures. *Journal of Parasitology* 90, 841–844.
- Jansen F, Dorny P, Gabriël S, Dermauw V, Johansen MV and Trevisan C** (2021) The survival and dispersal of *Taenia* eggs in the environment: what are the implications for transmission? A systematic review. *Parasites & Vectors* 14, 88.
- Kapel CMO, Torgerson PR, Thompson RCA and Deplazes P** (2006) Reproductive potential of *Echinococcus multilocularis* in experimentally infected foxes, dogs, raccoon dogs and cats. *International Journal for Parasitology* 36, 79–86.
- Keilbach NM, de Aluja AS and Sarti-Gutierrez E** (1989) A programme to control taeniasis-cysticercosis (*T. solium*): experiences in a Mexican village. *Acta Leidensia* 57, 181–189.
- Kern P** (2010) Clinical features and treatment of alveolar echinococcosis. *Current Opinion in Infectious Diseases* 23, 505.
- Kubina S, Costa D, Favennec L, Gargala G, Rousseau A, Villena I, La Carbona S and Razakandrainibe R** (2021) Detection of infectious *Cryptosporidium parvum* oocysts from lamb's lettuce: CC-qPCR's intake. *Microorganisms* 9, 215.
- Lass A, Szostakowska B, Myjak P and Korzeniewski K** (2015) The first detection of *Echinococcus multilocularis* DNA in environmental fruit, vegetable, and mushroom samples using nested PCR. *Parasitology Research* 114, 4023–4029.
- Lass A, Szostakowska B, Myjak P and Korzeniewski K** (2016) Fresh fruits, vegetables and mushrooms as transmission vehicles for *Echinococcus multilocularis* in highly endemic areas of Poland: reply to concerns. *Parasitology Research* 115, 3637–3642.
- Lass A, Szostakowska B, Myjak P and Korzeniewski K** (2017) Detection of *Echinococcus multilocularis* DNA in fruit, vegetable, and mushroom samples collected in the non-endemic territory of the Pomerania province and comparison of the results with data from rural areas of the neighbouring highly endemic Warmia-Masuria Province, Poland. *Acta Parasitologica* 62, 459–465.
- Lass A, Szostakowska B, Kontogeorgos I, Korzeniewski K, Karamon J, Sulima M and Karanis P** (2019) First detection of *Echinococcus multilocularis* in environmental water sources in endemic areas using capsule filtration and molecular detection methods. *Water Research* 160, 466–474.
- Lass A, Ma L, Kontogeorgos I, Xueyong Z, Li X and Karanis P** (2020) Contamination of wastewater with *Echinococcus multilocularis* – possible implications for drinking water resources in the QTP China. *Water Research* 170, 115334.
- Laws GF** (1968) Physical factors influencing survival of taeniid eggs. *Experimental Parasitology* 22, 227–239.
- Lawson JR and Gemmell MA** (1985) The potential role of blowflies in the transmission of taeniid tapeworm eggs. *Parasitology* 91, 129–143.
- Lee PS, Dong MH, Yan XL, He TY, Yu SF, Wee SL and Wilson J** (2023) Blowfly-derived mammal DNA as mammal diversity assessment tool: determination of dispersal activity and flight range of tropical blowflies. *Biodiversity Data Journal* 11, e108438.
- Lightowlers MW, Jensen O, Fernandez E, Iriarte JA, Woollard DJ, Gauci CG, Jenkins DJ and Heath DD** (1999) Vaccination trials in Australia and Argentina confirm the effectiveness of the EG95 hydatid vaccine in sheep. *International Journal for Parasitology* 29, 531–534.
- Malkamäki S, Näreaho A, Lavikainen A, Oksanen A and Sukura A** (2019) A new SYBR green real-time PCR assay for semi-quantitative detection of *Echinococcus multilocularis* and *Echinococcus canadensis* DNA on bilberries (*Vaccinium myrtillus*). *Food and Waterborne Parasitology* 17, e00068.
- Malkamäki S, Oksanen A, Näreaho A and Sukura A** (2022) Dispersal of taeniid eggs: experimental faecal contamination of forest environment followed by DNA detection in wild berries. *Food and Waterborne Parasitology* 27, e00152.
- Martinez MJ, de Aluja AS and Gemmell M** (2000) Failure to incriminate domestic flies (Diptera: muscidae) as mechanical vectors of taenia eggs (Cyclophyllidae: Taeniidae) in rural Mexico. *Journal of Medical Entomology* 37, 489–491.
- Massolo A, Simoncini A and Romig T** (2022) The 'bridge effect' by intermediate hosts may explain differential distributions of *Echinococcus* species. *Trends in Parasitology* 38, 501–512.
- Matoff K and Kolev G** (1964) The role of the hairs, muzzle and paws of *Echinococcus* dogs in the Epidemiology of *Echinococcosis*. *Zeitschrift Fur Tropenmedizin Und Parasitologie* 15, 452–460.
- Matsuo K, Inaba T and Kamiya H** (2003) Detection of *Echinococcus multilocularis* eggs by centrifugal flotation technique: Preliminary survey of soil left in the ferryboats commuting between Hokkaido Island, where *E. multilocularis* is endemic, and mainland Japan. *Japanese Journal of Infectious Diseases* 56, 118–119.
- Mori K, Imamura A, Hirayama I and Minamoto T** (2023) Detection of *Echinococcus multilocularis* in repurposed environmental DNA samples from river water. *PeerJ* 11, e15431.
- M'rad S, Chaabane-Banaoues R, Lahmar I, Oumaima H, Mezhoud H, Babba H and Oudni-M'rad M** (2020) Parasitological contamination of vegetables sold in Tunisian retail markets with helminth eggs and protozoan cysts. *Journal of Food Protection* 83, 1104–1109.
- Nagy A, Ziadinov I, Schweiger A, Schnyder M and Deplazes P** (2011) Hair coat contamination with zoonotic helminth eggs of farm and pet dogs and foxes. *Berliner Und Munchener Tierarztliche Wochenschrift* 124, 503–511.
- Liu B, Xue C, Wang X, Wu W, Han S, Wu W, National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention (Chinese Center for Tropical Diseases Research); NHC Key Laboratory of Parasite and Vector Biology; WHO Collaborating Centre for Tropical Diseases; National Center for International Research on Tropical Diseases, Shanghai, China, and Institute for Infectious Disease and Endemic Disease Control, Beijing Center for Disease Prevention and Control, Beijing, China** (2022) The first detection of *Echinococcus Granulosus* DNA in residents' hands, dogs' hair, and soil in highly endemic region of Echinococcosis — Naqu City, Xizang Autonomous Region, China, 2020. *China CDC Weekly* 4, 982–985.
- Nosik AP** (1952) Resistance of the oncospheres of *Echinococcus granulosus* to some physical and chemical factors. *Sbornik Trudov Kharkovskogo Veterinarnogo Instituta* (21), 304–311.
- Okolo MI** (1986) Prevalence and public health implications of *Echinococcus granulosus* in rural dogs in Eastern Nigeria. *International Journal of Zoonoses* 13, 19–24.
- Oksanen A, Siles-Lucas M, Karamon J, Possenti A, Conraths FJ, Romig T, Wysocki P, Mannocci A, Mipatrini D, La Torre G, Boufana B and Casulli A** (2016) The geographical distribution and prevalence of *Echinococcus multilocularis* in animals in the European Union and adjacent countries: a systematic review and meta-analysis. *Parasites & Vectors* 9, 519.
- Ortiz Pineda C, Temesgen TT and Robertson LJ** (2020) Multiplex quantitative PCR analysis of strawberries from Bogotá, Colombia, for contamination with three parasites. *Journal of Food Protection* 83, 1679–1684.
- Otero-Abad B, Armua-Fernandez MT, Deplazes P, Torgerson PR and Hartnack S** (2017) Latent class models for *Echinococcus multilocularis* diagnosis in foxes in Switzerland in the absence of a gold standard. *Parasites & Vectors* 10, 612.
- Possenti A, Manzano-Román R, Sánchez-Ovejero C, Boufana B, La Torre G, Siles-Lucas M and Casulli A** (2016) Potential risk factors associated with human cystic echinococcosis: systematic review and meta-analysis. *PLoS Neglected Tropical Diseases* 10, e0005114.
- Raymond S and St. Clair CC** (2023) Urban magpies frequently feed on coyote scats and may spread an emerging zoonotic tapeworm. *EcoHealth* 20, 441–452.
- Robertson LJ, Troell K, Woolsey ID and Kapel CMO** (2016) Fresh fruit, vegetables, and mushrooms as transmission vehicles for *Echinococcus*

- multilocularis* in Europe: inferences and concerns from sample analysis data from Poland. *Parasitology Research* **115**, 2485–2488.
- Romig T, Deplazes P, Jenkins D, Giraudoux P, Massolo A, Craig PS, Wassermann M, Takahashi K and de la Rue M (2017) Ecology and life cycle patterns of *Echinococcus* Species. In Thompson RCA, Deplazes P and Lymbery AJ (eds), *Advances in Parasitology Echinococcus and Echinococcosis, Part A*, Vol. **95**. Academic Press, pp. 213–314. <https://doi.org/10.1016/bs.apar.2016.11.002>
- Rousseau A, Escotte-Binet S, La Carbona S, Dumètre A, Chagneau S, Favennec L, Kubina S, Dubey JP, Majou D, Bigot-Clivot A, Villena I and Aubert D (2019) *Toxoplasma gondii* oocyst infectivity assessed using a sporocyst-based cell culture assay combined with quantitative PCR for environmental applications. *Applied and Environmental Microbiology* **85**, e01189–19.
- Saelens G, Robertson L and Gabriël S (2022) Diagnostic tools for the detection of taeniid eggs in different environmental matrices: a systematic review. *Food and Waterborne Parasitology* **26**, e00145.
- Sánchez Thevenet P, Jensen O, Mellado I, Torrecillas C, Raso S, Flores ME, Minvielle MC and Basualdo JA (2003) Presence and persistence of intestinal parasites in canine fecal material collected from the environment in the Province of Chubut, Argentine Patagonia. *Veterinary Parasitology* **117**, 263–269.
- Sánchez Thevenet P, Alvarez HM, Torrecillas C, Jensen O and Basualdo JA (2020) Dispersion of *Echinococcus granulosus* eggs from infected dogs under natural conditions in Patagonia, Argentina. *Journal of Helminthology* **94**, e29.
- Schiller EL (1955) Studies on the helminth fauna of Alaska. XXVI. Some observations on the cold-resistance of eggs of *Echinococcus sibiricensis* Rausch and Schiller, 1954. *The Journal of Parasitology* **41**, 578–582.
- Serra E, Masu G, Chisu V, Cappai S, Masala G, Loi F and Piseddu T (2022) Environmental contamination by *Echinococcus* spp. Eggs as a risk for human health in educational farms of Sardinia, Italy. *Veterinary Sciences* **9**, 143.
- Shaikenov BS, Torgerson PR, Mathis A, Deplazes P, Rysmukhambetova AT and Massenov B (2004) Short report: the use of a polymerase chain reaction to detect *Echinococcus granulosus* (G1 Strain) eggs in soil samples. *The American Journal of Tropical Medicine and Hygiene* **71**, 441–443.
- Shcherbakov OV, Aghayan SA, Gevorgyan HSh, Abgaryan TA, Gevorgyan RH, Jiménez-Meléndez A and Robertson LJ (2024) Preliminary investigations of parasite contamination of water sources in Armenia. *Food and Waterborne Parasitology* **34**, e00221.
- Siebert S and Neumann S (2022) Wind-borne dispersion of *Echinococcus multilocularis* eggs – a flight model. *Journal of Helminthology* **96**, e45.
- Slepnev NK (1982) The survival of *Echinococcus granulosus* oncospheres under the influence of chemical and physical factors. Trudy Belorusskogo Nauchno Issledovatel' Skogo Instituta Eksperimental' Noi Veterinarij Veterinarnaya Nauka Proizvodstvu **19**, 96–98.
- Sweatman GK and Williams RJ (1963) Survival of *Echinococcus granulosus* and *Taenia hydatigena* eggs in two extreme climatic regions of New Zealand. *Research in Veterinary Science* **4**, 199–218.
- Szostakowska B, Lass A, Kostyra K, Pietkiewicz H and Myjak P (2014) First finding of *Echinococcus multilocularis* DNA in soil: Preliminary survey in Varmia-Masuria Province, northeast Poland. *Veterinary Parasitology* **203**, 73–79.
- Temesgen TT, Robertson LJ and Tysnes KR (2019) A novel multiplex real-time PCR for the detection of *Echinococcus multilocularis*, *Toxoplasma gondii*, and *Cyclospora cayentanensis* on berries. *Food Research International* **125**, 108636.
- Temesgen TT, Stigum VM and Robertson LJ (2022) Surveillance of berries sold on the Norwegian market for parasite contamination using molecular methods. *Food Microbiology* **104**, 103980.
- Thevenet PS, Jensen O, Drut R, Cerrone GE, Grenóvero MS, Alvarez HM, Targovnik HM and Basualdo JA (2005) Viability and infectiousness of eggs of *Echinococcus granulosus* aged under natural conditions of inferior arid climate. *Veterinary Parasitology* **133**, 71–77.
- Thomas LJ and Babero BB (1956) Observations on the infectivity of *Echinococcus* eggs obtained from foxes (*Alopex lagopus* Linn.) on St. Lawrence Island, Alaska. *The Journal of Parasitology* **42**, 659.
- Thompson RCA and Eckert J (1982) The production of eggs by *Echinococcus multilocularis* in the laboratory following in vivo and in vitro development. *Zeitschrift Für Parasitenkunde Parasitology Research* **68**, 227–234.
- Torgerson PR (2016) Fresh fruits, vegetables, and mushrooms as transmission vehicles for *Echinococcus multilocularis*. *Parasitology Research* **115**, 4447–4448.
- Torgerson PR, Pilkington J, Gulland FM and Gemmill MA (1995) Further evidence for the long distance dispersal of taeniid eggs. *International Journal for Parasitology* **25**, 265–267.
- Torgerson PR, Schweiger A, Deplazes P, Pohar M, Reichen J, Ammann RW, Tarr PE, Halkik N and Müllhaupt B (2008) Alveolar echinococcosis: from a deadly disease to a well-controlled infection. Relative survival and economic analysis in Switzerland over the last 35 years. *Journal of Hepatology* **49**, 72–77.
- Torgerson PR, Devleeschauwer B, Praet N, Speybroeck N, Willingham AL, Kasuga F, Rokni MB, Zhou X-N, Fèvre EM, Sripa B, Gargouri N, Fürst T, Budke CM, Carabin H, Kirk MD, Angulo FJ, Havelaar A and de Silva N (2015) World Health Organization estimates of the global and regional disease burden of 11 foodborne parasitic diseases, 2010: a data synthesis. *PLOS Medicine* **12**, e1001920.
- Torgerson PR, Robertson LJ, Enemark HL, Foehr J, van der Giessen JWB, Kapel CMO, Klun I and Trevisan C (2020) Source attribution of human echinococcosis: a systematic review and meta-analysis. *PLoS Neglected Tropical Diseases* **14**, e0008382.
- Trachsel D, Deplazes P and Mathis A (2007) Identification of taeniid eggs in the faeces from carnivores based on multiplex PCR using targets in mitochondrial DNA. *Parasitology* **134**, 911–920.
- Umhang G, Bastien M, Renault C, Faisse M, Caillet C, Boucher J-M, Hormaz V, Poule M-L and Boué F (2017) A flotation/sieving method to detect *Echinococcus multilocularis* and *Toxocara* spp. eggs in soil by real-time PCR. *Parasite* **24**, 28.
- Umhang G, Bastien F, Cartet A, Ahmad H, van der Ark K, Berg R, Bonelli P, Davidson RK, Deplazes P, Deksne G, João Gargate M, van der Giessen J, Jamil N, Jokelainen P, Karamon J, M'rad S, Maksimov P, Oudni-M'rad M, Muchaamba G, Oksanen A, Pepe P, Poule M-L, Rinaldi L, Samorek-Pieróg M, Santolamazza F, Santoro A, Santucci C, Saarma U, Schnyder M, Villena I, Wassermann M, Casulli A and Boué F (2024) Detection of Em, Eg sl and other Taeniidae species in lettuces and berries from Europe, Tunisia and Pakistan: two multicenter studies from the MEmE project. *bioRxiv* 2024.06.10.598207. <https://doi.org/10.1101/2024.06.10.598207>
- Vargas-Calla A, Gomez-Puerta LA, Pajuelo MJ, García HH and Gonzalez AE (2018) Molecular detection of Taeniid eggs in beetles collected in an area endemic for *Taenia solium*. *The American Journal of Tropical Medicine and Hygiene* **99**, 1198–1200.
- Veit P, Bilger B, Schad V, Schäfer J, Frank W and Lucius R (1995) Influence of environmental factors on the infectivity of *Echinococcus multilocularis* eggs. *Parasitology* **110**, 79–86.
- Wachira TM, Macpherson CN and Gathuma JM (1991) Release and survival of *Echinococcus* eggs in different environments in Turkana, and their possible impact on the incidence of hydatidosis in man and livestock. *Journal of Helminthology* **65**, 55–61.
- Wang Q, Zhong B, Yu W, Zhang G, Budke CM, Liao S, He W, Chen F, Xu K, Xie F, Danbazeli, Wang Q, Yang L, Huang Y, Li R, Yao R, Giraudoux P and Craig PS (2021) Assessment of a 10-year dog deworming programme on the transmission of *Echinococcus multilocularis* in Tibetan communities in Sichuan Province, China. *International Journal for Parasitology* **51**, 159–166.
- WHO (2015) WHO estimates of the global burden of foodborne diseases: foodborne diseases burden epidemiology reference group 2007–2015. Available at <https://www.who.int/publications-detail-redirect/9789241565165> (accessed 11 March 2024).
- Ziadinov I, Mathis A, Trachsel D, Rysmukhambetova A, Abdyjaparov TA, Kuttubaev OT, Deplazes P and Torgerson PR (2008) Canine echinococcosis in Kyrgyzstan: using prevalence data adjusted for measurement error to develop transmission dynamics models. *International Journal for Parasitology* **38**, 1179–1190.