

On the von Neumann entropy of certain quantum walks subject to decoherence[†]

CHAOBIN LIU and NELSON PETULANTE

*Department of Mathematics, Bowie State University, 14000 Jericho Park Road,
Bowie, Maryland 20715, U.S.A.*

Email: {cliu;npetulante}@bowiestate.edu

Received 29 March 2010; revised 15 July 2010

In this paper, we consider a discrete-time quantum walk on the N -cycle governed by the condition that at every time step of the walk, the option persists, with probability p , of exercising a projective measurement on the coin degree of freedom. For a bipartite quantum system of this kind, we prove that the von Neumann entropy of the total density operator converges to its maximum value. Thus, when influenced by decoherence, the mutual information between the two subsystems corresponding to the space of the coin and the space of the walker must eventually diminish to zero. Put plainly, any level of decoherence greater than zero forces the system to become completely ‘disentangled’ eventually.

1. Introduction

A quantum walk (QW) is a reversible process that is commonly described as the quantum-mechanical analogue of a classical random walk. In recent times, quantum walks have attracted extensive attention, mainly for their value as potential sources of new algorithms (Kempe 2003; Ambainis 2003; Kendon 2006; Venegas-Andraca 2008; Konno 2008).

Like classical random walks, quantum walks are classified into two main types: discrete-time quantum walks (Nayak and Vishwanath 2000; Ambainis *et al.* 2001; Aharonov *et al.* 2001) and continuous-time quantum walks (Fahri and Gutmann 1998; Childs *et al.* 2002). Both types exhibit similar dynamical properties, and the continuous type can be obtained from the discrete type through a suitable limiting process (Strauch 2006; Childs 2010).

Whereas continuous-time quantum walks can be modelled in terms of a single position Hilbert space \mathcal{H}_p , discrete-time quantum walks cannot be modelled except in terms of a bipartite system involving the tensor product of the position Hilbert space \mathcal{H}_p and an auxiliary coin Hilbert space \mathcal{H}_c . The conditional shift operator, which governs the itinerary of the walker, induces entanglement between the degree of freedom of the coin and the spatial degree of freedom of the walker.

To capture the full computational power inherent in any quantum-mechanical process, it is essential to be able to model the phenomenon of quantum entanglement. Recently,

[†] The authors acknowledge, with due gratitude, the support provided to us by the NSF-funded HBCU-UP/BETTER Project at Bowie State University.

several important studies of quantum entanglement in discrete-time quantum walks have been completed, from both a numerical perspective (Carneiro *et al.* 2005; Maloyer and Kendon 2007; Venegas-Andraca and Bose 2009) and an analytical perspective (Abal *et al.* 2006; Annabestani *et al.* 2010a).

In recent years, several schemes have been proposed for the implementation of quantum walks in realistic media (Travaglione and Milburn 2002; Dur *et al.* 2002; Sanders *et al.* 2003; Du *et al.* 2003; Ryan *et al.* 2005; Eckert *et al.* 2005; Zou *et al.* 2006; van Hoogdalem and Blaauboer 2009). However, any attempt to implement a quantum system in a physical channel must take into consideration the critical issue of ‘decoherence’, whereby the idiosyncratic features of a quantum system succumb to macroscopic (classical) manifestations. Various mathematical models of decoherence in discrete-time QWs have been investigated both numerically and analytically (Kendon and Tregenna 2002; Brun *et al.* 2003a; Brun *et al.* 2003b; Kendon and Tregenna 2003a; Kendon and Tregenna 2003b; Romanelli 2005; Kosik *et al.* 2006; Richter 2007; Zhang 2008; Banerjee *et al.* 2008; Shikano *et al.* 2010; Liu and Petulante 2010; Annabestani *et al.* 2010b).

In this paper, as in Brun *et al.* (2003a; 2003b), we adopt a specific model of decoherence, which might best be described as follows. At each time step of the quantum walk, an observer stands ready to perform a projective measurement on the coin degree of freedom. The probability of performing a measurement is given by a fixed parameter p , called the ‘decoherence rate’.

There is not much evidence in the current literature of any attempts to provide a precise formulation of the relationship between entanglement and decoherence. A notable exception is the numerical study given in Maloyer and Kendon (2007). However, the measure of entanglement employed in Maloyer and Kendon (2007), which they called the *negativity* measure, differs considerably from the measure of entanglement adopted in this paper.

Here, we utilise the concept of von Neumann entropy to quantify the information content of the various components of the quantum walk system, including the mutual information shared between its subsystems (coin and position). This approach enables us to provide a precise formulation of the measure of entanglement between the subsystems. In the presence of any non-zero level of decoherence, the von Neumann entropy associated with the total probability density of the system tends to its maximum value, implying the total collapse of entanglement between subsystems.

2. Quantum walks on the N -cycle subject to decoherence

We begin by defining the elements, as formulated in Liu and Petulante (2010), of a quantum random walk on the N -cycle – the definitions and corresponding notation are analogous to those outlined in Liu and Petulante (2010). Let t denote the number of time steps from the moment the discrete quantum walk is launched on the N -cycle. The temporal evolution of the system is modelled by $\psi_t = U^t \psi_0$, where ψ_0 is the initial state and $U = S(\mathbb{I} \otimes U_c)$ is the evolution operator on the Hilbert space $\mathcal{H} = \mathcal{H}_N \otimes \mathcal{H}_2$. Here U_c denotes the coin operator on the coin subspace \mathcal{H}_2 spanned by an orthonormal basis $\{|j\rangle, j = 1, 2, \dots\}$, and S denotes the conditional shift operator on the position subspace \mathcal{H}_N

spanned by an orthonormal basis $\{|x\rangle, x \in \mathbb{Z}_N\}$. Thus, a typical state ψ in \mathcal{H} may be expressed as $\psi = \sum_x \sum_j \psi(x, j)|x\rangle \otimes |j\rangle$.

For a QW launched with initial state ψ_0 , the probability $P(x, t)$ of finding the walker at the position $x \in \mathbb{Z}_N$ at time t is given by the standard formula $P(x, t) = \text{Tr}(|x\rangle\langle x|\rho(t))$, where the time-dependent density operator $\rho(t)$ is defined by

$$\rho(t) = \psi_t \psi_t^\dagger = U^t |\psi_0\rangle\langle\psi_0| U^{\dagger t}. \tag{1}$$

During the evolution of the quantum walk, the history of decoherence-inducing events, if any, including any acts of measurement, may be modelled by the probabilistic option of applying to the coin degree of freedom, at each time step of the walk, a *unital* family of operators $\{\hat{A}_n\}_{0 \leq n \leq v}$, jointly satisfying the condition

$$\sum_{0 \leq n \leq v} \hat{A}_n^\dagger \hat{A}_n = I. \tag{2}$$

Accordingly, when adjusted for decoherence, the density operator of the system acts on the probability density function ρ via the formula

$$\rho(t + 1) = \sum_{0 \leq n \leq v} U \hat{A}_n \rho(t) \hat{A}_n^\dagger U^\dagger. \tag{3}$$

For simplicity, and without loss of generality, we will assume from this point onward that every quantum walk under consideration is launched from position $|0\rangle$ in coin state ψ_0 . Upon applying Fourier transformations to all elements of the QW system, the formulation of the density operator assumes the following form:

$$\rho(t) = \frac{1}{N} \sum_k \sum_{k'} |k\rangle\langle k'| \otimes \mathcal{L}_{kk'}^t |\psi_0\rangle\langle\psi_0|, \tag{4}$$

where

$$|k\rangle = \frac{1}{\sqrt{N}} \sum_x \exp(i2\pi xk/N) |x\rangle$$

and where the so-called ‘decoherence super-operator’ $\mathcal{L}_{kk'}$ is defined by the formula:

$$\begin{aligned} \mathcal{L}_{kk'} |\psi_0\rangle\langle\psi_0| &= \sum_n U_c(k) \hat{A}_n |\psi_0\rangle\langle\psi_0| \hat{A}_n^\dagger U_c^\dagger(k') \\ &= \begin{bmatrix} A_{11}(1, k, k') & A_{12}(1, k, k') \\ A_{21}(1, k, k') & A_{22}(1, k, k') \end{bmatrix}, \end{aligned} \tag{5}$$

in which

$$U_c(k) = \begin{bmatrix} e^{-\frac{2\pi ik}{N}} & 0 \\ 0 & e^{\frac{2\pi ik}{N}} \end{bmatrix} U_c. \tag{6}$$

Note that all essential features of the quantum walk, including all peculiarities of quantum behaviour connoted by the words ‘decoherence’ and ‘entanglement’, are fully encoded in the anatomy of the super-operator $\mathcal{L}_{kk'}$. This is what enables us to study the level of quantum entanglement in quantum walks subject to decoherence.

After t iterations, let the elements of the 2×2 matrix $\mathcal{L}_{kk'}^t|\psi_0\rangle\langle\psi_0|$ be denoted by

$$\mathcal{L}_{kk'}^t|\psi_0\rangle\langle\psi_0| = \begin{bmatrix} A_{11}(t, k, k') & A_{12}(t, k, k') \\ A_{21}(t, k, k') & A_{22}(t, k, k') \end{bmatrix}. \tag{7}$$

In terms of the standard basis for the $2N$ -dimensional Hilbert space $\mathcal{H} = \mathcal{H}_N \otimes \mathcal{H}_2$, the density operator $\rho(t)$ is given by

$$\rho(t) = \sum_{1 \leq x, y \leq N} \sum_{1 \leq j, l \leq 2} P_{xyjl}(t) |x\rangle\langle y| \otimes |j\rangle\langle l|, \tag{8}$$

where

$$P_{xyjl}(t) = \frac{1}{N} \sum_k \sum_{k'} \langle x|k\rangle\langle k'|y\rangle A_{jl}(t, k, k'). \tag{9}$$

In terms of the above definitions, the probability $P(x, t)$ of finding the walker at position x at time t is given by

$$\begin{aligned} P(x, t) &= \text{Tr}(|x\rangle\langle x|\rho(t)) \\ &= \frac{1}{N} \sum_k \sum_{k'} \langle x|k\rangle\langle k'|x\rangle \text{Tr}(\mathcal{L}_{kk'}^t|\psi_0\rangle\langle\psi_0|) \\ &= \frac{1}{N} \sum_k \sum_{k'} \langle x|k\rangle\langle k'|x\rangle (A_{11}(t, k, k') + A_{22}(t, k, k')) \\ &= P_{xx11}(t) + P_{xx22}(t). \end{aligned} \tag{10}$$

To avoid unpleasant complications and to permit us more easily to illustrate our approach to the analysis of a QW on the N -cycle subject to decoherence-inducing influences, we will confine our attention in this paper to a specific model. Let $\beta \in (0, \frac{\pi}{2})$ and $k \in \{0, 1, \dots, N - 1\}$. To serve as the coin operator of the system, we choose

$$U_c(\beta) = \begin{bmatrix} \cos \beta & \sin \beta \\ \sin \beta & -\cos \beta \end{bmatrix}, \tag{11}$$

whose Fourier dual is given by

$$U_c(\beta, k) = \begin{bmatrix} e^{-\frac{i2\pi k}{N}} \cos \beta & e^{-\frac{i2\pi k}{N}} \sin \beta \\ e^{\frac{i2\pi k}{N}} \sin \beta & -e^{\frac{i2\pi k}{N}} \cos \beta \end{bmatrix}. \tag{12}$$

Note that when $\beta = \frac{\pi}{4}$, the operator given by Equation 12 is none other than the Hadamard coin operator.

By the same token, we specialise to the following choice of three ($v = 2$) operators to serve as the unital family $\{\hat{A}_n\}_{0 \leq n \leq v}$ of decoherence-inducing operators on the coin degree of freedom, as in Equation 2:

$$\begin{aligned} \hat{A}_0 &= \sqrt{1 - p} \sigma_0 \\ \hat{A}_1 &= \frac{\sqrt{p}}{2} (\sigma_0 + \sigma_z) \\ \hat{A}_2 &= \frac{\sqrt{p}}{2} (\sigma_0 - \sigma_z) \end{aligned}$$

where $0 \leq p \leq 1$ and σ_0 and σ_z are the Pauli matrices. The level of decoherence induced by these operators is reflected by the value of p , which is called the *decoherence rate*. Specifically, the QW evolves as if the state of the coin is measured at each time step with probability p . Thus, when $p = 0$, the QW evolves as a purely coherent quantum process. At the other extreme, when $p = 1$, the QW behaves exactly like a classical random walk.

Now let $\mathbf{L}(\mathbb{C}^2)$ denote the Hilbert space of all 2×2 complex matrices with inner product given by

$$\langle M_1, M_2 \rangle \equiv \text{tr}(M_1^\dagger M_2). \tag{13}$$

Lemma 1. Let \mathcal{S} be a superoperator on the Hilbert space $\mathbf{L}(\mathbb{C}^2)$ defined by

$$\mathcal{S} = \sum_{n=0}^2 U_1 \hat{A}_n \cdot \hat{A}_n^\dagger U_2 : B \mapsto \sum_{n=0}^2 U_1 \hat{A}_n B \hat{A}_n^\dagger U_2,$$

where U_1, U_2 are 2×2 unitary matrices and $B \in \mathbf{L}(\mathbb{C}^2)$. Then $\langle \mathcal{S}B, \mathcal{S}B \rangle \leq \langle B, B \rangle$. In particular, $\langle \mathcal{S}B, \mathcal{S}B \rangle = \langle B, B \rangle$ for all $B \in \mathbf{L}(\mathbb{C}^2)$ if and only if the decoherence rate $p = 0$.

Proof. See Liu and Petulante (2010). □

An immediate corollary of this lemma, and one that is essential to our analysis, is the fact that $|\lambda| \leq 1$ for every eigenvalue λ of \mathcal{S} . To justify this assertion, suppose B_λ is an eigenvector of \mathcal{S} belonging to λ . Then $\langle \mathcal{S}B_\lambda, \mathcal{S}B_\lambda \rangle = \langle \lambda B_\lambda, \lambda B_\lambda \rangle = |\lambda|^2 \langle B_\lambda, B_\lambda \rangle$. But since, according to the lemma, $\langle \mathcal{S}B, \mathcal{S}B \rangle \leq \langle B, B \rangle$, we see that $|\lambda| \leq 1$.

We will now cast our reasoning in terms of the super-operator $\mathcal{L}_{k,k'}$, which maps $\mathbf{L}(\mathbb{C}^2)$ to $\mathbf{L}(\mathbb{C}^2)$. If we choose the Pauli matrices $\sigma_0, \sigma_x, \sigma_y$ and σ_z as a basis for $\mathbf{L}(\mathbb{C}^2)$, then, in terms of this basis, the 4×4 matrix representation of $\mathcal{L}_{k,k'}$ is given by

$$\mathcal{L}_{k,k'} = \begin{bmatrix} c^- & iqs^- \sin 2\beta & 0 & is^- \cos 2\beta \\ 0 & -qc^+ \cos 2\beta & qs^+ & c^+ \sin 2\beta \\ 0 & -qs^+ \cos 2\beta & -qc^+ & s^+ \sin 2\beta \\ is^- & qc^- \sin 2\beta & 0 & c^- \cos 2\beta \end{bmatrix} \tag{14}$$

where, for brevity, we have set $q = 1 - p$ and

$$\begin{aligned} c^+ &= \cos \frac{2\pi(k' + k)}{N} \\ s^+ &= \sin \frac{2\pi(k' + k)}{N} \\ c^- &= \cos \frac{2\pi(k' - k)}{N} \\ s^- &= \sin \frac{2\pi(k' - k)}{N}. \end{aligned}$$

After a somewhat tedious, but not very difficult calculation, we arrive at the following explicit formula for the characteristic polynomial $f(\lambda)$ of $\mathcal{L}_{k,k'}$:

$$\begin{aligned}
 f(\lambda) &= \det(\lambda \mathbb{I}_4 - \mathcal{L}_{k,k'}) \\
 &= \lambda^4 \\
 &\quad + (1 + \cos 2\beta)(qc^+ - c^-)\lambda^3 \\
 &\quad + ((1 + q^2)\cos 2\beta - 2qc^+c^-(1 + \cos 2\beta))\lambda^2 \\
 &\quad + q(1 + \cos 2\beta)(c^+ - qc^-)\lambda \\
 &\quad + q^2.
 \end{aligned}
 \tag{15}$$

The following proposition summarises some basic attributes of the eigenvalues of $\mathcal{L}_{k,k'}$.

Proposition 1. Suppose $0 < p < 1$. Let a typical eigenvalue of $\mathcal{L}_{k,k'}$ be denoted by λ . Then:

- (i) $\|\lambda\| \leq 1$.
- (ii) If $\|\lambda\| = 1$, then $\lambda = \pm 1$.
- (iii) $\lambda = 1$ when and only when $k = k'$, in which case the algebraic multiplicity of $\lambda = 1$ is 1.
- (iv) $\lambda = -1$ when and only when $|k' - k| = \frac{N}{2}$, in which case the algebraic multiplicity of $\lambda = -1$ is 1.

Proof. See Appendix A. □

Proposition 1 enables us to specify the long-term behaviour of the matrix components of the total density operator given by Equation 8.

Proposition 2. For the matrix defined in Equation 7, the following assertions hold:

- (i) Suppose $k = k'$:

If $j = l$,

$$\lim_{t \rightarrow \infty} A_{jl}(t, k, k') = \frac{1}{2}.$$

If $j \neq l$,

$$\lim_{t \rightarrow \infty} A_{jl}(t, k, k') = 0.$$

- (ii) Suppose $|k - k'| = N/2$:

If $j = l$,

$$\lim_{t \rightarrow \infty} (-1)^t A_{jl}(t, k, k') = \frac{1}{2}.$$

If $j \neq l$,

$$\lim_{t \rightarrow \infty} A_{jl}(t, k, k') = 0.$$

(iii) Suppose $|k - k'| \neq N/2$ and $\neq 0$:

Then, for all combinations of j, l , we have

$$\lim_{t \rightarrow \infty} A_{jl}(t, k, k') = 0.$$

Proof. See Appendix B. □

In view of Proposition 2, the behaviour as $t \rightarrow \infty$ of the operator in Equation 8 can be specified as follows.

Theorem 1. For a quantum walk on the N -cycle, let the total density operator $\rho(t)$ be defined as in Equation 8. Then:

(i) Suppose N is odd:

If $x = y$ and $j = l$,

$$\lim_{t \rightarrow \infty} P_{xyjl}(t) = \frac{1}{2N}.$$

If $x \neq y$ or $j \neq l$,

$$\lim_{t \rightarrow \infty} P_{xyjl}(t) = 0.$$

(ii) Suppose N is even:

If $x = y$ and $j = l$ and if $t - x$ is even,

$$\lim_{t \rightarrow \infty} P_{xyjl}(t) = \frac{1}{N}.$$

Otherwise,

$$\lim_{t \rightarrow \infty} P_{xyjl}(t) = 0.$$

Thus, in every case, as $t \rightarrow \infty$, the operator $\rho(t)$, viewed as a $2N \times 2N$ matrix, converges to a diagonal matrix. If N is odd, the diagonal elements all converge to $1/2N$. If N is even, the diagonal elements converge in an alternating pattern to $1/N$ or 0 .

Proof. See Appendix C. □

As noted in Zurek (2003), decoherence on a quantum system is manifested through its density matrix by the vanishing of the off-diagonal elements. In the context of a coin-driven quantum walk on the N -cycle, this is precisely what Theorem 1 asserts. Note that the off-diagonal elements of the density matrix are precisely the elements that represent the quantum correlations (also known as entanglement) between the coin subsystem and the position subsystem. Unsurprisingly, at least for quantum walks, decoherence turns out to be practically synonymous with ‘disentanglement’. A more precise elaboration of the relationship between decoherence and entanglement is deferred to the next section.

For the position distribution $P(x, t)$ (see Equation 10), Theorem 1 has an immediate corollary, which echoes a similar result given in Liu and Petulante (2010).

Corollary 1. Suppose a quantum walk on the N -cycle is launched from the origin with initial coin state $|\psi_0\rangle$ and with decoherence rate $p > 0$. If N is odd, $P(x, t)$ converges to $1/N$ on all nodes of the cycle. If N is even, $P(x, t)$ converges to $2/N$ on the supporting nodes of the cycle and to 0 on the non-supporting nodes of the cycle.

3. Entanglement versus decoherence

Since a purely coherent QW is a reversible process, the von Neumann entropy of the total density operator is invariant with respect to time. If the QW is launched in a pure state, then it will continue to evolve in a pure state and the entropy of the reduced density operator on the coin subsystem can serve as a measure of its degree of entanglement relative to the subsystem of the walker (Carneiro *et al.* 2005; Abal *et al.* 2006; Venegas-Andraca and Bose 2009; Annabestani *et al.* 2010a).

Unlike the purely coherent case, a quantum walk, when subject to decoherence, evolves in a mixture of states. If the ‘decoherence rate’ is non-zero, the von Neumann entropy of the total density operator is no longer invariant. Using a very simple argument, it can be shown that the entropy of the total density operator is a strictly increasing function of time. The following two facts constitute a basis for the argument:

- (a) projective measurements increase entropy; and
- (b) the entropy is a concave function of its inputs.

Thus, to measure the level of quantum entanglement in a QW subject to non-zero decoherence, the von Neumann entropy must be considered separately for each of the subsystems as well as for the total system.

The von Neumann entropy of a quantum system A , denoted $S(A)$, is a measure of the uncertainty implied by the multitude of potential outcomes, as reflected by its density matrix $\rho(A)$. By definition, $S(A) = S(\rho(A)) = -\text{Tr}(\rho \ln \rho)$.

For a composite system with two components A and B , the joint entropy of their conjunction, denoted by $S(A, B)$, is defined by the formula $S(A, B) = -\text{Tr}(\rho^{AB} \ln \rho^{AB})$, where ρ^{AB} is the density matrix of the composite quantum system AB .

A good measure of the level of quantum entanglement between the two components A and B is the so-called mutual information $S(A : B)$ defined by the formula $S(A : B) = S(A) + S(B) - S(A, B)$.

We continue to confine our attention to the case of a quantum walk on the N -cycle subject to decoherence on the coin degree of freedom. Our main objective in the rest of this paper is to show that whenever the decoherence rate $p > 0$, the mutual information between the subsystem of the coin and the subsystem of the walker must eventually diminish to 0.

The following Lemma, which is due to Watrous, is essential to our reasoning.

Lemma 2 (Watrous 2008). Let \mathcal{X} denote a complex Euclidean space and $\text{Pos}(\mathcal{X})$ denote the set of positive semidefinite operators defined on \mathcal{X} with norm

$$\|\rho\|_{\text{tr}} = \text{Tr} \left(\sqrt{\rho^\dagger \rho} \right).$$

Then, with respect to this norm, the von Neumann entropy $S(\rho)$ is continuous at every point $\rho \in \text{Pos}(\mathcal{X})$.

Proof. See Watrous (2008). □

The operator norm $\|\cdot\|_{\text{tr}}$, which appears in the statement of Lemma 2, is known as the Schatten 1-norm or sometimes simply as the ‘trace norm’.

Theorem 2. Suppose the QW is launched on the N -cycle with initial coin state $|\psi_0\rangle$ and decoherence rate $p > 0$. Let $\rho(t)$ denote the time-dependent density operator of the overall system. If N is odd, then $\lim_{t \rightarrow \infty} S(\rho(t)) = 1 + \log_2 N$. If N is even, then $\lim_{t \rightarrow \infty} S(\rho(t)) = \log_2 N$.

Proof. See Appendix D. □

A basic fact from information theory is that for any operator ρ defined on a Hilbert space of dimension d , the maximum value of the entropy $S(\rho)$ is $\log_2 d$. This implies that the limiting entropy values given by Theorem 2 are actually maximal, for both even and odd values of the cycle length N . To see this, note that the overall Hilbert space $\mathcal{H}_2 \times \mathcal{H}_N$, over which the quantum walk evolves, is of dimension $2N$. This easily explains why, when N is odd, the entropy is maximal. However, when N is even, exactly half of the nodes of the cycle (every second one) are necessary and sufficient for completely determining the position distribution of the walker’s itinerary. So, when N is even, the quantum walk evolves over a Hilbert space that is effectively of dimension N .

Finally, we consider separately the long-term trend of the entropies associated with the reduced density operators of the two constituent subsystems (coin and position) and the mutual information between them.

For the subsystem associated with the coin, the time-dependent *reduced* density operator $\rho_c(t)$ is given by $\rho_c(t) = \text{trace}_w(\rho(t))$, where the subscript w signifies exclusion or ‘tracing out’, relative to the overall system density operator $\rho(t)$, of the walker’s degrees of freedom. Similarly, for the subsystem associated with the walker, the time-dependent *reduced* density operator $\rho_w(t)$ is given by $\rho_w(t) = \text{trace}_c(\rho(t))$, where the subscript c signifies exclusion or ‘tracing out’, relative to the overall system density operator $\rho(t)$, of the coin’s degrees of freedom.

The following theorem summarises our main finding.

Theorem 3. Suppose a quantum walk is launched on the N -cycle with initial coin state $|\psi_0\rangle$ and decoherence rate $p > 0$. Let $\rho_c(t)$ and $\rho_w(t)$ denote the time-dependent reduced density operators associated with the subsystems of the coin and the walker, respectively. Then the long-term trend of the mutual information between the coin subsystem and the walker subsystem is given by $\lim_{t \rightarrow \infty} S(\rho_c(t) : \rho_w(t)) = 0$.

Proof. See Appendix E. □

Finally, it can be shown, without much difficulty, that as $t \rightarrow \infty$, the entropy values of the reduced density operators $\rho_c(t)$ and $\rho_w(t)$ each converges to its maximum value. To do this, it is enough to employ some elementary calculus based on Theorem 2, together with the following basic facts:

- (1) $S(\rho_c(t)) \leq 1$;
- (2) $S(\rho_w(t)) \leq \ln N$ when N is odd; and
- (3) $S(\rho_w(t)) \leq \ln(N/2)$ when N is even.

Corollary 2. Suppose a quantum walk is launched on the N -cycle with initial coin state $|\psi_0\rangle$ and decoherence rate $p > 0$. Then:

- (i) $\lim_{t \rightarrow \infty} S(\rho_c(t)) = 1$.
- (ii) If N is odd, then $\lim_{t \rightarrow \infty} S(\rho_w(t)) = \ln N$.
- (iii) If N is even, then $\lim_{t \rightarrow \infty} S(\rho_w(t)) = \ln(N/2)$.

4. Conclusions and further questions

The model of decoherence used in this article is only one of several prevalent in the current literature. It would be interesting to investigate how quantum entanglement responds to other models of decoherence, and not just for quantum walks on the N -cycle, but for quantum walks over other kinds of topological networks as well. We speculate that, in every case, decoherence serves to erase quantum entanglement between the subsystems of a given quantum system. This implies that, as noted in Zurek (2003), the progressive disappearance of entanglement should be accompanied by the corresponding disappearance of the off-diagonal elements of the density operators. Ultimately, the density operators should become indistinguishable from diagonal matrices.

The natural question arising in this context is how long does it take for a quantum system subject to decoherence to reach its stationary state devoid of any entanglement? To address this question, we propose a definition of *decoherence time* analogous to the measure of *mixing time* used in the literature. Let ρ_∞ denote the limiting (stationary) density operator of the quantum system and $\|\cdot\|_{\text{tr}}$ denote the trace norm as defined above. Then, for every $\epsilon > 0$, we define

$$D(\epsilon) = \min \{ \tau \mid \forall t > \tau : \|\rho(t) - \rho_\infty\|_{\text{tr}} < \epsilon \}. \tag{16}$$

To estimate $D(\epsilon)$, it would suffice to have good control over the eigenvalues and eigenvectors of the superoperator $\mathcal{L}_{kk'}^t$. Unfortunately, even for very simple systems, the task of specifying the eigenvalues and eigenvectors of the superoperator $\mathcal{L}_{kk'}^t$ can be a challenge. But the rewards of success would be worth the effort. Among other things, it would permit us to compare mixing time and decoherence time, since the relationship could prove quite interesting.

Appendix A. Proof of Proposition 1

Proof of (i). $\mathcal{L}_{k,k'}$ is a special case of the superoperator \mathcal{S} in Lemma 1, according to which, the moduli of all eigenvalues of $\mathcal{L}_{k,k'}$ are less than or equal to unity. □

Proof of (ii). Suppose $e^{i\theta}$ is a non-real eigenvalue of $\mathcal{L}_{k,k'}$, where θ is a real number. Then the conjugate $e^{-i\theta}$ also must be an eigenvalue and $e^{-i\theta} \neq e^{i\theta}$. Hence

$$f(\lambda) = (\lambda - e^{i\theta})(\lambda - e^{-i\theta})[\lambda^2 + a\lambda + (1 - p)^2]$$

for some $a \in \mathbb{C}$. For brevity, let $q = 1 - p$. Comparing corresponding coefficients of both sides of Equation 15, we obtain the following system of equations:

$$\begin{aligned} a - 2 \cos \theta &= (1 + \cos 2\beta) (qc^+ - c^-) \\ 1 + q^2 - 2a \cos \theta &= -2qc^+c^- + (1 + q^2 - 2qc^+c^-) \cos 2\beta \\ a - 2q^2 \cos \theta &= (1 + \cos 2\beta) (qc^+ - q^2c^-). \end{aligned} \tag{17}$$

After some elementary algebraic manipulations, we infer that

$$1 + q^2 = -q(1 + \cos 2\beta) \cos \frac{2\pi(k' + k)}{N} \cos \frac{2\pi(k' - k)}{N},$$

which is impossible since the modulus of the left-hand side is strictly greater than the modulus of the right-hand side. More precisely, note that the modulus of the right-hand side is strictly less than $2q$ which, in turn is less than $1 + q^2$. This contradiction implies that any unit eigenvalue of $\mathcal{L}_{k,k'}$ must be real. \square

Proof of (iii). $\lambda = 1$ is an eigenvalue of $\mathcal{L}_{k,k'}$ if and only if

$$f(1) = (1 + \cos 2\beta) \left(1 - \cos \frac{2\pi(k' - k)}{N} \right) \left[1 + 2(1 - p) \cos \frac{2\pi(k' + k)}{N} + (1 - p)^2 \right] = 0,$$

if and only if

$$1 - \cos \frac{2\pi(k' - k)}{N} = 0,$$

which implies $k' = k$. Moreover, since

$$f'(1) = (1 - \cos 2\beta)[1 - (1 - p)^2] \neq 0,$$

the algebraic multiplicity of $\lambda = 1$ is 1. \square

Proof of (iv). $\lambda = -1$ is an eigenvalue of $\mathcal{L}_{k,k'}$ if and only if

$$f(-1) = (1 + \cos 2\beta) \left(1 + \cos \frac{2\pi(k' - k)}{N} \right) \left[1 - 2(1 - p) \cos \frac{2\pi(k' + k)}{N} + (1 - p)^2 \right] = 0$$

if and only if

$$1 + \cos \frac{2\pi(k' - k)}{N} = 0,$$

which implies

$$|k' - k| = \frac{N}{2}.$$

In this case, since

$$f'(-1) = (1 - \cos 2\beta)[(1 - p)^2 - 1] \neq 0,$$

the algebraic multiplicity of $\lambda = -1$ is 1. \square

Appendix B. Proof of Proposition 2

Proof. Viewed as a 2×2 matrix, $\mathcal{L}_{k,k'}^t |\psi_0\rangle\langle\psi_0|$ is a linear combination of the Pauli matrices $\sigma_0, \sigma_x, \sigma_y$ and σ_z with corresponding weights given by

$$W(t, k, k') = \mathcal{L}_{k,k'}^t \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \end{bmatrix},$$

where the column vector $[\alpha_1, \alpha_2, \alpha_3, \alpha_4]^T$ represents $|\psi_0\rangle\langle\psi_0|$ with $\alpha_1 = \frac{1}{2}$ for all choices of $|\psi_0\rangle$.

Proof of (i). For brevity, let

$$\begin{aligned} q &= 1 - p \\ \tilde{c} &= \cos \frac{4\pi k}{N} \\ \tilde{s} &= \sin \frac{4\pi k}{N}. \end{aligned}$$

When $k = k'$, we have

$$\mathcal{L}_{k,k} = \begin{bmatrix} 1 & 0 \\ 0 & Q_0 \end{bmatrix}, \tag{18}$$

where

$$Q_0 = \begin{bmatrix} q\tilde{c} \cos 2\beta & q\tilde{s} & \tilde{c} \sin 2\beta \\ -q\tilde{s} \cos 2\beta & -q\tilde{c} & \tilde{s} \sin 2\beta \\ q \sin 2\beta & 0 & \cos 2\beta \end{bmatrix}. \tag{19}$$

Therefore

$$\mathcal{L}_{k,k}^t = \begin{bmatrix} 1 & 0 \\ 0 & Q_0^t \end{bmatrix}, \tag{20}$$

in terms of which, the four components of $\mathcal{L}_{k,k}^t |\psi_0\rangle \langle \psi_0|$ can be expressed as follows:

$$\begin{aligned} A_{11} &= \frac{1}{2} + \epsilon_1 \\ A_{22} &= \frac{1}{2} - \epsilon_1 \\ A_{12} &= \overline{A_{21}} = \epsilon_2 \end{aligned}$$

where ϵ_1 and ϵ_2 are linear combinations of the elements of the matrix Q_0^t . By Proposition 2, the eigenvalues of Q_0 all possess moduli strictly less than 1. Therefore $Q_0^t \rightarrow 0$ as $t \rightarrow \infty$, from which we have assertion (i) of the proposition. □

Proof of (ii). Using similar reasoning to the above, when $|k - k'| = N/2$, we have

$$\mathcal{L}_{k,k'}^t = \begin{bmatrix} (-1)^t & 0 \\ 0 & Q_1^t \end{bmatrix}, \tag{21}$$

where Q_1 is the 3×3 matrix adjoint to the leading entry c^- of the 4×4 matrix $\mathcal{L}_{k,k'}$ in Equation 14. The conclusion follows by a line of reasoning analogous to that used in the proof of part (i). □

Proof of (iii). If $|k - k'| \neq N/2$ and $\neq 0$, then, by Proposition 1, the modulus of every eigenvalue of $\mathcal{L}_{k,k'}$ is strictly less than 1. Thus $\mathcal{L}_{k,k'}^t \rightarrow 0$ as $t \rightarrow \infty$, which proves assertion (iii). □

This completes the proof of Proposition 2 □

Appendix C. Proof of Theorem 1

Proof of (i). For this part we assume N is odd. We consider three sub-cases:

(a) $x = y$ and $j = l$:

By Equation 9, we have

$$\begin{aligned}
 N^2 P_{xxjj}(t) &= \sum_{k=k'} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k') + \sum_{k \neq k'} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k') \\
 &= \sum_{k=0}^{N-1} A_{jj}(t, k, k) + \sum_{k \neq k'} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k').
 \end{aligned}
 \tag{22}$$

By Proposition 2, If $k = k'$, the term $A_{jj}(t, k, k)$ in (22) converges to $\frac{1}{2}$. If $k \neq k'$, the term $A_{jj}(t, k, k')$ in (22) converges to zero. So

$$\lim_{t \rightarrow \infty} P_{xyjl}(t) = \frac{1}{2N}.$$

(b) $x \neq y$ and $j = l$:

By Equation 9,

$$N^2 P_{xyjj}(t) = \sum_{k=0}^{N-1} e^{\frac{2\pi i k(x-y)}{N}} A_{jj}(t, k, k) + \sum_{k \neq k'} e^{\frac{2\pi i k(x-y)}{N}} A_{jj}(t, k, k').
 \tag{23}$$

As in the previous case, Proposition 2 enables us conclude that

$$\lim_{t \rightarrow \infty} P_{xyjl}(t) = 0.$$

(c) $j \neq l$:

In this case, by Proposition 2, $A_{jl}(t, k, k') \rightarrow 0$ as $t \rightarrow \infty$. Since $P_{xyjl}(t)$ is a linear combination of $A_{jl}(t, k, k')$ for $k, k' = 0, 1, 2, \dots, N - 1$ and $j, l = 1, 2$, it follows that

$$P_{xyjl}(t) = \frac{1}{N} \sum_k \sum_{k'} \langle x|k \rangle \langle k'|y \rangle A_{jl}(t, k, k') \rightarrow 0$$

as $t \rightarrow \infty$. □

Proof of (ii). In this part N is even. We consider the same three sub-cases as in part (i):

(a) $x = y$ and $j = l$:

Suppose $x = y$ and $j = l$. By Equation 9,

$$\begin{aligned}
 N^2 P_{xxjj}(t) &= \sum_{k=k'} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k') + \sum_{|k-k'|=\frac{N}{2}} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k') \\
 &\quad + \sum_{|k-k'| \neq \frac{N}{2}, 0} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k') \\
 &= \sum_{k=0}^{N-1} A_{jj}(t, k, k) + \sum_{|k-k'|=\frac{N}{2}} (-1)^x A_{jj}(t, k, k') \\
 &\quad + \sum_{|k-k'| \neq \frac{N}{2}, 0} e^{\frac{2\pi i x(k-k')}{N}} A_{jj}(t, k, k').
 \end{aligned}
 \tag{24}$$

By Proposition 2, the first and third summands of Equation 24 converge to $N/2$ and 0, respectively, and the second summand converges to

$$(-1)^{x-t} \frac{N}{2}.$$

It follows that

$$\lim_{t \rightarrow \infty} P_{xxjj}(t) = \frac{1}{N}$$

when $x - t$ is even and

$$\lim_{t \rightarrow \infty} P_{xxjj}(t) = 0$$

when $x - t$ is odd.

(b) $x \neq y$ and $j = l$:

By Equation 9,

$$\begin{aligned} N^2 P_{xyjj}(t) &= \sum_{k=k'} e^{\frac{2\pi i k(x-y)}{N}} A_{jj}(t, k, k') + \sum_{|k-k'|=\frac{N}{2}} e^{\frac{2\pi i x k}{N}} e^{\frac{-2\pi i y k'}{N}} A_{jj}(t, k, k') \\ &+ \sum_{|k-k'| \neq \frac{N}{2}, 0} e^{\frac{2\pi i x k}{N}} e^{\frac{-2\pi i y k'}{N}} A_{jj}(t, k, k') \\ &= \sum_{k=0}^{N-1} e^{\frac{2\pi i k(x-y)}{N}} A_{jj}(t, k, k) + \sum_{|k-k'|=\frac{N}{2}} (-1)^x e^{\frac{2\pi i k'(x-y)}{N}} A_{jj}(t, k, k') \\ &+ \sum_{|k-k'| \neq \frac{N}{2}, 0} e^{\frac{2\pi i (xk-yk')}{N}} A_{jj}(t, k, k'). \end{aligned} \tag{25}$$

By Proposition 2, $A_{jj}(t, k, k)$ converges to $1/2$ for all k , which implies that the first summand of Equation 25 converges to 0 as $t \rightarrow \infty$. In the third summand, all of the terms $A_{jj}(t, k, k')$ converge to 0. That just leaves us to evaluate the second summand, which can be reconstituted as

$$\frac{(-1)^{x-t}}{2} \sum_{|k-k'|=\frac{N}{2}} e^{\frac{i2\pi k'(x-y)}{N}} (-1)^t A_{jj}(t, k, k'). \tag{26}$$

By Proposition 2,

$$\lim_{t \rightarrow \infty} (-1)^t A_{jj}(t, k, k') = \frac{1}{2}.$$

Thus the third summand also converges to 0 as $t \rightarrow \infty$, and we conclude that

$$\lim_{t \rightarrow \infty} P_{xyjj}(t) = 0.$$

(c) $j \neq l$:

The assertion is proved in exactly the same way as in the corresponding case in part (i), when N is odd – see part(i)(c) above. □

Appendix D. Proof of Theorem 2

Proof. Let $\rho(\infty)$ denote the diagonal matrix

$$\text{diag} \left(\frac{1}{2N}, \frac{1}{2N}, \dots, \frac{1}{2N} \right).$$

If N is odd, Theorem 1 implies that

$$\lim_{t \rightarrow \infty} \|\rho(t) - \rho(\infty)\|_{\text{tr}} = 0.$$

By Lemma 2, we have

$$\lim_{t \rightarrow \infty} S(\rho(t)) = S(\rho(\infty)) = 1 + \ln N.$$

When N is even, we can prove the assertion

$$\lim_{t \rightarrow \infty} S(\rho(t)) = \ln N$$

in the same way. □

Appendix E. Proof of Theorem 3

Proof. When combined with Theorem 2, the inequality

$$S(\rho_c(t), \rho_w(t)) \leq S(\rho_c(t)) + S(\rho_w(t)) \leq 1 + \ln N$$

implies that

$$\lim_{t \rightarrow \infty} S(\rho_c(t), \rho_w(t)) = 1 + \ln N.$$

Thus,

$$\lim_{t \rightarrow \infty} [S(\rho_c(t)) + S(\rho_w(t))] = 1 + \ln N.$$

And since

$$S(\rho_c(t) : \rho_w(t)) = S(\rho_c(t)) + S(\rho_w(t)) - S(\rho_c(t), \rho_w(t)),$$

we have

$$\begin{aligned} \lim_{t \rightarrow \infty} S(\mathcal{H}_N : \mathcal{H}_c) &= \lim_{t \rightarrow \infty} S(\rho_c(t) : \rho_w(t)) \\ &= \lim_{t \rightarrow \infty} [S(\rho_c(t)) + S(\rho_w(t))] - \lim_{t \rightarrow \infty} S(\rho_c(t), \rho_w(t)) \\ &= 0. \end{aligned} \quad \square$$

References

Abal, G., Siri, R., Romanelli, A. and Donangelo, R (2006) Quantum walk on the line: entanglement and non-local initial conditions. *Physical Review A* **73** 042302 (Erratum: *Physical Review A* **73** 069905(E)).

Aharanov, D., Ambainis, A., Kempe, J. and Vazirani, U. (2001) Quantum walks on graphs. In: *Proceedings of the 33rd Annual ACM Symposium on Theory of Computing*, ACM 50–59.

Ambainis, A. (2003) Quantum walks and their algorithmic applications. *Int. J. Quantum Information* **1** (4) 507–518.

- Ambainis, A., Bach, E., Nayak, A., Vishwanath, A and Watrous, J. (2001) One-dimensional quantum walks. In: *Proceedings of the 33rd Annual ACM Symposium on Theory of Computing*, ACM 37–49.
- Annabestani, M., Abolhasani, M.R. and Abal, G. (2010a) Asymptotic entanglement in a two-dimensional quantum walk. *J. Phys. A: Math. Theor.* **43** 075301.
- Annabestani, M., Akhtarshenas, S.J. and Abolhassani, M.R. (2010b) Decoherence in one-dimensional Quantum Walk. *Physical Review A* **81** 032321.
- Banerjee, S., Srikanth, R., Chandrashekar, C.M. and Rungta, P. (2008) Symmetry-noise interplay in a quantum walk on an n -cycle. *Physical Review A* **78** 052316.
- Brun, T.A., Carteret, H.A. and Ambainis, A. (2003a) The quantum to classical transition for random walks. *Physical Review Letters* **91** (13) 130602.
- Brun, T.A., Carteret, H.A. and Ambainis, A. (2003b) Quantum random walks with decoherent coins. *Physical Review A* **67** 032304.
- Carneiro, I., Loo, M., Xu, X., Girerd, M., Kendon, V. and Knight, P.L. (2005) Entanglement in coined quantum walks on regular graphs. *New J. Phys.* **7** 156.
- Childs, A.M. (2010) On the relationship between continuous- and discrete-time quantum walk. *Communications in Mathematical Physics* **294** 581–603.
- Childs, A.M., Fahri, E and Gutmann, S. (2002) An example of the difference between quantum and classical random walks. *Quant. Inf. Proc.* **1** 35.
- Du, J., Li, H., Xu, X., Shi, M., Wu, J., Zhou, X. and Han, R. (2003) Experimental implementation of the quantum random-walk algorithm. *Physical Review A* **67** 042316.
- Dur, W., Raussendorf, R., Kendon, V. and Briegel, H.J. (2002) Quantum walks in optical lattices. *Physical Review A* **66** 052319.
- Eckert, K., Mompert, J., Birkel, G. and Lewenstein, M. (2005) One- and two-dimensional quantum walks in arrays of optical traps. *Physical Review A* **72** 012327.
- Fahri, E and Gutmann, S. (1998) Quantum computation and decision trees. *Physical Review A* **58** 915.
- Kempe, J. (2003) Quantum random walks – an introductory overview. *Contemp. Phys.* **44** (4) 307–327.
- Kendon, V. (2006) Decoherence in quantum walks – a review. *Mathematical Structures in Computer Science* **17** 1169–1220.
- Kendon, V. and Tregenna, B. (2002) Decoherence in a quantum walk on a line. In: Shapiro, J.H. and Hirota, O. (eds.) *Quantum Communication, Measurement & Computing (QCMC-02)*, Rinton Press 463.
- Kendon, V. and Tregenna, B. (2003a) Decoherence in discrete quantum walks (available at arXiv:quant-ph/0301182).
- Kendon, V. and Tregenna, B. (2003b) Decoherence is useful in quantum walks. *Physical Review A* **67** 042315.
- Konno, N. (2008) Quantum walks. In: Franz, U. and Schurmann, M. (eds.) *Quantum Potential Theory. Springer-Verlag Lecture Notes in Mathematics* **1954** 309–452.
- Kosik, J., Buzek, V. and Hillery, M. (2006) Quantum walks with random phase shifts. *Physical Review A* **74** 022310.
- Liu, C. and Petulante, N. (2010) Quantum walks on the N -cycle subject to decoherence on the coin degree of freedom. *Physical Review E* **81** 031113.
- Maloyer, O and Kendon, V. (2007) Decoherence versus entanglement in coined quantum walks. *New Journal of Physics* **9** 87.
- Nayak, A and Vishwanath, A. (2000) Quantum Walk on the Line (available at arXiv:quant-ph/0010117).

- Richter, P. (2007) Quantum speedup of classical mixing processes. *Physical Review A* **76** 042306.
- Romanelli, A., Siri, R., Abal, G., Auyuanet, A. and Donangelo, R. (2005) Decoherence in the quantum walk on the line. *Physica A* **347** 137–152.
- Ryan, C.A., Laforest, M., Boileau, J.C. and Laflamme, R. (2005) Experimental implementation of a discrete-time quantum random walk on an NMR quantum-information processor. *Physical Review A* **72** 062317.
- Sanders, B., Bartlett, S., Tregenna, B. and Knight, P. (2003) Quantum quincunx in cavity quantum electrodynamics. *Physical Review A* **67** 042305.
- Shikano, Y., Chisaki, K., Segawa, E. and Konno, N. (2010) Emergence of randomness and arrow of time in quantum walks. *Physical Review A* **81** 062129.
- Strauch, F. (2006) Connecting the discrete- and continuous-time quantum walks. *Physical Review A* **74** 030301(R).
- Travaglione, B.C. and Milburn, G.J. (2002) Implementing the quantum random walk. *Physical Review A* **65** 032310.
- van Hoogdalem, K. A. and Blaauboer, M. (2009) Implementation of the quantum-walk step operator in lateral quantum dots. *Physical Review B* **80** 125309.
- Venegas-Andraca, S.E. (2008) *Quantum Walks for Computer Scientists*, (Synthesis Lectures on Quantum Computing), Morgan and Claypool Publishers.
- Venegas-Andraca, S.E. and Bose, S. (2009) Quantum Walk-based Generation of Entanglement Between Two Walkers (available at quant-ph/0901.3946).
- Watrous, J. (2008) Theory of Quantum Information, Lecture notes from Fall 2008, Institute for Quantum Computing, University of Waterloo, Canada.
- Zhang, K. (2008) The limiting distribution of decoherent quantum random walks. *Physical Review A* **77** 062302.
- Zou, X., Dong, Y., and Guo, G. (2006) Optical implementation of one-dimensional quantum random walks using orbital angular momentum of a single photon. *New J. Phys.* **8** 81.
- Zurek, W.H. (2003) Decoherence and the transition from quantum to classical – Revisited. In: Duplantier, B., Raimond, J.-M. and Rivasseau, V. (eds.) *Decoherence Poincaré Seminar 2005*, *Progress in Mathematical Physics*, Birkhuser Verlag 1–31. (Eprint available at arXiv:quant-ph/0306072.)