

cambridge.org/psm


## Correspondence

**Cite this article:** Liu RT (2023). Correspondence: Psychological medicine. *Psychological Medicine* **53**, 284–285. <https://doi.org/10.1017/S0033291721002841>

Received: 20 June 2021  
Accepted: 27 June 2021  
First published online: 2 August 2021

**Author for correspondence:**

Richard T. Liu,  
E-mail: [rliupsych@gmail.com](mailto:rliupsych@gmail.com)

Richard T. Liu 

Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, USA

## To the Editor:

I would like to thank Sally McManus and colleagues for this opportunity to engage in a correspondence regarding this important topic of non-suicidal self-injury (NSSI) epidemiology.

I agree with several points shared by McManus and colleagues. Prevalence of several clinical phenomena, such as suicide and depression, have been observed to change at the population level over time (Hedegaard, Curtin, & Warner, 2020; Mojtabai, Olfson, & Han, 2016). McManus and colleagues provide important and unique data that similarly suggest shifts in prevalence of NSSI with time (McManus et al., 2019).

The observation that a relatively older data set was used in my paper is also absolutely correct. The assertion that the year of data collection was not presented in the introduction, however, is not; it is mentioned in the introduction, followed by a citation of McManus, Meltzer, Brugha, and Bebbington (2009), and it is mentioned again in the methods section. Beyond indicating the date of data collection, care was taken to specify that the data were from the 2007 Adult Psychiatric Morbidity Survey (APMS), with the aforementioned citation provided, to avoid potential confusion with the 2000 and 2014 surveys of the same name. The 2007 survey data were used because they remain, to my knowledge, the most recent epidemiological data publicly available that would allow for addressing the aims of my paper. The limitations of these data were mentioned in the discussion section, as were the uniquely valuable insight these data nonetheless provide regarding the phenomenology of NSSI. There is also precedence for such use of older epidemiological data. The National Comorbidity Survey Adolescent Supplement serves as a prominent example. Data collection was from February 2001 to January 2004. Yet, several important studies on prevalence and/or psychiatric comorbidity have been published after a similarly long period of time, or even longer (e.g. Burstein, Beesdo-Baum, He, & Merikangas, 2014; Nock et al. 2013; Paksarian et al. 2020; Platt, Keyes, McLaughlin, & Kaufman, 2019). This reflects the unique value that such epidemiological data continue to provide.

As for being the first epidemiological study of NSSI, were such a claim made of the current study, I agree that it would be indefensible. This claim, however, was never made. Indeed, the second paragraph of the introduction focused entirely on several prior epidemiological studies of this phenomenon. Moreover, in the introduction, I made a qualified statement: my intent was to offer ‘the first epidemiological study to present (i) fine-grained data on sociodemographic and psychiatric correlates of NSSI, (ii) evaluations of NSSI in relation to suicidal ideation and suicide attempts, and (iii) correlates of medical and psychiatric treatment use specifically for NSSI’ (Liu, in press), rather than the first NSSI epidemiological study of any kind. I will briefly elaborate on each of the three qualifiers in turn. First, most, if not all, epidemiological studies have provided some data on sociodemographic correlates. In my study, however, I was interested in presenting unique findings relating to a broad range of sociodemographic characteristics, including sex, race/ethnicity, age, education, marital status, income, poverty, and parental involvement during childhood. Also, to my knowledge, no prior epidemiological study has examined psychiatric disorders in relation to NSSI, outside of a notable recent study with college students (Kiekens et al., 2018; although here the focus was on the provisional DSM-5 NSSI disorder rather than NSSI occurrence).<sup>†1</sup> Second, to my knowledge, no prior epidemiological study has examined NSSI as a predictor of suicidal ideation and behavior. Third, regarding medical and psychiatric treatment for NSSI, McManus et al. (2019) combined the two in presenting some valuable findings on temporal changes over time and individual characteristics as predictors. In contrast, I sought to observe the important distinction between treatment for physical injury from NSSI (e.g. receiving stitches) and treatment for it as a mental health concern (e.g. receiving psychotherapy for NSSI), and as detailed in my paper, doing so yielded notably different patterns of findings for each.

Finally, I would like to share my view that the primary aims of McManus and colleagues’ recent study (McManus et al., 2019) and mine are rather complementary. The former leverages the APMS data sets to provide unique insight on population-level temporal trends over a 15-year period, whereas the latter sought to present an in-depth snapshot characterizing socio-demographic and clinical phenomena associated with NSSI. These two studies therefore

<sup>†</sup>The notes appear after the main text.

collectively provide a more complete understanding of NSSI than would be the case in a single study by itself.

## Notes

1 Of note, Kiekens et al. (in press) have a new study just recently published online in this journal, examining psychiatric disorders associated with NSSI occurrence in the same population.

## References

- Burstein, M., Beesdo-Baum, K., He, J. P., & Merikangas, K. R. (2014). Threshold and subthreshold generalized anxiety disorder among US adolescents: Prevalence, sociodemographic, and clinical characteristics. *Psychological Medicine*, 44, 2351–2362. <https://doi.org/10.1017/S0033291713002997>.
- Hedegaard, H., Curtin, S. C., & Warner, M. (2020). Suicide mortality in the United States, 1999–2018. *NCHS Data Brief*, 362, 1–8.
- Kiekens, G., Hasking, P., Bruffaerts, R., Alonso, J., Auerbach, R. P., Bantjes, J., ... Kessler, R. C. (in press). Non-suicidal self-injury among first-year college students and its association with mental disorders: Results from the World Mental Health International College Student (WMH-ICS) initiative. *Psychological Medicine*.
- Kiekens, G., Hasking, P., Claes, L., Mortier, P., Auerbach, R. P., Boyes, M., ... Bruffaerts, R. (2018). The DSM-5 nonsuicidal self-injury disorder among incoming college students: Prevalence and associations with 12-month mental disorders and suicidal thoughts and behaviors. *Depression and Anxiety*, 35, 629–637. <https://doi.org/10.1002/da.22754>.
- Liu, R. T. (in press). The epidemiology of non-suicidal self-injury: Lifetime prevalence, sociodemographic and clinical correlates, and treatment use in a nationally representative sample of adults in England. *Psychological Medicine*. <https://doi.org/10.1017/S003329172100146X>.
- McManus, S., Gunnell, D., Cooper, C., Bebbington, P. E., Howard, L. M., Brugha, T., ... Appleby, L. (2019). Prevalence of non-suicidal self-harm and service contact in England, 2000–14: Repeated cross-sectional surveys of the general population. *The Lancet. Psychiatry*, 6, 573–581. [https://doi.org/10.1016/S2215-0366\(19\)30188-9](https://doi.org/10.1016/S2215-0366(19)30188-9).
- McManus, S., Meltzer, H., Brugha, T., & Bebbington, P. (2009). *Adult psychiatric morbidity in England, 2007: Results of a household survey*. NHS Information Centre for Health and Social Care.
- Mojtabai, R., Olfson, M., & Han, B. (2016). National trends in the prevalence and treatment of depression in adolescents and young adults. *Pediatrics*, 138, e20161878. <https://doi.org/10.1542/peds.2016-1878>.
- Nock, M. K., Green, J. G., Hwang, I., McLaughlin, K. A., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2013). Prevalence, correlates, and treatment of lifetime suicidal behavior among adolescents: Results from the national comorbidity survey replication adolescent supplement. *JAMA Psychiatry*, 70, 300–310. <https://doi.org/10.1001/2013.jamapsychiatry.55>.
- Paksarian, D., Rudolph, K. E., Stapp, E. K., Dunster, G. P., He, J., Mennitt, D., ... Merikangas, K. R. (2020). Association of outdoor artificial light at night with mental disorders and sleep patterns among US adolescents. *JAMA Psychiatry*, 77, 1266–1275. <https://doi.org/10.1001/jamapsychiatry.2020.1935>.
- Platt, J. M., Keyes, K. M., McLaughlin, K. A., & Kaufman, A. S. (2019). Intellectual disability and mental disorders in a US population representative sample of adolescents. *Psychological Medicine*, 49, 952–961. <https://doi.org/10.1017/S0033291718001605>.