

Challenges and opportunities for quantifying greenhouse gas emissions through dairy cattle research in developing countries

Research Reflection

Cite this article: Munidasa S, Eckard R, Sun X, Cullen B, McGill D, Chen D and Cheng L (2021). Challenges and opportunities for quantifying greenhouse gas emissions through dairy cattle research in developing countries. *Journal of Dairy Research* **88**, 3–7. <https://doi.org/10.1017/S0022029921000182>

Received: 28 September 2020

Revised: 16 January 2021

Accepted: 25 January 2021

Keywords:

Emission estimation; livestock production; low-and medium-income countries; sustainability

Author for correspondence:

Long Cheng,
Email: long.cheng@unimelb.edu.au

Sineka Munidasa¹, Richard Eckard², Xuezhao Sun^{3,4}, Brendan Cullen²,
David McGill², Deli Chen² and Long Cheng¹

¹Faculty of Veterinary and Agricultural Sciences, Dookie Campus, The University of Melbourne, Victoria 3647, Australia; ²Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, Victoria 3010, Australia; ³The Innovation Centre of Ruminant Precision Nutrition and Smart and Ecological Farming, Jilin Agricultural Science and Technology University, Jilin City, Jilin Province, China and ⁴Jilin Inter-regional Cooperation Centre for the Scientific and Technological Innovation of Ruminant Precision Nutrition and Smart and Ecological Farming, Jilin City, Jilin Province, China

Abstract

The global dairy sector is facing the challenge of reducing greenhouse gas (GHG) emissions whilst increasing productivity to feed a growing population. Despite the importance of this challenge, many developing countries do not have the required resources, specifically funding, expertise and facilities, for quantifying GHG emissions from dairy production and research. This paper aims to address this challenge by discussing the magnitude of the issue, potential mitigation approaches and benefits in quantifying GHG emissions in a developing country context. Further, the paper explores the opportunities for developing country dairy scientists to leverage resources from developed countries, such as using existing relevant GHG emission estimation models. It is clear that further research is required to support developing countries to quantify and understand GHG emissions from dairy production, as it brings significant benefits including helping to identify and implement appropriate mitigation strategies for local production systems, trading carbon credits and achieving the nationally determined contribution obligations of the Paris Agreement.

Dairy production has been a crucial part of agricultural and food systems around the world for centuries. There is an increasing need to better understand and quantify the impact of worldwide dairy systems on climate change as dairy production emits greenhouse gases, such as methane (CH₄) and nitrous oxide (N₂O) (FAO and GDP, 2018). Enteric CH₄ production accounts for nearly 60% of GHG emissions from dairy cattle (FAO and GDP, 2018), and greenhouse gas emissions represent a large inefficiency in dairy production because these gases represent a loss of energy and nitrogen (a fundamental component of protein) from the dairy cattle, which otherwise can be used for production gains (Eckard and Clark, 2020). Hence, quantifying and managing GHG emissions from dairy cattle is essential to support the sustainable development of dairying. Moreover, accurate quantification will guide the setting of future national and international policies to meet the goals set in the Paris Agreement 2015 by 195 countries (FAO and GDP, 2018).

Dairy sectors in developing countries are dominated by smallholder dairy production systems. In these systems, farming households tend to keep 2–5 animals per herd (Staal *et al.*, 2016) and in the majority of cases are an integrated crop-livestock farming system, where animal feeding is made up of crop residues in combination with a cut and carry system for fodders (in more resource rich areas) through to free ranging systems (in resource poor areas). These systems tend to rely on family labour and produce milk for both household consumption and, where possible, sale into the market. In many cases, milk in developing countries comes from these smallholder systems. For instance, 85% of total Kenya milk production comes from smallholders (Gerosa and Skoet, 2013) and in Pakistan 95% of milk consumed is from local and unpackaged milk sources produced by smallholder farmers (Zia *et al.*, 2011). Intensification of the dairy system is common in developing countries as a response to increased demand for milk. However, traditional smallholder systems still remain dominant (Staal *et al.*, 2016). Developing countries contribute to a large proportion of global milk production. For instance, Asia, which is primarily made up of developing countries, accounts for 42% (360 million tonnes) of the world's milk production (FAO, 2019). Many of the developing countries' dairy research programmes and projects have little capacity for quantifying GHG emissions due to resource and expertise limitations. Thus, the main goal of this Research Reflection was to explore challenges and opportunities for quantifying GHG through dairy research in developing countries.

Importance of dairy production for food security, livelihood and economy in developing countries

Global demand for food is expected to more than double by 2050 as a result of human population growth and an increase in living standards, particularly in developing countries (Rojas-Downing *et al.*, 2017). Demand for milk is expected to rise by 48% from 2005 to 2050 (Alexandratos and Bruinsma, 2012), with 14% of the world's population benefiting from income generated from dairy production (Hemme and Otte, 2010). This proportion increases dramatically in developing country dairy sectors, where more than 50% of the world's milk-producing cattle are located (FAOSTAT, 2018). Developing countries are increasingly demanding dairy products, while developed countries are traditionally high in dairy product consumption (Gerber *et al.*, 2013).

Dairy production is often more than just income generation for rural farming families. It contributes to livelihoods, food security and ultimately helps to achieve a number of the United Nations Sustainable Development Goals in the developing countries (Reisinger and Clark, 2018). Since dairying plays an important role in economic and social wellbeing in developing countries, maintaining sustainable dairy production is crucial. However, expansion of these dairy production systems may negatively impact their environmental sustainability, largely due to GHG emissions from dairy production systems. Therefore, quantifying and mitigating GHG emissions from dairy cattle is an essential component to support the sustainable development of dairying in developing countries.

Importance of understanding and quantifying greenhouse gas emissions from dairy production through research in developing countries

According to Tubiello (2011), three-quarters of global agricultural GHG emissions are produced by developing countries. The livestock industry is a key part of global agriculture production and it contributes to about 15% of global anthropogenic emissions. From this, dairy cattle production accounts for approximately 30% of the livestock sector's emissions globally (Gerber *et al.*, 2013). In 2015, dairy production added 1,711.8 million tonnes of carbon dioxide equivalent (CO₂e) GHG to the atmosphere (FAO and GDP, 2018).

Emission can be expressed as both an absolute emission (per hectare, per cow) and emission intensity (emission per unit of product) term (Reisinger and Ledgard, 2013). The least productive countries may have the highest emission intensities (per unit of milk). However, there may be a low emission per hectare depending on the stocking rate and relatively low GHG emission per cow. The trade-off between these metrics can be complicated. For example, Eckard *et al.* (2010) modelled and compared the pasture-based dairy systems in terms of low-quality and high-quality diets by adjusting the stocking rates using the DairyMod, a mechanistic whole-farm systems model. Their analysis showed that feeding high-quality pasture reduced emission intensity (kg CO₂e per litre milk) by 19%, however, emission per hectare increased by 26% due to the higher stocking rates compared to low quality pasture feeding. This result shows the importance of having different metrics to quantifying emissions as it gives a different dimension of the issue. Further, these metrics are important for different policy contexts. For example, absolute emission can be used for determining the contribution of a country to gross emission, while emission intensity is more

useful in benchmarking production efficiency between systems, regions and countries (Reisinger and Ledgard, 2013).

It is interesting to note that North American and South Asian dairy cattle produced 116.6 kg CH₄ and 9867 kg milk per head per year and 62.1 kg CH₄ and 1388 kg milk per head per year, respectively (FAO and GDP, 2018). These values indicate a major difference in their GHG emission rates and milk production performances due to production system differences. Similarly, these values highlight the considerable differences in GHG emission intensities (emission per unit of product) between developed (North American) and developing countries (South Asian). Further, when it comes to emission intensity measures, a recent study showed that developed country dairy production had a lower emission intensity (averaged 1.4 kg CO₂e per kg fat-and protein- corrected milk (FPCM)) compared with developing country dairy production (averaged 5.4 CO₂e per kg FPCM) (FAO and GDP, 2018).

An important consideration in the developing country context is that dairy cattle in these systems play multiple functions on the smallholder farms. In these systems, dairy cattle are not just a source of milk, but they are also a source of draught power, store of capital, insurance for crop failure and provide nutrient cycling for cropping (Tricarico *et al.*, 2020). This multifunctional nature of the contribution needs to be taken into account in determining their actual emissions contribution. For instance, a case-study of mixed smallholder milk production systems conducted in Kenya emphasized that including multiple functions of dairy cattle has a strong impact on estimation outcomes of emission intensity of dairying (Weiler *et al.*, 2014).

Increasing productivity has a risk of reducing some of the non-food use of dairying, for instance, draught power. However, the benefit as a store of capital, insurance for crop failure and nutrients for cropping will still apply. Increasing productivity could bring more revenue for the farmers and will increase the benefit of insurance for crop failure, as it will provide more economic opportunities to ensure the improved livelihood of farmers. Further, milk will bring more benefits than the non-food uses for farm families and the rural communities, as it fulfils the human nutrition requirement by providing high-quality protein and other micronutrients. For example, the deficit for milk in Ethiopia is approximately 4.5 billion litres per year (Tricarico *et al.*, 2020). This can be addressed by increasing the productivity of the systems, which will ultimately reduce the GHG emission intensity of the system as well.

Currently, many developing countries use default Intergovernmental Panel on Climate Change (IPCC) Tier 1 method to estimate the GHG emissions, which adopts a fixed emission value/factor per animal as there is no local available data to practice Tier 2 calculations. This leads to incorrect estimations of emissions, as Tier 1 method does not account for local livestock factors such as milk production, feed quality and environmental conditions. A recent study on CH₄ emissions from selected African cattle production showed approximately 40% overestimation of emission when IPCC Tier 1 method was used than using method considered dairy cattle intake (Goopy *et al.*, 2018). This demonstrates clearly the need to more accurately measure and/or estimate GHG emissions from dairy cattle in developing countries. Further, lack of activity data (e.g. animal number and information about different dairy production systems in developing countries) is another major problem associated with the estimation of GHG emissions. Therefore, developing countries use other generalized data sources, for instance, FAOSTAT. This

will lead to over/underestimation of GHG emissions. Therefore, more specific data sources are required to fill these activity data gaps to support adopting higher inventories in developing countries.

Overall, accurately quantifying GHG emissions from dairy cattle through more research in developing countries, across a range of relevant production systems, is important for identifying GHG emissions by size and source. This in turn; (1) allows scientists to explore locally relevant GHG emissions management and mitigation strategies, (2) provides evidence for producers or the dairy industry to trade in carbon credits on voluntary markets, providing the potential for income diversification and (3) supports the transparent process of reaching and reporting nationally determined contribution obligations set in the Paris Agreement (IIED, 2019).

There are a number of management practices which can be implemented for mitigating GHG emission and/or GHG emissions intensity. These include nutritional intervention, herd size, manure management, housing systems and energy efficiency (including renewables) (Grossi *et al.*, 2019). Mitigation potential of GHG emissions using these strategies is highly diverse among farms and regions depending on different factors. For example, diversity of the production systems, environmental conditions, availability of the technology, economic incentives and institutional frameworks influence the degree of mitigation possible (FAO, 2010). However, while dairy systems in developing countries have potential to improve production and efficiency, it is not as simple as adopting those technologies and practices used in developed countries, where evidence is mounting that these more intensive systems may not be environmentally sustainable (Weiler *et al.*, 2014). Therefore, greater investment in dairy research in developing countries is required to address these complex situations and build more sustainable systems.

Challenges for quantifying greenhouse gas emissions through dairy research in developing countries

During the last few decades, quantifying and managing environmental impacts has been recognized as an important part of dairy research (Cheng, 2020). However, the amount of research of this sort that is funded in developing countries is limited compared to that within developed countries. Such differences are likely a reflection of food security being a major pressing priority in many of the developing countries, but also that government funds for research in developing countries are generally at a lower level than in developed countries (Gaffney *et al.*, 2019). Further, lack of expertise and facilities are two major constraints associated with quantifying GHG emissions through dairy research in developing countries.

There are a number of methods developed for quantifying GHG emissions from dairy cattle under research conditions. For example, enteric CH₄ emissions from dairy cattle can be measured by respiration chamber, sulphur hexafluoride tracer technique, ventilated hood /face mask and using the Greenfeed system (Hammond *et al.*, 2016; Zhao *et al.*, 2020). However, all these methods require substantial facilities and expertise to accurately quantify CH₄ emissions. While respiration chambers are regarded as a gold standard for CH₄ measurements, the system is costly to build and operate and as a result has historically only been used in developed countries (e.g. New Zealand, Australia, Belgium, Denmark and Germany: Pinares-Patiño and Waghorn, 2018), although chambers are now available in Brazil, China and Kenya.

The lack of relevant expertise and facilities in developing countries is also evident in a lack of publications in this area. The six countries with the largest dairy populations are India, Brazil, Sudan, Pakistan, China and Ethiopia. These countries account for nearly 40% of the total world dairy cattle population (AHDB Dairy, 2019). Using these individual country names together with methane and dairy, a search of the PubMed database returned only 4 publications between the year 2010 and 2020. These studies were conducted in China (2), Brazil (1) and India (1). In contrast, during the same period, scientists from Australia produced 9 publications, Canada published 13 publications and a similar case was found from other major developed countries. This simple comparison is an indicator that these developing countries are not quantifying dairy cattle GHG emissions often, which is a combination of lack of research funding, relevant facilities, expertise and environmental impacts being a lower research priority.

Opportunities for quantifying greenhouse gas emissions through dairy research in developing countries

Opportunities exist to allow scientists in developing countries to measure and estimate GHG emissions from dairy cattle. Firstly, open-access publications are increasingly available online to allow these scientists to learn about the latest dairy cattle GHG emissions research. Secondly, developed countries are offering support under the Paris Agreement for capacity building, transferring technology and financial support for minimizing GHG emissions in developing countries (FAO, 2016). The Australia Centre for International Agricultural Research (ACIAR) was established to support capability building in developing countries, including GHG emissions quantification and mitigation. Thirdly, there are many international organizations, individually and collaboratively, involved in supporting the development of skills and expertise of livestock GHG emissions management, including the Global Research Alliance on Agricultural Greenhouse Gases (GRA), the International Livestock Research Institute and Food and Agriculture Organization (FAO). These are good opportunities for dairy scientists in developing countries to leverage expertise and facilities from developed countries and enhance their understandings of GHG emissions and locally appropriate mitigation options.

Adopting existing greenhouse gas emission estimation models

There is a range of approaches that can be used to estimate GHG emissions from dairy cattle, ranging from the use of emissions factors based on animal numbers (analogous to IPCC Tier 1 methods; IPCC, 2006) to mechanistic models (Tier 3) that simulate the processes involved (Rotz, 2018). The simple emission factor approach requires minimal data collection but may not reflect the environmental conditions well. The more complex modelling approaches require more data (e.g. rumen function, physiological stage and rate of feed passage for enteric CH₄ emission estimation using process-based mechanistic models) and may provide more accurate estimates of emissions (Rotz, 2018). Data requirements for Tier 2 methods are intermediate between the two approaches. The Tier 2 methods use detailed information on sub-groups of animals along with estimates of feed intake based on live weight, live weight gain, milk yield, feed digestibility and other factors that reflect the local management practices (Wilkes and van Dijk, 2018).

Table 1. Dairy cattle enteric methane production estimation models, that include a tropical dataset

Model	Sample size	Region	Breed	Accuracy/precision	Reference
CH ₄ production (MJ/day) = 0.063 × GEI	Dairy cattle = 220, Beef cattle = 813	Temperate and tropical	Holstein–Friesian, Angus, Brahman, Shorthorn	$R^2 = 0.93, P < 0.001$	Charmley <i>et al.</i> (2016)
CH ₄ production (g/day) = 20.7 × DMI				$R^2 = 0.92, P < 0.001$	
CH ₄ production (MJ/day) = 3.35 + 0.047 × GEI	Lactating cattle = 284	Tropical	Girolando, Gir and Holstein	$R^2 = 0.55, P < 0.05$	Ribeiro <i>et al.</i> (2020)
CH ₄ production (MJ/day) = 4.15 + 0.822 × DMI				$R^2 = 0.50, P < 0.05$	
CH ₄ production (MJ/day) = 1.29 + 0.878 × DMI	Dairy and beef cattle = 830	Tropical	Mainly Zebu type	$R^2 = 0.70, P < 0.001$	Patra (2017)

GEI, Gross energy intake (MJ per day); DMI, Dry matter intake (kg per day).

While there is substantial potential to draw on the existing estimation models from developed countries to quantify GHG emissions in developing countries through dairy research, selection of the suitable modelling approaches requires careful consideration of the availability of local data and experimental conditions (dietary and animal characteristics). The production systems in developed countries and developing countries are different in numerous ways such as feeding practices, climatic conditions and animal characteristics (breeds, production capacity, heat tolerance). Dairy systems in developed countries are characterized normally by larger herd size, high yielding breeds and, often, confinement-based production systems either annually or seasonally. Further, these systems rely on stored forages, purchased grains and concentrates. However, there are some developed country production systems which are more grazing-based, for example, Australia, New Zealand and Ireland. In contrast, the smallholder systems of developing countries have lower-yielding, but more locally adapted dairy cattle breeds. Dairy systems of developing countries are mostly based on locally produced grasses and crop residues (e.g. corn stover, cereal straw). For instance, in India, dairy cattle diets are commonly based on approximately 61% crop residues and green fodders, 19% of by-products and 20% concentrates (FAO, IDF and IFCN, 2014). Therefore, direct application of GHG estimation models from developed countries may lead to over/underestimation of GHG emissions (FAO, IDF and IFCN, 2014). Local research to validate and/or adapt the approaches will be helpful to increase the estimation accuracy/precision. For instance, Ribeiro *et al.* (2020) evaluated the estimation precision and accuracy of the existing enteric CH₄ estimation models from the literature for the tropical conditions and recommended the model proposed by Charmley *et al.* (2016) be used for national inventories in tropical countries based on high concordance with local data (Table 1).

Use of proxies or markers to indicate greenhouse gas emissions

Many proxies have been developed to estimate the enteric CH₄ emission. These proxies, indicators or indirect traits, vary from simple and low-cost measures to complex and expensive methods (Negussie *et al.*, 2017). The ideal proxies for estimating GHG through research in developing countries should be cost-effective and require minimal expertise and facilities. Although there are plenty of proxies to estimate enteric CH₄ emission, the methods that use rumen samples, faeces, urine and blood are common,

but these samples are not always easy to obtain without intrusive procedures.

Negussie *et al.* (2017) published a systematic review on potential CH₄ proxies including feed intake and feeding behaviour, rumen function, metabolites and microbiome, milk production and composition, hindgut and faeces, and measurements at the level of the whole animal. According to the review, milk mid-infrared spectroscopy (MIR) and milk fatty-acid profile are promising methods to derive proxies for CH₄ estimation. Notably, Bougouin *et al.* (2019) found that milk fatty-acids have better potential to estimate enteric CH₄ production when combined with other variables (e.g. dry matter intake, body weight) rather than using only milk fatty-acids. However, the milk fatty-acid based method is expensive, which makes it less practical to use under many developing countries conditions.

There is an interesting perspective to consider GHG emissions model data proxies. Models rely on animal activity data, such as regional animal numbers. Sometimes regional animal numbers may not be available, but the district veterinary officer often knows how many vaccines to order each year. This may be a proxy to estimate animal numbers. Researchers could also use local information on household structures to estimate the number of cows per household in a region. Further, measuring dry matter intake to estimate enteric CH₄ emission is challenging. However, animal productivity data (e.g. milk production per day), can use as a proxy to estimate feed dry matter intake of the animal. These proxies are mainly important for inventory purposes. Overall, future exploration of potential proxies to estimate GHG emissions through dairy research is an essential step to support sustainable dairy production development in developing countries.

In conclusion, the dairy industry globally faces increasing pressure to reduce GHG emissions. There is a great potential for developing countries to contribute to this reduction, as a large proportion of dairy cattle are part of the farming systems in these countries with higher emission intensities compared to dairy sectors in developed countries. The multi-functional value of dairy cattle in subsistence agriculture is important to be considered when it comes to the development of sustainable dairy production and research in developing countries. The research capacity to quantify GHG emissions in developing countries is limited. Despite this, local expertise and facilities could be supported with application of appropriate research technologies from developed countries to help understand the source and scale of GHG emissions from dairy cattle in the different farming systems.

References

- AHDB Dairy (2019) World Dairy Cow Numbers. Available at <https://dairy.ahdb.org.uk/resources-library/market-information/farming-data/world-cow-numbers/#.Xvv2AG5uI2w> (Accessed 01.07.2020).
- Alexandratos N and Bruinsma J (2012) *World Agriculture Towards 2030/2050: The 2012 Revision*. ESA Working paper No. 12-03. Rome: Food and Agriculture Organization of the United Nations.
- Bougouin A, Appuhamy JRN, Ferlay A, Kebreab E, Martin C, Moate PJ, Benchaar C, Lund P and Eugène M (2019) Individual milk fatty acids are potential predictors of enteric methane emissions from dairy cows fed a wide range of diets: approach by meta-analysis. *Journal of Dairy Science* **102**, 10616–10631.
- Charmley ESRO, Williams SRO, Moate PJ, Hegarty RS, Herd RM, Oddy VH, Reyenga P, Staunton KM, Anderson A and Hannah MC (2016) A universal equation to predict methane production of forage-fed cattle in Australia. *Animal Production Science* **56**, 169–180.
- Cheng L (2020) The need for consideration of environmental implications in developing countries' dairy research. *Journal of Dairy Research* **87**, 143–144.
- Eckard RJ and Clark H (2020) Potential solutions to the major greenhouse-gas issues facing Australasian dairy farming. *Animal Production Science* **60**, 10–16.
- Eckard RJ, Grainger C and De Klein CAM (2010) Options for the abatement of methane and nitrous oxide from ruminant production: a review. *Livestock Science* **130**, 47–56.
- FAO (2010) Greenhouse gas emissions from the dairy sector. *A Life Cycle Assessment*. pp 89–91. Eds Gerber P, Vellinga T, Opio C, Henderson B and Steinfeld H, Rome: Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/3/k7930e/k7930e00.pdf> (accessed 05. 12. 2020).
- FAO (2016) *The Agricultural Sectors in Nationally Determined Contributions (NDCs)*. Rome: Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/3/a-i6400e.pdf> accessed on 28. 07. 2020
- FAO (2019) *Dairy Market Review*. Rome: Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/3/ca3879en/ca3879en.pdf> (accessed 10. 09. 2020).
- FAO and GDP (2018) *Climate Change and the Global Dairy Cattle Sector – The Role of the Dairy Sector in a Low-Carbon Future*. Rome: Food and Agriculture Organization of the United Nations and Global Dairy Platform. Available at <http://www.fao.org/3/CA2929EN/ca2929en.pdf> (accessed 30. 07. 2020).
- FAO, IDF and IFCN (2014) *World Mapping of Animal Feeding Systems in the Dairy Sector*. Rome: Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/3/a-i3913e.pdf> (accessed 28.11. 2020).
- FAOSTAT (2018) *Statistical Databases*. Rome: Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/faostat/en/#data/QA> (accessed 27. 08. 2020).
- Gaffney J, Challenger M, Califf K and Harden K (2019) Building bridges between agribusiness innovation and smallholder farmers: a review. *Global Food Security* **20**, 60–65.
- Gerber PJ, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, Faluccci A and Tempio G (2013) *Tackling Climate Change Through Livestock: A Global Assessment of Emissions and Mitigation Opportunities*. Rome, Italy. Food and Agriculture Organization of the United Nations (FAO)
- Gerosa S and Skoet J (2013) Milk availability: current production and demand and medium term outlook, chapter 2, pp 24–25. In Muehlhoff E, Bennett and McMahon D (eds), *Milk and Dairy Products in Human Nutrition*. Rome, Italy. Food and Agriculture Organization of the United Nations (FAO).
- Goopy JP, Onyango AA, Dickhoefer U and Butterbach-Bahl K (2018) A new approach for improving emission factors for enteric methane emissions of cattle in smallholder systems of East Africa – Results for Nyando, Western Kenya. *Agricultural Systems* **161**, 72–80.
- Grossi G, Goglio P, Vitali A and Williams AG (2019) Livestock and climate change: impact of livestock on climate and mitigation strategies. *Animal Frontiers* **9**, 69–76.
- Hammond KJ, Crompton LA, Bannink A, Dijkstra J, Yáñez-Ruiz DR, O'Kiely P, Kebreab E, Eugène MA, Yu Z, Shingfield KJ and Schwarm A (2016) Review of current in vivo measurement techniques for quantifying enteric methane emission from ruminants. *Animal Feed Science and Technology* **219**, 13–30.
- Hemme T and Otte J (2010) *Status and Prospects for Smallholder Milk Production: A Global Perspective*. Rome, Italy. Food and Agriculture Organization of the United Nations (FAO)
- IIED (2019) *Briefing. Meeting the Enhanced Transparency Framework: What Next for the LDCs?* Copenhagen, Denmark. The International Institute for Environment and Development. Available at <https://pubs.iied.org/pdfs/17730IIED.pdf> (accessed 21. 08. 2020).
- IPCC (2006) Guidelines for national greenhouse gas inventories. Volume 4, Agriculture, forestry and other land use. Chapter 10, Emissions from Livestock and manure management. Intergovernmental Panel on Climate Change. Available at https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_10_Ch10_Livestock.pdf (accessed 23. 07. 2020).
- Negussie E, de Haas Y, Dehareng F, Dewhurst RJ, Dijkstra J, Gengler N, Morgavi DP, Soyeurt H, van Gastelen S, Yan T and Biscarini F (2017) Invited review: large-scale indirect measurements for enteric methane emissions in dairy cattle: a review of proxies and their potential for use in management and breeding decisions. *Journal of Dairy Science* **100**, 2433–2453.
- Patra AK (2017) Prediction of enteric methane emission from cattle using linear and non-linear statistical models in tropical production systems. *Mitigation and Adaptation Strategies for Global Change* **22**, 629–650.
- Pinares-Patiño C and Waghorn G (2018) *Technical Manual on Respiration Chamber Designs*. Wellington, New Zealand: Ministry of Agriculture and Forestry. Available at <https://globalresearchalliance.org/wp-content/uploads/2018/02/LRG-Manual-Facility-BestPract-Sept-2018.pdf> (accessed 03. 07. 2020).
- Reisinger A and Ledgard S (2013) Impact of greenhouse gas metrics on the quantification of agricultural emissions and farm-scale mitigation strategies: a New Zealand case study. *Environmental Research Letters* **8**, 025019.
- Reisinger A and Clark H (2018) How much do direct livestock emissions actually contribute to global warming? *Global Change Biology* **24**, 1749–1761.
- Ribeiro RS, Rodrigues JPP, Maurício RM, Borges ALCC, Reis R, Berchielli TT, Valadares Filho SC, Machado FS, Campos MM, Ferreira AL and Júnior RG (2020) Predicting enteric methane production from cattle in the tropics. *Animal: An International Journal of Animal Bioscience* **14**, 1–15.
- Rojas-Downing MM, Nejadhashemi AP, Harrigan T and Woznicki SA (2017) Climate change and livestock: impacts, adaptation, and mitigation. *Climate Risk Management* **16**, 145–163.
- Rotz CA (2018) Symposium review: modelling greenhouse gas emissions from dairy farms. *Journal of Dairy Science* **101**, 6675–6690.
- Staal S, Ahuja V, Hemme T and Sharma VP (2016) Dairy Economics and Policy: Focus on Asia (No. 2016-02). Dairy Asia Working Paper. Available at <http://www.dairyasia.org/file/Dairy-Asia-Working-Paper-4-on-Dairy-Genetics-10-2-17.pdf> (accessed 02. 12. 2020).
- Tricarico JM, Kebreab E and Wattiaux MA (2020) MILK Symposium review: sustainability of dairy production and consumption in low-income countries with emphasis on productivity and environmental impact. *Journal of Dairy Science* **103**, 9791–9802.
- Tubiello FN (2011) *Linking Climate Change Financing and Sustainability: Implications for Agriculture*. Rome: Food and Agriculture Organization of the United Nations (FAO).
- Weiler V, Udo HM, Viets T, Crane TA and De Boer IJ (2014) Handling multifunctionality of livestock in a life cycle assessment: the case of smallholder dairying in Kenya. *Current Opinion in Environmental Sustainability* **8**, 29–38.
- Wilkes A and van Dijk S (2018) Tier 2 inventory approaches in the livestock sector: a collection of agricultural greenhouse gas inventory practices. Available at https://globalresearchalliance.org/wp-content/uploads/2018/12/Livestock-Tier-2-collection_Final_181130.pdf (accessed 02. 08. 2020).
- Zhao Y, Nan X, Yang L, Zheng S, Jiang L and Xiong B (2020) A review of enteric methane emission measurement techniques in ruminants. *Animals* **10**, 1004.
- Zia UE, Mahmood T and Ali MR (2011) *Dairy Development in Pakistan*. Rome: Food and Agriculture Organization of the United Nations (FAO). Available at <http://www.fao.org/3/a-al750e.pdf> (accessed 15. 01. 2021).