

Increased white matter integrity of posterior cingulate gyrus in the evolution of post-traumatic stress disorder

Zhang L, Li W, Shu N, Zheng H, Zhang Z, Zhang Y, He Z, Hou C, Li Z, Liu J, Wang L, Duan L, Jiang T, Li L. Increased white matter integrity of posterior cingulate gyrus in the evolution of post-traumatic stress disorder.

Objective: Functional imaging studies of post-traumatic stress disorder (PTSD) have shown an increased activation of posterior cingulate gyrus (PCG) of the brain. The aim of this study was to explore white matter integrity of PCG in PTSD subjects.

Methods: White matter integrity, as determined from fractional anisotropy (FA) value using diffusion tensor imaging, was assessed for PCG in subjects with and without PTSD from a severe mine accident. All subjects were also measured by the PTSD Checklist Civilian Version (PCL-C), the State-Trait Anxiety Inventory (STAI), the logical memory subtest and the visual reproduction subtest of the Wechsler Memory Scale-Revised in China. Sixteen PTSD subjects (8 subjects in each group) in the longitudinal study and 13 PTSD subjects as well as 14 non-PTSD controls in the cross-sectional case–control study were respectively recruited.

Results: In the longitudinal study, subjects with PTSD showed increased FA values in left PCG during the follow-up scan. In the cross-sectional study, FA values in bilateral PCG in PTSD subjects were higher than controls. Within the PTSD group ($n = 13$), FA values in the left PCG correlated positively with logical memory and negatively with PCL-C intrusion and STAI-trait (STAI-t) subscores. FA values in right PCG correlated negatively with STAI-t and STAI-state subscores.

Conclusion: These findings suggest that alterations of white matter integrity in PCG link to mnemonic and affective processing in PTSD over the long-term follow-up period.

Li Zhang^{1,*}, Weihui Li^{1,*}, Ni Shu², Huirong Zheng^{1,3}, Zhijun Zhang⁴, Yan Zhang¹, Zhong He⁵, Cailan Hou^{1,3}, Zexuan Li¹, Jun Liu⁵, Lifeng Wang¹, Lian Duan¹, Tianzi Jiang², Lingjiang Li^{1,6}

¹Mental Health Institute, The Second Xiangya Hospital of Central South University, Changsha, China; ²National Laboratory of Pattern Recognition, Automation Institute, Chinese Academy of Sciences, Beijing, China; ³Guangdong Mental Health Institute, Guangdong General Hospital, Guangdong Academy of Medical Sciences, Guangzhou, China; ⁴Department of Neurology, Affiliated Zhongda Hospital and Institute of the Neuropsychiatry of Southeast University, Nanjing, China; ⁵Department of Radiology, The Second Xiangya Hospital of Central South University, Changsha, China; and ⁶Chinese University of Hong Kong, Hong Kong, China

*Both authors contributed equally.

Keywords: anxiety; diffusion tensor imaging; memory; post-traumatic stress disorder

Lingjiang Li, Mental Health Institute, The Second Xiangya Hospital of Central South University, No. 139 Renmin Middle Road, Changsha 410011, Hunan, China.
Tel: +8673185292157;
Fax: +867315360086;
E-mail: llj2920@163.com

Accepted for publication May 15, 2011

Introduction

Post-traumatic stress disorder (PTSD) is characterised by specific symptoms, including intrusive thoughts, hyperarousal and avoidance, numbing, poor concentration and difficulty in explicitly recalling aspects of traumatic event. Several brain imaging studies have reported various structural and functional abnormalities of brain in subjects with PTSD (1–5). Most consistent findings are activation failure in medial prefrontal cortex, hippocampus and enhanced activation in amygdala in PTSD (6–11).

Additionally, studies in PTSD have shown that posterior cingulate gyrus (PCG) functionally related to aforementioned regions (12) is implicated in a neural circuit of stress (13).

It is thought that PCG, which consists of Brodmann areas 23, 29, 30 and 31, plays a critical role in memory, visuospatial orientation, monitoring eye movements and formation and retention of conditioned fear response (14,15). Notably, PCG has reciprocal connections with prefrontal cortex, anterior cingulate gyrus (ACG) and limbic system (16), from which PCG transmits information

to posterior neocortical association areas (14,15). Lesions in PCG in animals resulted in an impairment of spatial learning and memory function, therefore rats failed to perform the place navigation task (17,18). Similarly, Katayama et al. (19) have reported that PCG infarction in a woman leads to a failure in memorising a new route which likely results from a loss of directional memory over wide areas. Additionally, from an animal model of PTSD it appears that there is a significant degree of cortex specificity in memory impairment following underwater trauma (20).

Researches using diffusion tensor imaging (DTI) to explore dysfunction of white matter networks about PCG in PTSD were rare. DTI is a developed magnetic resonance imaging (MRI) technique that can provide information about white matter microstructural integrity *in vivo* (21,22). Fractional anisotropy (FA) value derived from DTI is measured by magnitude and direction of water diffusion (23). Previous DTI studies concerning PTSD have found abnormalities of white matter integrity in ACG and PCG. Abe et al. (24) reported increased FA values in left ACG, which correlated positively with symptom severity in victims with PTSD. However, Kim et al. (25) found that FA values in left ACG were lower in PTSD subjects than in healthy controls and they correlated negatively with severity. Kim et al. (26) also showed decreased FA values in multiple subregions of left cingulum bundle, especially in its anterior portion. There was only one published DTI study revealing lower FA values in right PCG in subjects with PTSD rather than in those without PTSD (27). Differences in types of trauma, illness duration, asymmetrical parameters in imaging processing and analysis, comorbid psychiatric disorders, including major depressive disorder, alcohol dependence and generalised anxiety disorder, could account for discrepant findings.

On 8 June 2005, a severe coal mine accident occurred in Zijiang Coal Mine in Hunan Province, China. Twenty-two miners lost their lives and the other 112 miners were rescued after more than 10 h of the ordeal in the darkness (7). The epidemiological data were obtained, and the survivors diagnosed with and without PTSD were scanned by neuroimaging techniques at 2, 10 and 24 months post-trauma, respectively. These techniques consisted of functional MRI (fMRI), three dimensions and DTI. Our group (7) has recently examined the neural correlates of PTSD using trauma-related imagery adapted to fMRI. In that study, subjects with acute PTSD at 2 months post-trauma exhibited increased activation in left PCG and decreased activation in bilateral middle frontal gyri when presented with trauma-relevant pictures versus neutral pictures. The PTSD group

also showed decreased activation in right frontal gyrus while performing memory performance. The recent findings suggested neurophysiological alterations and memory performance deficit in acute PTSD.

This study presents the results of a longitudinal study of white matter integrity in PTSD and data were obtained at 10 and 24 months post-trauma. We hypothesised that PTSD subjects would exhibit increased white matter integrity in PCG, consistent with increased activation in PCG in PTSD reported in prior functional neuroimaging studies (1,6,28–31). We have not found any prior research assessing longitudinal changes in PTSD using DTI. Survivors in this life-threatening coal mining accident experienced simultaneously and had high homogeneity in demographic background, which offered a distinct advantage in reducing the impact of confounding factors.

Materials and methods

Subjects

PTSD and matched control subjects were chosen from survivors of the coal mine accident and assessed according to the Structured Clinical Interview for Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV). Eighteen individuals who met the DSM-IV criteria of current PTSD were recruited into the study at 10 months post-trauma. At 24 months post-trauma, three original PTSD subjects dropped out of the study and the other seven people did not meet the PTSD diagnostic criteria. Thus the remaining 8 original PTSD subjects, along with 5 new people who met the criteria of current PTSD and 14 controls exposed to the same accident were recruited at 24 months post-trauma. Exclusion criteria for both groups included any history of head injury, any significant medical or neurological conditions, comorbid psychiatric disorders, substance abuse or dependence and mental retardation. During the study's 14-month follow-up period, subjects were excluded from the study if they experienced another traumatic event. All subjects were males and dextral, free of medication and without excessive head movement. Informed consent was written after the procedures had been fully explained. This study was approved by the Ethics Committee of the Second Xiangya Hospital of Central South University, China.

Instruments

Severity of illness was assessed by the PTSD Checklist Civilian Version (PCL-C), which is a 17 question, self-report measure used to evaluate PTSD symptoms from the DSM-IV criteria (32). Three subscores from

PCL-C were calculated corresponding to three PTSD symptom clusters: intrusion, avoidance and hyperarousal. Anxiety symptoms were assessed by the State-Trait Anxiety Inventory (STAI) (33). Depressive symptoms were assessed by the Beck Depression Inventory (BDI) (34). This self-report inventory consists of 20 items to assess state anxiety and another 20 items to assess trait anxiety. The logical memory subtest and the visual reproduction subtest of Wechsler Memory Scale-Revised in China were used to evaluate the short-term memory of survivors (7).

Data acquisition and processing

DTI was performed using a standard head coil on a 1.5T-Tesla General Electric scanner (Twin-speed, Milwaukee, WI, USA). Cushions were placed around the subjects' head to minimise head movement. Each volume consisted of 30 contiguous axial slices. Single-shot echo planar imaging with alignment of the anterior commissure–posterior commissure plane was undertaken, using the following parameters: repetition time = 12 000 ms, echo time = 107 ms, acquisition matrix = 128×128 , field of view = 24×24 , excitation number = 5, slice thickness = 4 mm and no gap. The diffusion sensitising gradients were applied along 13 non-collinear directions ($b = 1000 \text{ s/mm}^2$), together with an acquisition without diffusion weighting ($b = 0 \text{ s/mm}^2$) (35).

Three pairs of eigenvalues ($\lambda_1, \lambda_2, \lambda_3$) and eigen vectors were derived by diagonalisation of the diffusion tensor matrix. Subsequently, the FA value was calculated on a voxel-by-voxel basis according to the equation in Basser's study (36).

The method of imaging processing was similar to that described previously (35,37). Parametric images of FA and $b = 0$ were calculated with the DTI-Studio version 2.40 (H. Jiang, S. Mori; Department of Radiology, Johns Hopkins University, Baltimore, MD, USA) and transformed from Digital Imaging and Communications in Medicine (DICOM) format to analyse format for further processing using statistical parametric mapping (SPM2; Wellcome Department of Imaging Neuroscience, London, UK), implementing on Matlab 6.5 (MathWorks, Sherbon, MA, USA). For each subject, the b_0 image was normalised to the standard Montreal Neurological Institute (MNI) space using SPM2, and then the transformation matrix was applied to the FA map in order to normalise the map to the standard MNI space. All the images were resampled with a voxel size of $2 \times 2 \times 2 \text{ mm}^3$. The normalised FA maps were smoothed with an 8-mm full-width at half-maximum isotropic Gaussian kernel to decrease spatial noise, and a mean image (FA template) was created.

ROI analysis

As extensive alterations were observed in PCG in PTSD subjects, this region was brought into our central attention. Region of interest (ROI) analysis was performed to identify the FA value of the region. The colour-coded FA maps of DTI were used to outline different white matter fibre systems (38). In Colour Map-0, the axial line was located in the genu–splenium of the corpus callosum in the middle sagittal plane. The coronal line was taken from the medial surface of splenium of corpus callosum to the lateral surface in the middle sagittal plane and every movement of coronal line represents 2 mm from the medial surface. Different coronal plane follows every movement of coronal line in the middle sagittal plane. Two ROIs were located in bilateral PCG from each coronal plane. The FA value was obtained from Colour Map-0. Then, the mean FA values of bilateral PCG in different coronal planes (eight coronal planes for each individual) were calculated.

All measurements were performed by one rater (L. Z.) without the knowledge of subjects' identity. The intrarater reliability was established by rating five subjects randomly sampled from the whole subject group; the interrater reliability was also established by independent ratings of five subjects by two skilled raters who were familiar with brain anatomy (L. Z. and Yin Yan). Before the intraclass correlation coefficients (ICCs) were calculated, raters practised on another set of brains. The ICCs were 0.8969 for the left PCG and 0.9722 for the right PCG.

Statistical analysis

Intergroup [PTSD subjects at 10 months post-trauma ($n = 8$) vs. PTSD subjects at 24 months post-trauma; PTSD subjects at 24 months post-trauma ($n = 13$) vs. controls ($n = 14$)] differences in mean FA values were examined with paired t -test and two sample t -test, respectively. They were performed in a voxel-by-voxel manner. A statistical threshold of $p < 0.005$ (height threshold, uncorrected) (39) as well as an extent threshold of cluster size >50 voxels (400 mm^3) were considered to be statistically significant.

Paired t -test and independent t -test were also used in demographic and clinical assessment. Pearson correlation analysis was utilised to evaluate the correlations between mean FA values of each ROI and clinical variables, which involved subscores of PCL-C and STAI, logical memory and visual reproduction scores. A level of $p < 0.05$ (two tailed) was considered statistically significant. SPSS 13.0 (SPSS Inc, Chicago, IL, USA) for Windows was used for the computations.

Results

General information

Results relating to the PTSD groups (10 months post-trauma vs. 24 months post-trauma) in the longitudinal study were shown in Table 1. Significant differences were evident between two groups in PCL-C, STAI, BDI, logical memory and visual reproduction scores, and PCL-C intrusion, hyperarousal, STAI-state (STAI-s) and STAI-t subscores. There were no significant differences in PCL-C avoidance subscores between two groups. Results relating to the PTSD and control groups (24 months post-trauma) in the cross-sectional case-control study were presented in Table 2. As shown in the table, there were no significant differences in age, educational level, working depth underground, STAI-s subscores, logical memory and visual reproduction scores between the two groups. Significant differences were evident in PCL-C, STAI, BDI scores, PCL-C intrusion, avoidance, hyperarousal and STAI-t subscores.

Group comparison

In the longitudinal study, compared with the 10 months post-trauma, PTSD subjects at 24 months post-trauma showed significant increased FA values in the left PCG (Fig. 1) and decreased FA values in the right transverse temporal gyrus, bilateral temporal sub-gyri, left superior temporal gyrus, right prefrontal gyrus, right superior frontal gyrus, right medial frontal gyrus, right middle frontal gyrus, right frontal sub-gyrus and left cuneus (Table 3). In the cross-sectional study, compared with the control group, areas with higher FA values in the PTSD

Table 2. General information of the PTSD and control groups in the cross-sectional study

	PTSD group (n = 13)		Control group (n = 14)		t	p
	Mean	SD	Mean	SD		
Age (years)	37.54	3.69	40.86	5.20	-1.898	0.069
Educational level (years)	7.85	2.51	9.07	1.82	-1.460	0.157
Working depth underground (m)	-350.58	42.08	-330.43	46.77	-1.147	0.263
PCL-C	55.07	12.95	38.64	11.73	3.460	0.002**
PCL-C intrusion	16.77	3.39	11.29	4.07	3.788	0.001**
PCL-C avoidance	20.54	6.40	13.57	4.16	3.378	0.002**
PCL-C hyperarousal	17.76	4.18	13.78	4.82	2.284	0.031*
STAI	108.23	9.47	97.21	11.83	2.657	0.014*
STAI-s	52.07	5.78	48.42	6.82	1.493	0.148
STAI-t	56.15	4.86	48.78	6.23	3.407	0.002**
BDI	32.15	8.00	21.07	12.61	2.702	0.012*
Logical memory	10.92	5.11	10.43	3.72	0.290	0.78
Visual reproduction	8.92	3.71	8.21	3.24	0.530	0.601

BDI, Beck Depression Inventory; PCL-C, PTSD Checklist Civilian Version; PTSD, post-traumatic stress disorder; STAI, State-Trait Anxiety Inventory; STAI-s, STAI-state; STAI-t, STAI-trait.

*p < 0.05, **p < 0.01.

group were identified with bilateral PCG (Fig. 2), right precuneus, right parietal sub-gyrus, left middle temporal gyrus (Table 4). Regions with lower FA values in the PTSD group were not found.

Correlation

In the cross-sectional study, FA values of the left PCG in PTSD (n = 13) correlated positively with the logical memory scores (r = 0.61, t = 2.55, p =

Table 1. General information of the PTSD groups in the longitudinal study

	PTSD group (10 months post-trauma, n = 8)		PTSD group (24 months post-trauma, n = 8)		t	p
	Mean	SD	Mean	SD		
PCL-C	64.25	6.16	57.00	8.50	2.742	0.029*
PCL-C intrusion	19.88	2.36	16.75	1.67	3.751	0.007**
PCL-C avoidance	20.63	3.78	20.88	5.08	-0.182	0.861
PCL-C hyperarousal	21.88	2.42	19.38	2.97	2.887	0.023*
STAI	87.75	14.07	106.00	9.32	3.059	0.009**
STAI-s	43.63	8.72	53.25	8.33	2.258	0.040*
STAI-t	44.13	7.38	52.75	5.78	2.604	0.021*
BDI	46.38	3.34	28.13	7.06	7.412	0.000**
Logical memory	3.88	1.73	12.25	5.15	4.363	0.002**
Visual reproduction	2.88	4.55	8.38	4.24	2.501	0.025*

BDI, Beck Depression Inventory; PCL-C, PTSD Checklist Civilian Version; PTSD, post-traumatic stress disorder; STAI, State-Trait Anxiety Inventory; STAI-s, STAI-state; STAI-t, STAI-trait.

*p < 0.05, **p < 0.01.




Fig. 1. Increased FA in left PCG (the orange show) in PTSD at 24 months compared to 10 months post-trauma in the longitudinal study. The colour bar represents the T score. FA, fractional anisotropy; PCG, posterior cingulate gyrus; PTSD, post-traumatic stress disorder.

Table 3. Significant difference about FA value in PTSD subjects in the longitudinal study

Region	L/R	Voxel	Z score	MNI coordinates		
				x	y	z
Greater increase						
Posterior cingulate gyrus	L	99	3.53	-12	-44	4
Greater reduction						
Transverse temporal gyrus	R	533	4.55	40	-30	10
Temporal sub-gyrus	R	533	4.24	34	-32	16
	L	886	3.80	-46	-12	-22
Superior temporal gyrus	L	131	3.67	-40	-36	6
Prefrontal gyrus	R	310	3.18	56	8	10
Superior frontal gyrus	R	736	3.50	34	52	20
Medial frontal gyrus	R	10 638	4.23	10	62	22
Middle frontal gyrus	R	310	3.27	45	4	42
Frontal sub-gyrus	R	310	3.71	46	4	22
Cuneus	L	169	3.77	-12	-78	10

FA, fractional anisotropy; L, left side; MNI, Montreal Neurological Institute; R, right side.

Region displayed are for $p < 0.005$, cluster size >50 voxels.

0.027) and negatively with PCL-C intrusion ($r = -0.586$, $t = -2.398$, $p = 0.035$) as well as STAI-t subscores ($r = -0.605$, $t = -2.518$, $p = 0.029$). FA values of the right PCG correlated negatively with STAI-s ($r = -0.580$, $t = -2.362$, $p = 0.038$) and STAI-t subscores ($r = -0.630$, $t = -2.691$, $p = 0.021$) (Fig. 3). FA values of bilateral PCG showed no significant correlations with any other clinical variables in PTSD. No correlations were observed in controls.

In the longitudinal study, FA values of the left PCG in the PTSD group ($n = 8$) at 10 months post-trauma correlated positively with the STAI-s subscores only ($r = 0.773$, $R^2 = 0.5978$, $t = 2.986$, $p = 0.024$). Meanwhile, FA values of the left PCG in the PTSD group ($n = 8$) at 24 months post-trauma correlated negatively with PCL-C intrusion sub-

Table 4. Significant difference about FA value between the PTSD and control groups in the cross-sectional study

Region	L/R	Voxel	Z score	MNI coordinates		
				x	y	z
PTSD group > control group						
Posterior cingulate gyrus	R	114	2.84	12	-56	28
	L	66	3.33	-8	-44	30
Precuneus	R	114	3.40	8	-60	40
Parietal sub-gyrus	R	88	3.73	40	-38	36
Middle temporal gyrus	L	69	4.27	-56	-54	-2
PTSD group < control group						
None	-	-	-	-	-	-

FA, fractional anisotropy; L, left side; MNI, Montreal Neurological Institute; PTSD, post-traumatic stress disorder; R, right side.

Region displayed are for $p < 0.005$, cluster size >50 voxels.

scores only ($r = -0.749$, $R^2 = 0.56$, $t = -2.765$, $p = 0.033$).

Discussion

Using DTI approach, we found increased FA values in the left PCG over a 14-month follow-up period in PTSD subjects. To our knowledge, this is the first longitudinal study describing morphological changes of white matter over time through the whole brain of PTSD. Although there is little previous information concerning white matter connection of PCG in PTSD, some follow-up studies showed glucose hypometabolism (40,41) and greater grey matter loss in PCG (42) in patients with amnesic mild cognitive impairment relative to controls. These results suggest that PCG has dynamic structural and metabolic alterations in the course of diseases.

We also observed white matter abnormalities in bilateral PCG in the cross-sectional case-control study. Similar to our findings, one prior study (43)




Fig. 2. Higher FA in right PCG (a) and left PCG (b) (the orange show) in the PTSD group compared to the control group in cross-sectional study. FA, fractional anisotropy; PCG, posterior cingulate gyrus; PTSD, post-traumatic stress disorder.




Fig. 3. Correlations between FA values of left PCG in PTSD and (a) logical memory score ($r = 0.61$, $t = 2.55$, $p = 0.027$), (b) PCL-C intrusion subscores ($r = -0.586$, $t = -2.398$, $p = 0.035$, data from two of the victims were overlapping), (c) STAI-t subscores ($r = -0.605$, $t = -2.518$, $p = 0.029$; correlations between FA values of right PCG in PTSD), (d) STAI-s subscores ($r = -0.580$, $t = -2.362$, $p = 0.038$) and (e) STAI-t subscores ($r = -0.630$, $t = -2.691$, $p = 0.021$). FA, fractional anisotropy; PCG, posterior cingulate gyrus; PCL-C, PTSD Checklist Civilian Version; PTSD, post-traumatic stress disorder; STAI, State-Trait Anxiety Inventory; STAI-s, STAI-state; STAI-t, STAI-trait.

reported that subjects with panic disorder (PD) exhibited increased white matter FA values in the right PCG. Our result is also congruent with the findings of functional neuroimaging studies, reporting increased regional cerebral blood flow (rCBF) in bilateral PCG in subjects with PTSD relative to controls (6,28,29). In addition, a structural neuroimaging study exhibited grey matter density increasing significantly in the right PCG in rape victims with PTSD compared with healthy controls (44). We therefore propose that dysfunction in PCG may contribute to the mechanism of neuropathology in PTSD.

Another major finding was that increased FA values in the left PCG in PTSD correlated negatively with intrusive symptom, which often manifested as a rapid succession of intrusive memory (45). This pattern of correlation is consistent with functional evidence that increased activation in PCG is involved in intrusive or traumatic memory processing in subjects with PTSD (30,31,46–51). These studies rely on trauma-related scripts, sounds and pictures that can elicit intrusive or traumatic memory in PTSD (52,53). Furthermore, elevated FA values of PCG in PTSD subjects with improvement of intrusive symptom in the longitudinal study suggested that it may be a protective strategy for PCG to prevent the deterioration of intrusive recollection in PTSD.

More interestingly, our follow-up study showed that anxiety symptoms including state anxiety and trait anxiety increased in severity of PTSD. These findings supported the high rates of comorbid anxiety disorders or symptoms in full PTSD (54,55) or

sub-threshold PTSD (56). Our results also showed that increased white matter integrity in the left PCG in PTSD correlated negatively with trait anxiety. The same phenomenon happened between right PCG and trait anxiety as well as state anxiety. Unfortunately, our results were inconsistent with the study that increased FA values in right PCG in PD patients correlated positively with severity of anxiety symptoms (43). Another study also showed that right PCG correlated positively with trait anxiety (57) in healthy subjects during fear processing. In addition, Bench et al. (58) used a positron emission tomography (PET) approach to have found state anxiety associated with increased rCBF in PCG in depressive patients. Furthermore, from PET scans significant reductions in serotonin 1A receptor which played a crucial role in the pathophysiology of affected disorders were observed in PCG in individuals with anxiety disorders (59–61). Many factors involving different types of mental illnesses and methods of imaging data processing as well as analysis could result in these inconsistent consequences. Taken together, however, these findings support hypothesis that PCG is implicated in affective regulation (62–65), which would contribute to the pathogenesis of anxiety spectrum disorders including PTSD and so on. We speculated that it might become an important target of early intervention for PTSD.

The limitations of this study are the small sample size and the risk of a type I error (uncorrected).

Moreover, there was no control group at 10-month post-trauma. In addition, the findings of this study showed that quite many other brain regions (especially in frontal lobe and temporal lobe) might undergo specific change in PTSD, which would be discussed elsewhere due to limitation of space.

In spite of these limitations, using the voxel-based method with a relatively strict restriction of $p < 0.005$ and cluster size >50 voxels, we presented evidence for possible alterations of FA value in PTSD subjects, which suggests that white matter pathology may occur late in the course of illness.

In conclusion, these findings suggest that alterations of white matter integrity in PCG link to mnemonic and affective processing in individuals with PTSD over a long-term follow-up period. The results reveal that PTSD is associated with structural plastic changes to brain white matter in the evolution of illness. Further studies can use DTI to explore whether such changes are progressive over the course of PTSD and the relationship between the abnormalities and symptom severity, memory tests. We also would like to explore changes of white matter integrity in people with lifetime PTSD but not currently in the evolution of this illness.

Acknowledgements

We thank our subjects for their readiness to engage in this study and the Department of Radiology for providing scan time and technical assistant. We would also like to thank the study staff at the Mental Health Institute for their diligent work and essential contributions to the recruitment of participants and collection of data. We thank the fund support from the National Natural Science Foundation of China (30830046 to L. L.), the National Science and Technology Program of China (2007BAI17B02 to L. L.), the National 973 Program of China (2009CB918303 to L. L. and 2007CB512308 to Z. Z.), Program of Chinese Ministry of Education (20090162110011 to L. L.) and National Hi-Tech Research and Development Program of China (863 program: 2008AA02Z413 to Z. Z.).

References

- LANIUS RA, FREWEN PA, GIROTTI M, NEUFELD RW, STEVENS TK, DENSMORE M. Neural correlates of trauma script-imagery in posttraumatic stress disorder with and without comorbid major depression: a functional MRI investigation. *Psychiatry Res* 2007;**155**:45–56.
- LANIUS RA, BLUHM R, LANIUS U, PAIN C. A review of neuroimaging studies in PTSD: heterogeneity of response to symptom provocation. *J Psychiatr Res* 2006;**40**:709–729.
- FONZO GA, SIMMONS AN, THORP SR, NORMAN SB, PAULUS MP, STEIN MB. Exaggerated and disconnected insular-amygdalar blood oxygenation level-dependent response to threat-related emotional faces in women with intimate-partner violence posttraumatic stress disorder. *Biol Psychiatry* 2010;**68**:433–441.
- SCHUFF N, ZHANG Y, ZHAN W et al. Patterns of altered cortical perfusion and diminished subcortical integrity in

- posttraumatic stress disorder: an MRI study. *Neuroimage* 2011;**54**:S62–68.
- LANDRE L, DESTRIEUX C, BAUDRY M et al. Preserved subcortical volumes and cortical thickness in women with sexual abuse-related PTSD. *Psychiatry Res* 2010;**183**:181–186.
- BREMNER JD, NARAYAN M, STAIB LH, SOUTHWICK SM, MCGLASHAN T, CHARNEY DS. Neural correlates of memories of childhood sexual abuse in women with and without posttraumatic stress disorder. *Am J Psychiatry* 1999;**156**:1787–1795.
- HOU C, LIU J, WANG K et al. Brain responses to symptom provocation and trauma-related short-term memory recall in coal mining accident survivors with acute severe PTSD. *Brain Res* 2007;**1144**:165–174.
- ARMONY JL, CORBO V, CLEMENT MH, BRUNET A. Amygdala response in patients with acute PTSD to masked and unmasked emotional facial expressions. *Am J Psychiatry* 2005;**162**:1961–1963.
- BREMNER JD. Effects of traumatic stress on brain structure and function: relevance to early responses to trauma. *J Trauma Dissociation* 2005;**6**:51–68.
- BREMNER JD. Neuroimaging in posttraumatic stress disorder and other stress-related disorders. *Neuroimaging Clin N Am* 2007;**17**:523–538.
- BRUNETTI M, SEPEDE G, MINGOIA G et al. Elevated response of human amygdala to neutral stimuli in mild post traumatic stress disorder: neural correlates of generalized emotional response. *Neuroscience* 2010;**168**:670–679.
- VAN DEN HEUVEL M, MANDL R, LUIGJES J, HULSHOFF POL H. Microstructural organization of the cingulum tract and the level of default mode functional connectivity. *J Neurosci* 2008;**28**:10844–10851.
- BREMNER JD. Functional neuroimaging in post-traumatic stress disorder. *Expert Rev Neurother* 2007;**7**:393–405.
- MEUNIER M, JAFFARD R, DESTRADE C. Differential involvement of anterior and posterior cingulate cortices in spatial discriminative learning in a T-maze in mice. *Behav Brain Res* 1991;**44**:133–143.
- VOGT BA, FINCH DM, OLSON CR. Functional heterogeneity in cingulate cortex: the anterior executive and posterior evaluative regions. *Cereb Cortex* 1992;**2**:435–443.
- MADDOCK RJ, BUONOCORE MH. Activation of left posterior cingulate gyrus by the auditory presentation of threat-related words: an fMRI study. *Psychiatry Res* 1997;**75**:1–14.
- SUTHERLAND RJ, WHISHAW IQ, KOLB B. Contributions of cingulate cortex to two forms of spatial learning and memory. *J Neurosci* 1988;**8**:1863–1872.
- WHISHAW IQ, MAASWINKEL H, GONZALEZ CL, KOLB B. Deficits in allothetic and idiothetic spatial behavior in rats with posterior cingulate cortex lesions. *Behav Brain Res* 2001;**118**:67–76.
- KATAYAMA K, TAKAHASHI N, OGAWARA K, HATTORI T. Pure topographical disorientation due to right posterior cingulate lesion. *Cortex* 1999;**35**:279–282.
- RICHTER-LEVIN G. Acute and long-term behavioral correlates of underwater trauma – potential relevance to stress and post-stress syndromes. *Psychiatry Res* 1998;**79**:73–83.
- BASSER PJ, MATTIELLO J, LEBIHAN D. MR diffusion tensor spectroscopy and imaging. *Biophys J* 1994;**66**:259–267.
- MOSELEY ME, COHEN Y, KUCHARCZYK J et al. Diffusion-weighted MR imaging of anisotropic water diffusion in cat central nervous system. *Radiology* 1990;**176**:439–445.

23. BEAULIEU C. The basis of anisotropic water diffusion in the nervous system - a technical review. *NMR Biomed* 2002;**15**:435–455.
24. ABE O, YAMASUE H, KASAI K et al. Voxel-based diffusion tensor analysis reveals aberrant anterior cingulum integrity in posttraumatic stress disorder due to terrorism. *Psychiatry Res* 2006;**146**:231–242.
25. KIM MJ, LYOO IK, KIM SJ et al. Disrupted white matter tract integrity of anterior cingulate in trauma survivors. *Neuroreport* 2005;**16**:1049–1053.
26. KIM SJ, JEONG DU, SIM ME et al. Asymmetrically altered integrity of cingulum bundle in posttraumatic stress disorder. *Neuropsychobiology* 2006;**54**:120–125.
27. WANG HH, ZHANG ZJ, TAN QR et al. Psychopathological, biological, and neuroimaging characterization of posttraumatic stress disorder in survivors of a severe coal mining disaster in China. *J Psychiatr Res* 2010;**44**:385–392.
28. SACHINVALA N, KLING A, SUFFIN S, LAKE R, COHEN M. Increased regional cerebral perfusion by 99mTc hexamethyl propylene amine oxime single photon emission computed tomography in post-traumatic stress disorder. *Mil Med* 2000;**165**:473–479.
29. BREMNER JD, STAIB LH, KALOUPEK D, SOUTHWICK SM, SOUFER R, CHARNEY DS. Neural correlates of exposure to traumatic pictures and sound in Vietnam combat veterans with and without posttraumatic stress disorder: a positron emission tomography study. *Biol Psychiatry* 1999;**45**:806–816.
30. FREWEN PA, LANIUS RA, DOZOIS DJ et al. Clinical and neural correlates of alexithymia in posttraumatic stress disorder. *J Abnorm Psychol* 2008;**117**:171–181.
31. WHALLEY MG, RUGG MD, SMITH AP, DOLAN RJ, BREWIN CR. Incidental retrieval of emotional contexts in post-traumatic stress disorder and depression: an fMRI study. *Brain Cogn* 2009;**69**:98–107.
32. PETERLIN B, TIETJEN G, MENG S, LIDICKER J, BIGAL M. Post-traumatic stress disorder in episodic and chronic migraine. *Headache* 2008;**48**:517–522.
33. BARNES L, HARP D, JUNG W. Reliability generalization of scores on the Spielberger state-trait anxiety inventory. *Educ Psychol Meas* 2002;**62**:603.
34. BECK AT. Depression: causes and treatment. Philadelphia: University of Pennsylvania Press, 1967.
35. MA N, LI L, SHU N et al. White matter abnormalities in first-episode, treatment-naive young adults with major depressive disorder. *Am J Psychiatry* 2007;**164**:823–826.
36. BASSER PJ, MATTIELLO J, LEBIHAN D. Estimation of the effective self-diffusion tensor from the NMR spin echo. *J Magn Reson* 1994;**103**:247–254.
37. HAO Y, YAN Q, LIU H, XU L, XUE Z, SONG X. Schizophrenia patients and their healthy siblings share disruption of white matter integrity in the left prefrontal cortex and the hippocampus but not the anterior cingulate cortex. *Schizophr Res* 2009;**114**:128–135.
38. ASSAF Y, PASTERNAK O. Diffusion tensor imaging (DTI)-based white matter mapping in brain research: a review. *J Mol Neurosci* 2008;**34**:51–61.
39. READING S, YASSA M, BAKKER A et al. Regional white matter change in pre-symptomatic Huntington's disease: a diffusion tensor imaging study. *Psychiatry Res* 2005;**140**:55–62.
40. VILLAIN N, FOUQUET M, BARON J et al. Sequential relationships between grey matter and white matter atrophy and brain metabolic abnormalities in early Alzheimer's disease. *Brain* 2010;**133**:3301–3314.
41. FOUQUET M, DESGRANGES B, LANDEAU B et al. Longitudinal brain metabolic changes from amnesic mild cognitive impairment to Alzheimer's disease. *Brain* 2009;**132**:2058.
42. CHETELAT G, LANDEAU B, EUSTACHE F et al. Using voxel-based morphometry to map the structural changes associated with rapid conversion in MCI: a longitudinal MRI study. *Neuroimage* 2005;**27**:934–946.
43. HAN DH, RENSHAW PF, DAGER SR et al. Altered cingulate white matter connectivity in panic disorder patients. *J Psychiatr Res* 2008;**42**:399–407.
44. SUI SG, WU MX, KING ME et al. Abnormal grey matter in victims of rape with PTSD in Mainland China: a voxel-based morphometry study. *Acta Neuropsychiatrica* 2010;**22**:118–126.
45. TUERK PW, GRUBAUGH AL, HAMNER MB, FOA EB. Diagnosis and treatment of PTSD-related compulsive checking behaviors in veterans of the Iraq war: the influence of military context on the expression of PTSD symptoms. *Am J Psychiatry* 2009;**166**:762–767.
46. BREMNER JD, VYTHILINGAM M, VERMETTEN E et al. Neural correlates of declarative memory for emotionally valenced words in women with posttraumatic stress disorder related to early childhood sexual abuse. *Biol Psychiatry* 2003;**53**:879–889.
47. SOUZA MM, MELLO E SOUZA T, VINADE ER et al. Effects of posttraining treatments in the posterior cingulate cortex on short- and long-term memory for inhibitory avoidance in rats. *Neurobiol Learn Mem* 2002;**77**:202–210.
48. DRIESSEN M, BEBLO T, MERTENS M et al. Posttraumatic stress disorder and fMRI activation patterns of traumatic memory in patients with borderline personality disorder. *Biol Psychiatry* 2004;**55**:603–611.
49. LANIUS RA, WILLIAMSON PC, DENSMORE M et al. The nature of traumatic memories: a 4-T FMRI functional connectivity analysis. *Am J Psychiatry* 2004;**161**:36–44.
50. GEUZE E, VERMETTEN E, RUF M, DE KLOET CS, WESTENBERG HG. Neural correlates of associative learning and memory in veterans with posttraumatic stress disorder. *J Psychiatr Res* 2008;**42**:659–669.
51. MASAKI Y, NISHIKAWA T, IKEJIRI Y et al. Neural substrates of emotionally valenced episodic memory: a PET study using film stimuli. *Psychiatry Clin Neurosci* 2006;**60**:S46–S51.
52. LANIUS R. Dr. Lanius replies. *Am J Psychiatry* 2004;**161**:1927.
53. HULL A. Neuroimaging findings in post-traumatic stress disorder: systematic review. *Br J Psychiatry* 2002;**181**:102–110.
54. GARIEBALLA SS, SCHAUER M, NEUNER F et al. Traumatic events, PTSD, and psychiatric comorbidity in forensic patients – assessed by questionnaires and diagnostic interview. *Clin Pract Epidemiol Ment Health* 2006;**2**:7.
55. MAYOU R, BRYANT B, EHLERS A. Prediction of psychological outcomes one year after a motor vehicle accident. *Am J Psychiatry* 2001;**158**:1231–1238.
56. MARSHALL RD, OLFSO M, HELLMAN F, BLANCO C, GUARDINO M, STRUENING EL. Comorbidity, impairment, and suicidality in subthreshold PTSD. *Am J Psychiatry* 2001;**158**:1467–1473.
57. ETKIN A, KLEMENHAGEN KC, DUDMAN JT et al. Individual differences in trait anxiety predict the response of the

- basolateral amygdala to unconsciously processed fearful faces. *Neuron* 2004;**44**:1043–1055.
58. BENCH C, FRISTON K, BROWN R, FRACKOWIAK R, DOLAN R. Regional cerebral blood flow in depression measured by positron emission tomography: the relationship with clinical dimensions. *Psychol Med* 1993;**23**:579–590.
 59. NEUMEISTER A, BAIN E, NUGENT AC et al. Reduced serotonin type 1A receptor binding in panic disorder. *J Neurosci* 2004;**24**:589–591.
 60. KIM J, GORMAN J. The psychobiology of anxiety. *Clin Neurosci Res* 2005;**4**:335–347.
 61. SPINDELEGGER C, LANZENBERGER R, WADSAK W et al. Influence of escitalopram treatment on 5-HT 1A receptor binding in limbic regions in patients with anxiety disorders. *Mol Psychiatry* 2009;**14**:1040–1050.
 62. ETKIN A, PITTENGER C, POLAN H, KANDEL E. Toward a neurobiology of psychotherapy: basic science and clinical applications. *J Neuropsychiatry Clin Neurosci* 2005;**17**:145.
 63. PHILLIPS M, MARKS I, SENIOR C et al. A differential neural response in obsessive–compulsive disorder patients with washing compared with checking symptoms to disgust. *Psychol Med* 2000;**30**:1037–1050.
 64. OCHSNER KN, LUDLOW DH, KNIERIM K et al. Neural correlates of individual differences in pain-related fear and anxiety. *Pain* 2006;**120**:69–77.
 65. TILLFORS M, FURMARK T, MARTEINSDOTTIR I et al. Cerebral blood flow in subjects with social phobia during stressful speaking tasks: a PET study. *Am J Psychiatry* 2001;**158**:1220–1226.